



Universiteit
Leiden
The Netherlands

Modeling evolution from gas to young massive star clusters

Polak, B.; Mac Low, M.-M.; Klessen, R.; Appel, S.M.; Cournoyer-Cloutier, C.; Lewis, S.; ... ; Bolly, C.

Citation

Polak, B., Mac Low, M. -M., Klessen, R., Appel, S. M., Cournoyer-Cloutier, C., Lewis, S., ...
McMillan, S. (2023). Modeling evolution from gas to young massive star clusters. *Bulletin Of
The American Astronomical Society*, 207.04. Retrieved from
<https://hdl.handle.net/1887/3719326>

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)
Downloaded from: <https://hdl.handle.net/1887/3719326>

Note: To cite this publication please use the final published version (if applicable).

Bulletin of the AAS • Vol. 55, Issue 6 (AAS242 Abstracts)

Modeling Evolution from Gas to Young Massive Star Clusters

**Brooke Polak¹ Mordecai-Mark Mac Low² Ralf Klessen¹
Sabrina M. Appel³ Claude Cournoyer-Cloutier⁴ Sean Lewis⁵
Aaron Tran⁶ Martijn Wilhelm⁷ Simon Portegies Zwart⁷ Simon Glover¹
Stephen McMillan⁵**

¹Universität Heidelberg, ²American Museum of Natural History, ³Rutgers University,
⁴McMaster University, ⁵Drexel University, ⁶Columbia University, ⁷Leiden University

Published on: Jul 01, 2023

URL: <https://baas.aas.org/pub/2023n6i207p04>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Despite their pervasive influence, many aspects of star cluster formation are poorly understood because they are difficult to model. Simulation of their formation requires modeling a multitude of physical processes, from magnetohydrodynamics (MHD) to stellar feedback and gravitational dynamics, on scales from binary orbits to the widths of molecular clouds. To address these issues, we use Torch: a code employing the Astrophysical Multipurpose Software Environment (AMUSE) framework to couple stellar evolution and N-body codes with the MHD code FLASH. We include star formation based on sampling of the initial mass function after sink particle accretion, ray-tracing for ionizing and non-ionizing radiation, and mechanical feedback in the form of winds and supernovae. Torch originally required excessive computational resources to model giant molecular clouds (GMCs) with masses exceeding $10^5 M_{\odot}$, as it could not evolve the hundreds of thousands of stars that form in more massive clouds. This left a considerable gap, as the most massive GMCs in our galaxy exceed $10^6 M_{\odot}$. To enable Torch to model more massive GMCs, we integrate the N-body code PeTar to more efficiently handle the stellar dynamics. We present results from new Torch simulations of star clusters forming from clouds of mass 10^4 , 10^5 , and $10^6 M_{\odot}$. The most massive GMC forms over 100,000 stars within the first million years, making it one of the largest star cluster formation models resolving individual stars ever computed. We compare the global properties of each forming cluster over time and consider which remain constant as cloud mass is increased. We present the star formation histories and the time evolution of the gas properties, cluster mass, half-mass radius, and stellar velocity dispersion for each cloud.