



Universiteit
Leiden
The Netherlands

Measuring what matters: using claims data to evaluate healthcare outcomes and volume-outcome relationships

Schepens, M.H.J.

Citation

Schepens, M. H. J. (2024, February 29). *Measuring what matters: using claims data to evaluate healthcare outcomes and volume-outcome relationships*. Retrieved from <https://hdl.handle.net/1887/3719801>

Version: Publisher's Version


License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3719801>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 3

Does centralization of radical prostatectomy reduce the incidence of postoperative urinary incontinence?



Maike H.J. Schepens
Miranda L. van Hooff
Onno van der Galiën
Cathelijne M.P. Ziedses des Plantes
Diederik M. Somford
Pim J. van Leeuwen
Martijn B. Busstra
Sjoerd Repping
Michel W.J.M. Wouters
Jacques van Limbeek

Abstract

Background: On the basis of previous analyses of the incidence of urinary incontinence (UI) after radical prostatectomy (RP), the hospital RP volume threshold in the Netherlands was gradually increased from 20 per year in 2017, to 50 in 2018 and 100 from 2019 onwards.

Objective: To evaluate the impact of hospital RP volumes on the incidence and risk of UI after RP (RP-UI).

Design, setting, and participants: Patients who underwent RP during 2016–2020 were identified in the claims database of the largest health insurance company in the Netherlands. Incontinence was defined as an insurance claim for ≥ 1 pads/d.

Outcome measurements and statistical analysis: The relationship between hospital RP volume (HV) and RP-UI was assessed via multivariable analysis adjusted for age, comorbidity, postoperative radiotherapy, and lymph node dissection.

Results and limitation: RP-UI incidence nationwide and by RP volume category did not decrease significantly during the study period, and 5-yr RP-UI rates varied greatly among hospitals (19–85%). However, low-volume hospitals (≤ 120 RPs/yr) had a higher percentage of patients with RP-UI and higher variation in comparison to high-volume hospitals (>120 RPs/yr). In comparison to hospitals with low RP volumes throughout the study period, the risk of RP-UI was 29% lower in hospitals shifting from the low-volume to the high-volume category (>120 RPs/yr) and 52% lower in hospitals with a high RP volume throughout the study period (>120 RPs/yr for 5 yr).

Conclusions: A focus on increasing hospital RP volumes alone does not seem to be sufficient to reduce the incidence of RP-UI, at least in the short term. Measurement of outcomes, preferably per surgeon, and the introduction of quality assurance programs are recommended.

Patient summary: In the Netherlands, centralization of surgery to remove the prostate (RP) because of cancer has not yet improved the occurrence of urinary incontinence (UI) after surgery. Hospitals performing more than 120 RP operations per year had better UI outcomes. However, there was a big difference in UI outcomes between hospitals.

1 Introduction

Prostate cancer (PCa) is the most common cancer diagnosis among men.¹ The average 10-yr survival rate for patients with localized PCa is high and radical prostatectomy (RP) offers only a modest survival benefit in comparison to observation in selected patients^{2–4}. RP may result in severe functional side effects, such as urinary incontinence (UI) and sexual dysfunction.⁵ The risk of UI is higher after RP than after other treatments such as external beam radiotherapy and brachytherapy.⁶ Factors that may influence UI after RP (RP-UI) are well established.⁷ Certain biological and patient factors, including older age, higher body mass index, pre-existing lower urinary tract symptoms (LUTS), lower membranous urethral length (MUL), and functional bladder changes, have a negative impact on RP-UI.⁸ In addition to patient and tumor characteristics, the RP volume per surgeon plays a role in the variation in outcomes for this procedure.^{9–11}

Several studies have shown a positive correlation between favorable perioperative and functional outcomes and the hospital annual RP volume.^{12,13} Although some countries have specialized PCa centers with high patient volumes, many European countries have not yet implemented policies to centralize RP.⁹ In the Netherlands, a centralization policy for RPs has emerged. A significant driver for this centralization was our previous Dutch nationwide study based on claims data, which demonstrated that the risk of RP-UI was 30% lower in hospitals conducting more than 100 RPs/yr in comparison to hospitals performing fewer RPs.¹⁴ These results led to a stepwise increase in the national hospital volume standard for RP from 20 RPs/yr up to 2017, to 50 in 2018 and 100 from 2019 onwards. The rationale behind this increase was predicated on the assumption that an increase in the minimum annual RP volume would yield better functional and oncological outcomes.

The aim of the present population-based study was to use claims data to evaluate the impact of hospital RP volumes on the incidence and risk of RP-UI.

2 Patients and methods

2.1 Study approval

Data collection and analyses for this study were performed under the strict privacy rules and regulations of the Dutch government and health insurance companies. Patients included in the analyses could not be identified, so no informed consent or ethics approval was necessary.

2.2 Study aims

The aims of the study were to assess (1) the trend for RP volumes per hospital, (2) the

incidence of RP-UI at a national level, (3) variations in RP-UI incidence among hospitals, and (4) the risk of RP-UI.

2.3 Primary outcome

The primary outcome for the study was RP-UI. We defined UI as in our previous study:¹⁴ (1) the use of one or more incontinence pads per day 12–15 mo after RP and/or (2) surgery for RP-UI within 15 mo after RP.

2.4 Study design

2.4.1 RP volumes per hospital

Publicly available data from the website of the Dutch Health Institute¹⁵ were used to analyze the annual RP volume per hospital.

2.4.2 Incidence of RP-UI at a national level

A historical cohort of PCa patients undergoing RP between January 1, 2016 and December 31, 2020 and with claims for incontinence pads or surgery was identified from the claims database of the largest Dutch health insurer (Zilveren Kruis). Patients meeting the following criteria were excluded from the study: those who switched to another health insurer, deceased patients, patients with claims data indicating the use of incontinence pads 1–4 mo before RP, and patients insured by de Friesland, an insurance company label whose reimbursement policy did not provide full coverage for incontinence pads.

UI was analyzed by dividing the total number of patients with claims for incontinence pads by the total number of RP patients for each year (frequency and percentage). Differences were compared using a χ^2 test.

2.4.3 Variation in RP-UI incidence among hospitals

The average 5-yr UI incidence rate per hospital was first determined and then the average number of RPs performed annually at each hospital was calculated for two time periods: 2016–2018 (period 1) and 2019–2020 (period 2). Second, we recalibrated the optimal cutoff level for low-volume (LV) and high volume (HV) hospitals (see below). Third, hospitals were categorized as follows: (1) LV₁-LV₂ = LV in both time periods, (2) LV₁-HV₂ = LV in period 1 and HV in period 2; and (3) HV₁-HV₂ = HV in both time periods. For each hospital volume category, the RP-UI rates for period 1 and period 2 were calculated. Patients who underwent RP in hospitals that discontinued RP from 2019 onwards were excluded.

2.4.4 Risk of RP-UI

The impact of hospital RP volume was evaluated by developing an (explanatory) multivariable logistic regression model.¹⁶ The incidence of RP-UI was the primary outcome (dependent variable) and was dichotomized.

For all patients, data on patient characteristics (age at RP and comorbidity), treatment (RP date, lymph node dissection at RP, radiotherapy within 15 mo after RP, and hospital), and RP-UI outcome were collected from claims databases. Comorbidity was based on the ICD-10 code and specified as a diagnosis for each patient (eg, endocrine, heart, nervous system diseases). Case-mix variables were included as independent variables to adjust the effect of hospital volume category on RP-UI incidence.

Models were developed and the optimal threshold value was determined as follows. First, all potential confounders for which data were available (hospital volume [LV₁-LV₂, LV₁-HV₂, and HV₁-HV₂], age [continuous], radiotherapy [yes vs no], lymph node dissection [yes vs no], interaction variable between radiotherapy and lymph node dissection [yes vs no], and comorbidities [per diagnosis: yes vs no]) were separately tested in multivariate backwards analysis ($p < 0.05$) against the primary outcome (RP-UI). The variables were included as case-mix (independent) variables to adjust the effect of hospital volume category on RP-UI on the basis of an a priori hypothesized causal relationship.^{7,17} We used lymph node dissection as a proxy for tumor stage.


Second, relevant and significant confounders ($p < 0.05$) were included in the multivariable logistic regression model. Third, we recalibrated the cutoff value of 100 RPs/yr to distinguish LV and HV hospitals from our previous study¹⁴ by performing sensitivity analyses using predefined cutoffs of 100, 110, 120, 125, and 130 RPs/yr. All analyses were performed in SAS v9.4 (SAS Institute, Cary, NC, USA).

3 Results

3.1 RP volumes per hospital

The median annual RP volume per hospital increased from 51 in 2016 to 136 in 2020 and 161 in 2021 (Fig. 1). The annual RP volume varied widely among hospitals and increased over time as a limited number of HV hospitals emerged. The number of hospitals performing RP decreased from 37 in 2016 to 19 in 2020.


Figure 1 Boxplots of the number of radical prostatectomy (RP) procedures per hospital per year showing the median, range, interquartile range, and number of hospitals performing RP.



In total, data for 3843 patients who underwent RP during 2016–2020 were retrieved from the database. Data for four groups of patients were excluded (Fig. 2). The study population for analysis of RP-UI incidence at the national level included 2663 primary RP patients (mean age 66.7 yr, standard deviation 6.0).

At a national level, RP-UI incidence varied between 30.6% and 34.9% over the study period (Fig. 3). The RP-UI rate was 33.6% during 2016–2018 and decreased to 31.5% in 2019–2020 (difference not significant). The overall RP-UI incidence rate for 2016–2020 was 32.6%.

Figure 3 Mean proportion of patients with urinary incontinence 12–15 mo after radical prostatectomy (RP) by year. Differences in the rate of urinary incontinence after RP at a national level were not significant ($\chi^2 = 1.38, p > 0.05$).



3.3 Incidence of RP-UI among hospitals


Patient and treatment characteristics and outcomes per hospital for the subpopulation of 2376 patients (Fig. 2) are shown in Table 1. On the basis of goodness of fit and optimal dispersion of the number of hospitals among categories, the cutoff for LV versus HV in the analysis was set at 120 RPs/yr (Supplementary Table 1).

There was considerable variation in the 5-yr mean RP-UI incidence rate per hospital, ranging from 19.4% to 84.6%. Notably, the LV₁-LV₂ group showed wide variation. However, even in the HV₁-HV₂ group, significant hospital variation was observed, ranging from 19.4% to 44.7%.

LV₁-LV₂ hospitals had the highest mean RP-UI incidence (43.1%; Table 2) and the highest variation ($\pm 30.2\%$). HV₁-HV₂ hospitals consistently had the lowest RP-UI incidence, with

3.2 RP-UI incidence at a national level

Figure 2 Flow chart of the study populations for assessing the incidence of RP-UI at a national level, variations in RP-UI incidence among hospitals, and the risk of RP-UI.



RP = radical prostatectomy; RP-UI = urinary incontinence after RP.

a rate of 28.2% ($\pm 11.6\%$) in 2016–2018 that decreased to 27.9% ($\pm 12.1\%$) in 2019–2020. LV₁-HV₂ hospitals had lower RP-UI rates than LV₁-LV₂ hospitals and higher rates than HV₁-HV₂ hospitals, at 38.3% ($\pm 8.4\%$) in 2016–2018 and 33.6% ($\pm 6.3\%$) in 2019–2020. In all hospital volume categories, RP-UI incidence did not decrease significantly from the first period to the second period.

Table 1 RP volume category, patient characteristics, and RP-UI incidence by hospital during the study period (2016–2020)

Hospital	Volume category	Mean age (yr)	Mean proportion of patients (%)		
			RT	LND	RP-UI
1	LV ₁ -LV ₂	68.0	0	61.5	46.2
2	LV ₁ -LV ₂	68.6	7.7	69.2	84.6
3	LV ₁ -LV ₂	66.5	17.4	8.7	82.6
4	LV ₁ -LV ₂	67.5	8.9	20.4	33.1
5	LV ₁ -LV ₂	65.7	1.8	41.1	53.6
6	LV ₁ -LV ₂	64.9	9.0	25.6	36.1
7	LV ₁ -HV ₂	69.0	7.1	47.3	32.1
8	LV ₁ -HV ₂	67.0	13.8	44.8	37.9
9	LV ₁ -HV ₂	66.0	4.8	38.1	36.5
10	LV ₁ -HV ₂	66.1	6.0	44.6	41.0
11	LV ₁ -HV ₂	67.1	5.1	19.3	35.2
12	LV ₁ -HV ₂	67.3	17.6	54.1	39.2
13	LV ₁ -HV ₂	67.9	5.2	31.0	32.8
14	HV ₁ -HV ₂	66.4	7.7	50.7	32.9
15	HV ₁ -HV ₂	65.9	4.6	40.7	19.4
16	HV ₁ -HV ₂	66.4	4.0	50.7	44.7
17	HV ₁ -HV ₂	67.1	6.8	54.4	26.6
18	HV ₁ -HV ₂	66.9	5.9	53.5	23.3

HV = high RP volume; HV₁ = HV during 2016–2018; HV₂ = HV during 2019–2020; LV = low RP volume; LV₁ = LV during 2016–2018; LV₂ = LV during 2019–2020; LND = lymph node dissection; RP = radical prostatectomy; RT = radiotherapy; RP-UI = urinary incontinence after RP.

Table 2 Percentage of patients with RP-UI by hospital RP volume category

Hospital volume category	Mean RP-UI incidence \pm SD (%)		Hospitals	Patients
	2016-2018	2019-2020	(n)	(n)
LV ₁ -LV ₂	43.1 (± 27.0)	40.6 (± 30.2)	6	395
LV ₁ -HV ₂	38.3 (± 8.4)	33.6 (± 6.3)	7	653
HV ₁ -HV ₂	28.2 (± 11.6)	27.9 (± 12.1)	5	1328

HV = high RP volume (>120 RPs/yr); HV₁ = HV during 2016-2018; HV₂ = HV during 2019-2020; LV = low RP volume (≤ 120 RPs/yr); LV₁ = LV during 2016–2018; LV₂ = LV during 2019–2020; RP = radical prostatectomy; SD = standard deviation; RP-UI = urinary incontinence after RP.

3.4 Relationship between hospital volume and RP-UI risk

The final multivariable model showed that hospital volume category was independently correlated with RP-UI incidence. More precisely, patients undergoing RP in HV₁-HV₂ hospitals were 52% less likely to experience RP-UI than patients in LV₁-LV₂ hospitals (adjusted odds ratio [aOR] 0.48, 95% confidence interval [CI] 0.38–0.62; Table 3), while patients undergoing RP in LV₁-HV₂ hospitals were 29% less likely to suffer from RP-UI (aOR 0.71, 95% CI 0.55–0.92). Age (aOR 1.05, 95% CI 1.03–1.06) and lymph node dissection (aOR 1.34, 95% CI 1.12–1.61) also contributed to the model (Table 3). None of the other variables significantly improved this core model.

Table 3 Multivariable logistic regression and adjusted odds ratio for the incidence of urinary incontinence after RP^a

Parameter	Effect estimate \pm SE (%)	<i>p</i> value	Odds ratio (95% CI)
Age at RP	0.05 \pm 0.01	<.0001	1.05 (1.03 - 1.06)
Lymph node dissection	0.28 \pm 0.09	0.003	1.34 (1.12 - 1.61)
Radiotherapy ^b	0.26 \pm 0.17	0.91	1.294 (0.921 - 1.81)
HV1-HV2 vs LV1-LV2	-0.36 \pm 0.06	<.0001	0.48 (0.38 - 0.62)
LV1-HV2 vs LV1-LV2	0.01 \pm 0.07	0.88	0.71 (0.55 - 0.92)
Intercept	-3.85 \pm 0.53	<.0001	

CI = confidence interval; HV = high RP volume; HV1 = HV during 2016–2018; HV2 = HV during 2019–2020; LV = low RP volume; LV1 = LV during 2016–2018; LV2 = LV during 2019–2020; RP = radical prostatectomy; SE = standard error.

a For the final model, total concordance was 61.3%, with $\chi^2 = 8.20$ (Hosmer and Lemeshow test; *p* = 0.42).

b Not included in the final model.

Although the risk of RP-UI was lower for patients in HV hospitals, the RP-UI incidence at a national level did not decrease significantly after centralization. This might be attributable to other factors, such as an increase in the age of patients undergoing RP over the study period. The proportion of patients aged >70 yr was 30.5% in 2016–2018 and 37.6% in 2019–2020, and our risk model demonstrated that the risk of RP-UI increases with age. Other unmeasured factors, such as volume per surgeon and tumor stage, may have also contributed to the nonsignificant reduction in RP-UI at a national level. Further studies are needed to evaluate the impact of these factors and to study other functional and oncological outcomes over time.

4.1 Studies on the impact of hospital and surgeon volumes on RP-UI

Our study revealed an overall annual RP-UI rate of 33% for a continence definition of <1 pad/d. Relevant (single center) studies in the literature have reported RP-UI rates of approximately 10%.^{14,18–25} Many of these studies defined continence as “no leak” or “no pads” (Table 4). Our results are closest to findings reported by Haglind et al¹⁸ from a multicenter study in Sweden (*n* = 1847) in which the RP-UI rate was 20–21%. The comparatively higher national RP-UI rate (33%) in our study may be explained by study differences, as the study by Haglind et al did not encompass all hospitals, used patient-reported questionnaires, followed a prospective design, and excluded patients aged >75 yr.¹⁸

4 Discussion

Implementation of volume thresholds of 50 RPs/yr in 2018 and 100 RPs/yr from 2019 onwards in the Netherlands led to a notable reduction in the number of hospitals conducting RPs (from 37 to 19). There was also a significant increase in the median number of RPs performed annually per hospital (from 51 to 136). At a national level, RP-UI rates decreased marginally from 33.6% to 31.5%. The wide variation in hospital 5-yr RP-UI rates (19–85%) may have contributed to this nonsignificant decrease.

After adjustment for relevant confounding factors, hospital volume was the most predictive factor for RP-UI. Patients who underwent RP in hospitals that transitioned from LV to HV had 29% lower likelihood of experiencing RP-UI than patients in LV1-LV2 hospitals. Patients who underwent RP in hospitals with consistently high RP volumes throughout the study period had 52% lower risk of RP-UI. The proportion of patients undergoing RP in a HV hospital (>120 RPs/yr) increased from 33.5% in 2016 to 86.9% in 2020.¹⁵

Table 4 Overview of peer-reviewed evidence on 12-mo UI following radical prostatectomy

Reference and study period	Cohort	Patients (n)	Age (yr)	UI at 12 mo (%)	DoC	Data used	Surgical method
Zorn [19] 2003-2005	Single center	300	59.4	10	0 pads	NR	RARP
Shikanov [20] 2003-2008	Single center	380	58.1	18	<1 pad	Open interview	RARP
Patel [21] 2008-2009	Single surgeon	1100	58	2.6	0 pads	NR	RARP
Novara [22] 2005-2009	Single center	308	61.6	10	No leak	ICIQ-UI	RARP
Haglund [18] 2008-2011	14 centers	1847	63	21.3	<1 pad	Standardized questionnaire	RARP
Haglund [18] 2008-2011	14 centers	778	63	20.2	<1 pad	Standardized questionnaire	RRP
Coughlin [23] 2010-2015	Single center	157	35-70	10	0 pads	NR	RARP/RRP
Schepens [14] 2014-2015	Nationwide	1590	65	26.0	<1 pad	Claims data	NR
Sauer [24] 2014-2018	Single center	133	65	17	≤ 1 dry pad	Standardized questionnaire	RARP/RRP
Lee [25] 2004-2015	Single center	1691	≤ 70	11.4	< 1 pad	PROs at assessment	RARP/RRP
Lee [25] 2004-2015	Single center	610	> 70	18.5	< 1 pad	PROs at assessment	RARP/RRP
Present study 2016-2020	30% of national population	2663	66.8	32.6	<1 pad	Claims data	NR

DoC = definition of continence; ICIQ-UI = International Consultation on Incontinence Questionnaire-Urinary Incontinence; NR = not reported; PROs = patient-reported outcomes; RARP = robot-assisted radical prostatectomy; RRP = retropubic radical prostatectomy; UI = urinary incontinence.

4.2 Studies on the impact of hospital and surgeon volumes on RP-UI

Our previous nationwide study revealed that the risk of RP-UI was 30% lower for patients undergoing RP in hospitals with an annual volume of >100 RPs.¹⁴ The current study, in which we recalibrated the threshold to 120 RPs/yr, confirms this volume–RP-UI relationship. However, our study could not identify the optimal volume threshold for RP, as the volume threshold was chosen for methodological reasons. Another study confirmed that a high annual caseload (>200 RPs/year) was associated with lower RP-UI incidence.¹⁰ However, Nossiter et al²⁶ found no difference in RP-UI between LV and HV hospital cohorts in the UK, which might be explained by lower hospital RP volumes in the UK than in the Netherlands and/or the use of patient questionnaires in their study.

Our explanatory model demonstrated that hospital volume and age are important factors for RP-UI risk, but not all the variance in RP-UI incidence could be explained. A systematic review of the association between volume and outcome after RP revealed that both higher hospital volume and higher surgeon volume improved outcomes such as postoperative UI.¹⁰ Other studies confirmed that an increase in surgeon volume—but not hospital volume— was associated with improvement in outcomes such as long-term postoperative UI.^{10,27} Begg et al²⁷ concluded that surgeons who performed relatively poorly with respect to postoperative complications also performed poorly with respect to late urinary complications, and long-term RP-UI varied from 16% for high-volume to 20% for low-volume surgeons.

4.3 Studies on variation in RP-UI outcomes

Heterogeneity in RP-UI outcomes has been described^{28,29} but few studies have reported on the underlying factors. A Swedish population-based study concluded that the most important factor influencing RP-UI heterogeneity was surgeon experience, which accounted for 42% of the heterogeneity observed.³⁰ High-volume surgeons generally improve their technique via experience.³¹ However, even within a group of high-volume surgeons, substantial variation in outcome has been observed.³² Performance per surgeon could not be evaluated in the present study.

We found wide variation in the incidence of RP-UI among hospitals. While some HV centers reported remarkably high RP-UI rates, the highest variation was observed in LV hospitals. Therefore, it is possible that surgeon-volume relationships, which we could not measure, contributed in part to the variation in RP-UI rates we observed. Further studies should evaluate which additional measures are needed to reduce variation and to identify the optimal volume threshold per hospital and per surgeon.

4.4 Studies on reducing the risk of RP-UI

As observed for other cancer surgeries, simply increasing the minimum volume standard without also measuring outcomes in quality assurance programs (QAPs) may not be sufficient to improve the quality of care.³³ QAPs are structured programs in which health

care employees critically review the outcomes of their patients and continuously analyze and discuss these results in order to improve outcomes. Cathcart et al³⁴ evaluated the effects of a QAP for RP and found that 3-mo UI scores improved significantly for all but one surgeon who had low RP-UI rates at study initiation. Improving patient outcomes via short quality-improvement cycles is easier in settings in which a higher volume of patients is treated. For a low volume of patients, it can take years to measure a difference in outcomes after a change in practice.

We expect that at a national level the risk of RP-UI can be further reduced by (1) measuring outcomes at a per-surgeon level, (2) increasing the focus on audit and feedback via QAPs, and (3) applying per-surgeon RP volume thresholds.

4.5 Study strengths

The primary strength of our study lies in its novelty, as the first study to assess the trend in RP-UI incidence over time at a national level, interhospital variations, and the impact of hospital RP volumes on the incidence of RP-UI. The study strength stems from the extensive and representative nature of the data used. The source for our data was the database of the largest Dutch health insurance company, covering 27% of all RPs performed in the Dutch population during the study period. Most other studies on RP-UI are based on a single center or even a single surgeon. Patients insured by Zilveren Kruis are representative of the entire Dutch population³⁵, so the current results can be extrapolated to the total Dutch population.

4.6 Study limitations

Several limitations should be mentioned. First, we could not include all the variables identified in previous research as potential confounders, such as body mass index, LUTS, MUL, surgical technique, surgeon volume, and surgeon experience.^{11,27,36} Second, it was not possible to verify whether patients actually used the incontinence pads for which they claimed reimbursement. A recent study comparing RP-UI incidence derived from two sources (patient-reported outcome measure vs claims data) revealed comparable rates, with claims data slightly underestimating actual RP-UI rates.³⁷ Therefore, we do not suspect that the claims database as the source for the study data severely affected our results. Third, no distinction between laparoscopic and robot-assisted surgical approaches was possible. However, surgical approach was not associated with RP-UI incidence in a recent study¹⁷ and almost all RPs in the Netherlands were performed with robotic assistance during the study period.

5 Conclusions

The hospital RP volume threshold in the Netherlands has increased and the number of hospitals performing RP has thus decreased. Therefore, the average number of RPs per hospital increased. In all hospital volume categories, RP-UI incidence did not decrease significantly over the study period. However, LV1-LV2 hospitals had the highest percentage of patients with RP-UI and the highest variation.

In comparison to LV1-LV2 hospitals, the risk of RP-UI was 29% lower for patients undergoing RP in LV1-HV2 hospitals (≤ 120 RPs/yr during 2016-2018 and >120 RPs/yr during 2019-2020), and 52% lower for patients undergoing RP in HV1-HV2 hospitals (>120 RPs/yr for 5 yrs).

However, at a national level the RP-UI incidence did not significantly decrease. This could be influenced by the increasing age of patients at RP, the large variation in RP-UI rates per hospital, the absence of QAPs, and variables that could not be included in this study, such as surgeon experience and surgeon volume. Measurement of outcomes at a per-surgeon level and a greater focus on audit and feedback are recommended to improve outcomes relevant for patients.

Acknowledgements

We thank Ferdinand Roelfsema and Barbara Kolff van Oosterwijk for critically reviewing the manuscript.

6 Supplementary Table 1

Prediction model best RP volume cutoff based on goodness of fit characteristics and dispersion of number of hospitals for different volume cutoff values

Volume cut-off value	C-statistic	Hosmer Lemeshow test Goodness of fit Pr> Chi square	Number of hospitals per category		
			LV ₁ - LV ₂	LV ₁ - HV ₂	HV ₁ - HV ₂
100	0.602	0.517	3	4	12
110	0.606	0.185	3	7	7
120	0.614	0.487	6	7	5
125	0.616	0.514	6	8	4
130	0.614	0.373	9	5	4

We chose a volume cutoff of 120 RP per hospital per year. We chose the cutoff point based on a combination of optimal C-statistic, Hosmer Lemeshow test and an optimal distribution of hospitals per category. We did not choose for an optimal distribution of patients per category as this could distort the findings, because of the very high numbers of patients in few high-volume hospitals.

References

- 1 Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2021. *CA Cancer J Clin* 2021;71:7–33.
- 2 Wilt TJ, Vo TN, Langsetmo L, et al. Radical prostatectomy or observation for clinically localized prostate cancer: extended follow-up of the Prostate Cancer Intervention Versus Observation Trial (PIVOT). *Eur Urol* 2020;77:713–24.
- 3 Neal DE, Metcalfe C, Donovan JL, et al. Ten-year Mortality, disease progression, and treatment-related side effects in men with localised prostate cancer from the ProtecT randomised controlled trial according to treatment received. *Eur Urol* 2020;77:320–30.
- 4 Hamdy FC, Donovan JL, Lane JA, et al. Fifteen-year outcomes after monitoring, surgery, or radiotherapy for prostate cancer. *N Engl J Med* 2023;388:1547–58.
- 5 Vernooij RWM, Cremers RGHM, Jansen H, et al. Urinary incontinence and erectile dysfunction in patients with localized or locally advanced prostate cancer: a nationwide observational study. *Urol Oncol* 2020;38:735.e17–e25.
- 6 Albkri A, Girier D, Mestre A, Costa P, Droupy S, Chevrot A. Urinary incontinence, patient satisfaction, and decisional regret after prostate cancer treatment: a French national study. *Urol Int* 2018;100:50–6.
- 7 Lardas M, Grivas N, Debray TPA, et al. Patient- and tumour-related prognostic factors for urinary incontinence after radical prostatectomy for nonmetastatic prostate cancer: a systematic review and meta-analysis. *Eur Urol Focus* 2022;8:674–89.
- 8 Heesakkers J, Farag F, Bauer RM, Sandhu J, De Ridder D, Stenzl A. Pathophysiology and contributing factors in postprostatectomy incontinence: a review. *Eur Urol* 2017;71:936–44.
- 9 Van den Broeck T, Oprea-Lager D, Moris L, et al. A systematic review of the impact of surgeon and hospital caseload volume on oncological and nononcological outcomes after radical prostatectomy for nonmetastatic prostate cancer. *Eur Urol* 2021;80:531–45.
- 10 Leow JJ, Leong EK, Serrell EC, et al. Systematic review of the volumeoutcome relationship for radical prostatectomy. *Eur Urol Focus* 2018;4:775–89.
- 11 Thompson JE, Egger S, Böhm M, et al. Superior quality of life and improved surgical margins are achievable with robotic radical prostatectomy after a long learning curve: a prospective singlesurgeon study of 1552 consecutive cases. *Eur Urol* 2014;65:521–31.
- 12 Groeben C, Koch R, Baunacke M, Wirth MP, Huber J. High volume is the key for improving in-hospital outcomes after radical prostatectomy: a total population analysis in Germany from 2006 to 2013. *World J Urol* 2017;35:1045–53.
- 13 Gershman B, Meier SK, Jeffery MM, et al. Redefining and contextualizing the hospital volume-outcome relationship for robot-assisted radical prostatectomy: implications for centralization of care. *J Urol* 2017;198:92–9.
- 14 Schepens MHJ, Ziedses des Plantes CMP, Somford DM, et al. Hoe vaak incontinentie na radicale prostatectomie? Incidence of incontinence after radical prostatectomy using claims-based data. *Ned Tijdschr Geneesk* 2018;162:D2294.
- 15 Health Institute Netherlands. Prostate carcinoma indicators. Available at: <https://www.zorginzicht.nl/kwaliteitsinstrumenten/porstaatacarcinoom-indicatoren>.

- 16 Sainani KL. Explanatory versus predictive modeling. *PM R* 2014;6:841–4.
- 17 Baunacke M, Schmidt ML, Thomas C, et al. Long-term functional outcomes after robotic vs. retropubic radical prostatectomy in routine care: a 6-year follow-up of a large German health services research study. *World J Urol* 2020;38:1701–9.
- 18 Haglind E, Carlsson S, Stranne J, et al. Urinary incontinence and erectile dysfunction after robotic versus open radical prostatectomy: a prospective, controlled, nonrandomised trial. *Eur Urol* 2015;68:216–25.
- 19 Zorn KC, Gofrit ON, Orvieto MA, Mikhail AA, Zagaja GP, Shalhav AL. Robotic-assisted laparoscopic prostatectomy: functional and pathologic outcomes with interfascial nerve preservation. *Eur Urol* 2007;51:755–62.
- 20 Shikanov SA, Zorn KC, Zagaja GP, Shalhav AL. Trifecta outcomes after robotic-assisted laparoscopic prostatectomy. *Urology* 2009; 74:619–23.
- 21 Patel VR, Coelho RF, Chauhan S, et al. Continence, potency and oncological outcomes after robotic-assisted radical prostatectomy: early trifecta results of a high-volume surgeon. *BJU Int* 2010;106:696–702.
- 22 Novara G, Ficarra V, D'elia C, et al. Evaluating urinary continence and preoperative predictors of urinary continence after robot assisted laparoscopic radical prostatectomy. *J Urol* 2010; 184:1028–33.
- 23 Coughlin GD, Yaxley JW, Chambers SK, et al. Robot-assisted laparoscopic prostatectomy versus open radical retropubic prostatectomy: 24-month outcomes from a randomised controlled study. *Lancet Oncol* 2018;19:1051–60.
- 24 Sauer M, Tennstedt P, Berliner C, et al. Predictors of short and long term urinary incontinence after radical prostatectomy in prostate MRI: significance and reliability of standardized measurements. *Eur J Radiol* 2019;120:108668.
- 25 Lee YJ, Jung JW, Lee S, et al. Contemporary trends in radical prostatectomy and predictors of recovery of urinary continence in men aged over 70 years: comparisons between cohorts aged over 70 and less than 70 years. *Asian J Androl* 2020;22:280–6.
- 26 Nossiter J, Morris M, Cowling TE, et al. Hospital volume and outcomes after radical prostatectomy: a national population-based study using patient-reported urinary continence and sexual function. *Prostate Cancer Prostat Dis* 2023;26:264–70.
- 27 Begg CB, Riedel ER, Bach PB, et al. Variations in morbidity after radical prostatectomy. *N Engl J Med* 2002;346:1138–44.
- 28 Vickers A, Savage C, Bianco F, et al. Cancer control and functional outcomes after radical prostatectomy as markers of surgical quality: analysis of heterogeneity between surgeons at a single cancer center. *Eur Urol* 2011;59:317–22.
- 29 Carlsson S, Berglund A, Sjoberg D, et al. Effects of surgeon variability on oncologic and functional outcomes in a population-based setting. *BMC Urol* 2014;14:25.
- 30 Nyberg M, Sjoberg DD, Carlsson SV, et al. Surgeon heterogeneity significantly affects functional and oncological outcomes after radical prostatectomy in the Swedish LAPPRO trial. *BJU Int* 2021;127:361–8.
- 31 Prystowsky JB, Bordage G, Feinglass JM. Patient outcomes for segmental colon resection according to surgeon's training, certification, and experience. *Surgery* 2002;132:663–70.
- 32 Bianco Jr FJ, Riedel ER, Begg CB, Kattan MW, Scardino PT. Variations among high volume surgeons in the rate of complications after radical prostatectomy: further evidence that technique matters. *J Urol* 2005;173:2099–103.
- 33 Simunovic M, Urbach D, Major D, et al. Assessing the volumeoutcome hypothesis and region-level quality improvement interventions: pancreas cancer surgery in two Canadian provinces. *Ann Surg Oncol* 2010;17:2537–44.
- 34 Cathcart P, Sridhara A, Ramachandran N, Briggs T, Nathan S, Kelly J. Achieving quality assurance of prostate cancer surgery during reorganisation of cancer services. *Eur Urol* 2015;68:22–9.
- 35 van der Lee APM, Önsesveren I, Wierdsma AI, et al. The impact of antipsychotic formulations on time to medication discontinuation in patients with schizophrenia: a Dutch registry-based retrospective cohort study. *CNS Drugs* 2021;35:451–60.
- 36 Auffenberg GB, Qi J, Dunn RL, Linsell S, et al. Evaluation of patient and surgeon-specific variations in patient-reported urinary outcomes 3 months after radical prostatectomy from a statewide improvement collaborative. *JAMA Surg* 2021;156:e206359.
- 37 Baas D, Reitsma J, van Gerwen L, et al. P055: Correlation of healthcare insurance claims data for absorbing pads with PROMs as measure for urinary incontinence one year after radical prostatectomy. Presented at the 14th European Multidisciplinary Congress on Urological Cancers (EMUC 2022). https://www.postersessiononline.eu/173580348_eu/congresos/EMUC2022/aula/-P_55_EMUC2022.pdf.