



Universiteit
Leiden
The Netherlands

Executive functioning in antisocial behavior: a multi-level systematic meta-analysis

Jansen, J.M.; Franse, M.E.

Citation

Jansen, J. M., & Franse, M. E. (2024). Executive functioning in antisocial behavior: a multi-level systematic meta-analysis. *Clinical Psychology Review*, 109.
doi:10.1016/j.cpr.2024.102408

Version: Publisher's Version

License: [Creative Commons CC BY-NC 4.0 license](https://creativecommons.org/licenses/by-nc/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3721637>

Note: To cite this publication please use the final published version (if applicable).



Review

Executive functioning in antisocial behavior: A multi-level systematic meta-analysis

Jochem Milan Jansen^{a,b,*}, Melanie Elisabeth Franse^{c,d}

^a Institute for Criminal Law & Criminology, Faculty of Law, Leiden University, Leiden, Netherlands

^b Inforsa, Arkin, Amsterdam, The Netherlands

^c Netherlands Institute of Forensic Psychiatry and Psychology, Department of Science and Education, Herman Gorterstraat 5, 3511EW Utrecht, Netherlands

^d Institute of Education and Child Studies, Faculty of Social Sciences, Leiden University, Wassenaarseweg 52, 2333AK Leiden, Netherlands



ARTICLE INFO

Keywords:

Executive functioning

Antisocial behavior

Neuropsychology

Inhibition

Working Memory

Shifting

ABSTRACT

Neurobiological information – including executive functioning – is increasingly relevant for forensic clinical practice, as well as for the criminal justice system. Previous meta-analyses report that antisocial populations show impaired performance on executive functioning tasks, but these meta-analyses are outdated, have limitations in their methodological approach, and are therefore in need of an update. The current multi-level meta-analysis including 133 studies (2008-2023) confirms impaired performance in executive functioning ($d = .42$), but studies are heterogeneous. Several moderator analyses showed that neuropsychological test used, type of executive function component, and control group characteristics moderated the overall effect. Specifically, matching psychiatric problems in the non-antisocial control group eliminated any differences in executive functioning between groups. No moderation effects were found for assessment quality, hot or cold executive functions, and various population characteristics. These results could indicate that the assessment of executive functioning in antisocial populations may be less relevant for recidivism risk assessment than thought, although this should first be assessed in prospective longitudinal studies. Executive functioning could potentially be used to identify or screen for individuals with certain treatment needs or be used as a responsivity factor, especially in disorders which are often underdiagnosed in criminal justice settings.

1. Introduction

Neurobiological information is increasingly relevant for forensic clinical practice, as well as for the criminal justice system (Cheng, O'Connell, & Wormith, 2019). Neurobiological information may improve risk assessment (de Ruigh et al., 2021; Haarsma et al., 2020; Norman, Polaschek, & Starkey, 2023), and predict treatment completion in forensic psychiatric populations (Cornet, van der Laan, Nijman, Tollenaar, & de Kogel, 2015; Van der Sluys et al., 2020). One specific neurobiological domain which is increasingly assessed in this context is executive functioning (Haarsma et al., 2020; Norman et al., 2023), most often defined as (higher order) cognitive processes used to perform goal oriented, goal directed or future oriented actions, behaviors or responses (Baggetta & Alexander, 2016). Some authors propose that the nature of criminal responsibility can be reduced to EF (Hirstein, Siffert, & Fagan, 2018), and impaired executive functioning (EF) has been implicated in interpersonal problems (Sprague, Verona, Kalkhoff, & Kilmer, 2011),

physical health (Hall, Elias, & Crossley, 2006), and many psychiatric disorders, including substance abuse (Ersche et al., 2012) and emotion regulation difficulties (Fernandes, Wright, & Essau, 2023). Therefore, EF may be a transdiagnostic and/or risk factor for emotional, behavioral and psychotic disorders (Wade, Zeanah, Fox, & Nelson, 2020), all of which are prevalent within the judicial context.

Despite (custodial) sentences and forensic psychiatric treatment, world-wide recidivism remains at a relatively high and stable 20-60% reconviction rate within 2 years after release (Yukhnenko, Sridhar, & Fazel, 2019). EF could possibly provide (additional) information on recidivism risk and could guide forensic mental health care in order to reduce recidivism (Haarsma et al., 2020; Norman et al., 2023), but this requires a more specific understanding of the relationship between EF and antisocial behavior (ASB). This is necessary in order to assess the (potential) usability and feasibility of EF within a criminal justice setting. Most published research on this relationship uses a group comparison approach, including antisocial populations and non-

* Corresponding author at: Institute for Criminal Law & Criminology, Faculty of Law, Leiden University, Steenschuur 25, 2311ES Leiden, Netherlands.

E-mail address: j.m.jansen@law.leidenuniv.nl (J.M. Jansen).

offending control groups. Previous meta-analyses report that antisocial populations show impaired performance on EF tasks (Morgan & Lilienfeld, 2000; Ogilvie, Stewart, Chan, & Shum, 2011), but they are outdated, have limitations in their methodological approach and are therefore in need of an update.

1.1. Executive functioning

There is debate about the definition and operationalization of EF (see Baggetta and Alexander (2016) for a review). Studies generally agree that EF is a multidimensional construct, but disagree on the exact number of components and conceptualization of EF. For example, in a review of 106 studies, a total of 39 different components or processes of EF and 48 different theoretical models of EF were identified (Baggetta & Alexander, 2016). The most referenced model is proposed by Miyake et al. (2000) and includes three different components of EF; updating working memory, shifting between task sets, and inhibition of prepotent thoughts or actions. These functions correlate with one another, suggesting so called unity (e.g. they share an underlying ability), but are also separable, which indicates a degree of diversity (Friedman & Miyake, 2017; Miyake et al., 2000). According to this unity/diversity model, EF consists of three different components which share an underlying core ability, known as common EF. This model of shared and unique processes underlying EFs is supported by various neuroimaging studies (Saylik, Williams, Murphy, & Szameitat, 2022; Smolker, Friedman, Hewitt, & Banich, 2018). A similar model was proposed by Diamond (2013), and includes working memory, shifting, and inhibition, but does not include common EF or updating specifically. Instead, (Diamond) suggests that these three components work together in order to perform higher-order EFs such as planning, reasoning, or problem solving. Based on the review on EF conducted by Baggetta and Alexander (2016), we conclude there is most agreement on three separable EF components: working memory (span and updating), shifting between task sets, and inhibition of prepotent thoughts or actions.

Regarding the conceptualization of EF, age is a complicating factor, since confirmatory factor analyses show that different unity/diversity models of EF exist across the life span (Karr et al., 2018). EF seems to differentiate from a more unidimensional construct in children to both unidimensional and distinguishable constructs in adolescents and young adults (Karr et al., 2018). In this process of differentiation, inhibition and working memory seem to develop into distinguishable constructs earlier than shifting (Huizinga et al., 2006; Senn et al., 2004). It is proposed that this differentiation of EF is a direct reflection of cortical areas – which initially function non-specifically – becoming increasingly specialized during development, through activation, interactions, and experiences (Bardikoff & Sabbagh, 2017). From young adulthood to older adulthood, EF seems to dedifferentiate again, with greater unidimensionality of EF in older adults (Karr et al., 2022). Because of these differences in the unity/diversity of EF across the life span, it is important to assess the effects of age in the relationship between EF and ASB.

1.2. Assessment of executive functioning

Besides the conceptualization of EF, Baggetta and Alexander (2016) also reviewed how EFs are assessed, and report a wide variety in assessments. A total of 11 different batteries were reported, which are sets of (sub)tests or scales that measure different aspects of EF. Such batteries can either be performance based, or based on behavioral ratings by participants themselves or others (e.g. teacher, parent, staff). Additionally, 109 different neuropsychological tasks were identified, many of which were only reported once ($n=56$). Most commonly used tasks include the Stroop task, Digit span and the Go/No-Go task, but 27% of these tasks assessed multiple EF processes. For example, the Stroop task was used to measure inhibition, cognitive control, working memory, attention and overall/central executive functioning (Baggetta & Alexander, 2016).

This issue is known as the task impurity problem, which indicates that a certain task or outcome assesses or operates on a number of different executive (and/or non-executive) components. This severely hinders the interpretability of outcomes from such tasks, since it often remains unclear if task impairment is associated with task specific or common EF impairments (or another process all together). Snyder, Miyake, and Hankin (2015) describe that a specific outcome of an EF task often consists of variance explained by (1) task specific EF, (2) common EF, (3) non-EF processes, and (4) measurement error. The task impurity problem can be alleviated by using multiple measures of each EF component and extracting a latent variable constituting the task-specific EF of interest (Friedman et al., 2008). Unfortunately, this is a time consuming approach and therefore not always feasible. An alternative is a thorough task and task-outcome selection, since some tasks provide outcomes which are more closely related to specific EF (Snyder et al., 2015). For example, more traditional EF tasks (including the Stroop task and the Trail Making Test) measure both common and specific EF components and are sometimes considered too coarse to answer questions about specific EF. Some authors have provided guidelines or lists where the quality (including sensitivity to specific EF components) of certain task outcomes is assessed (see Op den Kelder, Van den Akker, Geurts, Lindauer, and Overbeek (2018) or Snyder et al. (2015)). Unfortunately, there is little consensus on when a certain task outcome is deemed to be of high quality, but in theory more specific EF measures should provide a more detailed answer on which processes are implicated in antisocial behavior.

Another complicating factor in the assessment of EF is that EF problems may manifest themselves differently in emotional and/or motivationally salient situations (*Dolan & Lennox, 2013). This idea is in line with studies showing that antisocial behavior is often associated with emotionally salient situations, such as reactive aggression (Bertsch, Florange, & Herpertz, 2020) or committing crime under the influence of peer pressure (Sijtsema & Lindenberg, 2018). Employing EF in these situations has been described as “hot” – as opposed to non-emotional “cold” – EF (Zelazo, 2020). Studies attempt to assess “hot” and “cold” EFs by varying task features (Salehinejad, Ghanavati, Rashid, & Nitsche, 2021). For example, a Go/No-Go task with neutral stimuli is used (i.e. symbols) for the assessment of “cold” EF, whereas emotionally salient images are used as stimuli for the assessment of “hot” EF (Salehinejad et al., 2021). Since ASB is more apparent in emotionally salient situations, it is expected that antisocial populations experience more problems with EF in “hot” situations, resulting in a larger EF impairment in “hot” tasks compared to “cold” tasks.

1.3. Antisocial behavior

ASB is a complex (social) construct, which has proven to be difficult to conceptualize within a single theoretical framework (Rutter, 2003). One proposed operationalization delineates antisocial behavior into three different categories: clinical psychiatric diagnosis, violation of legal and social norms, and aggressive or violent behavior (Ogilvie et al., 2011). Clinical diagnoses most frequently associated with antisocial behavior include oppositional defiant disorder (ODD), conduct disorder (CD), antisocial personality disorder (ASPD), and psychopathy/Callous Unemotional (CU)-traits. Both ODD and CD are sometimes referred to as Disruptive Behavioral Disorders (DBD). Legal operationalizations are related to the violation of social norms and include criminality or delinquency. This is often based on official records, such as the presence of a criminal record or being currently detained. Finally, physically aggressive or violent individuals are also considered antisocial.

1.4. Previous meta-analyses

So far two meta-analyses have assessed the difference in EF between antisocial and non-antisocial control groups (Morgan & Lilienfeld, 2000; Ogilvie et al., 2011). In both meta-analyses, antisocial individuals

generally scored worse on neuropsychological measures of EF than non-antisocial controls, with a medium effect-size ($d=.62$; (Morgan & Lilienfeld, 2000); $d=.44$; (Ogilvie et al., 2011)), but the included studies showed considerable heterogeneity. Several moderator analyses were reported, e.g. for the neuropsychological tests which were used and participant characteristics. The definition of antisocial behavior appeared to influence the overall effect size, since both meta-analyses reported lowest effect sizes for antisocial personality disorder, (ASPD; $d=.08$; $d=.19$) highest for criminality ($d=1.09$; $d=.61$), and psychopathy was somewhere in the middle ($d=.29$; $d=.42$). A meta-regression analysis with moderators revealed that diagnosis of ADHD in the ASB group negatively affected their EF performance, whereas age and sex did not (Ogilvie et al., 2011). These non-significant moderation effects for age and sex were reported in the meta-analysis by Morgan and Lilienfeld (2000) as well.

Although there is some evidence that specific characteristics may interact in explaining the association between ASB and EF, neither meta-analysis included moderation analyses with interactions between study characteristics. Individual studies support the notion of such interactions, for example, children with a higher levels of conduct problems (CP) and callous-unemotional (CU)-traits may display better EF performance compared to children in a low-CU/high-CP group, and low-CU/low-CP group (Graziano et al., 2022). Additionally, better EF was associated with more externalizing behavior in children with high CU-traits, but lower EF was not associated with externalizing behaviour in children with low CU-traits (de Graaf, Bolhuis, Cecil, White, & van Dongen, 2023). Finally, adolescents with both high-CP and high CU-traits reported higher violence and substance use, with the effects being stronger in youth with higher levels of executive control (i.e. inhibition). These studies in children therefore suggest that impaired EF is primarily a predictor for ASB in children with high CU-traits. Numerous other possible interactions could explain variance between studies and will be explored in this meta-analysis.

Unfortunately, neither of the meta-analyses used a theoretically based model of EF, but instead performed moderator analyses for the specific measure which was used to assess EF. Ogilvie et al. (2011) reported variability, with largest effects sizes for the self-ordered pointing task ($d=.83$; common EF/working memory), porteus maze test ($d=.71$; common EF/planning), delayed match to sample ($d=.59$; common EF/working memory) and the Go/No-Go task ($d=.56$; common EF/inhibition). Although Morgan and Lilienfeld (2000) assessed only a few neuropsychological tests, largest effect sizes were also reported for the maze test ($d=.80$; common EF/planning).

Both meta-analyses did not assess the relationship between EF and ASB in young children (<12 years of age), although much research exists on this topic. For example, a recent meta-analysis of prospective longitudinal studies indicates poor EF predicts future conduct problems (Yang et al., 2022). Although it is complicated to compare ASB in young children (i.e. children with a diagnosis of DBD) to adults (i.e. convicted criminals), it is unclear whether the difference in EF performance in ASB populations compared to controls varies with age. Based on the unity/diversity model - which proposes there are different models of EF throughout the life span - it could be expected that EF is also differently related to ASB throughout life. A recent meta-analysis found that children (aged 3-18) with DBD (described by the authors as an early manifestations of antisocial behavior) scored worse on EF than typically developing controls, with a small effect size for working memory ($d=-.26$), a small to medium effect size for inhibition ($d=-.30$ to $d=-.45$) and a small effect size for shifting ($d=-.31$) (Figueiredo, Ramiao, Barroso, & Barbosa, 2023). Another meta-analysis on the relationship between EF and DBD with a lower mean age (3-6 years) found somewhat smaller effect sizes, with small effect sizes for working memory ($d=.15$), inhibition ($d=.22$), and shifting ($d=.13$) (Schoemaker, Mulder, Dekovic, & Matthys, 2013). Compared to the meta-analyses from Morgan and Lilienfeld (2000) and Ogilvie et al. (2011) - where the youngest included participants were 14 years - smaller effect sizes were found in the meta-

analyses that included younger children. The current meta-analysis therefore incorporates (young) child and adult ASB populations in order to assess whether the association between EF and ASB varies with age.

Antisocial populations are characterized by high prevalence of ADHD, substance use disorders, psychosis/schizophrenia and other psychiatric problems (Fazel, Hayes, Bartellas, Clerici, & Trestman, 2016). It is unclear to what extent impairments in EF in antisocial groups are associated with the ASB or with underlying psychiatric symptoms characterized by impaired EF. Some studies indeed indicate that when ADHD children with and without CD are compared, no differences are found in EF (*Noordermeer et al., 2020). Although Ogilvie et al. (2011) showed that antisocial groups with ADHD perform worse than antisocial groups without ADHD, they did not assess whether the non-offending control groups included ADHD participants as well. The authors do suggest that it is possible that EF impairments are not specific to ASB, but that it could be associated with psychological, emotional and behavioral problems in general. Unfortunately, this hypothesis was not assessed in either of the meta-analyses, and it therefore remains unclear to what extent EFs are associated with ASB when analyses are controlled for comorbid psychological, emotional, and behavioral problems.

Although both meta-analyses conclude there is a robust relationship between impairments in EF and ASB, there are several theoretical and methodological limitations to consider. For example, although both meta-analyses assessed differences in specific measures used to assess EF, they lack a theoretically based assessment of EF. It is therefore unclear whether one EF component is more strongly associated with ASB than another. Additionally, traditional neuropsychological test are generally developed to assess large deviations in EF, whereas more recently developed tests are more sensitive to smaller deviations (Snyder et al., 2015). The meta-analysis by Ogilvie et al. (2011) used the 'extreme groups method', including only the largest effect size of each study. A disadvantage of this methodology is that it can cause inflation of the effect sizes in the meta-analysis. Both meta-analyses used a grand mean effect size per study, lumping together effect sizes for the association between antisocial groups and EF. Neither of the previous meta-analyses performed a multi-level meta-analysis which enables the researcher to include all effect sizes from a single study without violating the independence of effect sizes assumption (Fernandez-Castilla et al., 2020).

1.5. Current Study

The current meta-analysis aims to provide an updated assessment of the difference between antisocial populations and controls in EF, and to evaluate whether these differences vary between (1) EF components, (2) neuropsychological test used, (3) hot and cold EF, (4) EF assessment quality, (5) population characteristics of the antisocial and non-offending control groups, and (6) explorative interaction effects between these characteristics. A multi-level analysis will be performed, thereby addressing several limitations of previous meta-analyses.

2. Methods

This meta-analysis was reported following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) recommendations (Page et al., 2021). A PRISM research protocol was created before the study, but was not registered.

2.1. Search

A systematic search of the literature was performed on April 12th 2023, in four electronic databases: Criminal Justice Abstracts, PsychINFO, PubMed, and Web of Science. Within this search, there were no language or geographical restrictions, but there were restrictions for publication date (2008 and thereafter) and species (human only). After

consultation with an experienced librarian, the following terms and keywords have been used (see Appendix 1 for the specific search terms and filters used in each database): (1) Study population terms: antisocial, antisocial personality disorder, conduct disorder, oppositional defiant disorder, psychopath*, delinquent*, crim*, aggress*, violent*, offen*, assault*, unlawful*, cybercrim*, abus*, rule break*, inmate, prison*, jail*, incarcerat*, detain*, juvenile. (2) Outcome measure terms: executive function*, frontal function*, cognitive control, executive dysfunction*, shifting, inhibition, updating, working memory, planning, emotion* regulation, affect regulation, reappraisal. (3) In order to limit irrelevant results, all studies needed to contain at least one of the following terms: brain, neuro*, or cogni*. This search resulted in 14,831 unique articles.

2.2. In/Exclusion criteria

To be included in the meta-analysis, the studies needed to meet the following criteria: (1) ASB was operationalized as a clinical diagnosis that is related to ASB (Conduct Disorder, Oppositional Defiant Disorder, Antisocial Personality Disorder and/or Psychopathy) or by official records of criminality, delinquency and/or violent behavior. Diagnosis or ASB based on self-report measures or questionnaires was not sufficient. (2) EF was measured by a neuropsychological tasks assessing (updating) working memory, shifting, inhibition or higher-order EF. Implicit measures of EF and EF measured via self-report or questionnaires were not

included. (3) In line with previous meta-analyses, the antisocial group was compared to a non-antisocial control group on EF. The non-antisocial control group had to be free from any form of brain damage. (4) The outcome measures were sufficient to calculate effect sizes, for example means and standard deviations, *t*-values, *F*-values, *p*-values, and/or *r*-values.

2.3. Study selection

With the use of ASReview (Van De Schoot et al., 2021), an active learning software for meta-analytic screening, both authors independently assessed the identified studies for inclusion/exclusion, based on title and abstract. Based on text analysis ASReview presents the record that the machine deems most likely to be relevant first. Following a heuristic approach, the researchers stopped when they marked 200 consecutive articles as irrelevant because it is unlikely that any relevant studies remain present in the rest of the dataset. We chose 200 as a safe choice, whereas in literature often 50 or 100 is used (Ros, Bjarnason, & Runeson, 2017). This resulted in an inter-rater reliability of 98% between both authors. When the researchers were inconclusive about the eligibility of a study, they discussed the eligibility together until consensus was reached. This study selection based on title and abstract resulted in 337 possibly eligible studies. Following the inclusion and exclusion criteria, 133 articles of the initial 337 were included in the meta-analysis. The reasons for exclusion of the articles were: (1) no




Fig. 1. Flowchart. *The authors of these studies were contacted twice, but without any response.

sufficient operationalization of ASB ($n = 72$), (2) no non-ASB control group ($n = 58$), (3) no sufficient measure of EF ($n=36$), (4) missing required information ($n = 22$), (5) no quantitative study ($n = 8$), (6) and other reasons ($n = 8$), see [Figure 1](#).

2.4. Data collection and coding procedure

The following data was extracted from each study: (1) general study characteristics, (2) eligibility criteria, (3) information about the population and setting, (4) study methods, (5) study outcomes and results, (6) and key conclusions and limitations. The authors of articles that met the inclusion criteria but did not report sufficient information were contacted via email multiple times before exclusion.

Each outcome of each study was assessed for which specific EF component was measured; updating, working memory, shifting between task sets and inhibition of prepotent thoughts or actions. Since updating working memory and maintaining information in working memory (e.g. memory span), are often considered separate components ([Baggetta & Alexander, 2016](#)), they were both included separately. Additionally, higher order EFs – including planning, decision making and problem solving – were also included and labelled as Higher order EF. If different versions of neuropsychological tests were used, they were recoded into one specific test, for example, different versions of the Stroop test (word version, emotional version, children version, etc.) were all relabelled as Stoop test. Additionally, all outcomes where emotionally or motivationally salient stimuli were used, were considered “hot” EF, whereas all outcomes acquired under neutral stimuli were considered measuring “cold” EF.

Several participant characteristics were coded, including mean age of the ASB group and percentage of male participants. The ASB group was coded as being violent when specifically noted in the manuscript (e.g. conviction for violent crime, interpersonal violence perpetration, or sexual offending), and as non-violent otherwise. Additionally, dummy variables were created for all psychiatric disorders for both control and ASB groups separately when specifically assessed and described in the manuscripts. This indicates that studies which did not assess psychiatric diagnoses or offender characteristics were all labeled as non-diagnosed or non (sexually) violent.

Each outcome of each study was assessed for quality (low, medium or high) by JMJ. The assessment was based on previous work by [Op den Kelder et al., 2018](#) and [Snyder et al. \(2015\)](#), and extended for outcomes which had not been previously evaluated. All individual study outcomes were assessed on a case-by-case basis, and looked at whether EF assessment was confounded by assessment of speed or other EF elements and the level of cognitive load of the measures. For example, the Trail Making Test consists of part A and part B. Part B provides a measure of shifting, whereas part A provides a measure of (sustained) attention. Nevertheless sustained attention influences performance during part B. Therefore, a more specific and qualitatively better outcome measure would be to subtract the score on part A from part B. In this example, those effect sizes based on part B were scored of medium quality ([Op den Kelder et al. \(2018\)](#)), whereas a subtracted score (B-A) was scored as high quality. Since higher order EFs recruit several executive functions, they are generally scored as low quality. A full list of all outcomes per task and their quality assessment can be found in [Appendix 4](#).

2.4.1. Risk of bias assessment

Risk of bias was assessed using the Newcastle – Ottawa Quality Assessment Scale for case control studies ([Wells et al., 2000](#)), adapted slightly for the purpose of this study (see appendix 5). The instrument includes eight criteria, with a star assigned each time criteria are met (maximum nine stars). The first 10 studies were joint rated (IRR = 87%, disagreements resolved via discussion).

2.5. Analysis

2.5.1. Data structure

Most studies reported multiple effect sizes for multiple EFs. Since this violates the independence of effect sizes assumption of meta-analysis, a three-level meta-analysis with EF as a cross-classified random effects was conducted using the metafor package ([Viechtbauer, 2010](#)) in R-studio (version 2023.09.1). All extreme effect sizes were winsorized before analyses were performed, Cohen’s d was used as the measure of effect size, and p -values of $p < .05$ were considered statistically significant.

We found that the three-level model indeed provided a better fit compared to a two-level model with level 3 heterogeneity constrained to zero ($\chi^2_1 = 257.30$; $p < .001$). We also found that including EF as cross-classified random effects did not improve model fit ($\chi^2_1 = .00$; $p = 1.00$). Therefore, the final data structure used in subsequent analyses was a three level meta-analysis, modelling effect-sizes within studies. A trim and fill analysis was conducted to assess publication bias using a regular meta-analytic model, because the analysis is not available for a three-level meta-analysis ([Shi & Lin, 2019](#)).

2.5.2. Moderation analysis

Moderation analyses were conducted in order to assess whether the inclusion of a moderator could explain (some of) the heterogeneity in effect sizes. Separate analyses were conducted with categorical moderators concerning EF (components, hot vs cold, neuropsychological test used, and quality of assessment), and study population (ADHD DBD, schizophrenia/psychosis, psychopathy/CU-traits, SUD, or IED diagnosis in the ASB group, ASB group specified as violent or sexual offender, and diagnosis of the non-offending control group). Finally, age and sex were assessed as continuous moderators in separate analyses, and the effect of age group (adults vs youth subjects) were further investigated and reported in Appendix 6.

Exploratory interaction analyses were conducted in order to assess whether interaction between sample and or study characteristics could explain (some of the) heterogeneity in effect sizes. Such exploratory interaction analyses were conducted for: (1) diagnosis of ASPD and presence of psychopathic traits in adult samples, (2) diagnosis of DBD and presence of CU-traits in youth samples, (3) Psychopathy/CU-traits and hot vs cold EF, (4) ADHD diagnosis and hot vs cold EF, and (5) hot vs cold EF and violent vs not classified as violent ASB individuals. All analyses were only performed for cells containing at least 5 effect sizes.

3. Results

3.1. Study characteristics

A total of $n=133$ studies were incorporated into the analysis, resulting in $n=1238$ effect sizes. A total number of $n=9318$ ASB participants were included, and compared to $n=11738$ non-ASB controls. Participants were predominantly male in both ASB (88.92%) and non-antisocial control groups (85.75%), and mean age was similar in both groups (ASB: $M=23.01$ $SD=13.89$; Control $M=22.43$ $SD=13.40$). Studies in children all assessed ASB through diagnosis and not through official records. Several psychiatric diagnoses were identified in the ASB group, including: ADHD ($n_{effectsizes}=357$), DBD ($n_{effectsizes}=516$), Schizophrenia/psychotic ($n_{effectsizes}=30$), ASPD ($n_{effectsizes}=79$), CU-traits/Psychopathy ($n_{effectsizes}=37$), SUD ($n_{effectsizes}=22$), IED ($n_{effectsizes}=8$) or not diagnosed ($n_{effectsizes}=489$). In non-antisocial control groups, four diagnoses were identified: ADHD ($n_{effectsizes}=257$), SUD ($n_{effectsizes}=25$), and Schizophrenia/psychotic ($n_{effectsizes}=30$), and autism spectrum disorder ($n_{effectsizes}=3$). See [Appendix 3](#) for a table with study characteristics.

3.2. Risk of bias

Risk of bias assessment using the Newcastle – Ottawa Quality Assessment Scale for case control studies (Wells et al., 2000), shows that studies were generally of good quality, especially for definition of controls and ASB groups and representativeness of cases. There was higher risk of bias related to the way ascertainment of ASB was determined in both groups (i.e. when an antisocial group was assessed for ASB through official records, but controls were included based on self-report), for the non-response rate (which was often not well described), and for the comparability of ASB and control participants (which often differed in psychiatric diagnosis, age and/or gender), see Figure 2.

3.3. Three-level meta-analysis

The pooled Cohen’s *d* based on the three-level meta-analytic model was *d* = .42 (95%CI: .34-.50; *p* < .001). The estimated variance components were $\tau^2_{Level 3}$ = .17 and $\tau^2_{Level 2}$ = .20. This means that $I^2_{Level 3}$ = 40.5% of the total variation can be attributed to between-cluster, and $I^2_{Level 2}$ = 48.2% to within-cluster heterogeneity. The amount of variation, and the prediction interval of the main analysis (95% PI: -.78 – 1.63), reveal that the difference between antisocial groups and controls varies substantially. In order to assess the origins of this variability, several moderation analyses were conducted for: EF component, neuropsychological task used, hot versus cold EF, assessment quality, study population characteristics, and age and gender. The results of the trim and fill funnel plot did not reveal any filled in studies and therefore suggest little risk of publication bias (see Appendix 2)

3.4. Moderation analyses

3.4.1. Different executive function components

A three-level meta-analysis including EF component as a moderator showed that EF component did not moderate the pooled effect sizes $F_{(4, 1233)}=2.11$ *p*=.08. The effect sizes of all EF components differed from zero (see Figure 3), indicating that antisocial populations show impaired performance on all EFs. Although not statistically significant, the

impaired performance seems more evident for updating, see Figure 3.

3.4.2. Different neuropsychological tests

A three-level meta-analysis including neuropsychological test as a moderator showed it moderated the pooled effect sizes ($F_{(34, 1128)}=1.67$ *p*=.01), indicating that antisocial populations show more impaired performance on some neuropsychological tests compared to others, see Figure 4. Bonferroni corrected pairwise comparisons did not reveal any differences (all *p*>.15). Please note that specific tests with less than 5 effect sizes were excluded from the analyses.

3.4.3. Hot vs cold executive functioning

A three level meta-analysis including hot ($n_{effectsizes} = 168$) versus cold ($n_{effectsizes} = 1070$) EF as a moderator showed that hot and cold EF did not moderate the pooled effect sizes $F_{(1, 1236)}=1.25$ *p*=.26. Both hot (*d*=.48 *se*=.07 $t_{(1236)}=7.30$ *p*<.001) and cold (*d*=.41 *se*=.04, $t_{(1236)}=9.56$ *p*<.001) EF differed from zero. These results indicate that antisocial populations show impaired performance on EF, regardless of whether emotional or motivational stimuli were used or not.

3.4.4. Assessment quality

A three-level meta-analysis including assessment quality (low $n_{effectsizes} = 324$; medium $n_{effectsizes} = 338$; high $n_{effectsizes} = 576$) as a moderator showed quality did not moderate the pooled effect sizes, $F_{(2, 1235)}=.76$ *p*=.48. Whether an effect size was of low (*d*=.46 *se*=.05, $t_{(1235)}=8.79$ *p*<.001), medium (*d*=.40 *se*=.05, $t_{(1235)}=7.90$ *p*<.001) or high quality (*d*=.42 *se*=.05, $t_{(1235)}=9.30$ *p*<.001) did not make any difference in the pooled effect size, but all differed from zero. These results indicate that antisocial populations show impaired performance on EF regardless of the quality of the assessment of the task outcome

3.4.5. ASB operationalization / diagnoses

Studies included in this meta-analysis showed considerable variation regarding the characteristics of the included ASB participants, and results of moderation analyses showed a moderating effect for a diagnosis of schizophrenia/psychosis in the ASB group $F_{(1,1236)} = 10.15$ *p*=.002. ASB populations with a diagnosis of schizophrenia/psychosis showed no




Fig. 2. Risk of Bias assessment. This figure shows the results from the risk of bias assessment using the Newcastle Ottawa Scale. Green indicates the amount of studies which met criteria, yellow indicates when studies partially met criteria, and red indicates when studies did not meet criteria.




Fig. 3. Meta-analytic results including executive function components as moderating effect. Effect-size estimates are reported using Cohen's d .

significant impairment in EF ($d = -.03$ $p = .84$), whereas those without a diagnosis of schizophrenia/psychosis did ($d = .44$ $p < .001$). This result can likely be explained by the fact that all studies including an antisocial group with diagnosis of schizophrenia/psychosis used a non-antisocial control group which had a schizophrenia/psychosis diagnosis as well (see below).

There were no moderating effects for ADHD diagnosis $F_{(1,1236)} = 2.30$ $p = .13$, DBD diagnosis $F_{(1,1236)} = .54$ $p = .46$, ASPD diagnosis $F_{(1,1236)} = 2.61$ $p = .11$, SUD diagnosis $F_{(1,1236)} = 1.75$ $p = .19$, IED diagnosis $F_{(1,1236)} = .89$ $p = .34$, the presence of CU traits/psychopathy $F_{(1,1236)} = .72$ $p = .40$, whether ASB group specified as violent $F_{(1,1236)} = .01$ $p = .92$, specified as sexual offender $F_{(1,1236)} = 2.88$ $p = .09$, or whether ASB group was based on official records $F_{(1,1236)} = .03$ $p = .86$.

3.4.6. Non-offending control group with psychiatric diagnoses

A three-level meta-analysis including diagnoses of the non-offending control group did show a moderating effect $F_{(3, 1231)} = 30.58$ $p < .001$, indicating that studies using control groups with a psychiatric diagnosis differed in the reported effect sizes compared to studies who did not include control groups with a psychiatric diagnosis. Non-offending control groups with an ADHD diagnosis, substance use disorder, or schizophrenia/psychosis did not differ in EF compared to ASB groups – which were often also diagnosed with the same disorder (see Figure 5). A follow-up analyses showed a moderating effect of matching diagnosis in the ASB and non-offending control group $F_{(1,1233)} = 42.70$ $p < .001$, and revealed that ASB and non-offending control groups no longer differed in EF when accounting for underlying psychiatric disorders ($d = .06$ $se = .07$; $t_{(1233)} = .83$ $p = .41$). These results indicate that EF impairments in ASB, may – at least in part – be due to underlying mental health problems and not necessarily directly be related to the ASB.

3.4.7. Age and sex

A meta-regression, including age (of the ASB group) as a moderator indicated that age moderated the pooled effect size $F_{(1, 1234)} = 4.51$ $p = .03$, where differences in EF between ASB and non-antisocial control groups increased with age. Since EF components may develop differentially, a post-hoc moderation analyses including the interaction between age and EF component was performed, which revealed an interaction effect $F_{(4, 1226)} = 4.56$ $p = .001$. This significant interaction originates from a different relationship between age and both shifting and inhibition, where impairments in shifting (and not inhibition) seem to increase with age (see Figure 6). This interaction survived a Bonferroni correction for pairwise comparisons ($\Delta d = .013$ $p = .01$). A moderation analysis for sex, including the percentage of male ASB participants as moderator, did not reveal a statistically significant influence of sex on

the pooled effect size $F_{(1,1216)} = 2.35$ $p = .13$.

3.4.8. Exploratory interaction and post-hoc analyses

Interaction analysis for diagnosis of ASPD and presence of psychopathic traits in adult samples, did not reveal a statistically significant interaction $F_{(1, 514)} = .37$ $p = .54$, nor did the interaction analyses between Psychopathy/CUtraits and hot vs cold EF $F_{(1, 1234)} = .59$ $p = .44$, ADHD diagnosis and hot vs cold EF in youth $F_{(1,707)} = .01$ $p = .90$, or hot vs cold EF and violent ASB individuals $F_{(1,1234)} = .01$ $p = .73$. Several exploratory interaction analyses could not be conducted, because of empty cells in the data. For example, there were no studies in children with CU-traits but without a DBD diagnosis, and ADHD was not assessed in adults samples.

Finally, a post-hoc analysis was conducted to assess possible bias resulting from our coding procedure (see discussion). Results on this analysis of studies ($n = 11$, $n_{\text{effectsizes}} = 120$) which included DBD participants with and without ADHD, showed that ADHD did not increase EF impairment relative to the non-antisocial control group $F_{(1,118)} = 2.77$ $p = .10$.

4. Discussion

The current meta-analysis updates the available evidence for differences between antisocial populations and controls in EF using more advanced meta-analytic procedures, and by evaluating whether these differences vary between (1) EF components, (2) neuropsychological test used, (3) hot and cold EF, (4) EF assessment quality, (5) population characteristics of the antisocial and non-antisocial control groups. Additionally, (6) explorative interaction effects between these characteristics are conducted. Antisocial populations indeed showed impaired performance in EF (medium effect size $d = .42$), but studies are heterogeneous. Several moderator analyses showed that neuropsychological test used, age and other characteristics of both antisocial and non-offending control groups moderated the overall effect. Specifically, a non-antisocial control group with matching psychiatric problems eliminated any differences in EF between groups. No moderation effects were found for assessment quality, hot or cold EF, or various population characteristics (of the antisocial group), including diagnoses of DBD and ASPD. Whether the antisocial groups were defined through official records or not, or specified as violent or not also did not influence the results.

The reported impairment in EF in antisocial groups is in agreement with previous meta-analyses (Morgan & Lilienfeld, 2000; Ogilvie et al., 2011), although Morgan and Lilienfeld (2000) reported a larger effect size with a more select group of neuropsychological tests. In line with




Fig. 4. Meta-analytic results including neuropsychological test as moderating effect. Only neuropsychological tests which were used at least 5 times are included in this analysis. Effect-size estimates are reported using Cohen's d.

these previous meta-analyses we report high heterogeneity between studies, which indicates that the grand-mean effect should be interpreted with caution. It suggests that the effect sizes found in our meta-analyses did not derive from a single population of studies, which is most likely in the result of varying methodological approaches of the individual studies. In order to delineate this heterogeneity, several moderator analyses were conducted.

The use of different neuropsychological tests explains some of the heterogeneity, where antisocial groups experience the most problems on the Hayling test ($d=.86$), Spatial N-back ($d=.81$) and Porteus maze test ($d=.83$), whereas smaller differences were found using the Tower of London ($d=.15$), Object alternation test/delayed alternation test ($d=.06$) and the Delay discounting task ($d=.19$). Although these more extreme values were mostly for neuropsychological tests used 10 times or less, they are well aligned with the meta-analyses of Ogilvie et al. (2011), who – for example - report similar effect sizes for the Porteus maze test, and Tower of London task. Some differences were found as

well, as we report smaller effect sizes for the Go/No-Go task ($d=.56$ vs $d=.22$) and larger effect sizes for the stop signal task ($d=.42$ vs $d=.60$). Since we did not find moderating effects of the EF components, variability in neuropsychological tests is likely to reflect differences in task-instructions, stimuli and procedures.

Age was expected to have a moderating effect on the relationship between EF and ASB, because EFs continue to develop into late adolescence and smaller effect sizes were previously found for the relation between EF and ASB in children (Figueiredo et al., 2023; Schoemaker et al., 2013) compared to (young) adults (Morgan & Lilienfeld, 2000; Ogilvie et al., 2011). We indeed showed a moderating effect of age - where EF difficulties seem to increase with age - although this seems mostly driven by shifting, which component is thought to differentiate later in development. These results were corroborated when comparing studies in adults with studies in children (appendix 6). Although these results suggest that shifting is more weakly associated with ASB at a younger age, this effect was also expected for the other




Fig. 5. Meta-analytic results including control diagnosis as moderating effect. Effect-size estimates are reported using Cohen's *d*.




Fig. 6. Meta-analytic results for the interaction between age and both shifting and inhibition. Effect-size estimates are reported using Cohen's *d*.

constructs because lower effect sizes were found in younger participants for inhibition, shifting and working memory (Schoemaker et al., 2013). A possible explanation for not finding this effect in all EF components could be that only the mean age of participants per study was included, without correcting for age range. As a result, subtle differences between young children, adolescents, and older adults may have gone unnoticed. In addition, there is a significant gap in existing studies of EF and ASB with a mean age around young adulthood (18-26 years of age), while ASB is most common at this age and EF constructs are most distinguishable at this moment in the lifespan (Karr et al, 2022). Future studies should investigate age related differences in the association between EF components and ASB in order to more clearly assess these relationships.

Based on previous meta-analyses, it was expected that several characteristics of the ASB group would explain some of the heterogeneity between studies. The results show that heterogeneity was only explained when the ASB group included individuals with schizophrenia or psychosis, since these studies did not show any difference in EF between ASB and controls. These findings may be explained by the fact that all of these studies included a non-antisocial control group with schizophrenia or psychosis as well (see below). It is surprising that none of the other characteristics explained any of the heterogeneity, since previous meta-analysis did report that for example ASB participants

with ADHD report more EF impairments. This could be due to the fact that in our coding procedure studies that did not assess the presence of (for example) ADHD were coded as not having ADHD, since studies which screened for and consequently excluded ADHD participants were scarce. A post-hoc analysis on studies which included DBD participants with and without ADHD, showed that ADHD did not increase EF impairment relative to the non-antisocial control group. So although our coding procedure might have resulted in misclassified studies - which would cause an underestimation of the effect of ADHD diagnosis -, our post-hoc analysis suggests this bias is relatively small.

Although characteristics of the included ASB groups did not appear to explain heterogeneity between studies, but characteristics of control participants did. Our results indicate that the presence of a diagnosis in the non-offending control group (ADHD, substance use disorder, or psychosis/schizophrenia) completely abolished the difference in EF between the groups. A comparison between antisocial groups with non-antisocial control groups is based on the assumption that both groups are similar except for the ASB. Any differences in EF are therefore thought to be linked to the ASB. In practice, research shows that antisocial groups generally differ in many characteristics from controls, including socio-economic status, intelligence, or psychiatric problems. Our results show that a diagnosis (ADHD, substance use or schizophrenia/psychosis) in the non-antisocial control group abolished any difference with the

antisocial group. This suggests that impairments in EF could (at least in part) be related to underlying psychiatric problems - which are highly prevalent and underdiagnosed in antisocial populations (Buitelaar & Ferdinand, 2016; Fazel et al., 2016) - rather than being related to the ASB itself. This could indicate that the assessment of EF in ASB populations is less relevant for recidivism risk assessment, although this should first be assessed in longitudinal studies assessing the prospective and predictive value of EF for recidivism. The available evidence from such studies is currently inconclusive (Fine, Steinberg, Frick, & Cauffman, 2016; Ormachea et al., 2017), and reviews or meta-analyses have not yet been conducted. The results do indicate that EF could potentially be used to identify or screen for individuals with certain treatment needs or be used as a responsivity factor, especially in disorders which are often underdiagnosed in criminal justice settings.

It was expected that the extent of the impairments in EF would vary between the different components of EF, but our results suggest this is not the case since EF component did not moderate the overall effect size. Antisocial populations do appear to experience more difficulties with updating working memory. We expected that assessment quality would also moderate the overall effect size, because high quality assessments are more aimed at measuring specific EF components, minimalizing the influence of common EF. We did not find any moderating effects of assessment quality. One explanation for this result could be that antisocial groups show a general impairment in all EF components. An alternative explanation would be that impairment in underlying common EF results in (equally) reduced performance on the specific EF components. In order to assess whether some specific EF components are implicated more than others, studies should use multiple measures of each EF component and extract a latent variable constituting the task-specific EF of interest (Friedman et al., 2008).

4.1. Implications & directions for future research

Our results support previous studies reporting impairments in EF in antisocial groups, but also show that these differences could originate from underlying psychiatric problems rather than be directly related to ASB. Although we did not assess whether EF performance was predictive for recidivism or future ASB, our findings could suggest that EFs are not relevant for ASB and therefore for risk assessment because they relate to underlying psychiatric problems. On the other hand, it remains true that antisocial populations are characterized by higher prevalence of psychiatric problems and impaired EF, and that this could be related to or predictive of (future) antisocial behavior. Studies prospectively assessing the value of EF as predictor for future ASB are relatively scarce and show mixed results (Aharoni et al., 2013; Brassard & Joyal, 2022; Nikulina & Widom, 2019; Zijlmans et al., 2021). To our knowledge, reviews or meta-analyses on the predictive value of EF in ASB are currently unavailable. Since neurobiological information - including EF - is increasingly relevant for forensic clinical practice and risk assessment (de Ruigh et al., 2021; Haarsma et al., 2020; Norman et al., 2023), such reviews or meta-analyses are needed to delineate the association between EF, psychiatric problems and (future) ASB.

Antisocial groups are not more impaired on specific EF components, suggesting either a general impairment in antisocial groups in all EF components or that underlying common EF drives the difference between antisocial and non-antisocial control groups. We did not find any studies that used a latent variable approach, using multiple measures of each EF component to extract task-specific EF of interest (Friedman et al., 2008). These types of studies could provide a more definitive answer to the question whether impairments in specific EF components are implicated in ASB.

For future research on the relationship between EF and ASB, it is important more effort is made to match the controls more closely to the antisocial group in terms of socioeconomic status, education level and traumatic brain injury since these factors are highly associated with EF (Checa & Rueda, 2011; Jansen, 2020; Lawson, Hook, & Farah, 2018).

Many of the included studies did not take these differences between the ASB group and controls into account, which may lead to inflation of effect sizes.

4.2. Strengths & limitations

Our results were obtained using a multi-level analytic approach, enabling us to incorporate multiple effect sizes per study in our analysis. This is superior compared to using an "extreme groups method", which may cause an inflation of the resulting effect sizes of the meta-analysis. Nevertheless, we found similar results as previous meta-analyses, suggesting that the inflation of effect sizes was minimal.

In our analyses we assessed specific EF components and non-offending control group characteristics, both of which had not previously been analyzed or reported upon. Especially for the non-antisocial control group characteristics, we identified that underlying psychiatric problems may play a major role in explaining differences in EF performance between antisocial and non-antisocial control groups. Only 11% of the studies used an appropriate non-antisocial control group with a matching psychiatric diagnosis. Future studies should be aware of - and correct for - underlying psychiatric problems which could explain the often found difference in EF performance between antisocial and non-antisocial control groups.

Although several moderator and interaction analyses were conducted, many included studies did not report on any or all of the moderators. During data extraction all studies which did not report on - for example - violence were coded as "not specified as violent". Consequently, in the analyses we compared effect sizes from violent samples to samples which were not specified as violent. If individuals were not specified as violent, this does not necessarily mean that they are in fact non-violent, and it is likely that the comparison group includes violent individuals to some extent. The results of our moderation analyses might therefore under-estimate the true effect size for these moderators.

Studies using (only) self-report measures of ASB were excluded from the analyses. Some self-report measures are well validated, and in fact may identify a (now under-represented subgroup of) individuals which do exhibit ASB, but have not been caught. We have opted to exclude self-report measures because we wanted to adhere to the in/exclusion criteria used by the original two meta-analyses which excluded such self-report measures. Nevertheless, this should be noted as both a limitation of our study and as a possible avenue for further research. Finally - as both strength and limitation - all included effect sizes were scored on assessment quality by JMJ, because there is no full list of neuropsychological tests or agreed upon criteria for EF assessment quality. Previous studies were helpful, but did not include all measures reported in our selected studies (Op den Kelder et al., 2018; Snyder et al., 2015). Confounding influences of speed or other EF elements and the level of cognitive load of the measures were taken into account, but not all descriptions of the task procedures were clear on the (possible) level of interference. These quality labels should therefore be further validated and/or future studies should provide a golden standard for EF assessment.

4.3. Conclusion

The current meta-analysis updates the available evidence for differences between antisocial populations and controls in EF using more advanced meta-analytic procedures, and evaluates whether these differences vary between (1) EF components, (2) neuropsychological test used, (3) hot and cold EF, (4) EF assessment quality, and (5) population characteristics of the antisocial and non-offending control groups. Antisocial populations indeed show impaired performance in EF (medium effect size $d=.42$), but these differences could be explained by underlying psychiatric problems rather than by the exhibited ASB. No differences were found between hot and cold EF, but some neuropsychological test revealed greater differences between groups than others.

These results could indicate that the assessment of EF in ASB populations is less relevant for recidivism risk assessment, although this should first be assessed in prospective longitudinal studies. EF could potentially be used to identify or screen for individuals with certain treatment needs or be used as a responsivity factor, especially in disorders which are often underdiagnosed in criminal justice settings.

Declaration of competing interest

None.

Appendix A. Appendix 1

A.1. Search Criminal Justice Abstracts

(brain OR neuro* OR cogni*) AND ("executive function*" OR "frontal function*" OR "cognitive control" OR "executive dysfunction*" OR "shifting" OR "inhibition" OR "updating" OR "working memory" OR "planning" OR "verbal fluency" OR "delay gratification" OR "instant gratification" OR "impulsivity" OR "spatial working memory" OR "emotion regulation" OR "emotional regulation" OR "Affect regulation" OR "Reappraisal") AND ("conduct disorder" OR "oppositional defiant disorder" OR "antisocial*" OR "antisocial personality disorder" OR "psychopath" OR "psychopathy" OR "psychopaths" OR "delinquen*" OR "crim*" OR "aggress*" OR "violen*" OR "offen*" OR "assault*" OR "unlawful*" OR "cyberbully*" OR "bully*" OR Cybercrime*" OR "rule break*" OR "inmate*" OR "prison*" OR "jail" OR "incarcerat*" OR "detain*" OR "juvenile").

A.2. Search PsychInfo

(brain OR neuro* OR cogni*) AND ("executive function*" OR (DE "Executive Function") OR "frontal function" OR "cognitive control" OR (DE "Cognitive Control") OR "executive dysfunction*" OR (DE "Cognitive Impairment") OR "shifting" OR "inhibition" OR (DE "Inhibition (Personality)") OR "updating" OR "working memory" OR "planning" OR (DE "Planned Behavior") OR "verbal fluency" OR (DE "Verbal Fluency") OR "delay gratification" OR (DE "Delay of Gratification") OR "instant gratification" OR "impulsivity" OR (DE "Impulsiveness") OR "spatial working memory" OR "emotion regulation" OR "emotional regulation" OR (DE "Emotional Regulation") OR "Affect regulation" OR "Reappraisal") AND ("conduct disorder" OR (DE "Conduct Disorder") OR "oppositional defiant disorder" OR (DE "Oppositional Defiant Disorder") OR "antisocial*" OR (DE "Antisocial Behavior") OR "antisocial personality disorder" OR (DE "Antisocial Personality Disorder") OR "psychopath" OR "psychopathy" OR "psychopaths" OR (DE "Psychopathy") OR "delinquen*" OR (DE "Juvenile Delinquency") OR "crim*" OR (DE "Crime") OR (DE "Criminal Behavior") OR (DE "Criminal Conviction") OR (DE "Criminal Offenders") OR (DE "Criminal Record") OR "aggress*" OR (DE "Aggressive Behavior") " OR "violen*" OR (DE "Violence") OR "offen*" OR "assault*" OR "unlawful*" OR "cyberbully*" OR (DE "Cyberbullying") OR "bully*" OR (DE "Bullying") OR "25ybercrime*" OR (DE "Cybercrime") OR "rule break*" OR "inmate*" OR (DE "Prisoners") OR "prison*" OR "jail" OR (DE "Prisons") OR "incarcerat*" OR (DE "Incarceration") OR "detain*" OR "juvenile") NOT (Cancer OR Tumor OR metasta* OR rat* OR mouse OR mice OR rodent* OR fish OR zebra OR monkey*).


A.3. Search PubMed

(Psychiatry and Psychology Category[MeSH] OR neuro* OR cogni* OR brain) AND ((executive function[MeSH Terms] OR "executive function*" OR "frontal function*" OR "cognitive control" OR shifting OR inhibition OR "Inhibition, Psychological"[Mesh] OR updating OR "working memory" OR planning OR "emotion regulation" OR "emotional regulation" OR "Emotional Regulation"[Mesh] OR "Affect regulation" OR "Reappraisal")) AND (("conduct disorder" OR "oppositional defiant disorder" OR antisocial* OR "antisocial personality disorder" OR "Antisocial Personality Disorder"[Mesh] OR psychopath OR psychopaths OR psychopathy OR delinquen* OR crim* OR aggress* OR violen* OR offen* OR assault* OR unlawful* OR cybercrime* OR rule break* OR inmate* OR prison* OR jail OR incarcerat* OR detain* OR juvenile)) NOT ((meta-analysis[Filter] OR review[Filter] OR systematicreview[Filter])

A.4. Search Web of Science

(ALL=(brain OR neuro* OR cogni*)) AND (ALL= ("executive function*" OR "frontal function*" OR "cognitive control" OR "executive dysfunction*" OR "shifting" OR "inhibition" OR "updating" OR "working memory" OR "planning" OR "verbal fluency" OR "delay gratification" OR "instant gratification" OR "impulsivity" OR "spatial working memory" OR "emotion regulation" OR "emotional regulation" OR "Affect regulation" OR "Reappraisal")) AND (ALL=("conduct disorder" OR "oppositional defiant disorder" OR "antisocial*" OR "antisocial personality disorder" OR "psychopath" OR "psychopathy" OR "psychopaths" OR "delinquen*" OR "crim*" OR "aggress*" OR "violen*" OR "offen*" OR "assault*" OR "unlawful*" OR "26ybercrime*" OR "rule break*" OR "inmate*" OR "prison*" OR "jail" OR "incarcerat*" OR "detain*" OR "juvenile")) NOT (ALL=(Cancer OR Tumor OR metasta* OR rat* OR mouse OR mice OR rodent* OR fish OR zebra OR monkey*)) Refined by: PUBLICATION YEARS: (2023 OR 2022 OR 2021 OR 2011 OR 2020 OR 2010 OR 2019 OR 2009 OR 2018 OR 2008 OR 2017 OR 2016 OR 2015 OR 2014 OR 2013 OR 2012)

Appendix B. Appendix 2



Appendix 3

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
Antonini et al., 2015	Diagnosis	ODD	ADHD + ODD	33	9.44	72.72%	Memory Span Task	Working memory	Cold
			ADHD controls	67	(1.75)	75.76%	Wisconsin Card Sorting Task	memory maintenance	Cold
			Healthy controls	30	8.88	66.67%	Delay Discounting Task	Shifting	Hot
					9.00		Iowa Gambling Task	Choice impulsivity	Hot
				(1.80)			Affective decision making		
Frias-Armenta et al., 2011	Official records	Offender	Offenders	48	14.00	80.0%	Wisconsin Card Sorting Task	Shifting	Cold
			Healthy controls	27	(1.35)	-	Stroop Test	Inhibition	Cold
Baliouis et al., 2019	Official records + diagnosis	Offender, ASPD, Psychopathy	Offenders + ASPD	52	30.3	100%	Spatial Working Memory Task	Working memory	Cold
			Offenders + psychopathy	27	(8.9)	100%	Stockings of Cambridge	memory maintenance	Cold
			Healthy controls	20	(10.9)	100%	Intra/Extradimensional Shift Task	Planning	Hot
					33.9		Go/No-Go	Shifting	
				(10.7)			Inhibition		
Barkataki et al., 2008	Official records + diagnosis	Offender, ASPD	Offenders + ASPD	14	33.5	100%	Go/No-Go	Inhibition	Cold
			Offenders + schizophrenia	12	(10.45)	100%			
			Controls + schizophrenia	12	34.83				
			Healthy controls	14	(4.97)	100%			
				34.92					
				(7.60)					
				32.14					
				(7.75)					
Barlati et al., 2023	Official records	Offender	Violent offenders + schizophrenia	50	37.92	86%	Trail Making Test	Shifting	Cold
			Healthy controls	50	(11.14)	86%	Stroop Test	Inhibition	Cold
					37.70		Memory Span task	Working memory	Cold
				(11.08)		Tower of London	memory maintenance	Cold	
							Planning		
Barnett et al., 2009	Diagnosis	ODD, CD	ODD + ADHD	22	9.02	86.36%	Memory Span Task	Working memory	Cold
			CD + ADHD	20	(2.11)	80.0%	Spatial Working Memory Task	memory maintenance	Cold
			Controls + ADHD	23	8.24	73.91%	Delayed Match to	Working	
			Healthy controls	25	(1.64)	72.0%			

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
					8.23 (1.94)		Sample Spatial Recognition Task	memory maintenance	Cold
					8.81 (1.48)		Tower of London	Working memory maintenance	Cold
Becerra-García, 2015	Official records	Offender	Domestic offenders	10	42.00	100%	Trail Making Test	Planning	Cold
			Sex offenders	20	(8.48)	100%		Shifting	
			Violent offenders	9	37.55	100%		Working memory maintenance	
			Non-violent offenders	8	(9.27)	100%		Shifting	
			Healthy controls	31	30.22	100%		Shifting	
					(7.17)				
					40.88				
					(10.48)				
					38.45				
					(12.58)				
Becerra-García & Egan, 2014	Official records	Offender	Sex offenders (incestuous)	21	47.33	100%	Trail Making Test	Shifting	Cold
			Sex offenders	11	(7.92)	100%	Digit Span Backward	Working memory manipulation	Cold
			Healthy controls	28	49.09	100%			
					(13.38)				
					45.61				
					(8.77)				
Blum et al., 2017	Diagnosis	ASPD	ASPD	26	24.3	57.7%	Intra-Extradimensional Shift Task	Shifting	Cold
			Healthy controls	266	(3.8)	68.4%	Stop Signal Task	Inhibition	Cold
					22.8		Stockings of Cambridge	Planning	Cold
					(3.7)		Cambridge Gambling Task	Affective decision making	Hot
							Stroop Test	Inhibition	Hot
Borroni et al., 2015	Official records	Offender	Offenders	27	17.66	100%			Cold
			Healthy controls	27	(1.55)	100%			
			Healthy controls (age-paired)	27	17.11				
			Healthy controls (age- and education-paired)	27	(1.55)	100%			
					17.59				
					(1.69)				
Brænden et al., 2023	Diagnosis	ODD	ODD	23	9.9 (1.5)	58.0%	NEPSY-2 Design Fluency	Shifting	Cold
			Controls + ADHD	43	9.6 (1.8)	63.0%	NEPSY-2 Inhibition	Inhibition	Cold
							NEPSY-2 Inhibition	Shifting	Cold
							NEPSY-2 Word List Interference	Working memory maintenance	Cold
Bulgari et al., 2017	Official records	Offender	Offenders + schizophrenia	50	46.7	92.0%	Wisconsin Card Sorting Task	Shifting	Cold
			Controls + schizophrenia	37	(10.0)	86.5%	Iowa Gambling Task	Affective decision making	Hot
					49.2		Memory Span Task	Working memory maintenance	Cold
					(9.9)		Tower of London	Planning	
								Shifting	
Cantrell, 2008	Diagnosis	CD	CD	29	17.07	100%	Trail Making Test	Shifting	Cold
			CD + ADHD	20	(1.03)	100%	Stroop Test	Inhibition	Cold
			Healthy controls	59	16.75	100%	Wisconsin Card Sorting Task	Shifting	Cold
					(0.68)			Planning	Cold
					-		D-KEFS Tower Test		
Carter Leno et al., 2018	Diagnosis	ODD/CD	ODD/CD	26	12.79	83.72%	Go/No-Go Task	Inhibition	Cold
			Controls + ADHD	21	(1.61)	95.24%	Switch Task	Shifting	Cold
			Controls + ASS	41	12.98	65.38%			
			Healthy controls	43	(1.47)	58.54%			
					12.31				
					(1.62)				
					13.77				
					(1.08)				
Chamberlain et al., 2016	Diagnosis	ASPD	ASPD	17	23.8	58.8%	Intra/Extradimensional Shift Task	Shifting	Cold
			Healthy controls	229	(3.9)	61.6%	Stop Signal Task	Inhibition	Cold
					23.6		Cambridge Gambling Task	Affective decision making	Hot
					(3.1)		Spatial Working	Working memory	Cold

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
Coenen et al., 2022	Official records	Offender	Offenders Healthy controls	34 36	16.67 (1.21) 17.25 (1.00)	100% 100%	Memory Task	maintenance	
							Stockings of Cambridge	Planning	
							StopSignal task	Inhibition	Cold
							Stroop Test	Inhibition	Cold
							Digit span Backward	Working	Cold
							Spatial working memory	Working	Cold
							Trail Making Test	Memory	Cold
							Local-Global	Working	Cold
							Iowa Gambling task	Memory	Hot
								Shifting	
Cohen et al., 2010	Official records	Offender	Sex offenders Healthy controls	51 84	38.63 (12.2) 33.81 (9.7)	- -	Wisconsin Card	Shifting	Cold
							Sorting Task	Shifting	Cold
							Trail Making Test	Inhibition	Cold
							Stroop Test	Planning	Cold
							Porteus Maze Test		
Combalbert et al., 2016	Official records	Offender	Offenders Healthy controls	138 138	59.72 (8.02) 68.40 (8.03)	100% 100%	FAB Conflicting Information Task	Inhibition	Cold
Crippa et al., 2015	Diagnosis	ODD	ODD + ADHD Controls + ADHD Healthy controls	12 11 68	10.00 (1.65) 9.00 (1.67) 10.40 (2.04)	91.66% 90.90% 64.71%	Honk Test	Inhibition	Cold
							Battersea Multitask paradigm	Planning	Cold
Cubillo et al., 2023	Diagnosis	CD	CD Healthy controls	753 693	14.14 (2.48) 14.39 (2.29)	33.60% 38.39%	Go/No-go Task	Inhibition	Cold
							Go/No-go Task	Inhibition	Hot
Curtis et al., 2021	Diagnosis	Offender	Violent offenders + SUD Non-violent offenders + SUD Controls + SUD	69 58 63	37.01 (8.35) 38.03 (9.35) 43.54 (11.72)	85% 77% 60%	Memory Span task	Working	Cold
							Stroop Test	Memory	Cold
							Trail Making Test	Inhibition	Cold
								Shifting	
De Brito et al., 2013	Diagnosis	ASPD, Psychopathy	ASPD + Psychopathy ASPD Healthy controls	17 28 21	40.0 (9.0) 35.8 (8.4) 35.0 (8.2)	100% 100% 100%	Digit Span (overall)	Working	Cold
							Object Alternation	memory	Cold
							Test	manipulation	Hot
							Response Reversal Task	Shifting	Hot
							Cambridge Gambling Task	Affective decision making	
Delfin et al., 2020	Official records	Offender	Offenders Healthy controls	27 20	36.63 (9.85) 33.10 (11.82)	100% 100%	Go/No-go Task	Inhibition	Cold
Déry et al., 2009	Diagnosis	CD	CD Healthy controls	105 138	9.57 (2.04) 10.08 (1.72)	74.29% 73.19%	Stroop Test	Inhibition	Cold
							Trail Making Test	Shifting	Cold
							Wisconsin Card Sorting Task	Shifting	Cold
Dolan & Lennox, 2013	Diagnosis	CD	CD CD + ADHD Healthy controls	72 35 22	16.41 (0.68) 15.97 (0.96) 15.63 (1.50)	100% 100% 100%	Stockings of Cambridge	Planning	Cold
							Intra/Extradimensional Shift	Shifting	Cold
							Intra/Extradimensional Shift	Inhibition	Cold
							Go/No-go Task	Inhibition	Hot
							Card Playing Task	Choice impulsivity	Hot
Dolan, 2012	Diagnosis, official records	ASPD, offender	Offenders + ASPD (LP) Offenders + ASPD (MP) Offenders + ASPD (HP) Healthy controls	35 28 33 49	37.18 (10.48) 35.04 (10.12) 38.79 (11.42) 33.69 (10.24)	100% 100% 100% 100%	Stockings of Cambridge	Planning	Cold
							Go/No-go Task	Inhibition	Cold
							Intra/Extradimensional Shift	Shifting	Cold

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF	
Easton et al., 2008	Official records	Offender	Violent offenders + SUD	9	40 (9.9)	100%	Wisconsin Card Sorting Task	Shifting	Cold	
			Controls + SUD	9	40 (9.9)	100%	Digit Span Backward	Working memory	Cold	
			Healthy controls	7	40 (9.9)	100%	Stroop Test	memory manipulation	Cold	
								Continuous Performance Task	Inhibition	Cold
Enticott et al., 2008	Official records	Offender	Offenders + schizophrenia	18	36.11 (10.78)	72.22%	Trail Making Test	Inhibition	Cold	
			Healthy controls	18		66.67%	Iowa Gambling Task	Affective decision making	Cold	
Euler et al., 2014	Diagnosis	CD	CD	20	14.25 (1.52)	100%	Stroop Test	Inhibition	Cold	
			Healthy controls	20	14.15 (0.88)	100%				
Ezpeleta & Granero, 2015	Diagnosis	ODD	ODD	51	3.87 (0.30)	56.9%	Continuous Performance Task	Inhibition	Cold	
			ODD + ADHD	10	3.69 (0.31)	40.0%				
			Controls + ADHD	23	3.74 (0.33)	73.9%				
			Healthy controls	538	3.76 (0.33)	48.5%				
Fairchild et al., 2009	Diagnosis	CD	CD (adolescent-onset)	34	15.54 (0.90)	100%	Wisconsin Card Sorting Task	Shifting	Cold	
			CD (early-onset)	38	15.75 (0.75)	100%				
			Healthy controls	84	15.77 (0.82)					
Feilhauer et al., 2012	Official records	Offender	Offenders	53	16.11 (0.64)	100%	Go/No-go Task	Inhibition	Cold	
			Healthy controls	64	15.92 (0.63)	100%				
Fonseca-Parra & Rey-Anacona, 2013	Diagnosis	ODD	ODD	13	8.69 (2.06)	100%	Wisconsin Card Sorting Task	Shifting	Cold	
			Healthy controls	18	9.28 (1.87)	100%	Pyramid of Mexico	Planning	Cold	
Franke et al., 2019	Official records	Offender	Sex offenders	15	50.5 (11.4)	100%	Go/No-go Task	Inhibition	Cold	
			Healthy controls	15	48.1 (11.0)	100%	Tower of London	Planning	Cold	
Ginsberg et al., 2010	Official records	Offender	Offenders + ADHD	30	34.4	100%	Digit Span (overall)	Working memory	Cold	
			Controls + ADHD	20	33.4	100%	Continuous Performance Task	memory maintenance	Cold	
			Healthy controls	18	35.2	100%		Inhibition		
Glenn et al., 2017	Diagnosis	ODD, CD	ODD/CD	33	11.39 (2.03)	87.9%	Stop Signal Task	Inhibition	Cold	
			ODD/CD + ADHD	133	10.44 (1.93)	89.5%	Stockings of Cambridge	Planning	Cold	
			Controls + ADHD	99	10.60 (1.78)	85.9%				
Gobbi et al., 2020	Official records	Offender	Offenders	126	44.9 (10.0)	81.75%	Wisconsin Card Sorting Task	Shifting	Cold	
			Healthy controls	121	44.5 (11.2)	74.38%	Iowa Gambling Task	Affective decision making	Hot	
								Memory Span Task	Working memory	Cold
								Tower of London	memory maintenance	Cold
Gonzalez-Gadea et al., 2014	Official records	Offender	Offenders	30	16.67 (0.54)	100%	Go/No-go Task	Inhibition	Cold	
			Healthy controls	16	16.00 (0.63)	100%	Hayling Test	Inhibition	Cold	
								Digit Span Backward	Working memory	Cold
								Memory Span Task	memory manipulation	Cold
								FAB Conflicting Information Task	Working memory maintenance	Cold
Grant et al., 2012	Official records	Offender	Offenders	14	21.0 (3.1)	64.3%	Cambridge Gambling Task	Affective decision making	Hot	
			Healthy controls	95	20.0 (2.8)	68.4%	Spatial Working Memory Task	Working memory	Cold	
										Cold

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
							Intra/Extradimensional Shift Stop Signal Task	maintenance Shifting Inhibition	
Guan et al., 2015	Official records	Offender	Offenders Healthy controls	20 20	16.58 (0.77) 16.23 (0.71)	100% 100%	Go/No-go Task	Inhibition	Cold
Gunn et al., 2018	Diagnosis	ASPD	ASPD + SUD Controls + SUD Healthy controls	139 309 185	- - -	- - -	Delay Discounting Task	Choice impulsivity	Hot
Habermeyer et al., 2013	Official records	Offender	Pedophilic sex offenders Healthy controls	11 7	49.0 (12.5) 47.0 (8.6)	100% 100%	Go/No-go Task	Inhibition	Cold
Hanlon et al., 2013	Official records	Offender	Offenders + schizophrenia Controls + schizophrenia	7 7	31.00 (5.45) 27.60 (3.23)	100% 100%	Trail Making Test Wisconsin Card Sorting Task Digit Span Backward	Shifting Shifting Working memory manipulation	Cold Cold Cold
Herrero et al., 2018	Official records	Offender	Sex offenders Child Sex offenders Non-sex offenders Healthy controls	26 17 35 32	37.8 (8.87) 44.0 (11.5) 34.84 (8.17) 29.0 (2.37)	100% 100% 100% 100%	Letter Memory Task Number Letter Switch Simon Task	Updating Shifting Inhibition	Cold Cold Cold
Herrero et al., 2010	Official records	Offender	Offenders Healthy controls	24 32	33.6 (7.5) 29.0 (2.4)	100% 100%	Letter Memory Task Number Letter Switch Simon Task	Updating Shifting Inhibition	Cold Cold Cold
Hobson et al., 2011	Diagnosis	ODD, CD	ODD/CD Healthy controls	28 34	12.64 (1.98) 13.13 (1.99)	67.86% 73.53%	Go/No-go Task Stop Signal Task Continuous Performance Test Wisconsin Card Sorting Task Iowa Gambling Task	Inhibition Inhibition Shifting Affective decision making	Cold Cold Cold Hot
Hoppenbrouwers et al., 2013	Official records, diagnosis	Offender, psychopathy	Offenders + psychopathy Healthy controls	13 15	34.2 (9.2) 34.0 (9.9)	100% 100%	Letter Number Sequencing Task	Working memory manipulation	Cold
Hummer et al., 2015	Diagnosis	CD, ODD	CD/ODD CD/ODD + ADHD Healthy controls	14 19 33	15.3 (1.3) 15.4 (1.5) 15.3 (1.4)	57.0% 84.0% 73.0%	Stroop Test Digit Span (overall)	Inhibition Working memory maintenance	Cold Cold
Hummer et al., 2011	Diagnosis	CD, ODD	CD/ODD CD/ODD + ADHD Healthy controls	23 25 25	14.8 (1.3) 14.7 (1.2) 15.1 (1.4)	56.52% 76.0% 52.0%	Stroop Test Digit Span (overall) Continuous Performance Task	Inhibition Working memory maintenance Inhibition	Cold Cold Cold
Hwang et al., 2016	Diagnosis	CD/ODD	CD/ODD (low CU) CD/ODD (high CU) Healthy controls	17 18 28	14.78 (2.39) 14.56 (1.84) 12.88 (2.03)	70.0% 56.0% 54.0%	Stroop Test Stroop Test	Inhibition Inhibition	Cold Hot
Iria et al., 2012	Official records, diagnosis	Offender, psychopathy	Offenders + psychopathy Offenders Controls + psychopathy Healthy controls	25 37 12 39	40.76 (10.03) 38.70 (8.98) 37.75 (8.87) 37.87 (11.63)	100% 100% 100% 100%	Go/No-go Task	Inhibition	Hot
Iselin & DeCoster, 2009	Official records	Offender	Offenders (adolescent) Offenders (young)	44 41	15.70 (1.67) 20.86	100% 100%	Continuous Performance Task	Inhibition	Cold

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
			adult)	33	(1.47)	100%			
			Controls	35	14.52	100%			
			(adolescent)		(1.53)				
			Controls (young adult)		19.34				
					(2.93)				
Iselin & DeCoster, 2012	Official records	Offender	Offenders (adolescent)	41	15.70	100%	Stroop Test	Inhibition	Cold
			Offenders (young adult)	40	(1.67)	100%			
			Offenders (young adult)	32	20.86	100%			
			Controls (adolescent)	31	(1.47)	100%			
			Controls (adolescent)		14.48				
			Controls (young adult)		(1.54)				
			Controls (young adult)		19.34				
					(0.80)				
Jiang et al., 2016	Diagnosis	ODD	ODD	7	11.76	95.83%	Stroop Test	Inhibition	Cold
			ODD + ADHD	17	12.64	95.83%	Wisconsin Card	Shifting	Cold
			Controls + ADHD	24	12.17	91.67%	Sorting Task	Working memory	Cold
			Healthy controls	36	12.92	75.0%	Memory Span Task	memory maintenance	Cold
							Digit Span Backward	Working memory	Cold
							Stockings of Cambridge	Working memory	Cold
							Spatial Working Memory Task	manipulation	
								Planning	
								Working memory	
								maintenance	
Joyal et al., 2020	Official records	Offender	Sex offenders (child victim)	39	15.0	100%	Wisconsin Card	Shifting	Cold
			Sec offenders (peer victim)	15	(1.5)	100%	Sorting Task	Affective decision making	Hot
			Non-sex offenders	41	16.0	100%	Iowa Gambling Task	decision making	Cold
			Healthy controls	39	(1.7)	100%	Stop Signal Task	Inhibition	
					16.1				
					(1.1)				
					15.4				
					(1.7)				
Kallitsoglou, 2018	Diagnosis	CD	CD	26	7.6	78.0%	Tower of London	Planning	Cold
			CD + reading disorder	27	(0.41)	77.0%	Digit Span Backward	Working memory	Cold
			Controls + reading disorder	35	7.6	54.0%	Continuous Performance Task	memory manipulation	Cold
			Healthy controls	31	(0.34)	74.0%		Inhibition	
					7.5				
					(0.33)				
					7.7				
					(0.37)				
Kashiwagi et al., 2015	Official records	Offender	Offenders + schizophrenia	30	44.1	100%	Digit Span (overall)	Working memory	Cold
			Controls + schizophrenia	24	(11.5)	100%	Tower of London	memory maintenance	Cold
					40.3			Planning	
					(10.7)				
Kim et al., 2023	Diagnosis	ODD	ODD + ADHD	36	7.8 (1.9)	91.7%	Stroop Test	Inhibition	Cold
			Controls + ADHD	307	7.7 (1.8)	83.7%	Trail Making Test	Shifting	Cold
			Healthy controls	128	8.2 (2.5)	50.8%			
Kleine Deters et al., 2020	Diagnosis	CD, ODD	ODD	44	12.3	72.7%	Delayed Match to Sample	Working memory	Cold
			CD	48	(2.6)	87.5%		memory maintenance	
			Healthy controls	86	13.5	57.0%			
					(3.0)				
					13.3				
					(2.7)				
Krakowski et al., 2015	Official records	Offender	Offenders	16	41.7	93.7%	Go/No-go Task	Inhibition	Hot
			Healthy controls	22	(19.6)	77.3%	Switch Task	Shifting	Cold
					41.4				
					(9.5)				
Krischer et al., 2008	Official records	Offender	Offenders	33	18.12	0.0%	Self-ordered Pointing Task	Working memory	Cold
			Healthy controls	20	(1.05)	0.0%	Affective Self-ordered Pointing Task	memory maintenance	Hot
					17.05			Working memory	
					(0.69)			memory maintenance	
Lin & Gau, 2017	Diagnosis	CD, ODD	CD + ADHD	76	13.21	93.42%	Digit Span Backward	Working memory	Cold
			ODD + ADHD	133	(1.70)	79.70%	Spatial Working Memory	memory manipulation	Cold
			Controls + ADHD	273	12.01	78.39%	Memory	manipulation	Cold
			Healthy controls	347	(1.57)	58.06%	CANTAB Spatial Span	Working memory	
					12.18			memory maintenance	
					(1.66)				

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
					12.90 (2.05)			Working memory maintenance	
Lishak et al., 2021	Official records	Offender	Domestic violent offenders	54	37.07 (9.27)	100%	Go/No-go Task	Inhibition	Cold
			Domestic violent offenders (+ additional criminality)	56	36.43 (8.5)	100%	Digit span	Working memory	Cold
			Healthy controls	82		100%			
					37.77 (9.55)				
Liu et al., 2023	Diagnosis	CD, ODD	CD/ODD + ADHD	1026	10.6 (2.62)	83.92%	Trail Making Test	Shifting	Cold
			Controls + ADHD	1026	10.36 (2.55)	86.74%	Stroop Test	Inhibition	Cold
			Healthy controls	406		61.58%			
					9.9 (1.65)				
Luman et al., 2009	Diagnosis	ODD	ODD + ADHD	18	9.83 (1.5)	69.0%	Stop Signal Task	Inhibition	Cold
			Controls + ADHD	20	8.83 (1.42)	69.0%			
			Healthy controls	50	9.5 (1.25)	56.0%			
					31.5 (1.25)				
Majorek et al., 2009	Official records	Offender	Offenders + schizophrenia	33	31.5 (1.25)	97.97%	Wisconsin Card Sorting Task	Shifting	Cold
			Controls + schizophrenia	38	35.9 (11.6)	47.37%	BADS Zoo Test	Planning	Cold
			Healthy controls	29	37.0 (13.7)	34.48%			
					9.85 (1.91)				
Manfei Xu et al., 2017	Diagnosis	ODD	ODD	14	9.85 (1.91)	100%	Stockings of Cambridge	Planning	Cold
			ODD + ADHD	29	10.11 (1.74)	100%	Spatial Working Memory	Working memory	Cold
			Controls + ADHD	39	9.16 (1.82)	100%	CANTAB Spatial Span	Working memory	Cold
			Healthy controls	52	10.02 (2.10)	100%	Wisconsin Card Sorting Task	Working memory maintenance	Cold
					9.16 (1.82)		Digit Span Backward	Working memory manipulation	Cold
					28.78 (6.32)				
Marceau et al., 2008	Official records	Offender	Offenders	584	28.78 (6.32)	100%	Trail Making Test	Shifting	Cold
			Psychiatric controls	494	28.94 (7.09)	100%	Wisconsin Card Sorting Task	Shifting	Cold
			Healthy controls	132	27.92 (7.03)	100%			
					38.04 (8.62)				
Massau et al., 2017	Official records	Offender	Child sexual offender + pedophilia	45	38.04 (8.62)	100%	Stop Signal Task	Inhibition	Cold
			Child sexual offender – pedophilia	19		100%	Information Sampling Task	Inhibition	Cold
			Controls + pedophilia	45	40.26 (12.71)	100%	Intra/Extradimensional Shift	Shifting	Cold
			Healthy controls	49		100%	Stockings of Cambridge	Planning	Cold
					36.51 (9.46)		Memory Span Task	Working memory	Cold
					36.43 (6.70)				
Medrano Nava et al., 2022	Diagnosis	CD, ODD	ODD + ADHD	10	7-9	100%	BANFE Labyrinths	Planning	Cold
			ODD + ADHD + ADD	10	7-9	100%	BANFE Card Classification	Shifting	Cold
			ODD + ADHD + LD	8	7-9	100%	BANFE Card game	Affective decision making	Hot
			ODD + CD + ADHD + LD	9	7-9	100%	BANFE Hanoi	Decision making	Cold
			Controls + ADHD	10	7-9	100%	BANFE Visuospatial WM	Planning	Cold
			Controls + ADHD + ADD	8	7-9	100%	BANFE Stroop B	Working memory	Cold
					31.5 (10.02)				
Meier et al., 2012	Official records	Offender	Offenders	13	31.5 (10.02)	100%	Go/No-go Task	Inhibition	Cold
			Controls + ADHD	13	31.3 (9.73)	100%	Go/No-go Task	Inhibition	Hot
			Healthy controls	13	28.5 (5.49)	100%			

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
Mellentin et al., 2013	Diagnosis	ASPD	ASPD + SUD Controls + SUD Healthy controls	16 30 17	36.81 (8.60) 34.18 (8.01) 36.65 (10.62)	93.75% 66.67% 76.47%	Iowa Gambling Task	Affective decision making	Hot
Moeller et al., 2014	Diagnosis	IED	IED Controls + SUD Healthy controls	11 17 22	33.5 (7.1) 32.6 (6.4) 43.2 (6.5)	100% 100% 100%	Stroop Test	Inhibition	Cold
Molleman et al., 2021	Official records	Offender	Offenders Offenders (psychiatric) Healthy controls	57 119 66	31.2 38.9 36.1	100% 100% 100%	Stroop Test Continuous Performance Test	Inhibition Inhibition	Hot Cold
Molleman et al., 2022	Official Records	Offender	Offenders Offenders (psychiatric) Healthy controls	57 119 65	31.2 (8.5) 38.9 (9.2) 36.1 (11.8)	100% 100% 100%	Reversal learning task StopSignal task	Shifting Inhibition	Hot Cold
Munkvold et al., 2014	Diagnosis	ODD	ODD ODD + ADHD Controls + ADHD Healthy controls	10 15 50 160	9.3 (0.8) 9.7 (1.0) 9.6 (0.8) 9.4 (1.0)	70.0% 100% 78.0% 55.63%	Continuous Performance Test	Inhibition	Cold
Muscatallo et al., 2014	Official records	Offender	Offenders Healthy controls	147 150	17.07 (1.1) 17.29 (0.9)	100% 100%	Stroop Test Wisconsin Card Sorting Test	Inhibition Shifting	Cold Cold
*Närhi, Lehto-Salo, Ahonen, & Marttunen, 2010	Diagnosis	CD/ODD	CD/ODD Healthy controls	77 48	15.35 (1.08) 14.79 (0.81)	48.1% 35.4%	Trail Making Test Wisconsin Card Sorting Test Tower of Hanoi Digit Span Backward	Shifting Shifting Planning Working memory manipulation	Cold Cold Cold Cold
Neves & Pinho, 2018	Official records	Offender	Offenders Healthy controls	59 59	34.4 (7.9) 33.7 (8.9)	51.0% 49.0%	Stroop Test Porteus Maze Test	Inhibition Planning	Cold Cold
Nishinaka et al., 2016	Official records	Offender	Offenders Healthy controls	71 54	42.79 (11.92) 42.06 (11.43)	84.51% 88.89%	Two Back Task Iowa Gambling Task Groton Maze Learning Task Stop Signal Task Digit Span Backward Delay Discounting Task	Updating Affective decision making Planning Inhibition Working memory maintenance Choice impulsivity	Cold Hot Cold
Noordermeer et al., 2020	Diagnosis	ODD	ODD + ADHD Controls + ADHD Healthy controls	82 82 82	16.3 (3.1) 16.3 (3.0) 16.1 (3.3)	67.0% 67.0% 67.0%	Stop Signal Task Digit Span Backward Delay Discounting Task	Inhibition Working memory maintenance	Cold Cold Hot
Nordvall, Jonsson, & Neely, 2017	Official records	Offender	Offenders Healthy controls	31 40	17.0 (1.7) 17.5 (1.3)	51.61% 50.0%	Stroop Test Switch Task Verbal n-back Task Memory Span Task	Inhibition Shifting Updating Working memory maintenance	Cold Cold Cold Cold
Nordvall, Neely, & Jonsson, 2017	Official records	Offender	Offenders Healthy controls	37 39	17.7 (1.4) 17.3 (1.1)	62.16% 51.28%	Stroop Test Switch Task Verbal n-back Task Memory Span Task	Inhibition Shifting Updating Working memory maintenance	Cold Cold Cold Cold
Ostrosky et al., 2012	Official records	Offender	Offenders Healthy controls	82 76	44 (9.0) 42 (8.5)	100% 100%	Porteus Maze Test BANFE Visuospatial WM Task Stroop Test BANFE Subtraction Task BANFE Gambling Task Wisconsin Card Sorting Task BANFE Maze Tower of Hanoi	Planning Working memory maintenance Inhibition Working memory maintenance Affective decision making Shifting	Cold Cold Cold Cold Hot Cold Cold Cold Cold

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
Pajer et al., 2008	Diagnosis	CD	CD	52	16.5	0.0%	Stroop Test	Planning	Cold
			Healthy controls	41	(0.95)	0.0%	Digit Span (overall)	Inhibition	Cold
Palix et al., 2022	Official records	Offender	Violent offenders	22	39.27	100%	Go/No-Go	Working memory	Cold
			Healthy controls	24	(12.12)	100%	Wisconsin Card Sorting Task	Shifting	Cold
Pasion et al., 2018	Official records	Offender	Offenders	56	32	100%	Stroop Test	Inhibition	Cold
			Healthy controls	48	(11.6)	100%	Spatial N-back Task	Updating	Cold
Patiz & Bayraktar, 2023	Official records	Offender	Offenders	30	15.97	76.7%	Wisconsin Card	Shifting	Cold
			Victims	30	(1.35)	63.3%	Sorting Task	Planning	Cold
			Healthy controls	25	-	44.0%	Tower Test		
Pera-Guardiola et al., 2016	Official records, diagnosis	Offender, ASPD	Offenders + ASPD (HP)	31	33.45	100%	Wisconsin Card	Shifting	Cold
			Offenders + ASPD (LP)	13	(4.84)	100%	Sorting Task		
			Offenders + ASPD (LP)	24	39.93	100%			
			Healthy controls	24	(10.01)				
Perino et al., 2019	Official records	Offender	Offenders	24	16.2	50.0%	Go/No-go Task	Inhibition	Cold
			Healthy controls	24	(1.2)	50.0%			
Poon, 2020	Official records	Offender	Offenders	122	15.28	50.82%	Cambridge Gambling Task	Affective decision making	Hot
Prehn et al., 2013	Official records, diagnosis	Offender, ASPD	Offenders + ASPD	15	27.87	100%	Verbal N-back Task	Updating	Cold
			Healthy controls	17	(9.86)	100%	Verbal N-back Task	Updating	Hot
Price et al., 2013	Official records	Offender	Sex offenders	27	43.31	100%	Hayling Test	Inhibition	Cold
			Non-sex offenders	21	(9.82)	100%	Stroop Test	Inhibition	Hot
			Healthy controls	38	35.24	100%			
Price et al., 2014	Official records	Offender	Sex offenders	24	15.96	100%	Hayling Test	Inhibition	Cold
			Non-sex offenders	21	(1.27)	100%	Stroop Test	Inhibition	Hot
			Healthy controls	21	16.75	100%			
Qian et al., 2010	Diagnosis	ODD	ODD + ADHD	53	9.25	79.25%	Stroop Test	Inhibition	Cold
			Controls + ADHD	89	(1.79)	62.92%	Trail Making Test	Shifting	Cold
			Healthy controls	116	9.07	83.62%	Digit Span Backward	Working memory	Cold
Raszkwicz, 2010	Diagnosis	CD	CD	34	15.51	82.35%	Continuous	Planning	Cold
			Healthy controls	35	(0.82)	45.71%	Performance Test	Inhibition	Cold
Rhodes et al., 2012	Diagnosis	ODD	ODD	21	9.91	100%	Memory Span Task	Working memory	Cold
			ODD + ADHD	27	(1.94)	100%	CANTAB Spatial	memory	Cold
			Controls + ADHD	21	9.77	100%	Span Task	maintenance	Cold
			Healthy controls	26	(1.82)	100%	Verbal Executive	Working	Cold
			Function Task		9.47			memory	Cold
			Spatial Executive		(2.01)			maintenance	Cold
			Function Task		9.69			Working	Cold
Spatial Working		(1.46)			memory	Cold			
Memory Task					manipulation				
Delayed Match to					Working				
Sample Task					memory				
Spatial N-back Task					manipulation				
					Working				
					memory				
					maintenance				
					Working				

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
Romero-Martínez, Lila, & Moya-Albiol, 2021	Official records	Offender	Domestic offenders Healthy controls	89 39	40.10 (11.05)	100% 100%	Wisconsin Card Sorting Task Switch Task	memory maintenance Updating Shifting Shifting	Cold Cold
Romero-Martínez, Lila, Vitoria-Estruch, & Moya-Albiol, 2021	Official records	Offender	Domestic offenders Healthy controls	47 41	38.61 (11.40)	100% 100%	Trail Making Test Wisconsin Card Sorting Task	Shifting Shifting	Cold Cold
Romero-Martínez, Lila, & Moya-Albiol, 2021	Official records	Offender	Domestic offenders Healthy controls	51 39	41.72 (11.01)	100% 100%	CANTAB Spatial Span Cambridge Gambling Task	Working memory maintenance Affective decision making	Cold Hot
Romero-Martínez, Lila, Vitoria-Estruch, & Moya-Albiol, 2021	Official records	Offender	Intimate partner violence perpetrators + SUD Intimate partner violence perpetrators - SUD Healthy controls	104 120 82	40.08 (9.49) 40.07 (9.17)	100% 100% 100%	Digit Span Task Wisconsin Card Sorting Task Key Test	Working memory Shifting Planning	Cold Cold Cold
Rosburg et al., 2018	Official records	Offender	Child sex offenders Healthy controls	40 21	40.30 (10.59) 36.5 (10.3)	100% 100%	Go/No-go Task	Inhibition	Cold
Rubia et al., 2009	Diagnosis	CD, ODD	CD/ODD Controls + ADHD Healthy controls	13 20 20	12.9 (2.2) 13.2 (1.4) 14.0 (1.9)	100% 100% 100%	Simon Task	Inhibition	Cold
Rubia et al., 2008	Diagnosis	CD, ODD	CD/ODD Controls + ADHD Healthy controls	13 20 20	13.0 (1.0) 13.2 (1.5) 14.0 (2.0)	100% 100% 100%	Stop Signal Task	Inhibition	Cold
Rubia et al., 2010	Diagnosis	CD, ODD	CD/ODD Controls + ADHD Healthy controls	14 14 20	12.6 (2.3) 13.3 (1.1) 13.5 (1.9)	100% 100% 100%	Switch Task	Shifting	Cold
Saarinen et al., 2015	Diagnosis	CD, ODD	CD/ODD Healthy controls	26 26	10.1 (1.2)		Spatial N-back Task	Updating	Cold
Barbosa and Monteiro, 2008	Official records	Offender	Offenders Healthy controls	30 30	39.3 (9.89) 32.7 (11.8)	100% 100%	Wisconsin Card Sorting Task BADS Action Program BADS Zoo Test BADS Key Test BADS Modified Sex Elements Task	Shifting Planning Planning Planning	Cold Cold Cold Cold
Schiffer & Vonlaufen, 2011	Official records	Offender	Non-sex offenders Child sex offenders Child sex offenders + pedophilia Healthy controls	16 15 15 17	37.4 (9.1) 44.2 (7.9) 38.7 (8.9)	100% 100% 100% 100%	Wisconsin Card Sorting Task Go/No-go Task Trail Making Test Memory Span Task Tower of London	Shifting Inhibition Shifting Working memory maintenance Planning	Cold Cold Cold Cold Cold
Schiffer et al., 2014	Official records	Offender	Offenders + SUD Healthy controls	21 23	37.7 (10.2) 35.2 (8.2) 34.1 (8.9)	100% 100%	Stroop Test	Inhibition	Cold
Schoemaker et al., 2012	Diagnosis	CD, ODD	CD/ODD CD/ODD + ADHD	33 52	4.32 (0.69) 4.51	81.8% 82.7%	Go/No-go Task Delayed Gratification Task	Inhibition Choice impulsivity	Cold Hot

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
			Controls + ADHD	61	(0.57)	80.3%	Shape School Task	Inhibition	Hot
			Healthy controls	56	4.60	69.6%	Object Alternation Test	Shifting	Cold
Schoorl et al., 2016	Diagnosis	CD, ODD	CD/ODD	65	10.3	100%	Ultimatum Game (child version)	Affective decision making	Hot
			Healthy controls	38	(1.28)	100%			
Schoorl et al., 2018	Diagnosis	CD, ODD	CD/ODD	65	10.3	100%	Digit Span Backward	Working memory	Cold
			Healthy controls	32	(1.28)	100%	Shifting Attention Set Task	memory manipulation	Cold
					9.9		Digit Span Backward	Shifting	Hot
					(1.24)		Shifting Attention Set Task	Working memory manipulation	Hot
Schwenck et al., 2017	Diagnosis	CD, ODD	CD/ODD	19	13.81	100%	Iowa Gambling Task	Affective Decision Making	Hot
			Healthy controls	24	(1.56)	100%			
Seruca & Silva, 2016	Official records	Offender	Offenders	42	33.67	100%	Trail Making Test	Shifting	Cold
			Healthy controls	28	(8.66)	100%	Digit Span Backward	Working memory	Cold
					35.57		Stroop Test	memory maintenance	Cold
					(8.95)		Porteus Maze Test	Inhibition	Cold
Shuai et al., 2011	Diagnosis	CD, ODD	CD/ODD + ADHD	38	10.34	100%	Stroop Test	Inhibition	Cold
			Controls + ADHD + LD	38	(2.53)	100%	Trail Making Test	Shifting	Cold
			Controls + ADHD	76	10.38	100%	Tower of Hanoi	Planning	Cold
			Healthy controls	76	(2.56)	100%			
					10.24				
					(2.40)				
					10.21				
					(2.30)				
Stratton et al., 2018	Official records	Offender	Offenders + schizophrenia	25	37.12	96.0%	Trail Making Test	Shifting	Cold
			Controls + schizophrenia	25	(12.91)	92.0%	Wisconsin Card	Shifting	Cold
			Healthy controls	25	33.89	84.0%	Sorting Test		
					(5.74)				
					30.92				
					(6.55)				
Swann et al., 2009	Diagnosis, official records	ASPD, offender	Offenders + ASPD	34	38.7	100%	Delay Discounting Task	Choice impulsivity	Hot
			Healthy controls	30	(10.3)	100%	Continuous Performance Task	Inhibition	Cold
					31.5		Risky Choice Task		
					(9.5)				
Syngelaki et al., 2009	Official records	Offender	Offenders	102	16.33	100%		Affective decision making	Hot
			Healthy controls	83	15.77	100%			
Szczypiński et al., 2022	Diagnosis, official records	Offender	Sex offenders (to child)	13	43.69	100%	Go/No-go Task	Inhibition	Cold
			Healthy controls	18	(7.61)	100%			Hot
					35.06				
					(7.69)				
Tung & Chhabra, 2011	Official records	Offender	Offenders	40	15.7	100%	Stroop Test	Inhibition	Cold
			Healthy controls	40	15.0	100%			
Turner et al., 2020	Official records	Offender	Sex offenders (to child)	70	41.66	100%	Iowa Gambling Task	Affective decision making	Hot
			Sex offenders (to adult)	49	(13.85)	100%	Wisconsin Card		Cold
			Non-sex offenders	54	37.73	100%	Sorting Task	Shifting	Cold
			Healthy controls	73	(11.44)	100%	Tower of London	Planning	Cold
					32.72		Trail Making Test	Shifting	Cold
					(9.81)		Stroop Test	Inhibition	
					38.10				
					(12.97)				
Turner et al., 2018	Official records	Offender	Sex offenders (to child)	63	42.14	100%	Go/No-go Task	Inhibition	Hot
			Healthy controls	63	(13.19)	100%	Iowa Gambling Task	Affective decision making	Hot
					27.0		Game of Dice Task	Affective decision making	Hot
					(4.75)				
Urazán-Torres et al., 2013	Diagnosis	CD	CD	39	9.54	77.0%	Wisconsin Card	Shifting	Cold
			Healthy controls	39	(1.99)	49.0%	Sorting Task	Planning	Cold
					9.26		Pyramid of Mexico		
					(2.11)				
Uytun et al., 2017	Diagnosis	CD	CD + ADHD	10	-	100%	Wisconsin Card	Shifting	Cold
			Controls + ADHD	10	-	100%	Sorting Task	Inhibition	Cold
			Healthy controls	10	-	100%	Stroop Test		

(continued on next page)

Appendix 3 (continued)

Study	Assessment Method	ASB classification	Participant Groups	n	Mean Age (SD)	Males (%)	EF task	EF domain	Hot/Cold EF
Van der Meere et al., 2008	Diagnosis	CD	CD + IQ 70-75	21	12.08	100%	Continuous Performance Test	Inhibition	Cold
			Controls + IQ 70-75	19	(1.08)	100%			
Vilà-Balló et al., 2014	Official records	Offender	Offenders	17	18.3	100%	Stop Signal Task	Inhibition	Cold
			Healthy controls	17	(0.3)	100%			
Vila-Ballo et al., 2015	Official records	Offender	Offenders	14	18.43	100%	Wisconsin Card Sorting Task	Shifting	Cold
			Healthy controls	14	(1.16)	100%			
Vitoria-Estruch et al., 2018	Official records	Offender	Offender (IPV) + SUD	28	40.21	100%	Switch Task	Shifting	Cold
			Offender (IPV)	35	(11.90)	100%	Memory Span Task	Working memory	Cold
			Healthy controls	37	39.34	100%	Digit Span Backward	memory	Cold
					(9.83)		Wisconsin Card Sorting Task	maintenance	Cold
					41.75		BADS Zoo Test	Working memory	Cold
					(11.0)		BADS Key Test	manipulation	Cold
Vloet et al., 2011	Diagnosis	CD, ODD	CD/ODD + ADHD	17	12.5	100%	Go/No-go Task	Inhibition	Cold
			Healthy controls	17	(1.6)	100%	Go/No-go Task	Inhibition	Hot
Weidacker et al., 2022	Official records	Offender	Sex offenders + pedophilia	11	43.55	100%	Stroop Test	Inhibition	Cold
			Controls + pedophilia	8		100%			
			Healthy controls	10	33.25	100%			
Zhu et al., 2014	Diagnosis	ODD	ODD	11	11.5	100%	Stop Signal Task	Inhibition	Cold
Zhu et al., 2021	Diagnosis	CD, ODD	Healthy controls	10	11.7	100%	Stroop Test	Inhibition	Cold
			CD/ODD	26	11.1	68.2%			
Zou et al., 2013	Official records	Offender	CD/ODD + ADHD	22	11.7	57.7%	Stroop Test	Inhibition	Hot
			Controls + ADHD	30	12.7	76.7%			
			Healthy controls	20	12.8	75.00%			
			Offenders (violent)	107	16.5	100%			
		Offenders (non-violent)	107	(0.6)	100%	Intra-Extradi-dimensional Shift	Shifting	Cold	
		Healthy controls	107	(0.6)	100%	Stockings of Cambridge	Working memory	Cold	
				16.5		Spatial Working Memory Task	maintenance	Cold	
				(0.6)					

Appendix 4

Executive Function	Neuropsychological test	Outcome Measure	Quality	Freq
Higher-order EF	BADS action program	Overall performance	Low	1
Higher-order EF	BADS Key Test	Overall performance	Low	2
Higher-order EF	BADS Key Test	Strategy	Low	1
Higher-order EF	BADS modified six elements	Overall performance	Low	1
Higher-order EF	BADS Zoo Test	Overall performance	Low	5
Higher-order EF	BANFE Gambling task	% disadvantage decks	High	1
Higher-order EF	BANFE Gambling task	Overall performance	High	1
Higher-order EF	BANFE Labyrinths	Completion time	Low	8
Higher-order EF	BANFE Labyrinths	Planning time	Low	8
Higher-order EF	BANFE Maze	Completion time	Low	1
Higher-order EF	BANFE Maze	Planning error	Low	1
Higher-order EF	Battersea Multitask Paradigm	BMP coherence	Low	2
Higher-order EF	Battersea Multitask Paradigm	BMP performance	Low	2
Higher-order EF	Battersea Multitask Paradigm	BMP planning	Low	2
Higher-order EF	Cambridge Gambling task	Decision-making	High	3
Higher-order EF	Cambridge Gambling task	Decision-making	Medium	1
Higher-order EF	Cambridge Gambling task	Delay aversion	Medium	1
Higher-order EF	Cambridge Gambling task	Deliberation time	Low	3
Higher-order EF	Cambridge Gambling task	Deliberation time	Medium	1

(continued on next page)

Appendix 4 (continued)

Executive Function	Neuropsychological test	Outcome Measure	Quality	Freq
Higher-order EF	Cambridge Gambling task	Number of bankruptcies	Low	1
Higher-order EF	Cambridge Gambling task	Proportion bet	Medium	1
Higher-order EF	Cambridge Gambling task	Risk adjustment	High	3
Higher-order EF	Cambridge Gambling task	Risk adjustment	Medium	1
Higher-order EF	Cambridge Gambling task	Risk taking	Medium	1
Higher-order EF	Child version of the Ultimatum Game (UG)	Emotional decision making	Medium	1
Higher-order EF	D-KEFS Tower Test	Total achievement scaled score	Low	2
Higher-order EF	Game of Dice Task	Net score	High	1
Higher-order EF	Groton Maze Learning Task	Completion time	Low	1
Higher-order EF	Iowa Gambling task	Advantageous Deck Picks	High	1
Higher-order EF	Iowa Gambling task	Net score	High	18
Higher-order EF	Iowa Gambling task	Number of risky picks	High	4
Higher-order EF	Iowa Gambling task	Ratio good/bad deck	High	4
Higher-order EF	Iowa Gambling task	Risk cards percentage	High	8
Higher-order EF	Iowa Gambling task	Total punctuation	High	8
Higher-order EF	One Touch Stocking of Cambridge	Mean choices to correct	Low	2
Higher-order EF	One Touch Stocking of Cambridge	Problems solved on first choice	Low	2
Higher-order EF	Porteus Maze test	Number of mazes completed	Low	4
Higher-order EF	Porteus Maze test	Qualitative error score	Low	2
Higher-order EF	Porteus Maze test	Total number of errors	Low	2
Higher-order EF	Pyramid of Mexico	Design with the fewest possible moves	Low	1
Higher-order EF	Pyramid of Mexico	Number of moves	Low	1
Higher-order EF	Pyramid of Mexico	Total no of solved problems	Low	3
Higher-order EF	Risky Choice Task	Mean number of points	High	1
Higher-order EF	Stockings of Cambridge	Average moves	Low	14
Higher-order EF	Stockings of Cambridge	Initial thinking time	Low	4
Higher-order EF	Stockings of Cambridge	Problems solved in minimum moves	Low	21
Higher-order EF	Stockings of Cambridge	Subsequent thinking time	Low	4
Higher-order EF	Tower of Hanoi	Completion time	Low	15
Higher-order EF	Tower of Hanoi	Error steps	Low	3
Higher-order EF	Tower of Hanoi	Movements	Low	8
Higher-order EF	Tower of Hanoi	Points obtained	Low	1
Higher-order EF	Tower of Hanoi	Steps needed to complete	Low	5
Higher-order EF	Tower of London	Completion time	Low	7
Higher-order EF	Tower of London	Failed attempts	Low	9
Higher-order EF	Tower of London	Problems solved in minimum moves	Low	8
Higher-order EF	Tower of London	Total in excess of the minimum	Low	4
Higher-order EF	Tower of London	Total no of solved problems	Low	6
Higher-order EF	Tower of London	Tower of London	Low	1
Inhibition	Card Playing Task	Payoff	Low	2
Inhibition	Continuous performance test	Commission errors	High	37
Inhibition	Delay discounting task	Delay discounting k-values	High	6
Inhibition	Delay discounting task	SKIP longest delay	High	1
Inhibition	Delay discounting task	SKIP Shortest delay	High	1
Inhibition	Delay discounting task	TCIP Maximum consecutive delayed responses	High	1
Inhibition	Delay discounting task	TCIP Percent immediate responses	High	1
Inhibition	Delayed Gratification Task	Complied with rules	High	4
Inhibition	Delayed Gratification Task	Impulsive Choices	High	2
Inhibition	Delayed Gratification Task	Long-term Choices	High	2
Inhibition	Delayed Gratification Task	Payoff	High	2
Inhibition	FAB Conflicting information task	Total erroneous responses	Low	3
Inhibition	Go/No-Go	dPrime	High	5
Inhibition	Go/No-Go	Percentage correct no-go responses	High	33
Inhibition	Go/No-Go	Percentage errors of commission	High	43
Inhibition	Go/No-Go	Reaction time errors of commission	High	12
Inhibition	Go/No-Go	Reaction time errors of commission	Low	1
Inhibition	Go/No-Go	Total percentage correct	Low	2
Inhibition	Go/No-Go	Total reaction time	Low	3
Inhibition	Hayling test	Category B errors	Medium	2
Inhibition	Hayling test	Impairment	High	2
Inhibition	Hayling test	Overall performance	Medium	5
Inhibition	Honk Test	Change errors	Medium	2
Inhibition	Honk Test	Stop errors	High	2
Inhibition	Information Sampling Task	Total correct	High	8
Inhibition	Intra/Extradimensional shift	Stages completed	Medium	1
Inhibition	Negative priming	Negative priming ratio	High	1
Inhibition	NEPSY-2 Inhibition	Inhibition	Medium	1
Inhibition	Shape School task	Total percentage correct	Medium	4
Inhibition	Simon task	Errors (Incongruent)	Medium	8
Inhibition	Simon task	Interference score	High	12
Inhibition	Simon task	RT card III	Medium	6
Inhibition	StopSignal task	% errors	Low	3
Inhibition	StopSignal task	% errors	Medium	1
Inhibition	StopSignal task	Errors (Incongruent)	Medium	1
Inhibition	StopSignal task	Proportion successful stops	Medium	3
Inhibition	StopSignal task	SSRT	High	22

(continued on next page)

Appendix 4 (continued)

Executive Function	Neuropsychological test	Outcome Measure	Quality	Freq
Inhibition	StopSignal task	Stop latency	Low	1
Inhibition	StopSignal task	Stop latency	Medium	5
Inhibition	Stroop Test	Errors	Medium	8
Inhibition	Stroop Test	Errors (incongruent)	Medium	44
Inhibition	Stroop Test	Interference score	High	52
Inhibition	Stroop Test	RT (incongruent)	Medium	48
Inhibition	Stroop Test	Time	Medium	8
Inhibition	Stroop Test	Total	Medium	8
Shifting	Intra/Extradimensional shift	EDS errors	High	10
Shifting	Intra/Extradimensional shift	Pre-ED errors	Medium	9
Shifting	Intra/Extradimensional shift	Reversal errors	Medium	1
Shifting	Intra/Extradimensional shift	Stages completed	Medium	10
Shifting	Intra/Extradimensional shift	Total errors	Low	2
Shifting	Intra/Extradimensional shift	Total errors adjusted	Medium	6
Shifting	Local-Global	Interference score	High	1
Shifting	NEPSY-2 Design Fluency	Cognitive flexibility	Medium	1
Shifting	NEPSY-2 Inhibition	Switching	Medium	1
Shifting	Number Letter switch	Overall performance	High	2
Shifting	Number Letter switch	Switch cost	High	4
Shifting	Object alternation test (OAT)/delayed alternation test (DAT)	% correct responses	High	4
Shifting	Object alternation test (OAT)/delayed alternation test (DAT)	Number of errors	Low	2
Shifting	Response Reversal task	Number of errors	High	2
Shifting	Reversal learning task	Error switch	Medium	2
Shifting	Reversal learning task	Number of stages completed	Low	2
Shifting	Shifting attention set task	SWitch cost	High	2
Shifting	Switch Task	% correct responses	Low	3
Shifting	Switch Task	Correct rejections	Medium	1
Shifting	Switch Task	Error switch	Medium	2
Shifting	Switch Task	RT shift	High	1
Shifting	Switch Task	RT shift	Medium	3
Shifting	Switch Task	Switch cost	High	16
Shifting	Switch Task	Switch cost	Low	1
Shifting	Trail Making Test	TMT B	Medium	3
Shifting	Trail Making Test	TMT B-A	High	11
Shifting	Trail Making Test	TMT-B	Medium	40
Shifting	Trail Making Test	TMT-B - A	High	8
Shifting	Wisconsin Card Sorting Task	% correct responses	Low	4
Shifting	Wisconsin Card Sorting Task	Categories completed	Low	35
Shifting	Wisconsin Card Sorting Task	Completion time	Low	9
Shifting	Wisconsin Card Sorting Task	Conceptual responses	Low	5
Shifting	Wisconsin Card Sorting Task	Criteria perseverations	High	1
Shifting	Wisconsin Card Sorting Task	Failure to maintain set	High	7
Shifting	Wisconsin Card Sorting Task	global score	High	1
Shifting	Wisconsin Card Sorting Task	Hits	High	8
Shifting	Wisconsin Card Sorting Task	Overall result	High	1
Shifting	Wisconsin Card Sorting Task	Overall performance	High	1
Shifting	Wisconsin Card Sorting Task	Perseverative errors	High	44
Shifting	Wisconsin Card Sorting Task	Perseverations	High	8
Shifting	Wisconsin Card Sorting Task	Perseverative responses	Medium	21
Shifting	Wisconsin Card Sorting Task	Switch cost	High	2
Shifting	Wisconsin Card Sorting Task	Total correct	Low	4
Shifting	Wisconsin Card Sorting Task	Total errors	Low	35
Shifting	Wisconsin Card Sorting Task	Total trials	Low	2
Shifting	Wisconsin Card Sorting Task	Trials to complete 1st category	Low	2
Updating	Letter Memory Task	Correct recalls	High	4
Updating	Spatial N-back	Accuracy	High	7
Updating	Spatial N-back	dPrime	High	1
Updating	Spatial N-back	RT 2-back	Medium	1
Updating	Two Back Task, TWOB	Accuracy	High	1
Updating	Verbal n-back	Accuracy	High	6
Updating	Verbal n-back	RT 2-back	Medium	4
Updating	Verbal n-back	Total accuracy	Medium	2
Updating	Verbal n-back	Updating cost	High	3
Working Memory	Affective self-ordered pointing task	Errors	High	2
Working Memory	BANFE subtraction task	Completion time	Medium	2
Working Memory	BANFE subtraction task	Overall performance	Medium	2
Working Memory	BANFE visual working memory task	Completion time	Medium	1
Working Memory	BANFE visual working memory task	Overall performance	Medium	1
Working Memory	BANFE visual working memory task	Perseverative errors	Medium	1
Working Memory	BANFE Visuospatial WM	Level	Medium	8
Working Memory	BANFE visuospatial working memory task	Level	Medium	1
Working Memory	BANFE visuospatial working memory task	Order error	Medium	1
Working Memory	BANFE visuospatial working memory task	Perseverative errors	Medium	1
Working Memory	CANTAB Spatial Span	SSP Length	High	12
Working Memory	CANTAB Spatial Span	SSP Length	Medium	1
Working Memory	Continuous performance test	Commission errors	High	1

(continued on next page)

Appendix 4 (continued)

Executive Function	Neuropsychological test	Outcome Measure	Quality	Freq
Working Memory	Delayed match to sample	Accuracy	High	14
Working Memory	Digit span Backward	Digit span backwards	High	3
Working Memory	Digit span Backward	Digit span backwards	Medium	30
Working Memory	Digit span overall	Digit span overall	Low	10
Working Memory	Letter Number Sequencing (LNS)	LNS score	High	1
Working Memory	Memory Span task	Accuracy	High	9
Working Memory	Memory Span task	Accuracy	Medium	2
Working Memory	Memory Span task	Digit span	High	5
Working Memory	Memory Span task	Digit span (WISC) + Corsi Block test	Medium	2
Working Memory	Memory Span task	SSP Length	High	17
Working Memory	Memory Span task	SSP Length	Medium	1
Working Memory	Memory Span task	Strategy	High	4
Working Memory	Memory Span task	SWM between errors 4-8 boxes	Medium	8
Working Memory	NEPSY-2 Word List Interference	Interference score	Medium	1
Working Memory	Self-ordered pointing task	Errors	High	1
Working Memory	Spatial Executive function task	Accuracy	High	4
Working Memory	spatial recognition task	Accuracy	Medium	2
Working Memory	Spatial working memory	Errors	High	1
Working Memory	Spatial working memory	SSP Length	High	1
Working Memory	Spatial working memory	Strategy	High	16
Working Memory	Spatial working memory	SWM between errors 4-8 boxes	Medium	4
Working Memory	Spatial working memory	Total errors	Low	1
Working Memory	Spatial working memory	Total errors 4-8 boxes	High	16
Working Memory	Verbal Executive function task	Accuracy	High	4

Appendix C. Appendix 5:*C.1. Newcastle – Ottawa quality assessment scale case control studies*

Note: A study can be awarded a maximum of one star for each numbered item within the Selection and Exposure categories. A maximum of two stars can be given for Comparability.

Selection1) Is the case definition adequate?

- a) yes, from judicial registration data or diagnosis *
- b) no description

2) Representativeness of the cases

- a) consecutive or obviously representative series of cases *
- b) potential for selection biases or not stated

3) Selection of Controls

- a) community controls (or controls with matched diagnosis) *
- b) convenience sample
- c) no description

4) Definition of Controls

- a) no history of criminal behaviour or antisocial behaviour diagnosis *
- b) no description of source

Comparability1) Comparability of cases and controls on the basis of the design or analysis

- a) study controls for diagnosis * (for example; ADHD is characterised by significant impairments in executive functioning)
- b) study controls for both age and gender * (both known and reliable predictors of criminal behaviour / ASB diagnosis: Mean difference in age \leq 5.0, percentage gender difference $<5\%$).

Exposure1) Ascertainment of exposure

- a) secure record (eg judicial registration data, or diagnosis) *
- b) Self report
- c) No description

2) Same method of ascertainment for cases and controls

- a) yes *
- b) no

3) Non-Response rate

- a) same rate for both groups *
- b) non respondents described
- c) rate different and no designation

Appendix D. Appendix 6

Meta-analytic results assessing differences between adult and youth samples.

D.1. Different executive functions

The three-level meta-analysis including the interaction between and main effects of age group (adults vs children) and executive function as moderators showed a significant interaction effect $F(4,1234)=4.51$ $p=.001$. Although none of the Bonferroni corrected pairwise comparisons revealed a significant interaction (see main manuscript section 3.4.1.), separate analysis per age group showed that executive function did not moderate the pooled effect sizes in children $F(4, 709)=1.33$ $p=.26$, but did in adults $F(4,525)=6.19$ $p<.001$. Bonferroni corrected pairwise comparisons indicated shifting ($\Delta d=.24$ $p<.01$) and updating ($\Delta d=.77$ $p<.01$) were more impaired than inhibition, and that updating was more impaired than higher-order EF ($\Delta d=.68$ $p<.05$).

D.2. Different neuropsychological tests

A three-level meta-analysis including neuropsychological test, age group (adult vs youth) and their interaction as moderators showed no significant interaction effect ($F(22, 1108)=1.00$ $p=.46$, or main effect of age group ($F(1, 1130)=.49$ $p=.49$). Neuropsychological test did moderate the pooled effect sizes ($F(34, 1130)=2.07$ $p<.001$), but these results are discussed in the main manuscript (section 3.4.2).

D.3. Hot vs cold executive functioning

A three-level meta-analysis including hot vs cold EF, age group (adult vs youth) and their interaction as moderators showed no significant interaction effect ($F(1, 1240)=1.17$ $p=.46$), or main effects of age group ($F(1, 1241)=1.16$ $p=.28$) or hot vs cold EF ($F(1, 1241)=.49$ $p=.49$).

D.4. Assessment quality

A three-level meta-analysis including assessment quality (high, medium, low), age group (adult vs. youth) and their interaction as moderators showed no significant interaction effect ($F(2, 1238)=.14$ $p=.87$, nor main effects of assessment quality ($F(2, 1240)=.83$ $p=.44$) and age group ($F(1,1240)=1.14$ $p=.29$).

References

- Aharoni, E., Vincent, G. M., Harenski, C. L., Calhoun, V. D., Sinnott-Armstrong, W., Gazzaniga, M. S., & Kiehl, K. A. (2013). Neuroprediction of future rearrest. *PNAS*, *110*(15), 6223–6228. <https://doi.org/10.1073/pnas.1219302110>
- *Antonini, T. N., Becker, S. P., Tamm, L., & Epstein, J. N. (2015). Hot and cool executive functions in children with attention-deficit/hyperactivity disorder and comorbid oppositional defiant disorder. *Journal of the International Neuropsychological Society*, *21*(8), 584–595. <https://doi.org/10.1017/s1355617115000752>
- Baggetta, P., & Alexander, P. A. (2016). Conceptualization and operationalization of executive function. *Mind, Brain, and Education*, *10*(1), 10–33. <https://doi.org/10.1111/mbe.12100>
- *Baliouis, M., Duggan, C., McCarthy, L., Huband, N., & Völlm, B. (2019). Executive function, attention, and memory deficits in antisocial personality disorder and psychopathy. *Psychiatry Research*, *278*, 151–161. <https://doi.org/10.1016/j.psychres.2019.05.046>
- *Barbosa, M. F. S., & Monteiro, L. M. C. (2008). Recurrent criminal behavior and executive dysfunction. *The Spanish Journal of Psychology*, *11*(1), 259–265. <https://doi.org/10.1017/S1138741600004297>
- Bardikoff, N., & Sabbagh, M. (2017). The differentiation of executive functioning across development: Insights from developmental cognitive neuroscience. In N. Budwig, E. Turiel, & P. D. Zelazo (Eds.), *New perspectives on human development* (pp. 47–66). Cambridge University Press. <https://doi.org/10.1017/CBO9781316282755.005>
- *Barkataki, I., Kumari, V., Das, M., Sumich, A., Taylor, P., & Sharma, T. (2008). Neural correlates of deficient response inhibition in mentally disordered violent individuals. *Behavioral Sciences & the Law*, *26*(1), 51–64. <https://doi.org/10.1002/bsl.787>
- *Barlatti, S., Nibbio, G., Stanga, V., Giovannoli, G., Calzavara-Pinton, I., Necchini, N., Vita, A., & 1-10. (2023). Cognitive and clinical characteristics of offenders and non-offenders diagnosed with schizophrenia spectrum disorders: results of the Recoviwel observational study. *European Archives of Psychiatry and Clinical Neuroscience*, *273*(6), 1–10. <https://doi.org/10.1007/s00406-022-01510-9>
- *Barnett, R., Maruff, P., & Vance, A. (2009). Neurocognitive function in attention-deficit-hyperactivity disorder with and without comorbid disruptive behaviour disorders. *The Australian and New Zealand Journal of Psychiatry*, *43*(8), 722–730. <https://doi.org/10.1080/00048670903001927>
- *Becerra-García, J. A. (2015). Neuropsychology of domestic violence: A comparative preliminary study of executive functioning. *Medicine, Science and the Law*, *55*(1), 35–39. <https://doi.org/10.1177/0025802414525148>
- *Becerra-García, J. A., & Egan, V. (2014). Neurocognitive functioning and subtypes of child molesters: Poorer working memory differentiates incestuous from non-incestuous offenders. *Psychiatry, Psychology and Law*, *21*(4), 585–590. <https://doi.org/10.1080/13218719.2013.873974>
- Bertsch, K., Florange, J., & Herpertz, S. C. (2020). Understanding Brain Mechanisms of Reactive Aggression. *Current Psychiatry Reports*, *22*(12), 81. <https://doi.org/10.1007/s11920-020-01208-6>
- *Blum, A. W., Leppink, E. W., & Grant, J. E. (2017). Neurocognitive dysfunction in problem gamblers with co-occurring antisocial personality disorder. *Comprehensive Psychiatry*, *76*, 153–159. <https://doi.org/10.1016/j.comppsy.2017.05.004>
- *Borrani, J., Frías, M., Ortiz, X., García, A., & Valdez, P. (2015). Analysis of cognitive inhibition and flexibility in juvenile delinquents. *The Journal of Forensic Psychiatry & Psychology*, *26*(1), 60–77. <https://doi.org/10.1080/14789949.2014.971852>
- *Bränden, A., Coldevin, M., Zeiner, P., Stubberud, J., & Melinder, A. (2023). Executive function in children with disruptive mood dysregulation disorder compared to attention-deficit/hyperactivity disorder and oppositional defiant disorder, and in children with different irritability levels. *European Child & Adolescent Psychiatry*, *21*, 1–11. <https://doi.org/10.1007/s00787-023-02143-6>
- Brassard, M. L., & Joyal, C. C. (2022). Predicting forensic inpatient violence with odor identification and neuropsychological measures of impulsivity: A preliminary study. *Journal of Psychiatric Research*, *147*, 154–158. <https://doi.org/10.1016/j.jpsychires.2022.01.021>
- Buitelaar, N. J., & Ferdinand, R. F. (2016). ADHD Undetected in Criminal Adults. *Journal of Attention Disorders*, *20*(3), 270–278. <https://doi.org/10.1177/1087054712466916>
- *Bulgari, V., Iozzino, L., Ferrari, C., Picchioni, M., Candini, V., De Francesco, A., & de Girolamo, G. (2017). Clinical and neuropsychological features of violence in schizophrenia: a prospective cohort study. *Schizophrenia Research*, *181*, 124–130. <https://doi.org/10.1016/j.schres.2016.10.016>
- *Cantrell, C. M. (2008). *Executive functioning in adolescents with disruptive behavior disorders* The University of Tulsa. Tulsa.
- *Carter Leno, V., Chandler, S., White, P., Pickles, A., Baird, G., Hobson, C., & Simonoff, E. (2018). Testing the specificity of executive functioning impairments in adolescents with ADHD, ODD/CD and ASD. *European Child & Adolescent Psychiatry*, *27*, 899–908. <https://doi.org/10.1007/s00787-017-1089-5>
- *Chamberlain, S. R., Derbyshire, K. L., Leppink, E. W., & Grant, J. E. (2016). Neurocognitive deficits associated with antisocial personality disorder in non-treatment-seeking young adults. *Journal of the American Academy of Psychiatry and the Law Online*, *44*(2), 218–225. <https://doi.org/10.1007/s00787-017-1089-5>
- Checa, P., & Rueda, M. R. (2011). Behavioral and brain measures of executive attention and school competence in late childhood. *Developmental Neuropsychology*, *36*(8), 1018–1032. <https://doi.org/10.1080/87565641.2011.591857>
- Cheng, J., O'Connell, M. E., & Wormith, J. S. (2019). Bridging Neuropsychology and Forensic Psychology: Executive Function Overlaps With the Central Eight Risk and Need Factors. *The International Journal of Offender Therapy and Comparative Criminology*, *63*(4), 558–573. <https://doi.org/10.1177/0306624X18803818>
- *Coenen, E., Pleysier, S., & Put, J. (2022). Executive functions, self-control and juvenile delinquency. *Psychology, Crime & Law*, *28*(10), 1004–1023. <https://doi.org/10.1007/s00787-017-1089-5>
- *Cohen, L. J., Nesci, C., Steinfeld, M., Haeri, S., & Galynker, I. (2010). Investigating the relationship between sexual and chemical addictions by comparing executive function in pedophiles, opiate addicts and healthy controls. *Journal of Psychiatric Practice*, *16*(6), 405. <https://doi.org/10.1097/01.pra.0000390759.04581.7c>
- *Combalbert, N., Pennequin, V., Ferrand, C., Vandevyvere, R., Armand, M., & Geffray, B. (2016). Mental disorders and cognitive impairment in ageing offenders. *The Journal of Forensic Psychiatry & Psychology*, *27*(6), 853–866. <https://doi.org/10.1080/14789949.2016.1244277>
- Cornet, L. J., van der Laan, P. H., Nijman, H. L., Tollenaar, N., & de Kogel, C. H. (2015). Neurobiological factors as predictors of prisoners' response to a cognitive skills training. *Journal of Criminal Justice*, *43*(2), 122–132. <https://doi.org/10.1016/j.jcrimjus.2015.02.003>

- *Crippa, A., Marzocchi, G. M., Piroddi, C., Besana, D., Giribone, S., Vio, C., & Sora, M. L. (2015). An integrated model of executive functioning is helpful for understanding ADHD and associated disorders. *Journal of Attention Disorders*, 19(6), 455–467. <https://doi.org/10.1177/1087054714542000>
- *Cubillo, A., Tkalec, A., Oldenhof, H., Unteraehrer, E., Raschle, N., Kohls, G., & Stadler, C. (2023). Linking heart rate variability to psychological health and brain structure in adolescents with and without conduct disorder. *Frontiers in Psychiatry*, 14(1101064). <https://doi.org/10.3389/fpsy.2023.1101064>
- *Curtis, A., Gooden, J. R., Cox, C. A., Harries, T., Peterson, V., Enticott, P. G., & Manning, V. (2021). Neurocognitive functioning among people accessing an addiction neuropsychology clinic with and without a history of offending behaviour. *Psychiatry, Psychology and Law*, 28, 854–866. <https://doi.org/10.1177/1087054714542000>
- *De Brito, S. A., Viding, E., Kumari, V., Blackwood, N., & Hodgins, S. (2013). Cool and hot executive function impairments in violent offenders with antisocial personality disorder with and without psychopathy. *PLoS One*, e65566. <https://doi.org/10.1371/journal.pone.0065566>
- *Delfin, C., Ruzich, E., Wallinius, M., Björnsdotter, M., & Andiné, P. (2020). Trait disinhibition and NoGo event-related potentials in violent mentally disordered offenders and healthy controls. *Frontiers in Psychiatry*, 11, Article 577491. <https://doi.org/10.3389/fpsy.2020.577491>
- *Déry, M., Toupin, J., & Durette, C. (2009). Habilité́s cognitives et trouble des conduites précoce chez les filles. *Revue Québécoise De Psychologie*.
- Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135–168. <https://doi.org/10.1146/annurev-psych-113011-143750>
- *Dolan, M. (2012). The neuropsychology of prefrontal function in antisocial personality disordered offenders with varying degrees of psychopathy. *Psychological Medicine*, 42(8), 1715–1725. <https://doi.org/10.1017/S0033291711002686>
- *Dolan, M., & Lennox, C. (2013). Cool and hot executive function in conduct-disordered adolescents with and without co-morbid attention deficit hyperactivity disorder: relationships with externalizing behaviours. *Psychological Medicine*, 43(11), 2427–2436. <https://doi.org/10.1017/S0033291712003078>
- *Easton, C. J., Sacco, K. A., Neavins, T. M., Wupperman, P., & George, T. P. (2008). Neurocognitive performance among alcohol dependent men with and without physical violence toward their partners: a preliminary report. *The American Journal of Drug and Alcohol Abuse*, 34(1), 29–37. <https://doi.org/10.1080/00952990701764326>
- *Enticott, P. G., Ogloff, J. R., Bradshaw, J. L., & Fitzgerald, P. B. (2008). Cognitive inhibitory control and self-reported impulsivity among violent offenders with schizophrenia. *Journal of Clinical and Experimental Neuropsychology*, 30(2), 157–162. <https://doi.org/10.1080/13803390701290055>
- Ersche, K. D., Turton, A. J., Chamberlain, S. R., Muller, U., Bullmore, E. T., & Robbins, T. W. (2012). Cognitive dysfunction and anxious-impulsive personality traits are endophenotypes for drug dependence. *American Journal of Psychiatry*, 169(9), 926–936. <https://doi.org/10.1176/appi.ajp.2012.11091421>
- *Euler, F., Sterzer, P., & Stadler, C. (2014). Cognitive control under distressing emotional stimulation in adolescents with conduct disorder. *Aggressive Behavior*, 40(2), 109–119. <https://doi.org/10.1002/ab.21508>
- *Ezpeleta, L., & Granero, R. (2015). Executive functions in preschoolers with ADHD, ODD, and comorbid ADHD-ODD: Evidence from ecological and performance-based measures. *Journal of Neuropsychology*, 9(2), 258–270. <https://doi.org/10.1111/jnp.12049>
- *Fairchild, G., van Goozen, S. H., Stollery, S. J., Aitken, M. R., Savage, J., Moore, S. C., & Goodyer, I. M. (2009). Decision making and executive function in male adolescents with early-onset or adolescence-onset conduct disorder and control subjects. *Biological Psychiatry*, 66(2), 162–168. <https://doi.org/10.1016/j.biopsych.2009.02.024>
- Fazel, S., Hayes, A. J., Bartellas, K., Clerici, M., & Trestman, R. (2016). Mental health of prisoners: prevalence, adverse outcomes, and interventions. *Lancet Psychiatry*, 3(9), 871–881. [https://doi.org/10.1016/S2215-0366\(16\)30142-0](https://doi.org/10.1016/S2215-0366(16)30142-0)
- *Feilhauer, J., Cima, M., Korebrits, A., & Kunert, H. J. (2012). Differential associations between psychopathy dimensions, types of aggression, and response inhibition. *Aggressive Behavior*, 38(1), 77–88. <https://doi.org/10.1002/ab.20415>
- Fernandes, B., Wright, M., & Essau, C. A. (2023). The Role of Emotion Regulation and Executive Functioning in the Intervention Outcome of Children with Emotional and Behavioural Problems. *Children (Basel)*, 10(1), 139. <https://doi.org/10.3390/children10010139>
- Fernandez-Castilla, B., Jamshidi, L., Declercq, L., Beretvas, S. N., Onghena, P., & Van den Noortgate, W. (2020). The application of meta-analytic (multi-level) models with multiple random effects: A systematic review. *Behavior Research Methods*, 52(5), 2031–2052. <https://doi.org/10.3758/s13428-020-01373-9>
- Figueiredo, P., Ramiao, E., Barros, R., & Barbosa, F. (2023). Executive (dys)functions and ODD and CD: A systematic review and meta-analysis. *Neuropsychology*, 37(2), 133–156. <https://doi.org/10.1037/neu0000881>
- Fine, A., Steinberg, L., Frick, P. J., & Cauffman, E. (2016). Self-control assessments and implications for predicting adolescent offending. *Journal of Youth and Adolescence*, 45, 701–712. <https://doi.org/10.1007/s10964-016-0425-2>
- *Fonseca-Parra, L. F., & Rey-Anacona, C. A. (2013). Características neuropsicológicas de niños con trastorno negativista desafiante. *Revista de Psicopatología y Psicología Clínica*, 18(3), 233–244. <https://doi.org/10.5944/rppc.vol.18.num.3.2013.12923>
- *Franke, L., Seipel, S., Vasic, N., Streb, J., Nigel, S., Otte, S., & Dudeck, M. (2019). Neuropsychological profile of pedophilic child sexual offenders compared with an IQ-matched non-offender sample—Results of a pilot study. *International Journal of Law and Psychiatry*, 64, 137–141. <https://doi.org/10.1016/j.ijlp.2019.03.002>
- *Frias-Armenta, M., Valdez-Ramírez, P., Nava-Cruz, G., Figueredo, A. J., & Corral-Verdugo, V. I. (2011). Executive functions, attention and juvenile delinquency. In M. Frias-armenta, & V. Corral-verdugo (Eds.), *Bio-psycho-social Perspectives on Interpersonal Violence*. Nova Science Publishers.
- Friedman, N. P., & Miyake, A. (2017). Unity and diversity of executive functions: Individual differences as a window on cognitive structure. *Cortex*, 86, 186–204. <https://doi.org/10.1016/j.cortex.2016.04.023>
- Friedman, N. P., Miyake, A., Young, S. E., DeFries, J. C., Corley, R. P., & Hewitt, J. K. (2008). Individual differences in executive functions are almost entirely genetic in origin. *Journal of Experimental Psychology: General*, 137, 201. <https://doi.org/10.1037/0096-3445.137.2.201>
- *Ginsberg, Y., Hirvikoski, T., & Lindfors, N. (2010). Attention deficit hyperactivity disorder (ADHD) among longer-term prison inmates is a prevalent, persistent and disabling disorder. *BMC Psychiatry*, 10(1), 1–13. <https://doi.org/10.1186/1471-244x-10-112>
- *Glenn, A. L., Rimmel, R. J., Ong, M. Y., Lim, N. S., Ang, R. P., Threadgill, A. H., & Ooi, Y. P. (2017). Neurocognitive characteristics of youth with noncomorbid and comorbid forms of conduct disorder and attention deficit hyperactivity disorder. *Comprehensive Psychiatry*, 77, 60–70. <https://doi.org/10.1016/j.comppsy.2017.06.005>
- *Gobbi, E., Cotelli, M., Manenti, R., Ferrari, C., Macis, A., Bianconi, G., & de Girolamo, G. (2020). Neuropsychological features in patients with severe mental disorders and risk of violence: a prospective multicenter study in Italy. *Psychiatry Research*, 289, Article 113027. <https://doi.org/10.1016/j.psychres.2020.113027>
- *Gonzalez-Gadea, M. L., Herrera, E., Parra, M., Gomez Mendez, P., Baez, S., Manes, F., & Ibanez, A. (2014). Emotion recognition and cognitive empathy deficits in adolescent offenders revealed by context-sensitive tasks. *Frontiers in Human Neuroscience*, 8, 850. <https://doi.org/10.3389/fnhum.2014.00850>
- de Graaf, I. E., Bolhuis, K., Cecil, C. A. M., White, T. H., & van Dongen, J. D. M. (2023). Child Neuropsychological Functioning and Interpersonal Callousness as Predictors of Externalising Behaviour in Early Adolescence: A Prospective Population-based Study. *Research on Child and Adolescent Psychopathology*, 51(10), 1465–1480. <https://doi.org/10.1007/s10802-023-01091-8>
- *Grant, J. E., Chamberlain, S. R., Schreiber, L. R., & Odraug, B. L. (2012). Neurocognitive deficits associated with shoplifting in young adults. *Comprehensive Psychiatry*, 53(8), 1049–1055. <https://doi.org/10.1016/j.comppsy.2012.04.012>
- Graziano, P. A., Landis, T., Maharaj, A., Ros-Demarize, R., Hart, K. C., & Garcia, A. (2022). Differentiating preschool children with conduct problems and callous-unemotional behaviors through emotion regulation and executive functioning. *Journal of Clinical Child & Adolescent Psychology*, 51(2), 170–182. <https://doi.org/10.1080/15374416.2019.1666399>
- *Guan, M., Liao, Y., Ren, H., Wang, X., Yang, Q., Liu, X., & Wang, W. (2015). Impaired response inhibition in juvenile delinquents with antisocial personality characteristics: A preliminary ERP study in a Go/Nogo task. *Neuroscience Letters*, 603, 1–5. <https://doi.org/10.1016/j.neulet.2015.06.062>
- *Gunn, R. L., Gerst, K. R., Lake, A. J., & Finn, P. R. (2018). The effects of working memory load and attention refocusing on delay discounting rates in alcohol use disorder with comorbid antisocial personality disorder. *Alcohol*, 66, 9–14. <https://doi.org/10.1016/j.alcohol.2017.07.009>
- Haarsma, G., Davenport, S., White, D. C., Ormachea, P. A., Sheena, E., & Eagleman, D. M. (2020). Assessing risk among correctional community probation populations: Predicting reoffense with mobile neurocognitive assessment software. *Frontiers in Psychology*, 10, 296. <https://doi.org/10.3389/fpsyg.2019.02926>
- *Habermeier, B., Esposito, F., Händel, N., Lemoine, P., Kuhl, H. C., Klarhöfer, M., & Graf, M. (2013). Response inhibition in pedophilia: An fMRI pilot study. *Neuropsychobiology*, 68(4), 228–237. <https://doi.org/10.1159/000355295>
- Hall, P. A., Elias, L. J., & Crossley, M. (2006). Neurocognitive influences on health behavior in a community sample. *Health Psychology*, 25(6), 778–782. <https://doi.org/10.1037/0278-6133.25.6.778>
- Hanlon, R. E., Brook, M., Stratton, J., Jensen, M., & Rubin, L. H. (2013). Neuropsychological and intellectual differences between types of murderers: Affective/impulsive versus predatory/instrumental (premeditated) homicide. *Criminal Justice and Behavior*, 40(8), 933–948. <https://doi.org/10.1177/0093854813479779>
- *Herrero, O., Escorial, S., & Colom, R. (2010). Basic executive processes in incarcerated offenders. *Personality and Individual Differences*, 48(2), 133–137. <https://doi.org/10.1016/j.paid.2009.09.009>
- *Herrero, Ó., Escorial, S., & Colom, R. (2018). Rapists and child abusers share low levels in executive updating, but do not in fluid reasoning. *The European Journal of Psychology Applied to Legal Context*, 11(1), 1–7. <https://doi.org/10.5093/ejpalc2018a10>
- Hirstein, W., Sifferd, K. L., & Fagan, T. K. (2018). *Responsible brains: Neuroscience, law, and human culpability*. MIT Press. <https://doi.org/10.7551/mitpress/11262.001.0001>
- *Hobson, C. W., Scott, S., & Rubia, K. (2011). Investigation of cool and hot executive function in ODD/CD independently of ADHD. *Journal of Child Psychology and Psychiatry*, 52(10), 1035–1043. <https://doi.org/10.1111/j.1469-7610.2011.02454.x>
- *Hoppenbrouwers, S. S., De Jesus, D. R., Stirpe, T., Fitzgerald, P. B., Voineskos, A. N., Schutter, D. J., & Daskalakis, Z. J. (2013). Inhibitory deficits in the dorsolateral prefrontal cortex in psychopathic offenders. *Cortex*, 49(5), 1377–1385. <https://doi.org/10.1016/j.cortex.2012.06.003>
- Huizinga, M., Dolan, C. V., & van der Molen, M. W. (2006). Age-related change in executive function: Developmental trends and a latent variable analysis. *Neuropsychologia*, 44(11), 2017–2036. <https://doi.org/10.1016/j.neuropsychologia.2006.01.010>
- *Hummer, T. A., Kronenberger, W. G., Wang, Y., Dunn, D. W., Mosier, K. M., Kalnin, A. J., & Mathews, V. P. (2011). Executive functioning characteristics associated with ADHD comorbidity in adolescents with disruptive behavior

- disorders. *Journal of Abnormal Child Psychology*, 39, 11–19. <https://doi.org/10.1007/s10802-010-9449-3>
- *Hummer, T. A., Wang, Y., Kronenberger, W. G., Dunn, D. W., & Mathews, V. P. (2015). The relationship of brain structure to age and executive functioning in adolescent disruptive behavior disorder. *Psychiatry Research: Neuroimaging*, 231(3), 210–217. <https://doi.org/10.1016/j.psychres.2014.11.009>
- *Hwang, S., Nolan, Z. T., White, S. F., Williams, W. C., Sinclair, S., & Blair, R. J. R. (2016). Dual neurocircuitry dysfunctions in disruptive behavior disorders: emotional responding and response inhibition. *Psychological Medicine*, 46(7), 1485–1496. <https://doi.org/10.1017/s0033291716000118>
- *Iria, C., Barbosa, F., & Paixão, R. (2012). The identification of negative emotions through a go/no-go task: Comparative research in criminal and non-criminal psychopaths. *European Psychologist*, 17(4), 291–299. <https://doi.org/10.1027/1016-9040/a000101>
- *Iselin, A. M. R., & DeCoster, J. (2009). Reactive and proactive control in incarcerated and community adolescents and young adults. *Cognitive Development*, 24(2), 192–206. doi:10.1016/j.cogdev.2008.07.001.
- Iselin, A. M. R., & DeCoster, J. (2012). Unique relations of age and delinquency with cognitive control. *Journal of Adolescence*, 35(2), 367–379. <https://doi.org/10.1016/j.adolescence.2011.08.004>
- Jansen, J. (2020). Traumatic brain injury and its relationship to previous convictions, aggression, and psychological functioning in Dutch detainees. *Journal of Forensic Psychology Research and Practice*, 20(5), 395–412. <https://doi.org/10.1080/24732850.2020.1755923>
- *Jiang, W., Li, Y., Du, Y., & Fan, J. (2016). Emotional regulation and executive function deficits in unmedicated Chinese children with oppositional defiant disorder. *Psychiatry Investigation*, 13(3), 277. <https://doi.org/10.4306/pi.2016.13.3.277>
- *Joyal, C. C., Tardif, M., & Spearson-Goulet, J. A. (2020). Executive functions and social cognition in juveniles who have sexually offended. *Sexual Abuse*, 32(2), 179–202. <https://doi.org/10.1177/1079063218807487>
- *Kallitsoglou, A. (2018). Executive function variation in children with conduct problems: Influences of coexisting reading difficulties. *Journal of Learning Disabilities*, 51(2), 168–180. <https://doi.org/10.1177/0022219417691834>
- Karr, J. E., Areshenkoff, C. N., Rast, P., Hofer, S. M., Iverson, G. L., & Garcia-Barrera, M. A. (2018). The unity and diversity of executive functions: A systematic review and re-analysis of latent variable studies. *Psychological Bulletin*, 144(11), 1147–1185. <https://doi.org/10.1037/bul0000160>
- *Kashiwagi, H., Kuroki, N., Ikezawa, S., Matsushita, M., Ishikawa, M., Nakagome, K., & Ikeda, M. (2015). Neurocognitive features in male patients with schizophrenia exhibiting serious violence: a case control study. *Annals of General Psychiatry*, 14(1), 1–6. <https://doi.org/10.1186/s12991-015-0086-7>
- *Kim, H., Jung, E., Lee, T., Kim, S., & Kim, H. W. (2023). Impact of Comorbid Oppositional Defiant Disorder on the Clinical and Neuropsychological Characteristics of Korean Children With Attention-Deficit/Hyperactivity Disorder. *Psychiatry Investigation*, 20(10), 962. <https://doi.org/10.30773/pi.2023.0091>
- *Kleine Deters, R., Naaijen, J., Rosa, M., Aggensteiner, P. M., Banaschewski, T., Saam, M. C., & Dietrich, A. (2020). Executive functioning and emotion recognition in youth with oppositional defiant disorder and/or conduct disorder. *The World Journal of Biological Psychiatry*, 21(7), 539–551. <https://doi.org/10.1080/15622975.2020.1747114>
- *Krakowski, M. I., Foxe, J., de Sanctis, P., Nolan, K., Hoptman, M. J., Shope, C., & Czobor, P. (2015). Aberrant response inhibition and task switching in psychopathic individuals. *Psychiatry Research*, 229(3), 1017–1023. <https://doi.org/10.1016/j.psychres.2015.06.018>
- *Krischer, M., Sevecke, K., Wunram, H. L., Lehmkuhl, G., & Pukrop, R. (2008). The impact of emotional stimuli on working memory in female delinquent juveniles. *Zeitschrift für Kinder- und Jugendpsychiatrie und Psychotherapie*, 36(6), 401–410. <https://doi.org/10.1024/1422-4917.36.6.401>
- Lawson, G. M., Hook, C. J., & Farah, M. J. (2018). A meta-analysis of the relationship between socioeconomic status and executive function performance among children. *Developmental Science*, 21(2), Article e12529. <https://doi.org/10.1111/desc.12529>
- *Lin, Y. J., & Gau, S. S. F. (2017). Differential neuropsychological functioning between adolescents with attention-deficit/hyperactivity disorder with and without conduct disorder. *Journal of the Formosan Medical Association*, 116(12), 946–955. <https://doi.org/10.1016/j.fjma.2017.02.009>
- *Lishak, V., Scott, K. L., Dyson, A., & Milovanov, A. (2021). General criminality as a marker of heterogeneity in domestically violent men: Differences in trauma history, psychopathology, neurocognitive functioning, and psychophysiology. *Journal of Interpersonal Violence*, 36(17–18), 9623–9648. <https://doi.org/10.1177/0886260519858065>
- *Liu, N., Jia, G., Qiu, S., Li, H., Liu, Y., Wang, Y., Niu, H., Liu, L., & Qian, Q. (2023). Different executive function impairments in medication-naïve children with attention-deficit/hyperactivity disorder comorbid with oppositional defiant disorder and conduct disorder. *Asian Journal of Psychiatry*, 81, Article 103446. <https://doi.org/10.1016/j.ajp.2022.103446>
- *Luman, M., Van Noesel, S. J., Papanikolaou, A., Van Oostenbruggen-Scheffer, J., Veugelers, D., Sergeant, J. A., & Oosterlaan, J. (2009). Inhibition, reinforcement sensitivity and temporal information processing in ADHD and ADHD+ ODD: evidence of a separate entity? *Journal of Abnormal Child Psychology*, 37(8), 1123–1135. <https://doi.org/10.1007/s10802-009-9334-0>
- *Majorek, K., Wolfkühler, W., Küper, C., Saimel, N., Juckel, G., & Brüne, M. (2009). “Theory of mind” and executive functioning in forensic patients with schizophrenia. *Journal of Forensic Sciences*, 54(2), 469–473. <https://doi.org/10.1111/j.1556-4029.2008.00966.x>
- *Marceau, R., Meghani, R., & Reddon, J. R. (2008). Neuropsychological Assessment of Adult Offenders. *Journal of Offender Rehabilitation*, 47, 41–73. <https://doi.org/10.1080/1059670801940409>
- *Massau, C., Tenbergen, G., Kärgel, C., Weiß, S., Gerwinn, H., Pohl, A., & Schiffer, B. (2017). Executive functioning in pedophilia and child sexual offending. *Journal of the International Neuropsychological Society*, 23(6), 460. <https://doi.org/10.1017/s1355617717000315>
- *Medrano Nava, E., Flores-Lázaro, J. C., Nicolini Sánchez, H., & Juárez García, F. (2022). Effects of comorbidity on executive functions among children with ADHD, finding trends. *Applied Neuropsychology: Child*, 17, 13. <https://doi.org/10.1080/21622965.2022.2135440>
- *Meier, N. M., Perrig, W., & Koenig, T. (2012). Neurophysiological correlates of delinquent behaviour in adult subjects with ADHD. *International Journal of Psychophysiology*, 84(1), 1–16. <https://doi.org/10.1016/j.ijpsycho.2011.12.011>
- *Mellentin, A. I., Skot, L., Teasdale, T. W., & Habekost, T. (2013). Conscious knowledge influences decision-making differently in substance abusers with and without comorbid antisocial personality disorder. *Scandinavian Journal of Psychology*, 54(4), 292–299. <https://doi.org/10.1111/sjop.12054>
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. *Cognitive Psychology*, 41(1), 49–100. <https://doi.org/10.1006/cogp.1999.0734>
- *Moeller, S. J., Froböse, M. I., Konova, A. B., Misyrlis, M., Parvaz, M. A., Goldstein, R. Z., & Alia-Klein, N. (2014). Common and distinct neural correlates of inhibitory dysregulation: stroop fMRI study of cocaine addiction and intermittent explosive disorder. *Journal of Psychiatric Research*, 58, 55–62. <https://doi.org/10.1016/j.jpsychires.2014.07.016>
- *Molleman, P. W., Driessen, J. M. A., Schilder, C., Bulten, B. H., & Brazil, I. A. (2021). Behavioral inhibition and activation system factors in offenders and non-offenders. *International Journal of Forensic Mental Health*, 1-13. <https://doi.org/10.1080/14999013.2021.1947422>
- *Molleman, P. W., Molleman, L., Schilder, C., Bulten, B. H., & Brazil, I. A. (2022). Can measures of cognitive flexibility and inhibition distinguish forensic psychiatric inpatients from prisoners? *The Journal of Forensic Psychiatry & Psychology*, 33(3), 371–388. doi:11245.1/927b1429-57a8-4e0f-8481-bbed1816fc4d.
- Morgan, A. B., & Lilienfeld, S. O. (2000). A meta-analytic review of the relation between antisocial behavior and neuropsychological measures of executive function. *Clinical Psychology Review*, 20(1), 113–136. [https://doi.org/10.1016/s0272-7358\(98\)00096-8](https://doi.org/10.1016/s0272-7358(98)00096-8)
- *Munkvold, L. H., Manger, T., & Lundervold, A. J. (2014). Conners’ continuous performance test (CCPT-II) in children with ADHD, ODD, or a combined ADHD/ODD diagnosis. *Child Neuropsychology*, 20(1), 106–126. <https://doi.org/10.1080/09297049.2012.753997>
- *Muscatello, M. R. A., Scimeca, G., Pandolfo, G., Micò, U., Romeo, V. M., Mallamaci, D., & Bruno, A. (2014). Executive functions and basic symptoms in adolescent antisocial behavior: a cross-sectional study on an Italian sample of late-onset offenders. *Comprehensive Psychiatry*, 55(3), 631–638. <https://doi.org/10.1016/j.comppsy.2013.11.015>
- **Närhi, V., Lehto-Salo, P., Ahonen, T., & Marttunen, M. (2010). Neuropsychological subgroups of adolescents with conduct disorder. *Scandinavian Journal of Psychology*, 51(3), 278–284. <https://doi.org/10.1111/j.1467-9450.2009.00767.x>
- *Neves, D., & Pinho, M. S. (2018). Self-regulation and the specificity of autobiographical memory in offenders. *International Journal of Law and Psychiatry*, 57, 91–99. <https://doi.org/10.1016/j.ijlp.2018.01.005>
- Nikulina, V., & Widom, C. S. (2019). Higher Levels of Intelligence and Executive Functioning Protect Maltreated Children Against Adult Arrests: A Prospective Study. *Child Maltreatment*, 24(1), 3–16. <https://doi.org/10.1177/1077559518808218>
- *Nishinaka, H., Nakane, J., Nagata, T., Imai, A., Kuroki, N., Sakikawa, N., & Hashimoto, K. (2016). Neuropsychological impairment and its association with violence risk in Japanese forensic psychiatric patients: A case-control study. *PLoS One*, 11(1), Article e0148354. <https://doi.org/10.1371/journal.pone.0148354>
- *Noordermeer, S. D., Luman, M., Buitelaar, J. K., Hartman, C. A., Hoekstra, P. J., Franke, B., & Oosterlaan, J. (2020). Neurocognitive deficits in attention-deficit/hyperactivity disorder with and without comorbid oppositional defiant disorder. *Journal of Attention Disorders*, 24(9), 1317–1329. <https://doi.org/10.1177/1087054715606216>
- *Nordvall, O., Jonsson, B., & Neely, A. S. (2017). Self-reported and performance-based measures of executive functions in internet youth. *Psychology, Crime & Law*, 23(3), 240–253. <https://doi.org/10.1080/1068316X.2016.1239725>
- *Nordvall, O., Neely, A. S., & Jonsson, B. (2017). Self-reported impulsivity and its relation to executive functions in internet youth. *Psychiatry, Psychology and Law*, 24(6), 910–922. <https://doi.org/10.1080/13218719.2017.1327312>
- Norman, E. M., Polaschek, D. L. L., & Starkey, N. J. (2023). Executive function in individuals who are compliant and non-compliant with the conditions of a community-based sentence. *Psychiatry, Psychology and Law*, 30(2), 161–176. <https://doi.org/10.1080/13218719.2021.2003268>
- Ogilvie, J. M., Stewart, A. L., Chan, R. C. K., & Shum, D. H. K. (2011). Neuropsychological Measures of Executive Function and Antisocial Behavior: A Meta-Analysis. *Criminology*, 49(4), 1063–1107. <https://doi.org/10.1111/j.1745-9125.2011.00252.x>
- Op den Kelder, R., Van den Akker, A. L., Geurts, H. M., Lindauer, R. J. L., & Overbeek, G. (2018). Executive functions in trauma-exposed youth: a meta-analysis. *European Journal of Psychotraumatology*, 9(1), 1450595. <https://doi.org/10.1080/20008198.2018.1450595>
- Ormachea, P. A., Lovins, B. K., Eagleman, D. M., Davenport, S., Jarman, A., & Haarsma, G. (2017). The role of tablet-based psychological tasks in risk assessment.

- Criminal Justice and Behavior*, 44(8), 993–1008. <https://doi.org/10.1177/0093854817714018>
- *Ostrosky, F., Borja, K. C., Rebolgar, C. R., & Galván, K. X. D. (2012). Neuropsychological profiles of members of organized crime and drug-traffic organizations. *Research and Reports in Forensic Medical Science*, 2, 19–30. <https://doi.org/10.2147/RRFMS.S32352>
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., ... Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *International Journal of Surgery*, 88, Article 105906. <https://doi.org/10.1136/bmj.n71>
- *Pajer, K., Chung, J., Leininger, L., Wang, W., Gardner, W., & Yeates, K. (2008). Neuropsychological function in adolescent girls with conduct disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 47(4), 416–425. <https://doi.org/10.1097/chi.0b013e3181640828>
- *Palix, J., Gillespie, S. M., Abbiati, M., & Abu-Akel, A. (2022). Dynamics of parasymphatic activity in violent incarcerated offenders before, during, and in recovery from an emotional inhibition task. *Scientific Reports*, 12(1), 7126. <https://doi.org/10.1038/s41598-022-10872-y>
- *Pasion, R., Cruz, A. R., & Barbosa, F. (2018). Dissociable effects of psychopathic traits on executive functioning: insights from the triarchic model. *Frontiers in Psychology*, 9, 1713. <https://doi.org/10.3389/fpsyg.2018.01713>
- *Patz, B., & Bayraktar, S. (2023). Evaluation of neuropsychological characteristics and attention bias in juvenile offenders, juvenile victims, and juveniles who have not participated in the criminal justice system. *Frontiers in Psychology*, 14(1229044). <https://doi.org/10.3389/fpsyg.2023.1229044>
- *Pera-Guardiola, V., Batalla, I., Bosque, J., Kosson, D., Pifarré, J., Hernández-Ribas, R., & Cardoner, N. (2016). Modulatory effects of psychopathy on Wisconsin Card Sorting Test performance in male offenders with Antisocial Personality Disorder. *Psychiatry Research*, 235, 43–48. <https://doi.org/10.1016/j.psychres.2015.12.003>
- *Perino, M. T., Guassi Moreira, J. F., McCormick, E. M., & Telzer, E. H. (2019). Apples to apples? Neural correlates of emotion regulation differences between high- and low-risk adolescents. *Social Cognitive and Affective Neuroscience*, 14(8), 827–836. <https://doi.org/10.1093/scan/nsz063>
- *Poon, K. (2020). Evaluating dual-process theory of decision-making in Chinese delinquent adolescents. *Australian Psychologist*, 55(3), 257–268. <https://doi.org/10.1111/ap.12417>
- *Prehn, K., Schulze, L., Rossmann, S., Berger, C., Vohs, K., Fleischer, M., & Herpertz, S. C. (2013). Effects of emotional stimuli on working memory processes in male criminal offenders with borderline and antisocial personality disorder. *The World Journal of Biological Psychiatry*, 14(1), 71–78. <https://doi.org/10.3109/15622975.2011.584906>
- *Price, S. A., Beech, A. R., Mitchell, I., & Humphreys, G. W. (2013). Measuring deviant sexual interest using the emotional Stroop task. *Criminal Justice and Behavior*, 40(9), 970–987. <https://doi.org/10.1177/009385481347626>
- *Price, S. A., Beech, A. R., Mitchell, I., & Humphreys, G. W. (2014). Measuring deviant sexual interest in adolescents using the emotional Stroop task. *Sexual Abuse*, 26(5), 450–471. <https://doi.org/10.1177/1079063213495897>
- *Qian, Y., Shuai, L., Cao, Q., Chan, R. C., & Wang, Y. (2010). Do executive function deficits differentiate between children with attention deficit hyperactivity disorder (ADHD) and ADHD comorbid with oppositional defiant disorder? A cross-cultural study using performance-based tests and the behavior rating inventory of executive function. *The Clinical Neuropsychologist*, 24(5), 793–810. <https://doi.org/10.1080/13854041003749342>
- *Raszkievicz, S. R. (2010). *The clinical utility of the Conners' Continuous Performance Test in the evaluation of youth with conduct disorder*. Milwaukee, Wisconsin: Marquette University.
- *Rhodes, S. M., Park, J., Seth, S., & Coghill, D. R. (2012). A comprehensive investigation of memory impairment in attention deficit hyperactivity disorder and oppositional defiant disorder. *Journal of Child Psychology and Psychiatry*, 53, 128–137. <https://doi.org/10.1111/j.1469-7610.2011.02436.x>
- *Romero-Martínez, Á., Lila, M., & Moya-Albiol, L. (2021). Alexithymic traits are closely related to impulsivity and cognitive and empathic dysfunctions in intimate partner violence perpetrators: New targets for intervention. *Applied Neuropsychology: Adult*, 28(1), 71–79. <https://doi.org/10.1080/23279095.2019.1594233>
- *Romero-Martínez, Á., Lila, M., Vitoria-Estruch, S., & Moya-Albiol, L. (2021). Can attention and working memory impairments of intimate partner perpetrators explain their risky decision making? *Journal of Interpersonal Violence*, 36(11-12). <https://doi.org/10.1177/0886260518814263>. NP6492-NP6507.
- Ros, R., Bjarnason, E., & Runeson, P. (2017). A machine learning approach for semi-automated search and selection in literature studies. In *Proceedings of the 21st International Conference on Evaluation and Assessment in Software Engineering*.
- *Rosburg, T., Deuring, G., Boillat, C., Lemoine, P., Falkenstein, M., Graf, M., & Mager, R. (2018). Inhibition and attentional control in pedophilic child sexual offenders—An event-related potential study. *Clinical Neurophysiology*, 129(9), 1990–1998. <https://doi.org/10.1016/j.clinph.2018.06.029>
- *Rubia, K., Halari, R., Cubillo, A., Mohammad, A. M., Scott, S., & Brammer, M. (2010). Disorder-specific inferior prefrontal hypofunction in boys with pure attention-deficit/hyperactivity disorder compared to boys with pure conduct disorder during cognitive flexibility. *Human Brain Mapping*, 31(12), 1823–1833. <https://doi.org/10.1002/hbm.20975>
- *Rubia, K., Halari, R., Smith, A. B., Mohammad, M., Scott, S., & Brammer, M. J. (2009). Shared and disorder-specific prefrontal abnormalities in boys with pure attention-deficit/hyperactivity disorder compared to boys with pure CD during interference inhibition and attention allocation. *Journal of Child Psychology and Psychiatry*, 50(6), 669–678. <https://doi.org/10.1111/j.1469-7610.2008.02022.x>
- *Rubia, K., Halari, R., Smith, A. B., Mohammed, M., Scott, S., Giampietro, V., & Brammer, M. J. (2008). Dissociated functional brain abnormalities of inhibition in boys with pure conduct disorder and in boys with pure attention deficit hyperactivity disorder. *American Journal of Psychiatry*, 165(7), 889–897. <https://doi.org/10.1176/appi.ajp.2008.07071084>
- de Ruigh, E. L., Kleeven, A. T. H., Jansen, L. M., de Vries Robbé, M., Vermeiren, R. R., Mulder, E. A., & Popma, A. (2021). Predicting youth reoffending after incarceration: added value of protective factors and heart rate variability. *The Journal of Forensic Psychiatry & Psychology*, 32(4), 449–480. <https://doi.org/10.1080/14789949.2020.1860250>
- Rutter, M. (2003). Commentary: causal processes leading to antisocial behavior. *Developmental Psychology*, 39(2), 372–378. <https://doi.org/10.1037/0012-1649.39.2.372>
- *Saarinen, S., Fontell, T., Vuontela, V., Carlson, S., & Aronen, E. T. (2015). Visuospatial working memory in 7- to 12-year-old children with disruptive behavior disorders. *Child Psychiatry & Human Development*, 46(1), 34–43. <https://doi.org/10.1007/s10578-014-0449-3>
- Salehinejad, M. A., Ghanavati, E., Rashid, M. H. A., & Nitsche, M. A. (2021). Hot and cold executive functions in the brain: A prefrontal-cingular network. *Brain and Neuroscience Advances*, 5, 1–19. <https://doi.org/10.1177/23982128211007769>
- Saylik, R., Williams, A. L., Murphy, R. A., & Szameitat, A. J. (2022). Characterising the unity and diversity of executive functions in a within-subject fMRI study. *Scientific Reports*, 12(1), 8182. <https://doi.org/10.1038/s41598-022-11433-z>
- *Schiffer, B., Pawliczek, C., Mu, B., Forsting, M., Gizewski, E., Leygraf, N., & Hodgins, S. (2014). Neural mechanisms underlying cognitive control of men with lifelong antisocial behavior. *Psychiatry Research: Neuroimaging*, 222(1-2), 43–51. <https://doi.org/10.1016/j.pscychresns.2014.01.008>
- *Schiffer, B., & Vonlaufen, C. (2011). Executive dysfunctions in pedophilic and nonpedophilic child molesters. *The Journal of Sexual Medicine*, 8(7), 1975–1984. <https://doi.org/10.1111/j.1743-6109.2010.02140.x>
- *Schoemaker, K., Bunte, T., Wiebe, S. A., Espy, K. A., Deković, M., & Matthys, W. (2012). Executive function deficits in preschool children with ADHD and DBD. *Journal of Child Psychology and Psychiatry*, 53(2), 111–119. <https://doi.org/10.1111/j.1469-7610.2011.02468.x>
- Schoemaker, K., Mulder, H., Dekovic, M., & Matthys, W. (2013). Executive functions in preschool children with externalizing behavior problems: a meta-analysis. *Journal of Abnormal Child Psychology*, 41(3), 457–471. <https://doi.org/10.1007/s10802-012-9684-x>
- *Schoorl, J., van Rijn, S., de Wied, M., Van Goozen, S., & Swaab, H. (2016). Emotion regulation difficulties in boys with oppositional defiant disorder/conduct disorder and the relation with comorbid autism traits and attention deficit traits. *PLoS One*, 11(7), Article e0159323. <https://doi.org/10.1371/journal.pone.0159323>
- *Schoorl, J., van Rijn, S., de Wied, M., Van Goozen, S., & Swaab, H. (2018). Boys with oppositional defiant disorder/conduct disorder show impaired adaptation during stress: An executive functioning study. *Child Psychiatry & Human Development*, 49(2), 298–307. <https://doi.org/10.1007/s10578-017-0749-5>
- *Schwenck, C., Ciaramidaro, A., Selivanova, M., Tournay, J., Freitag, C. M., & Siniatchkin, M. (2017). Neural correlates of affective empathy and reinforcement learning in boys with conduct problems: fMRI evidence from a gambling task. *Behavioural Brain Research*, 320(75–84). <https://doi.org/10.1016/j.bbr.2016.11.037>
- Senn, T. E., Espy, K. A., & Kaufmann, P. M. (2004). Using path analysis to understand executive function organization in preschool children. *Developmental Neuropsychology*, 26(1), 445–464. https://doi.org/10.1207/s15326942dn2601_5
- *Seruca, T., & Silva, C. F. (2016). Executive functioning in criminal behavior: Differentiating between types of crime and exploring the relation between shifting, inhibition, and anger. *International Journal of Forensic Mental Health*, 15(3), 235–246. <https://doi.org/10.1080/14999013.2016.1158755>
- Shi, L., & Lin, L. (2019). The trim-and-fill method for publication bias: practical guidelines and recommendations based on a large database of meta-analyses. *Medicine (Baltimore)*, 98(23), Article e15987. <https://doi.org/10.1097/MD.00000000000015987>
- *Shuai, L., Chan, R. C., & Wang, Y. (2011). Executive function profile of Chinese boys with attention-deficit hyperactivity disorder: different subtypes and comorbidity. *Archives of Clinical Neuropsychology*, 26(2), 120–132. <https://doi.org/10.1093/acclin/acq101>
- Sijtsma, J. J., & Lindenberg, S. M. (2018). Peer influence in the development of adolescent antisocial behavior: Advances from dynamic social network studies. *Developmental Review*, 50(140–154). <https://doi.org/10.1016/j.dr.2018.08.002>
- Smolker, H. R., Friedman, N. P., Hewitt, J. K., & Banich, M. T. (2018). Neuroanatomical Correlates of the Unity and Diversity Model of Executive Function in Young Adults. *Frontiers in Human Neuroscience*, 12, 283. <https://doi.org/10.3389/fnhum.2018.00283>
- Snyder, H. R., Miyake, A., & Hankin, B. L. (2015). Advancing understanding of executive function impairments and psychopathology: bridging the gap between clinical and cognitive approaches. *Frontiers in Psychology*, 6, 328. doi:10.3389/fpsyg.2015.00328.
- Sprague, J., Verona, E., Kalkhoff, W., & Kilmer, A. (2011). Moderators and mediators of the stress-aggression relationship: executive function and state anger. *Emotion*, 11(1), 61–73. <https://doi.org/10.1037/a0021788>
- *Stratton, J., Cobia, D. J., Reilly, J., Brook, M., & Hanlon, R. E. (2018). Differences in neuropsychological functioning between homicidal and nonviolent schizophrenia samples. *Journal of Forensic Sciences*, 63(5), 1435–1443. <https://doi.org/10.1111/1556-4029.13750>
- *Swann, A. C., Lijffijt, M., Lane, S. D., Steinberg, J. L., & Moeller, F. G. (2009). Trait impulsivity and response inhibition in antisocial personality disorder. *Journal of Psychiatric Research*, 43(12). doi:1057-1063.

- *Syngelaki, E. M., Moore, S. C., Savage, J. C., Fairchild, G., & Van Goozen, S. H. (2009). Executive functioning and risky decision making in young male offenders. *Criminal Justice and Behavior*, 36(11), 1213–1227. <https://doi.org/10.1177/0093854809343095>
- *Szczypliński, J., Wypych, M., Krasowska, A., Wiśniewski, P., Kopera, M., Suszek, H., & Wojnar, M. (2022). Abnormal behavioral and neural responses in the right dorsolateral prefrontal cortex during emotional interference for cognitive control in pedophilic sex offenders. *Journal of Psychiatric Research*, 151, 131–135. <https://doi.org/10.1016/j.jpsychires.2022.04.012>
- *Tung, S., & Chhabra, N. (2011). A comparative study on the neuropsychological status of delinquent and non-delinquent boys. *International Journal of Culture and Mental Health*, 4(2), 121–127. <https://doi.org/10.1080/17542863.2010.530772>
- *Turner, D., Laier, C., Brand, M., Bockshammer, T., Welsch, R., & Rettenberger, M. (2018). Response inhibition and impulsive decision-making in sexual offenders against children. *Journal of Abnormal Psychology*, 127(5), 471. <https://doi.org/10.1037/abn0000359>
- *Turner, D., Roszyk, A., Szumski, F., & Rettenberger, M. (2020). Deviant sexual interests but not antisocial behaviors are associated with deficits in executive functioning in individuals convicted of sexual offenses against children. *Neuropsychology*, 34(8), 906. <https://doi.org/10.1037/neu0000703>
- *Urazán-Torres, G. R., Puche-Cabrera, M. J., Caballero-Forero, M., & Rey-Anacona, C. A. (2013). Cognitive and executive functions in colombian school children with conduct disorder: sex differences. *Revista Colombiana de Psiquiatría*, 42(4), 324–332. [https://doi.org/10.1016/S0034-7450\(13\)70029-5](https://doi.org/10.1016/S0034-7450(13)70029-5)
- *Uytun, M. C., Karakaya, E., Oztop, D. B., Gengec, S., Gumus, K., Ozmen, S., & Ozsoy, S. D. (2017). Default mode network activity and neuropsychological profile in male children and adolescents with attention deficit hyperactivity disorder and conduct disorder. *Brain Imaging and Behavior*, 11, 1561–1570. <https://doi.org/10.1007/s11682-016-9614-6>
- Van De Schoot, R., De Bruin, J., Schram, R., Zahedi, P., De Boer, J., Weijdema, F., ... Oberski, D. L. (2021). An open source machine learning framework for efficient and transparent systematic reviews. *Nature Machine Intelligence*, 3(2), 125–133. <https://doi.org/10.1038/s42256-020-00287-7>
- *Van der Meere, J., Van der Meer, D. J., Borger, N., & Pirila, S. (2008). Attention and impulse control in children with borderline intelligence with or without conduct disorder. *Perceptual and Motor Skills*, 107(3), 676–682. <https://doi.org/10.2466/pms.107.3.676-682>
- Van der Sluys, M. E., Zijlmans, J., Popma, A., Van der Laan, P. H., Scherder, E. J. A., & Marhe, R. (2020). Neurocognitive predictors of treatment completion and daytime activities at follow-up in multiproblem young adults. *Cognitive, Affective, & Behavioral Neuroscience*, 20(5), 1103–1121. <https://doi.org/10.3758/s13415-020-00822-4>
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36, 1–48. <https://doi.org/10.18637/jss.v036.i03>
- *Vila-Ballo, A., Cunillera, T., Rostan, C., Hdez-Lafuente, P., Fuentemilla, L., & Rodriguez-Fornells, A. (2015). Neurophysiological correlates of cognitive flexibility and feedback processing in violent juvenile offenders. *Brain Research*, 1610, 98–109. <https://doi.org/10.1016/j.brainres.2015.03.040>
- *Vilá-Balló, A., Hdez-Lafuente, P., Rostan, C., Cunillera, T., & Rodriguez-Fornells, A. (2014). Neurophysiological correlates of error monitoring and inhibitory processing in juvenile violent offenders. *Biological Psychology*, 102, 141–152. <https://doi.org/10.1016/j.biopsycho.2014.07.021>
- *Vitoria-Estruch, S., Romero-Martínez, A., Lila, M., & Moya-Albiol, L. (2018). Differential cognitive profiles of intimate partner violence perpetrators based on alcohol consumption. *Alcohol*, 70, 61–71. <https://doi.org/10.1016/j.alcohol.2018.01.006>
- *Vloet, T. D., Konrad, K., Herpertz-Dahlmann, B., & Kohls, G. (2011). The effect of social and monetary reward on inhibitory control in boys with hyperkinetic conduct disorder. *Zeitschrift für Kinder- und Jugendpsychiatrie und Psychotherapie*, 39(5), 341–349. <https://doi.org/10.1024/1422-4917/a000127>
- Wade, M., Zeanah, C. H., Fox, N. A., & Nelson, C. A. (2020). Global deficits in executive functioning are transdiagnostic mediators between severe childhood neglect and psychopathology in adolescence. *Psychological Medicine*, 50(10), 1687–1694. <https://doi.org/10.1017/S0033291719001764>
- *Weidacker, K., Kärge, C., Massau, C., Krueger, T. H. C., Walter, M., Ponseti, J., ... Schiffer, B. (2022). Interference inhibition in offending and non-offending pedophiles: A preliminary event-related fMRI study. *Neuropsychologia*, 173, Article 108301. <https://doi.org/10.1016/j.neuropsychologia.2022.108301>
- Wells, G. A., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., & Tugwell, P. (2000). The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. https://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
- *Xu, M., Jiang, W., Du, Y., Li, Y., & Fan, J. (2017). Executive function features in drug-naïve children with oppositional defiant disorder. *Shanghai Archives of Psychiatry*, 29(4), 228. <https://doi.org/10.11919/j.issn.1002-0829.216104>
- Yang, Y., Shields, G. S., Zhang, Y., Wu, H., Chen, H., & Romer, A. L. (2022). Child executive function and future externalizing and internalizing problems: A meta-analysis of prospective longitudinal studies. *Clinical Psychology Review*, 97, Article 102194. <https://doi.org/10.1016/j.cpr.2022.102194>
- Yukhnenko, D., Sridhar, S., & Fazel, S. (2019). A systematic review of criminal recidivism rates worldwide: 3-year update. *Wellcome Open Research*, 4(28), 28. <https://doi.org/10.12688/wellcomeopenres.14970.3>
- Zelazo, P. D. (2020). Executive Function and Psychopathology: A Neurodevelopmental Perspective. *Annual Review of Clinical Psychology*, 16(1), 431–454. <https://doi.org/10.1146/annurev-clinpsy-072319-024242>
- *Zhu, Y., Liu, L., Yang, D., Ji, H., Huang, T., Xue, L., & Fang, Y. (2021). Cognitive control and emotional response in attention-deficit/hyperactivity disorder comorbidity with disruptive, impulse-control, and conduct disorders. *BMC Psychiatry*, 21(1), 1–13. <https://doi.org/10.1186/s12888-021-03221-2>
- *Zhu, Y., Ying, K., Wang, J., Su, L., Chen, J., Lin, F., & Wang, S. (2014). Differences in functional activity between boys with pure oppositional defiant disorder and controls during a response inhibition task: a preliminary study. *Brain Imaging and Behavior*, 8(4), 588–597. <https://doi.org/10.1007/s11682-013-9275-7>
- Zijlmans, J., Marhe, R., Bevaart, F., Van Duin, L., Luijckx, M. A., Franken, L., ... Popma, A. (2021). The predictive value of neurobiological measures for recidivism in delinquent male young adults. *Journal of Psychiatry and Neuroscience*, 46(2), E271–E280. <https://doi.org/10.1503/jpn.200103>
- *Zou, Z., Meng, H., Ma, Z., Deng, W., Du, L., Wang, H., & Hu, H. (2013). Executive functioning deficits and childhood trauma in juvenile violent offenders in China. *Psychiatry Research*, 207(3), 218–224. <https://doi.org/10.1016/j.psychres.2012.09.013>