



Universiteit
Leiden
The Netherlands

Three-dimensional charge density wave in the dual heavy fermion system U Pt₂Si₂

Petkov, V.; Baumbach, R.; Milinda Abeykoon, A.M.; Mydosh, J.A.

Citation

Petkov, V., Baumbach, R., Milinda Abeykoon, A. M., & Mydosh, J. A. (2023). Three-dimensional charge density wave in the dual heavy fermion system U Pt₂Si₂. *Physical Review B*, 107(24). doi:10.1103/PhysRevB.107.245101

Version: Publisher's Version

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/3728696>

Note: To cite this publication please use the final published version (if applicable).

Three-dimensional charge density wave in the dual heavy fermion system UPt_2Si_2

Valeri Petkov^{1,*}, R. Baumbach^{2,3}, A. M. Milinda Abeykoon⁴, and J. A. Mydosh⁵

¹*Department of Physics, Central Michigan University, Mt. Pleasant, Michigan 48858, USA*

²*National High Magnetic Field Laboratory, Florida State University, Tallahassee, Florida 32310, USA*

³*Department of Physics, Florida State University, Tallahassee, Florida 32310, USA*

⁴*Photon Sciences Division, Brookhaven National Laboratory, Upton, New York 11973, USA*

⁵*Institute Lorentz, Leiden University, Leiden 2300RA, The Netherlands*



(Received 26 February 2023; accepted 15 May 2023; published 1 June 2023)

Heavy fermion liquids offer via their Kondo lattice diverse possibilities for exotic ground states. Using variable-temperature atomic pair distribution function analysis, we study the local atomic structure of the “dual” heavy fermion liquid UPt_2Si_2 , which exhibits antiferromagnetism consistent with localized $5f$ -electron states and transport properties characteristic to itinerant $5f$ -“ spd ” hybridized electron systems. We show that UPt_2Si_2 exhibits periodic lattice distortions (PLDs) involving both uranium and platinum atoms that are characteristic to three-dimensional charge density waves. The temperature evolution of the PLDs tracks that of the transport and magnetic properties, suggesting the presence of little-known $5f$ -electron-lattice interactions. We argue that PLDs in heavy fermion liquids in general, and in particular in UPt_2Si_2 , appear as new degrees of freedom that entangle competing electronic states and, as such, must be accounted for when their rich physics is considered.

DOI: [10.1103/PhysRevB.107.245101](https://doi.org/10.1103/PhysRevB.107.245101)

I. INTRODUCTION

Quantum materials with diverse interactions (lattice, charge, spin, and orbital) are at the nexus of efforts to produce novel states of matter. Among these materials, strongly correlated f -electron systems serve as an especially deep reservoir for novel behaviors, where examples include heavy-fermion Kondo lattices, unconventional superconductors, hidden order, hybridization-gap topological insulators, and Weyl-Kondo semimetals [1–8]. A large number of such systems belong to the ThCr_2Si_2 structure-type family, which is derived from the prototypical BaAl_4 structure [9–21]. Less well studied are systems that crystallize in the closely related CaBe_2Ge_2 -type structure, which, contrary to the ThCr_2Si_2 structure, lacks inversion symmetry along the c -axis. Recently, materials with this structure have attracted increased interest because of the discovery of (i) a possible topological superconductivity in CeRh_2As_2 and (ii) a prediction of gapless Weyl-Kondo nodal lines in CePt_2Si_2 and CeRh_2Ga_2 [22–29].

This led us to focus on UPt_2Si_2 (CaBe_2Ge_2 structure type), which had long been thought to exhibit prototypical localized f -electron behavior with crystal-electric field splitting of the localized uranium $5f$ -electron states and antiferromagnetic (AF) order at $T_N = 34$ K [30,31]. However, several recent studies have challenged this perspective by providing evidence for the presence of $5f$ orbitals—conduction “ spd ” electrons hybridization, similar to what is seen in other Kondo lattice materials [32–34]. This includes new evidence for coherence behavior in the electrical transport properties; the observation of high magnetic-field phase transitions suggesting the presence of Fermi surface instabilities due to

a breakdown of the Kondo coherent ground state; inelastic neutron-scattering measurements that are consistent with a dual, localized, itinerant $5f$ -electron character; and density functional theory calculations that favor a scenario where the $5f$ -electrons of uranium are mostly itinerant. Furthermore, x-ray diffraction (XRD) measurements revealed large anisotropic thermal factors for atoms occupying the Si-Pt-Si layers, which were interpreted as an indication of strong crystallographic disorder [35]. This disorder was proposed to be responsible for Anderson localization along the c -axis of the crystal lattice, resulting in an anisotropic resistivity. However, more recent studies connected this behavior to the emergence of charge density wave (CDW) order at an anomalously large temperatures $T_{\text{CDW}} = 315$ K [29].

Collectively, these behaviors reveal an unusual diversity of phenomena with a potential interplay between Kondo lattice hybridization, CDW, and AF orders. However, there remains uncertainty about the interrelationship between these diverse phenomena and the crystal structure, including the periodic lattice distortions (PLDs) known to accompany CDWs. To clarify these questions, we use variable-temperature atomic pair distribution function (PDF) analysis coupled with structure modeling to study lattice distortions in UPt_2Si_2 . We find that UPt_2Si_2 is a rare example of a ternary Fermi liquid where both metallic species are displaced from their position in the undistorted crystal lattice in a correlated manner. The displacements appear above T_{CDW} and evolve nonlinearly with diminishing temperature, increasingly disturbing the spatial coherence of the crystal lattice. Moreover, the deviations from linearity closely track changes in the electronic and magnetic properties, indicating the presence of a little-known strong interaction between lattice, electronic, and magnetic degrees of freedom in UPt_2Si_2 . Beyond this, our results call for similar local structure studies of other members of CaBe_2Ge_2 - and

*petko1vg@cmich.edu




FIG. 1. (a) Fragment from the crystal structure of UPt_2Si_2 featuring a sequence of alternating Pt(1)-Si(1)-Pt(1) and Si(2)-Pt(2)-Si(2) layers and uranium atoms (in green) positioned between the layers. Platinum atoms are in black and silicon atoms are in brown. (b) Resistivity of UPt_2Si_2 as a function of temperature. Arrows mark inflection points on the resistivity versus temperature curve corresponding to the onset of a charge density wave (T_{CDW}), change of conductivity type (T_{coh}), and emergence of antiferromagnetic (T_N) order. (c) Hysteresis curves (magnetization vs. external magnetic field) for UPt_2Si_2 measured at three different temperatures. (d) Magnetic susceptibility (T) for UPt_2Si_2 . The broad peak at $T_N = 37$ K reflect the emergence of an antiferromagnetic order with decreasing temperature.

ThCr_2Si_2 -type families of heavy fermion liquids, where models that neglect lattice distortions and local crystal symmetry breaking may not fully describe the observed fascinating phenomena.

II. EXPERIMENTAL

A. Sample preparation and properties characterization

Polycrystalline samples of UPt_2Si_2 were synthesized by arc melting the constituent elements (99.99% pure, lump form) in a 1:2:2 molar ratio of U:Pt:Si. The resulting boule was flipped and re-arc'd five times to ensure homogeneity. The final product was faceted, indicating the formation of large grains. In-house x-ray diffraction studies on powdered pieces showed that the as-cast UPt_2Si_2 possesses a CaBe_2Ge_2 -type structure (space group $P4/nmm$), shown in Fig. 1(a).

Specimens for electrical resistivity measurements were prepared by cleaving fragments from the main boule. This produces slabs of material, to which platinum wires were attached using a micro-spot-welding device. Standard four-wire electrical resistance $R(T)$ measurements were performed for $1.8 < T < 350$ K using a quantum design physical properties measurement system (PPMS). Results for the room-temperature normalized resistivity $R/R_{300\text{K}}$, shown Fig. 1(b), are consistent with earlier reports. In particular, $R/R_{300\text{K}}$ shows a strongly nonlinear evolution with

temperature, where clear inflection points are observed at $T_{\text{CDW}} = 315$ K and $T_N = 37$ K. Below T_{CDW} , $R/R_{300\text{K}}$ increases with decreasing temperatures and goes through a maximum at $T_{\text{coh}} = 150$ K. Such a phenomenon is typically observed with Kondo-lattice systems and is associated with the emergence of coherent hybridization between the f - and conduction electron states, leading to a reduction of the spin disorder scattering and low-temperature formation of a heavy Fermi liquid state. Finally, there is a rapid decrease at T_N , consistent with the further removal of the spin disorder scattering due to magnetic ordering, and a saturation toward a relatively large residual value of $R_{2\text{K}}/R_{300\text{K}} = 0.4$.

The magnetic properties were also studied using a PPMS. Results for the magnetic susceptibility $\chi(T)$ and magnetization as a function of magnetic field are shown in Figs. 1(c) and 1(d). The $\chi(T)$ data exhibit a Curie-Weiss behavior for $100 \text{ K} < T < 300 \text{ K}$, where fits to the data using the expression $\chi(T) = C/(T - \Theta)$ yield an effective (high-temperature) magnetic moment $\mu_{\text{eff}} = 3.20 \mu_B/\text{U}$ and $\Theta = -60$ K. That is to be compared with the low-temperature, AF-ordered magnetic moment of $1.7 \mu_B/\text{U}$ at 4.2 K observed by neutron-scattering experiments [36]. The difference between the uranium moments at high temperature ($3.2 \mu_B$) and 4.2 K ($1.7 \mu_B$) indicates the presence of a strong hybridization between the $5f$ -electrons of uranium and “ spd ” conduction electrons, leading to a Kondo-type reduction of




FIG. 2. (a) and (b) XRD intensity color maps for UPt_2Si_2 collected in the temperature range from 10 K to 400 K. The maps cover different ranges of Bragg angles. While the low-angle Bragg peaks shown in (a) do not seem to change much with temperature, the high-angle Bragg peaks shown in (b) are seen to show a complex temperature dependence. (c) Atomic PDF intensity color map for UPt_2Si_2 . The PDF peak at about 3 Å splits into two components below $T_{\text{CDW}} = 315$ K. The intensity of the peak at 4.25 Å suddenly starts increasing below 100 K (see the horizontal arrows). (d) Selected atomic PDFs for UPt_2Si_2 . Vertical arrows mark PDF features, i.e., atomic pair distances, that evolve markedly with temperature.

the local uranium moments. In principle, crystal electric field splitting of the $5f$ -electron multiplet could also result in a reduction of the low-temperature moment of uranium. The AF ordering, where the uranium magnetic moments are aligned along the c -axis of the crystal lattice, is seen as a strong decrease in $\chi(T)$ below T_N [Fig. 1(d)]. The nearly linear response of $M(H)$ at 5 K [Fig. 1(c)], i.e., within the ordered state, is also consistent with antiferromagnetism.

B. Synchrotron radiation studies

Synchrotron high-energy XRD experiments were conducted at the beamline 28-ID-1 at the National Synchrotron Light Source-II, Brookhaven National Laboratory using x rays with an energy of 74.46 keV ($\lambda = 0.1665$ Å). XRD data were collected while varying temperature between 10 K and 400 K in steps of 5 K. Experimental XRD patterns and atomic PDFs derived from the data using standard procedures [37] are summarized in Fig. 2.

III. STRUCTURE MODELING

A. Average crystal structure as a function of temperature

To reveal the evolution of the average crystal structure of UPt_2Si_2 with temperature, synchrotron XRD data

were subjected to Rietveld analysis using the software package FullProf [38]. Representative Rietveld fits based on a space group (SG) $P4/nmm$ model are shown in Figs. 3(a) and 3(b). The fits were successful and produced the lattice parameters and unit cell volume summarized in Figs. 3(c) and 3(d), and Fig. 3(e), respectively. The c lattice parameter and volume are seen to diminish smoothly with decreasing temperature. By contrast, the rate of decrease of the a lattice parameter with decreasing temperature changes markedly at T_{CDW} , indicating that the atomic displacements related to the emergent CDW are likely to occur in the basal planes of the crystal lattice.

B. Local atomic structure as a function of temperature

As shown in Fig. 2(b), higher angle Bragg peaks, which are extra sensitive to the local atomic structure, show an intricate evolution with temperature. Inspection of the respective atomic PDFs [Fig. 2(c)], which directly reflect frequently occurring atomic pair distances, shows that the peak positioned near 3.0 Å splits into two components below 350 K. The two components increasingly become separated from each other with decreasing temperature, and the intensity of the PDF peak at about 4.25 Å suddenly increases below 100 K. The profile of the PDF peak at about 5.5 Å also changes




FIG. 3. (a) and (b) Successful Rietveld fits to XRD patterns for UPt_2Si_2 . The fits are based on a tetragonal (SG $P4/nmm$) model as explained in the text. The goodness-of-fit factor R_{wp} for the fits are on the order of 9%. Experimental data are given as symbols, the computed data are given as red lines, and the residual difference (shifted for clarity) is given as a blue line. (c)–(e) Rietveld refined tetragonal lattice parameters and unit cell volume given as solid symbols. PDF refined lattice parameters and unit cell volume are also given as open symbols. Blue arrows mark inflection points on the lattice parameters versus temperature curves where CDW (T_{CDW}) and AF (T_N) orders emerge on cooling. The difference between Rietveld and PDF values indicates that, locally, the atomic arrangement in UPt_2Si_2 is more compressed in comparison to the average crystal structure. Error bars in (c)–(e) are close to the size of used symbols.

significantly with temperature. The evolution of interatomic distances in UPt_2Si_2 is well illustrated in Fig. 2(d), where selected atomic PDFs are shown. Here it is seen that, locally, the crystal lattice continuously evolves with temperature. To assess the evolution in more detail, we fit the experimental PDF data with a model based on the SG $P4/nmm$ structure using the Rietveld refined structure data as a starting point. Note that atomic PDF analysis is advantageous to Rietveld analysis when exploring local distortions of the crystal lattice because it takes into account both the diffuse and Bragg components of the diffraction data, while Rietveld analysis uses only the latter and, hence, is mainly sensitive to the average long-range crystal structure [39,40].

Fits including interatomic distances within two unit cells of UPt_2Si_2 (up to $\sim 20\text{ Å}$) are shown in Fig. 4. As can be seen in Figs. 4(a) and 4(b), several PDF features, including the sequence of well-defined PDF peaks positioned between 2 Å and 4 Å , and the cluster of overlapping PDF peaks at about 5.5 Å , i.e., the local atomic structure of UPt_2Si_2 , are not well reproduced by this model. Therefore, we allowed atoms in the tetragonal unit cell to move away from their position in the undistorted lattice, thus locally breaking the tetragonal lattice symmetry. The fits improved to an acceptable level, as demonstrated in Figs. 4(c) and 4(d), only when both Pt(2) and uranium atoms underwent a considerable displacement from their position in the undistorted lattice, while changes in the position of Pt(1), Si(1), and Si(2) atoms deteriorated the fits.

Changes in the position of Pt(2) atoms were expected because prior single crystal studies [29] indicated that they are associated with the CDW in UPt_2Si_2 emerging with decreasing temperature. A change in the position of uranium atoms was unexpected.

Fits including interatomic distances longer than 20 Å are shown in Fig. 5. As can be seen in Figs. 5(b) and 5(d), at 400 K , the lattice distortions in UPt_2Si_2 do not affect its long-range structure significantly. By contrast, at 10 K , the lattice distortions disturb the long-range structure over interatomic distances extending to about 35 Å , above which the average tetragonal crystal symmetry is recovered. Tetragonal lattice parameters and unit cell volume derived from the PDFs in Fig. 4 are compared with the Rietveld derived values in Figs. 3(c)–3(e). A fragment of the PDF-refined crystal structure of UPt_2Si_2 at 10 K and a projection of the fragment on the basal plane (ab) of the tetragonal lattice are shown in Fig. 6(a) and Fig. 6(b), respectively.

IV. DISCUSSION

As can be seen in Figs. 3(c)–3(e), the PDF refined lattice parameters and unit cell volume for UPt_2Si_2 are systematically smaller than the Rietveld refined values. This observation indicates that the local crystal structure is more densely packed in comparison with the bulk values. This may not come as a surprise because the CaBe_2Ge_2 structure type is




FIG. 4. (a) and (b) Unsuccessful and (c) and (d) successful fits to the low- r part of PDFs for UPt_2Si_2 obtained at two characteristic temperatures. The unsuccessful fits are based on a tetragonal (SG $P4/nmm$) model while the successful fits are based on a tetragonal (SG $P4/nmm$) model where the local symmetry is broken as explained in the text. The goodness-of-fit factor R_{wp} for the unsuccessful fits is on the order of 40% while that for the successful fits is on the order of 12%. Experimental data are given as symbols, computed data are given as red lines, and the residual difference (shifted for clarity) is given as a blue line.

relatively open, allowing the constituent atoms to change their positions to minimize the energy of CaBe_2Ge_2 structure-type compounds when they are subjected to external stimuli. Contrary to the case of Rietveld refined values, the temperature variation of the PDF refined values is uneven, showing inflection points at temperatures close to T_{CDW} and T_N . Moreover, the a lattice parameter counterintuitively increases and then barely evolves with temperature over a temperature region centered at T_{coh} [compare data in Figs. 1(b) and 3(d)]. Evidently, changes in the transport and magnetic properties of UPt_2Si_2 and those in its local atomic structure are strongly correlated.

Among all atoms in the unit cell, only the Pt(2) atoms in the Si(2)-Pt(2)-Si(2) planes and the uranium atoms positioned between the planes [see Fig. 6(a)] are significantly displaced from their positions in the undistorted lattice. For both atoms, the in-plane component of the displacement is much larger than the out-of-plane component. Also, the in-plane component of the displacements for Pt(2) atoms is larger than the in-plane component of uranium displacements. The opposite is true for the out-of-plane components. In particular, the in-plane displacement for Pt(2) atoms increases with diminishing temperature from a value of 0.04 Å at 400 K to a value of 0.15 Å at 10 K. By contrast, the Pt(2) out-of-plane displacement decreases from the 400 K value of 0.014 Å to the near-negligible value of 0.004 Å at 10 K. On the other hand, the in-plane displacement for uranium atoms decreases with decreasing temperature from a value of 0.07 Å at 400 K to a value of 0.03 Å at 10 K. At the same time, the out-of-plane displacement for uranium atoms

barely increases from its 400 K value of about 0.005 Å to 0.009 Å at 10 K. Notably the displacements for Pt(2) and that for uranium atoms concurrently evolve with diminishing temperature, clearly indicating the presence of significant $5d$ - $5f$ -electron hybridization in UPt_2Si_2 . Overall, the displacements of Pt(2) and uranium atoms follow a repetitive pattern, characteristic to PLDs known to accompany CDWs. In line with recent single-crystal studies [29], the PLDs are largely confined to the basal layers comprising Pt(2) atoms sandwiched between silicon planes. However, because both Pt(2) and uranium atoms are also considerably displaced in an out-of-plane direction, the CDW appearing in UPt_2Si_2 below 315 K is essentially a three-dimensional (3D) modulation of the crystal lattice. Signatures of the CDW transition are seen as a sudden increase in the resistivity [Fig. 1(b)] and wedge-like bump in the specific heat appearing between 305 K and 320 K [29,41].

From the structure data obtained in this and previous studies [29], the following picture for the relationship between lattice, electronic, and magnetic degrees of freedom in UPt_2Si_2 emerges. At temperatures well above T_{CDW} , Pt(2) and uranium atoms are already displaced from their positions in the undistorted tetragonal lattice. The displacements may be viewed as a precursor to the PLDs emerging below T_{CDW} [29]. Likely because the PLDs are short ranged near T_{CDW} , i.e., do not disturb the long-range structure (>20 Å) of UPt_2Si_2 significantly [see Figs. 5(b) and 5(d)], the emerged CDW appears commensurate with the underlying tetragonal crystal lattice. Nonetheless, the PLDs are large enough to increase the local atomic packing significantly in




FIG. 5. (a) Unsuccessful and (c) successful fit to the higher r part of the PDF for UPt_2Si_2 obtained at 10 K. The unsuccessful fit is based on a tetragonal (SG $P4/nmm$) model while the successful fit is based on a tetragonal (SG $P4/nmm$) model where the local symmetry is broken as explained in the text. The latter model also fits well the low- r part of the 10 K PDF data set, as shown in Fig. 4(c). (b) and (d) Near-equally successful fits to the high- r part of the PDF for UPt_2Si_2 obtained at 400 K. The fits are based on (b) an unmodified tetragonal (SG $P4/nmm$) model and (d) a tetragonal (SG $P4/nmm$) model where the local symmetry is broken as explained in the text. Evidently, at high temperatures, the lattice distortions in UPt_2Si_2 leave its long-range structure largely intact, i.e., they need not be necessarily evoked to explain it, which is not true for the shorter range structure [see Fig. 4(b)]. By contrast, at low temperatures, the lattice distortions significantly disturb the long-range structure of UPt_2Si_2 up to distances of about 35 Å [vertical broken line in (a)], above which the average tetragonal crystal symmetry is recovered (see the sharp drop in the fit residual). The goodness-of-fit factor R_{wp} for the unsuccessful fit in (a) is 28% while that for the successful fits is on the order of 12%. Experimental data are given as symbols, computed data are given as red lines, and the residual difference (shifted for clarity) is given as a blue line.

comparison to the average structure. The nonuniform change in the atomic displacements, related bonding distances, and local volume with temperature may be expected to render the thermal evolution of transport and magnetic properties nonlinear because the volume and electronic structure of heavy fermion liquids from the CaBe_2Ge_2 -type family are strongly coupled [10]. In particular, the slow increase in the electrical resistivity upon decreasing temperature below T_{CDW} can be associated with the gradual increase in the in-plane Pt(2) and out-of-plane uranium displacements, the latter of which reaches a plateau at T_{coh} . The in-plane displacements of uranium, U_{ab} , also reach a plateau at T_{coh} [(see Fig. 6(c)]. The subsequent decrease in the resistivity with decreasing temperature mirrors the decrease in the out-of-plane Pt(2) displacements. Then, the high value of the residual resistivity at very low temperatures can be attributed to residual electron-phonon scattering arising from the significant in-plane Pt(2) and out-of-plane uranium displacements. In addition, the local breaking of crystal symmetry will affect the crystal field splitting of the uranium 5*f*-electron multiplet and, hence, contributes to the observed changes in the magnetic moment of uranium atoms with temperature. Moreover, the periodic displacement of uranium atoms and related periodic modulation of the local crystal symmetry may modulate the AF

ordered pattern of uranium magnetic moments below T_N , as observed by resonant x-ray scattering experiments [8]. The observed irreversibility in the magnetic susceptibility below T_N can also be explained by the presence of local lattice distortions involving the magnetically active uranium atoms [42]. Finally, the coupled sharp increase and decrease of platinum and uranium atomic displacements below T_N signal an increased lattice instability, which may contribute to the complex magnetic phase diagram observed below 20 K in high magnetic fields [34]. The increased lattice instability is also demonstrated by the increased spatial extent of the PLDs distortions (~ 35 Å) at low temperatures [Figs. 5(a) and 5(c)]. The latter can be a major factor behind the observed [29] change of the character of the CDW from initially commensurate at high temperatures to distinctly incommensurate at low temperatures. More precisely, it could be that, at low temperatures, the incommensurate CDW structure of UPt_2Si_2 consists of commensurate CDW domains with a size of about 35 Å that are separated by discommensurations, e.g., narrow domain walls where the CDW phase changes fast [43]. Analysis of single crystal experiments suggests that the observed satellites signaling the emerged CDW order can indeed be regarded as overlapped patterns arising from such domains [29].




FIG. 6. (a) Fragment from the structure of UPt_2Si_2 at 10 K as derived from PDF analysis where Pt(2) atoms (red circles) appear displaced from their positions in the undistorted tetragonal structure (light-gray circles). Uranium atoms (solid green circles) also appear displaced from their positions in the undistorted lattice (light-green circles). By contrast, Si(1), Si(2), and Pt(1) are not displaced significantly from their positions in the undistorted lattice. (b) Projection of the displacements of Pt(2) and uranium atoms on the (ab) atomic plane. Blue arrows point in the direction of Pt(2) displacements. The brown dashed line, resembling a sine wave, emphasizes the periodicity in the Pt(2) atom displacements, forming a PLD. The same pertains to the green dashed line connecting the positions of displaced uranium atoms. (c) Amplitude of the atomic displacements of Pt(2) and uranium atoms from their positions in the undistorted lattice. Blue arrows mark inflection points on the displacement versus temperature curves where the CDW (T_{CDW}) and AF (T_N) orders set in. Black dashed lines highlight a range of temperatures, centered at about $T_{\text{coh}} = 150$ K, where, counterintuitively, the vertical and in-plane displacements of uranium atoms increase with diminishing temperature before reaching 100 K, and then remain unchanged down to 10 K. Error bars in (c) are close to the size of used symbols.

V. CONCLUSION

Magnetic, transport, and bonding phenomena in heavy fermion liquids such as UPt_2Si_2 are sensitive to the degree of hybridization between localized uranium $5f$ orbitals and “ spd ” conduction electrons, which, in turn, depends on the interatomic distances and crystal lattice symmetry. The spatial extent of the orbitals and their proximity to the Fermi level render these materials susceptible to external stimuli and give rise to competing crystallographic modifications, including local variations in the atomic positions, packing density, and volume. As our study shows, the variations in the atomic positions largely involve in-plane platinum and out-of-plane uranium displacement modes, thus forming an exotic 3D CDW below T_{CDW} . In addition, they progress nonlinearly with decreasing temperature, inevitably leading to concurrent nonlinear changes in both the magnetic and transport properties. The results suggest

that the marked flexibility of the crystal lattice must be accounted for in order to understand the rich physics of UPt_2Si_2 and, by extension, other heavy $5f$ fermion liquids in general.

ACKNOWLEDGMENTS

This work was supported by the US Department of Energy (DOE), Office of Science, Office of Basic Energy Sciences (Award No. DE-SC0021973) and used resources of the National Synchrotron Light Source at the Brookhaven National Laboratory provided by the DOE Office of Science (Contract No. DE-SC0012704). R.B. was supported by the National Science Foundation (NSF DMR-1904361). The National High Magnetic Field Laboratory is supported by the National Science Foundation (NSF DMR-1644779) and the State of Florida. Thanks are due to Dr. D. R. Tadiseti for the help with Rietveld analysis.

[1] I. Osborne and R. Coontz, Quantum wonderland, *Science* **319**, 1201 (2008).

[2] Q. Si and F. Steglich, Heavy fermions and quantum phase transitions, *Science* **329**, 1161 (2010).

- [3] P. Monthoux, D. Pines, and G. G. Lonzarich, Superconductivity without phonons, *Nature (London)* **450**, 1177 (2007).
- [4] C. Broholm, R. J. Cava, S. A. Kivelson, D. G. Nocera, M. R. Norman, and T. Senthil, Quantum spin liquids, *Science* **367**, 6475 (2020).
- [5] V. Martelli, A. Cai, E. M. Nica, and S. Paschen, Sequential localization of a complex electron fluid, *Proc. Natl. Acad. Sci. USA* **116**, 17701 (2019).
- [6] J. A. Mydosh, P. M. Oppeneer, and P. S. Riseborough, Hidden order and beyond: An experimental–theoretical overview of the multifaceted behavior of URu₂Si₂, *J Phys. Condens. Matter* **32**, 143002 (2020).
- [7] A. D. Huxley, Ferromagnetic superconductors, *Physica C* **514**, 368 (2015).
- [8] F. Kon, C. Tabata, K. Miura, R. Hibino, H. Hidaka, T. Yanagisawa, H. Nakao, and H. Amitsuka, Correlation between antiferromagnetic and charge-density-wave order in UPt₂Si₂ studied by resonant x-ray scattering, in *Proceedings of the International Conference on Strongly Correlated Electron Systems (SCES 2022)*, Amsterdam (SciPost Physics, 2022), https://scipost.org/preprints/scipost_202208_00025v1/.
- [9] M. Shatruk, ThCr₂Si₂ structure type: The “perovskite” of intermetallics, *J. Solid State Chem.* **272**, 198 (2019).
- [10] Y. Lai, J. Y. Chan, and R. E. Baumbach, Electronic landscape of the *f*-electron intermetallics with the ThCr₂Si₂ structure, *Sci. Adv.* **8**, eabp8264 (2022).
- [11] Y. Lai, K. Wei, G. Chappell, J. Diaz, T. Siegrist, P. J. W. Moll, D. Graf, and R. E. Baumbach, Tuning the structural and antiferromagnetic phase transitions in UCr₂Si₂: Hydrostatic pressure and chemical substitution, *Phys. Rev. Mater.* **4**, 075003 (2020).
- [12] H. Q. Yuan, F. M. Grosche, M. Deppe, C. Geibel, G. Sparn, and F. Steglich, Observation of two distinct superconducting phases in CeCu₂Si₂, *Science* **302**, 2104 (2003).
- [13] E. Schuberth, M. Tippmann, L. Steinke, S. Lausberg, A. Steppke, M. Brando, C. Krellner, C. Geibel, R. Yu, Q. Si, and F. Steglich, Emergence of superconductivity in the canonical heavy-electron metal YbRh₂Si₂, *Science* **351**, 485 (2016).
- [14] H. S. Jeevan, C. Geibel, and Z. Hossain, Quasiquartet crystal-electric-field ground state with possible quadrupolar ordering in the tetragonal compound YbRu₂Ge₂, *Phys. Rev. B* **73**, 020407 (2006).
- [15] Z. Ren, L. V. Pourovskii, G. Girit, G. Lapertot, A. Georges, and D. Jaccard, Giant Overlap Between the Magnetic and Superconducting Phases of CeAu₂Si₂ Under Pressure, *Phys. Rev. X* **4**, 031055 (2014).
- [16] S. Süllow, M. C. Aronson, B. D. Rainford, and P. Haen, Doniach Phase Diagram, Revisited: From Ferromagnet to Fermi Liquid in Pressurized CeRu₂Ge₂, *Phys. Rev. Lett.* **82**, 2963 (1999).
- [17] M. B. Fontes, M. A. Continentino, S. L. Bud’ko, M. El-Massalami, L. C. Sampaio, A. P. Guimarães, E. Baggio-Saitovitch, M. F. Hundley, and A. Lacerda, Physical properties of the Ce(Ru_{1-x}Fe_x)₂Ge₂ series, *Phys. Rev. B* **53**, 11678 (1996).
- [18] L. C. Gupta, D. E. MacLaughlin, C. Tien, C. Godart, M. A. Edwards, and R. D. Parks, Magnetic behavior of the Kondo-lattice system CeRu₂Si₂, *Phys. Rev. B* **28**, 3673 (1983).
- [19] K. Hiebl, C. Horvath, and P. Rogl, Magnetic behaviour of ternary silicides CeT₂Si₂ (T = Ru, Rh, Pd, OS, Ir, Pt) and boron substitution in Ce{Ru, Os}₂Si_{2-x}B_x, *J. Less Common Met.* **117**, 375 (1986).
- [20] M. Mihalik, M. Mihalik, and V. Sechovsky, Electrical transport and magnetism in CeFe₂Si₂ single crystal, *Physica B* **359**, 163 (2005).
- [21] H. Abe, K. Yoshii, and H. Kitazawa, Complex magnetic phase diagram of CeRu₂Ge₂, *Physica B* **312**, 253 (2002).
- [22] H. H. Lai, S. E. Grefe, S. Paschen, and Q. Si, Weyl-Kondo semimetal in heavy-fermion systems, *Proc. Natl. Acad. Sci. USA* **115**, 93 (2017).
- [23] S. Süllow, G. J. Nieuwenhuys, A. A. Menovsky, J. A. Mydosh, S. A. M. Mentink, T. E. Mason, and W. J. L. Buyers, Spin Glass Behavior in URh₂Ge₂, *Phys. Rev. Lett.* **78**, 354 (1997).
- [24] H. Ptasiwicz-Bak, Neutron diffraction study of magnetic ordering in UPd₂Si₂, UPd₂Ge₂, URh₂Si₂ and URh₂Ge₂, *J. Phys. F Met. Phys.* **11**, 1225 (1981).
- [25] M. Kuznietz, H. Pinto, H. Ettegui, and M. Melamud, Neutron-diffraction study of the magnetic structure of UCo₂Ge₂, *Phys. Rev. B* **40**, 7328 (1989).
- [26] T. Endstra, G. J. Nieuwenhuys, A. A. Menovsky, and J. A. Mydosh, Structural and magnetic properties of UCo₂Ge₂, *J. Appl. Phys.* **69**, 4816 (1991).
- [27] A. Gallagher, K. W. Chen, C. M. Moir, S. K. Cary, F. Kametani, N. Kikugawa, D. Graf, T. E. Albrecht-Schmitt, S. C. Riggs, A. Shekhter, and R. E. Baumbach, Unfolding the physics of URu₂Si₂ through silicon to phosphorus substitution, *Nat. Commun.* **7**, 10712 (2016).
- [28] Y. Dalichaouch, M. B. Maple, J. W. Chen, T. Kohara, C. Rossel, M. S. Torikachvili, and A. L. Giorgi, Effect of transition-metal substitutions on competing electronic transitions in the heavy-electron compound URu₂Si₂, *Phys. Rev. B* **41**, 1829 (1990).
- [29] J. Lee, K. Prokes, S. Park, I. Zaliznyak, S. Dissanayake, M. Matsuda, M. Frontzek, S. Stoupin, G. L. Chappell, R. E. Baumbach, C. Park, J. A. Mydosh, G. E. Granroth, and J. P. C. Ruff, Charge density wave with anomalous temperature dependence in UPt₂Si₂, *Phys. Rev. B* **102**, 041112 (2020).
- [30] G. J. Nieuwenhuys, Crystalline electric field effects in UPt₂Si₂ and URu₂Si₂, *Phys. Rev. B* **35**, 5260 (1987).
- [31] D. S. Grachtrup, M. Bleckmann, B. Willenberg, S. Süllow, M. Bartkowiak, Y. Skourski, H. Rakoto, I. Sheikin, and J. A. Mydosh, Field-induced phases in UPt₂Si₂, *Phys. Rev. B* **85**, 054410 (2012).
- [32] S. Elgazzar, J. Rusz, P. M. Oppeneer, and J. A. Mydosh, Electronic structure and Fermi surface of paramagnetic and antiferromagnetic UPt₂Si₂, *Phys. Rev. B* **86**, 075104 (2012).
- [33] D. S. Grachtrup, N. Steinki, S. Süllow, Z. Cakir, G. Zwacknagl, Y. Krupko, I. Sheikin, M. Jaime, and J. A. Mydosh, Magnetic phase diagram and electronic structure of UPt₂Si₂ at high magnetic fields: A possible field-induced Lifshitz transition, *Phys. Rev. B* **95**, 134422 (2017).
- [34] J. Lee, M. Matsuda, J. A. Mydosh, I. Zaliznyak, A. I. Kolesnikov, S. Süllow, J. P. C. Ruff, and G. E. Granroth, Dual Nature of Magnetism in a Uranium Heavy-Fermion System, *Phys. Rev. Lett.* **121**, 057201 (2018).
- [35] S. Süllow, A. Otop, A. Loose, J. Klenke, O. Prokhnenko, R. Feyerherm, R. W. A. Hendrikx, J. A. Mydosh, and H. Amitsuka, Electronic localization and two-dimensional metallic state in UPt₂Si₂, *J. Phys. Soc. Jpn.* **77**, 024708 (2008).

- [36] R. A. Steeman, E. Frikkee, S. A. M. Mentink, A. A. Menovsky, G. J. Nieuwenhuys, and J. A. Mydosh, Hybridisation effects in UPt_2Si_2 , *J. Phys. Condens. Matter* **2**, 4059 (1990).
- [37] C. L. Farrow, P. Juhás, J. W. Liu, D. Bryndin, E. S. Božin, J. Bloch, Th. Proffen, and S. J. L. Billinge, PDFfit2 and PDFgui: Computer programs for studying nanostructure in crystals, *J. Phys. Condens. Matter* **19**, 335219 (2007).
- [38] J. Rodriguez-Carvajal, Recent advances in magnetic structure determination by neutron powder diffraction, *Physica B* **192**, 55 (1993).
- [39] T. Egami and S. J. L. Billinge, *Underneath the Bragg Peaks: Structural Analysis of Complex Materials* (Pergamon, Amsterdam, 2003).
- [40] V. Petkov, in *Characterization of Materials*, edited by E. N. Kaufmann (Wiley, Hoboken, NJ, 2012).
- [41] M. Bleckmann, A. Otop, S. Sullow, R. Feyerherm, J. Klenke, A. Loose, R. W. A. Hendrikx, J. A. Mydosh, and H. Amitsuka, Structural properties, magnetic order and electronic transport in single crystalline UPt_2Si_2 , *J. Magn. Magn. Mater.* **322**, 2447 (2010).
- [42] A. Otop, F. J. Litterst, R. W. A. Hendrikx, J. A. Mydosh, and S. Sullow, Magnetic irreversibility in single crystalline UPt_2Si_2 , *J. Appl. Phys.* **95**, 6702 (2004).
- [43] H. J. Kim, C. D. Malliakas, A. T. Tomic, S. H. Tessmer, M. G. Kanatzidis, and S. J. L. Billinge, Local Atomic Structure and Discommensurations in the Charge Density Wave of CeTe_3 , *Phys. Rev. Lett.* **96**, 226401 (2006).