



Universiteit  
Leiden  
The Netherlands

## **Patient perspectives on indwelling urinary catheters and fluid balances after transsphenoidal pituitary surgery: a qualitative study**

Nollen, J.M.; Brunsveld-Reinders, A.H.; Peul, W.C.; Furth, W.R. van

### **Citation**

Nollen, J. M., Brunsveld-Reinders, A. H., Peul, W. C., & Furth, W. R. van. (2023). Patient perspectives on indwelling urinary catheters and fluid balances after transsphenoidal pituitary surgery: a qualitative study. *Bmj Open*, 13(3). doi:10.1136/bmjopen-2022-069598

Version: Publisher's Version

License: [Creative Commons CC BY-NC 4.0 license](https://creativecommons.org/licenses/by-nc/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3731566>

**Note:** To cite this publication please use the final published version (if applicable).

# BMJ Open Patient perspectives on indwelling urinary catheters and fluid balances after transsphenoidal pituitary surgery: a qualitative study

Jeanne-Marie Nollen <sup>1</sup>, Anja H Brunsveld-Reinders,<sup>2</sup> Wilco C Peul,<sup>1</sup> Wouter R van Furth<sup>1</sup>

**To cite:** Nollen J-M, Brunsveld-Reinders AH, Peul WC, *et al.* Patient perspectives on indwelling urinary catheters and fluid balances after transsphenoidal pituitary surgery: a qualitative study. *BMJ Open* 2023;**13**:e069598. doi:10.1136/bmjopen-2022-069598

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-069598>).

Received 28 October 2022  
Accepted 01 March 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Neurosurgery, Leiden University Medical Center, Leiden, Netherlands

<sup>2</sup>Department of Quality and Patient Safety, Leiden University Medical Center, Leiden, Netherlands

**Correspondence to**  
Jeanne-Marie Nollen;  
J.M.Nollen@lumc.nl

## ABSTRACT

**Objectives** To explore the perceptions and experiences of patients who underwent transsphenoidal pituitary gland and (para)sellar tumour surgery regarding indwelling urinary catheters (IDUCs) and the postoperative fluid balance.

**Design** Qualitative study using semistructured interviews based on the attitudes, social influence and self-efficacy model and expert knowledge.

**Participants** Twelve patients who underwent transsphenoidal pituitary gland tumour surgery and received an IDUC during or after surgery.

**Setting** One patient was interviewed in the endocrinology outpatient clinic and 11 patients were interviewed on the neurosurgery ward.

**Results** Five major themes emerged: (1) conflicting information and preoperative expectations, (2) IDUCs perceived as patient-friendly during bedrest, particularly for women, (3) little room for patients' opinions, (4) physical and emotional limitations and (5) fluid balance causes confusion. Information regarding IDUC placement and fluid balance given to patients both preoperatively and postoperatively did not meet their expectations, which led to confusion and uncertainty. The IDUC was perceived as preferable if bedrest was mandatory, preferred particularly by women. Patient could not mobilise freely due to the IDUC and felt ashamed, judged by others and dependent on nurses.

**Conclusions** This study provides insight into the challenges patients experience in relation to the IDUC and fluid balance. Perceptions on the necessity of an IDUC varied among patients and were influenced by both physical and emotional impediments. A clear, frequent and daily communication between healthcare professionals and patients to evaluate IDUC and fluid balance use is necessary to increase patient satisfaction.

## INTRODUCTION

To evaluate hospital care and the corresponding processes, patients' perspectives play a crucial role as they offer information that goes beyond the scope of regular hospital staff evaluations.<sup>1</sup>

Two frequently studied topics to gain insight in hospital care during the

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This qualitative study provides a broader understanding of challenges related to indwelling urinary catheters and fluid balance in relation to bed rest and diabetes insipidus.
- ⇒ Semistructured interviews were used to systematically explore set topics while allowing flexibility to explore participants' thoughts, feelings and beliefs.
- ⇒ Two researchers listened to the interviews and individually coded the papers, before discussing the results.
- ⇒ This study sample consisted of more female participants than male participants.

postoperative phase are indwelling urinary catheters (IDUCs) and fluid balances. While studies investigating fluid balances have focused primarily on accuracy and diagnostic value in critical care settings rather than focus on patient perspectives, patient experiences with and perceptions of IDUCs in the postoperative phase have been widely researched.<sup>2-4</sup> Patients have connected IDUCs with both infectious, including urinary tract infections (UTIs), and non-infectious problems, such as pain and discomfort.<sup>5</sup> These studies focused on general surgical populations despite literature indicating that patients' perspectives could be influenced by their specific illness and operation and that research should therefore keep the individual needs and specific situations in mind.<sup>6</sup>

One group of patients who are a useful source of information about IDUC and fluid balance experiences are patients who underwent transsphenoidal pituitary gland and (para)sellar tumour surgery. In the university hospital, IDUCs are not routinely placed during this surgery due to the relatively short operation time of 2–3 hours.<sup>7</sup> Despite this policy, IDUCs are frequently inserted postoperatively at the neurosurgical ward.

Two potential postoperative complications influence IDUC placement and the necessity of monitoring the fluid balance in this specific population. First, pituitary patients are at risk of developing the electrolyte disorder diabetes insipidus (DI).<sup>8</sup> Accurate monitoring of the fluid balance, every 3–6 hours postoperatively, is essential for the early detection of DI as well as the consideration of desmopressin therapy, which is the primary pharmacological treatment.<sup>9</sup> On the ward, nurses measure the urine volume in a urinal and patients use a personal fluid balance chart to register the fluid intake. As DI can occur in the 10 days following surgery, the fluid balance needs to be monitored after discharge.<sup>10</sup> IDUCs can contribute to a reliable fluid balance and are convenient for nurses when monitoring the urinary output.<sup>11 12</sup>

Second, to prevent post-transsphenoidal cerebrospinal fluid leakage, bed rest, with elevation of the head of bed at 30° for 24-hours, is a frequently occurring postoperative instruction which could influence the patient's ability to urinate.<sup>13 14</sup> Bed rest is identified as a risk factor for a retention bladder, which is defined as the inability to urinate despite a full bladder (>500 mL) and can lead to complications including UTIs and stretched bladder muscles.<sup>15 16</sup> If a patient develops a retention bladder, IDUCs are the primary intervention.<sup>17</sup>

Previous studies have explored pre-surgery and post-surgery symptom burden of DI and established the need for support before, during and after hospital admission.<sup>18 19</sup> However, to the best of our knowledge, patient perspectives regarding IDUCs and monitoring the fluid balance have not been studied in this specific patient population and setting despite having a major impact during the acute postoperative phase. Consequently, this study aims to explore the perspectives and experiences of patients who underwent transsphenoidal pituitary gland and (para)sellar tumour surgery regarding IDUCs and fluid balances on a neurosurgical ward.

## METHODS

### Study design

A qualitative study design was adopted which involved semistructured interviews with patients who underwent transsphenoidal pituitary gland and (para)sellar tumour surgery to explore their perceptions and experiences regarding IDUCs and the postoperative fluid balance.

### Setting and participants

The study was conducted in a 16-bed department of neurosurgery at a University Hospital in the Netherlands. Participants who underwent transsphenoidal pituitary gland and (para)sellar tumour surgery received an IDUC in the perioperative or postoperative period, and were aged >18 were approached face-to-face if they were admitted to the neurosurgical ward or by phone if they were discharged. Convenience sampling was used to approach 13 patients, 12 of which agreed to participate and 1 declined due to personal reasons. One patient was

## Box 1 Interview topics

1. How did patients experience the postoperative care on the neurosurgical ward?
  - Nursing care
  - Communication
  - Complications
  - Preoperative consultation in outpatient clinic
  - Experience with IDUC
  - Experience with fluid balance
2. How and to what extent was the patient involved in the decision to insert and remove the urinary catheter?
  - Preoperative information
  - Shared decision making
  - Role nurse/physician
  - Influence bedrest
  - Postoperative complications
3. How did patients experience the moment of IDUC insertion and removal?
  - Comfort
  - Physical situation
  - Time of day
  - Shared decision making
  - Nurse's role
  - Complications after removal
  - Fluid balance before and after removal
4. What was the patient's role in monitoring the fluid balance?
  - Bedpan/urinal
  - IDUC
  - Fluid balance chart
  - Patient participation
  - Collaboration with nurses
5. How did the IDUC affect mobilisation and interaction with caregivers/family members?
  - Stigma and feelings
  - Barriers

interviewed in the endocrinology outpatient clinic and 11 patients were interviewed on the neurosurgery ward. Data saturation was reached after 12 interviews which means that it is likely that no new information will arise during additional interviews.<sup>20</sup>

### Data collection

A semistructured interview guide was developed based on the attitudes, social influence and self-efficacy model and expert knowledge (Box 1). This model was deliberately chosen as it helps to elaborate on demonstrated health behaviours and accompanying motives.<sup>21</sup> Interviews were performed in Dutch.

Two pilot interviews were conducted. The topic list was adjusted two times based on the feedback of one test-participant and two participants who experienced difficulties explaining their role regarding IDUC removal. The audio-recorded interviews were held in a 3-month period, from mid-September until mid-November 2019, in a place and time that suited the participant. An oral summary was presented to each participant at the end of the interview to verify their story. Interviews were

conducted by an experienced neurosurgical nurse who was not involved in the care of the participating patients.

### Data analysis

The interviews were transcribed verbatim and analysed through thematic analysis.<sup>22</sup> Two researchers independently conducted the coding process and discussed the findings with one another. Transcripts were read and reread to become familiar with the data. During the first phase of coding, the data were segmented into meaningful parts. These parts were provided with summarising labels (codes). Subsequently, the codes were compared within and between transcripts by two researchers resulting in categories of codes on a more conceptual level. Finally, the created categories were described into themes. An iterative approach was adopted to enable continuous evaluation of the data.<sup>23</sup> The software programme Atlas.ti 8.4.15 was used to structure the process of data analysis.<sup>24</sup> Analysis was performed in Dutch and quotations were translated into English by a native speaker.

### Ethical considerations

All study procedures were in accordance with the declaration of Helsinki and the medical ethics committee of Academic Hospital approved the study protocol (N19.015).<sup>25</sup> Participants received an information sheet and an informed consent form prior to the interviews. All participants provided written informed consent. Furthermore, participants were asked for their permission to record the interview with a voice recorder.

### Patient and public involvement

The research question was developed by the researchers through their experience with the care for pituitary patients. Patients were not involved in the design and conduct of the study, the choice of outcome measures and recruitment for the study. Patients agreed with plans for dissemination of the results through scientific publication and education for nurses on the University hospital ward.

## RESULTS

### Patient characteristics

The sample included 12 patients (table 1) of which 83% (n=10) were women. The mean age of the participants was 55 years (range: 39–73 years). Four patients had an IDUC inserted during the operation. Eight patients had an IDUC inserted postoperatively on the ward as they developed a retention bladder. One patient who received an IDUC during the operation developed a retention bladder after IDUC removal which required re-catheterisation. The interviews had a duration of 23–58 min.

### Themes

Five major themes emerged: (1) conflicting information and preoperative expectations, (2) IDUCs perceived as patient-friendly during bedrest, particularly for women, (3) little room for patients' opinions, (4) physical and

**Table 1** Characteristics of study population (n=12)

	n (%)
Gender	
Male	10 (83)
Female	2 (17)
IDUC inserted during surgery	4 (33)
IDUC inserted on ward	9 (75)
Retention bladder	9 (75)
Bedrest	7 (58)
Diabetes insipidus	5 (42)
Cerebrospinal fluid leakage	1 (8)
	<b>Mean (min–max)</b>
Age	55 (39–73)
Length of hospital stay	4 (3–8)
Days IDUC inserted	2 (1–7)

IDUC, indwelling urinary catheter.

emotional limitations and (5) fluid balance causes confusion. Quotations are included to illustrate the text.

### Theme 1: conflicting information and preoperative expectations

During the preoperative consult, five patients were informed that they would not receive an IDUC during the operation, whereas the information booklet stated the opposite. Three patients stated that they did not discuss the IDUC during the consult and did not read the booklet prior to surgery, so therefore they were unaware of the possibility of an IDUC. Three participants expressed feeling indifferent towards receiving an IDUC as they trusted the medical staff to make the appropriate decision.

All participants received information during the preoperative consult on how to monitor the fluid balance after discharge; however, information on how to monitor the fluid balance during the hospital admission was provided to only four participants. Postoperatively, patients reported a large variation between nurses and their willingness to explain the fluid balance and having the patient monitor their input.

Two participants had undergone pituitary surgery in the past and were expecting to receive an IDUC based on their previous experiences. One participant was not content when she found out after her operation that she did not have an IDUC: 'I missed my IDUC. Because I had no discomfort from the IDUC the first time but I found it so dehumanising to urinate on the bedpan, especially because I was unable to empty my bladder and needed an IDUC because of that. In the end, there were four towels under me and I was completely covered in urine'.

The participants' preoperative attitudes toward IDUCs leaned towards the negative and were predominantly influenced by stigmas and stories told by their friends and families. One participant explained: 'I was so scared of

receiving and IDUC because I heard experiences from friends who had it (an IDUC) before and they said it hurts so badly to insert and remove it. So, after I heard all their terrible stories I thought no way I want an IDUC'. Another patient added: 'It is what we were taught by our parents in the old days. People were very dramatic about IDUCs; for me it is still a very sensitive subject. I was shocked when I found out I probably was getting one but there are more people in the hospital with one, I know that. But I have this image in my head of an elderly person in a wheelchair and then carrying around that bag... it makes you look so ill'.

#### Theme 2: IDUCs perceived as patient-friendly during bedrest, particularly for women

Eight female participants described their positive experiences with the IDUC in combination with postoperative bedrest. The general opinion was that providing a patient with an IDUC is more patient-friendly compared with having to use the bedpan. Ten out of the 12 participants felt that once the postoperative restriction mobility had ended, the IDUC had lost its added value.

Several complications associated with the bedpan were described. First, patients experienced a lack of privacy: 'In my room, one other patient was waiting for his operation, another person was waiting for his wife to come back from surgery. I'm sorry but I cannot urinate comfortably with others in the room. I couldn't urinate on the bedpan and I couldn't sit up straight in bed because I was on bedrest. The placement of the IDUC was an issue because they needed around six or seven attempts. It took almost 40 min before the IDUC was placed. Very painful and embarrassing for me. But when the IDUC was finally placed it was such a relief'. Second, using the bedpan was perceived as unsanitary: 'I had to urinate after the surgery but it was very difficult on the bedpan. I was so afraid that the urine would touch me or that I would wet my bed. It was so stressful and disgusting'. Third, participants felt dependent on nurses' schedules resulting in patients developing a retention bladder or having to try to control their bladder. Finally, bedpans were associated with physical discomfort.

Participants explained that the IDUC was generally promptly removed by a nurse once the mobility restriction had ended, which was usually around noon. Postponed removal was caused by nurses being too busy or the nurse's wish that the physiotherapist mobilised the patient beforehand. Postponed removal, at 06:00, made a strong impression on the patients: 'I was sleeping and it was very early in the morning and then she (the nurse) made a lot of noise, put all the lights on, pulled the IDUC out and that was it. While I was barely awake so I found that very uncomfortable'.

#### Theme 3: little room for patients' opinions

Patients had different perspectives on their role in the decision to insert or remove the IDUC. The four patients that had an IDUC inserted during surgery felt that they

were adequately informed sufficiently during the outpatient clinic consult. If an IDUC was required postoperatively, patients felt that nurses did not inform them adequately about their options and did not take their opinion into consideration.

The eight patients who did not receive an IDUC during the operation felt pressured by nurses to urinate promptly after their return to the ward, which generated stress and anxiety: 'I just woke up after the surgery and then they (nurses) checked how much fluid there was in my bladder and they said that it was too much. I had 1.2 liters of urine in my bladder and then I had 5 min to urinate, but I was still groggy from the surgery. After time was up they inserted an IDUC. It all went so fast. I just wished they had inserted the IDUC during the surgery' and 'I didn't really have an idea of what it would be like to have an IDUC. I never had one before and then all of a sudden they inserted one but they (nurses) didn't explain how they were going to do that, so that was very shocking to me. When I asked what was going to happen they explained a little bit but only after I asked for it. I just wish they told me earlier'. These eight patients wished they were involved more in the shared decision-making process.

#### Theme 4: physical and emotional limitations

The majority of the participants felt that an IDUC hinders mobilisation and reduces the need to be active since it makes mobilisation, especially to the bathroom, mostly redundant. One patient explained: 'All the hassle walking with the IDUC bag, I mean where do you put that thing. It limits my mobility so much. It really bothers me'. The increased strain on the tube when walking or turning over in bed led to discomfort and caused two patients to be scared that the IDUC might be disconnected and leak urine. Being dependent on nurses was also mentioned as a barrier to mobilise: 'I barely left my bed because then the nurse needed to help me and attach the IDUC to something. I didn't want to bother them (nurses) too much because they were so busy all the time'.

Reduced mobility was not experienced as bothersome by all participants: 'You feel it (the IDUC) pull and then you are afraid that it breaks so you have to be a bit careful, you cannot toss and turn in the bed. But lying still was no problem for me, I liked it'. A few participants felt uninformed by nurses and were left with questions about the postoperative mobilisation policy. One participant illustrated: 'I was happy lying in the bed but if no one says that you can walk you will stay in bed just because you don't know if you are even allowed to walk with an IDUC'.

Shame and fear of being judged for having an IDUC by nurses, other patients and visitors resulted in six participants to refrain from mobilising to areas outside their room and by trying to cover the IDUC: 'I think it is embarrassing to walk around with an IDUC. That's why I tried to cover up the bag with a cardigan or large trousers. I know I should not worry about that but I found the IDUC so distasteful to see'.

Since an IDUC is a foreign material, six patients who received an IDUC postoperatively experienced pain and discomfort when the IDUC was inserted. Patients complained of having bladder spasms, urine leaking next to the tube and feeling the need to urinate after the IDUC was inserted: 'I woke up during the night and I had a feeling of urinating but that was impossible because I had an IDUC. I found that very annoying'. After IDUC removal, three patients experienced a burning sensation when urinating which sometimes lasted for a couple of days.

Aside from physical discomfort, the interviews disclosed emotional strain caused by IDUCs. Four patients were afraid to develop a UTI as a result of the IDUC and these fears were confirmed by nurses. Before and shortly after the IDUC was removed, two patients were uncertain if their bladder could instantly regain its function and were worried that they could become incontinent. One participant explained: 'Just after the removal I was scared about what was going to happen. Did I have to run to the toilet every minute? At a certain point the IDUC gave me a feeling of peace because I didn't have to think about urinating. I was afraid that I needed to go to the bathroom six times each night and that I might be incontinent'.

#### Theme 5: fluid balance causes confusion

During hospital admission, only two participants monitored their fluid intake. The personal fluid balance chart was used simultaneously by the patient, nurses and hospital food service workers which led to confusion and deviating charts. One participant illustrated: 'I lost complete control of my input because some nurses wrote it down but other nurses didn't so it was very confusing to me. I didn't know if I was supposed to monitor my intake or not'. Participants also experienced difficulties with the fluid balance chart: 'I am always guessing how much ml is in one cup because the chart is difficult to understand. The nurses don't know either, they tell me different amounts per cup'.

Four participants voiced concerns regarding monitoring the fluid balance at home: 'the nurse monitored what was going in and out so of course I am starting to worry now that I am going home and have to do it myself. The nurses already worry if there is half a litre difference in the fluid balance and I really don't understand what all the fuss is about'. Ten patients would prefer more education on how to monitor the fluid balance as well as having the ability of guided practice.

Participants did not monitor the urinary output as they were not offered this option. Nine participants were willing to monitor their output during the hospital admission: 'I would like to monitor the output just so I know what is going on with my body. But I think it would be difficult to measure it on the day of the surgery since you are not feeling well then. but from day two on it would have been no problem for me'. Only one participant explicitly stated that she would find it disgusting to monitor the output during the hospital stay.

## DISCUSSION

The aim of this study was to explore patient perspectives regarding IDUCs and monitoring the fluid balance after transsphenoidal pituitary and (para)sellar surgery. Despite patients describing a broad range of physical and emotional limitations related to IDUCs, they were preferred under the condition of bedrest, especially by women. Our findings suggest that patients' experiences are largely influenced by the information they receive from healthcare professionals both before and during their hospital admission. Additionally, our study shows that despite patients being instructed to monitor the fluid intake, nurses take on responsibility for this task leaving the patient unprepared to monitor the fluid balance after discharge.

Most female participants were in favour of IDUC use during the period of mandatory bedrest due to negative experiences with the bedpan. Loss of privacy, dependency on nurses, embarrassment, physical discomfort and hygiene aspects, all described in previous research, contributed to patients preferring IDUCs instead of bedpans.<sup>26</sup>

This study confirms the importance of managing patients expectations and the consequences of patients receiving insufficient information.<sup>27</sup> The quality of patient information is an important factor related to patient-centred care as it contributes to increased patient participation.<sup>28 29</sup> Patients experienced negative effects including stress and confusion by receiving conflicting and too little information. Although it was not mentioned in this study by any of the participants, literature additionally reported that patients may question the competence of the healthcare professionals due to contradictory and incomplete information.<sup>30</sup>

Shared decision making was experienced as more present pre-operatively during scheduled consultations in contrast to acute situations, for example, a retention bladder, postoperatively on the ward. Patients felt pressured and overlooked by nurses. Literature acknowledges this phenomenon and states that shared decision making is influenced by the physical setting and variability of the illness and that therefore acute situations may lead to a healthcare provider-led approach.<sup>31</sup> This passive role assigned to patients postoperatively could be converted to an equal distribution of power between both parties through educational programmes for nurses and strategies (eg, decision flowcharts) that focus on increasing patients' decision-making capacity.<sup>32</sup>

This study highlights the need for patient involvement in clinical care during the hospital admission to ensure a safe transition from the hospital to the home setting. The lack of training and guidance during the postoperative period could be explained by nurses feeling hesitant to relinquish responsibility to patients as patient safety could be jeopardised.<sup>33</sup> Additionally, time constraints and the absence of a standardised educational protocol for nurses to train and educate patients could be of influence.<sup>34</sup> A practice environment where patients and their



relatives are trained to monitor both the fluid intake as well as the output to enable a gradual shift in responsibility, while still practicing in a safe and controlled setting, could strengthen patients' confidence.<sup>31 35</sup> To the best of our knowledge, no study has been conducted on such a specific educational programme.

Mobility challenges related to the IDUC, including prolonged time to ambulation (walking without the support of a nurse), immobility and discomfort, overlap with previous findings.<sup>36</sup> In this study, patients reported feeling dependent on nurses' directives which could have delayed the moment of mobilisation and thereby have a negative influence on the discharge date.<sup>37 38</sup>

We found that social influences, and stigmas could lead to embarrassment and fear of judgement from others. Although extensively described in long-term IDUC use, limited research has been conducted on the influence of social stigmas (eg, embarrassment) in hospital settings.<sup>39 40</sup>

The incidence of urinary retention in this study was 75% (9 out of 12), which does not fall in the reported incidence range of 5–70, and is significantly higher than the reported 5% in general surgical populations.<sup>15 41</sup> This high incidence could partly be explained by post-operative bedrest; however, additional influencing factors including perioperative fluids, concurrent diseases, duration of the surgery and perioperative medications were not reported since they were outside the scope of this study.<sup>42</sup> The results from this study could be different if the incidence of urinary retention, and subsequent catheterisation rate, were lower.

A major strength of this study is that a combination of patients who received an IDUC during and after the operation was interviewed. Due to this approach, a broad range of experiences and perspectives was gathered. In addition, by applying a code–recode procedure during the data analysis, the validity of the study increased.

A limitation of the study was the relatively small and specific patient population, in addition to this study being conducted in a single ward in a University hospital. However, we do feel that the results can be used for different patient groups who also require fluid balances. Additionally, the results provide information that could be used by others to obtain insight into the patient perspective and complicated dilemmas patients face during hospital admission. Second, interviews were conducted both on the ward and in the outpatient clinic. It could be possible that perspectives from the patient who was interviewed several days after discharge changed due to having time to reflect on their hospital admission.

Further research is necessary to assess the possibilities of patient involvement in monitoring the fluid balance during hospital admission. Furthermore, a nurse-led training programme should be developed and implemented on the ward to increase patient participation and build patients' confidence.

## CONCLUSION

IDUC placement and fluid balance measurements are important aspects of perioperative patient care after transsphenoidal pituitary gland and (para)sellar tumour surgery and have a major impact on the patient's overall evaluation. Patients who receive an IDUC during or after pituitary surgery experience a broad range of complications and are faced with a multitude of challenges related to communication and participation in care. In addition, insufficient information, predominantly provided by nurses, has a large impact on patient experiences and comprehension of the provided care. Patient involvement in both clinical care as well as shared decision making could be improved. Implementing an inpatient training programme to increase patient participation in clinical care is likely to be beneficial for the transition from the hospital to the home setting.

**Contributors** JMN contributed to conceptualisation, methodology, formal analysis, investigation, writing—original draft. AHB-R contributed to conceptualisation, methodology, formal analysis, writing: review & editing, supervision. WCP contributed to conceptualisation, writing: review & editing, supervision. WRvF contributed to conceptualisation, writing: review & editing, supervision and is the guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Consent obtained directly from patient(s).

**Ethics approval** This study involves human participants and was approved by Medisch-Ethische Toetsingscommissie Leiden, Den Haag, Delft (N19.015). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Data are available upon request.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iD

Jeanne-Marie Nollen <http://orcid.org/0000-0002-1241-5380>

## REFERENCES

- 1 Kleefstra S. *Hearing the patient's voice. The patient's perspective as outcome measure in monitoring the quality of hospital care.* 2016.
- 2 Safdar N, Codispoti N, Purvis S, *et al.* Patient perspectives on indwelling urinary catheter use in the hospital. *Am J Infect Control* 2016;44:e23–4.
- 3 Bhardwaj R, Pickard R, Carrick-Sen D, *et al.* Patients' perspectives on timing of urinary catheter removal after surgery. *Br J Nursing* 2012;21:S4–9.
- 4 Asfour HI. Fluid balance monitoring accuracy in intensive care units. *IOSR* 2016;05:53–62.
- 5 Saint S, Trautner BW, Fowler KE, *et al.* A multicenter study of patient-reported infectious and noninfectious complications associated with indwelling urethral catheters. *JAMA Intern Med* 2018;178:1078–85.
- 6 Sacristán JA, Aguarón A, Avendaño-Solá C, *et al.* Patient involvement in clinical research: why, when, and how. *Patient Prefer Adherence* 2016;10:631–40.

- 7 Zada G, Woodmansee WW, Iuliano S, *et al.* Perioperative management of patients undergoing transsphenoidal pituitary surgery. *Asian J Neurosurg* 2010;5:1–6.
- 8 Nayak P, Montaser AS, Hu J, *et al.* Predictors of postoperative diabetes insipidus following endoscopic resection of pituitary adenomas. *J Endocr Soc* 2018;2:1010–9.
- 9 Almalki MH, Ahmad MM, Brema I, *et al.* Management of diabetes insipidus following surgery for pituitary and suprasellar tumours. *Sultan Qaboos Univ Med J* 2021;21:354–64.
- 10 Lamas C, del Pozo C, Villabona C, *et al.* Clinical guidelines for management of diabetes insipidus and syndrome of inappropriate antidiuretic hormone secretion after pituitary surgery. *Endocrinol Nutr* 2014;61:e15–24.
- 11 Capatina C, Paluzzi A, Mitchell R, *et al.* Diabetes insipidus after traumatic brain injury. *J Clin Med* 2015;4:1448–62.
- 12 Schreckinger M, Szerlip N, Mittal S. Diabetes insipidus following resection of pituitary tumors. *Clin Neurol Neurosurg* 2013;115:121–6.
- 13 Elgamal E. CSF rhinorrhoea after transsphenoidal surgery. *Internet J Neurosurg* 2008;5 (1).
- 14 Prete A, Corsello SM, Salvatori R. Current best practice in the management of patients after pituitary surgery. *Ther Adv Endocrinol Metab* 2017;8:33–48.
- 15 Agrawal K, Majhi S, Garg R. Post-Operative urinary retention: review of literature. *WJA* 2019;8:1–12.
- 16 Alsaidi M, Guanjo J, Basheer A, *et al.* The incidence and risk factors for postoperative urinary retention in neurosurgical patients. *Surg Neurol Int* 2013;4:61.
- 17 Jackson J, Davies P, Leggett N, *et al.* Systematic review of interventions for the prevention and treatment of postoperative urinary retention. *BJS Open* 2019;3:11–23.
- 18 Waddle MR, Oudenhoven MD, Farin CV, *et al.* Impacts of surgery on symptom burden and quality of life in pituitary tumor patients in the subacute post-operative period. *Front Oncol* 2019;9:299.
- 19 Jakobsson Ung E, Olofsson A-C, Björkman I, *et al.* The pre- and postoperative illness trajectory in patients with pituitary tumours. *Endocr Connect* 2019;8:878–86.
- 20 Fusch PI, LRJTqr N. Are we there yet. *Data Saturation in Qualitative Research* 2015;20:1408.
- 21 Sheeran P, Maki A, Montanaro E, *et al.* The impact of changing attitudes, norms, and self-efficacy on health-related intentions and behavior: a meta-analysis. *Health Psychol* 2016;35:1178–88.
- 22 Cooper H, Camic PM, Long DL, *et al.* *Thematic analysis*. Washington: American Psychological Association, 2012.
- 23 Vaismoradi M, Jones J, Turunen H, *et al.* Theme development in qualitative content analysis and thematic analysis. *JNEP* 2016;6.
- 24 Friese S. *Qualitative data analysis with ATLAS. ti*: SAGE Publications Limited, 2019.
- 25 World Medical Association. World Medical association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013;310:2191–4.
- 26 Darbyshire D, Rowbotham D, Grayson S, *et al.* Surveying patients about their experience with a urinary catheter. *Int J Urol Nurs* 2016;10:14–20.
- 27 Clarke MA, Moore JL, Steege LM, *et al.* Health information needs, sources, and barriers of primary care patients to achieve patient-centered care: a literature review. *Health Informatics J* 2016;22:992–1016.
- 28 Scholl I, Zill JM, Härter M, *et al.* An integrative model of patient-centeredness—a systematic review and concept analysis. *PLoS One* 2014;9:e107828.
- 29 Vahdat S, Hamzehgardeshi L, Hessam S, *et al.* Patient involvement in health care decision making: a review. *Iran Red Crescent Med J* 2014;16:e12454.
- 30 Carpenter DM, Geryk LL, Chen AT, *et al.* Conflicting health information: a critical research need. *Health Expect* 2016;19:1173–82.
- 31 Truglio-Londrigan M, Slyer JT. Shared decision-making for nursing practice: an integrative review. *Open Nurs J* 2018;12:1–14.
- 32 Molina-Mula J, Gallo-Estrada J. Impact of nurse-patient relationship on quality of care and patient autonomy in decision-making. *Int J Environ Res Public Health* 2020;17:835.
- 33 Vaismoradi M, Jordan S, Kangasniemi M. Patient participation in patient safety and nursing input—a systematic review. *J Clin Nurs* 2015;24:627–39.
- 34 Oxelmark L, Ulin K, Chaboyer W, *et al.* Registered nurses' experiences of patient participation in hospital care: supporting and hindering factors patient participation in care. *Scand J Caring Sci* 2018;32:612–21.
- 35 Bombard Y, Baker GR, Orlando E, *et al.* Engaging patients to improve quality of care: a systematic review. *Implement Sci* 2018;13:98.
- 36 Mukantwari J, Omondi L, Ndateba I, *et al.* Indwelling urinary catheter use and associated outcomes among adult patients undergoing surgery. *Rwanda Journal of Medicine and Health Sciences* 2019;2:34.
- 37 Rojas-García A, Turner S, Pizzo E, *et al.* Impact and experiences of delayed discharge: a mixed-studies systematic review. *Health Expect* 2018;21:41–56.
- 38 Nollen JM, Pijnappel L, Schoones JW, *et al.* Impact of early postoperative indwelling urinary catheter removal: a systematic review. *J Clin Nurs* 8, 2022.
- 39 Wilde MH. Life with an indwelling urinary catheter: the dialectic of stigma and acceptance. *Qual Health Res* 2003;13:1189–204.
- 40 Oswald F, Young E, Denison F, *et al.* Staff and patient perceptions of a community urinary catheter service. *Int J Urol Nurs* 2020;14:83–91.
- 41 Dreijer B, Møller MH, Bartholdy J. Post-Operative urinary retention in a general surgical population. *Eur J Anaesthesiol* 2011;28:190–4.
- 42 Baldini G, Bagry H, Aprikian A, *et al.* Postoperative urinary retention: anesthetic and perioperative considerations. *Anesthesiology* 2009;110:1139–57.