



Universiteit
Leiden
The Netherlands

Feasibility to estimate mean systemic filling pressure with inspiratory holds at the bedside

Wijnberge, M.; Jansen, J.R.C.; Pinsky, M.R.; Klanderman, R.B.; Terwindt, L.E.; Bosboom, J.J.; ... ; Geerts, B.F.

Citation

Wijnberge, M., Jansen, J. R. C., Pinsky, M. R., Klanderman, R. B., Terwindt, L. E., Bosboom, J. J., ... Geerts, B. F. (2022). Feasibility to estimate mean systemic filling pressure with inspiratory holds at the bedside. *Frontiers In Physiology*, 13.
doi:10.3389/fphys.2022.1041730

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3736233>

Note: To cite this publication please use the final published version (if applicable).



OPEN ACCESS

EDITED BY
Peter Rasmussen,
University of Zurich, Switzerland

REVIEWED BY
Huaiwu He,
Peking Union Medical College Hospital
(CAMS), China
Songqiao Liu,
Southeast University, China

*CORRESPONDENCE
Marije Wijnberge,
m.wijnberge@amsterdamumc.nl
Alexander P. Vlaar,
a.p.vlaar@amsterdamumc.nl

SPECIALTY SECTION
This article was submitted to Clinical
and Translational Physiology,
a section of the journal
Frontiers in Physiology

RECEIVED 11 September 2022
ACCEPTED 25 October 2022
PUBLISHED 29 November 2022

CITATION
Wijnberge M, Jansen JRC, Pinsky MR,
Klanderman RB, Terwindt LE,
Bosboom JJ, Lemmers N, Vlaar AP,
Veelo DP and Geerts BF (2022),
Feasibility to estimate mean systemic
filling pressure with inspiratory holds at
the bedside.
Front. Physiol. 13:1041730.
doi: 10.3389/fphys.2022.1041730

COPYRIGHT
© 2022 Wijnberge, Jansen, Pinsky,
Klanderman, Terwindt, Bosboom,
Lemmers, Vlaar, Veelo and Geerts. This
is an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](#).
The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Feasibility to estimate mean systemic filling pressure with inspiratory holds at the bedside

Marije Wijnberge^{1,2*}, Jos R. C. Jansen³, Michael R. Pinsky⁴,
Robert B. Klanderman^{1,2}, Lotte E. Terwindt¹,
Joachim J. Bosboom^{1,2}, Nikki Lemmers¹, Alexander P. Vlaar^{2*},
Denise P. Veelo¹ and Bart F. Geerts⁵

¹Amsterdam UMC Location Academic Medical Center, Department of Anesthesiology, Amsterdam, Netherlands, ²Amsterdam UMC Location Academic Medical Center, Department of Intensive Care Medicine, Amsterdam, Netherlands, ³Leiden University Medical Center, Department of Intensive Care Medicine, Leiden, Netherlands, ⁴Department of Critical Care Medicine, University of Pittsburgh, Pittsburgh, PA, United States, ⁵Healthplus.ai, Amsterdam, Netherlands

Background: A decade ago, it became possible to derive mean systemic filling pressure (MSFP) at the bedside using the inspiratory hold maneuver. MSFP has the potential to help guide hemodynamic care, but the estimation is not yet implemented in common clinical practice. In this study, we assessed the ability of MSFP, vascular compliance (C_{sys}), and stressed volume (V_s) to track fluid boluses. Second, we assessed the feasibility of implementation of MSFP in the intensive care unit (ICU). Exploratory, a potential difference in MSFP response between colloids and crystalloids was assessed.

Methods: This was a prospective cohort study in adult patients admitted to the ICU after cardiac surgery. The MSFP was determined using 3–4 inspiratory holds with incremental pressures (maximum 35 cm H₂O) to construct a venous return curve. Two fluid boluses were administered: 100 and 500 ml, enabling to calculate V_s and C_{sys}. Patients were randomized to crystalloid or colloid fluid administration. Trained ICU consultants acted as study supervisors, and protocol deviations were recorded.

Results: A total of 20 patients completed the trial. MSFP was able to track the 500 ml bolus ($p < 0.001$). In 16 patients (80%), V_s and C_{sys} could be determined. V_s had a median of 2029 ml (IQR 1605–3164), and C_{sys} had a median of 73 ml mmHg⁻¹ (IQR 56–133). A difference in response between crystalloids and colloids was present for the 100 ml fluid bolus ($p = 0.019$) and in a post hoc analysis, also for the 500 ml bolus ($p = 0.010$).

Conclusion: MSFP can be measured at the bedside and provides insights into the hemodynamic status of a patient that are currently missing. The clinical feasibility of V_s and C_{sys} was judged ambiguously based on the lack of required hemodynamic stability. Future studies should address the clinical obstacles found in this study, and less-invasive alternatives to determine MSFP should be further explored.

Clinical Trial Registration: [ClinicalTrials.gov](#) Identifier NCT03139929.

KEYWORDS

hemodynamics, venous return, mean circulatory filling pressure, physiology, volume status, Guytonian model, vascular compliance

Introduction

At present, a decade after it became possible to estimate mean systemic filling pressure (MSFP) at the bedside, the parameter has not yet been implemented in (routine) clinical care. MSFP is considered the combined upstream pressure that drives blood flow into the right atrium, and MSFP allows the calculation of additional hemodynamic parameters such as the driving pressure for venous return (VRdp), stressed volume (Vs), and total systemic vascular compliance (Csys) (Maas et al., 2012a). Vs provides information on the effective circulating volume, a hemodynamic variable that is missing in current clinical practice. MSFP has helped to better understand the effects of vasopressors, propofol, and hyperoxia (Maas et al., 2013; de Wit et al., 2016; Helmerhorst et al., 2017; Adda et al., 2021). MSFP and the derived parameters could potentially be beneficial to guide hemodynamic care in patients admitted to the intensive care unit (ICU) (Rothe, 1993; Vos et al., 2020; Persichini et al., 2022).

MSFP can be determined in sedated and ventilated patients by extrapolating central venous pressure (CVP) versus cardiac output (CO) at different ventilatory plateau pressures during inspiratory holds (Wijnberge et al., 2018). Previous studies showed MSFP to predict fluid loading responsiveness (Guerin et al., 2015; Cooke et al., 2018). Although MSFP sounds promising, studies describing clinical guidance based on MSFP and the derived parameters are lacking (Vincent and Pinsky, 2018). Also, in previous MSFP studies (Keller et al., 2011; Maas et al., 2012b; Maas et al., 2012c), colloids were used, limiting the clinical translatability of results and feasibility, since in ICU patients, crystalloids are the preferred choice of fluids (Maas et al., 2012a).

In this study, our first aim was to assess the ability of MSFP, Csys, and Vs to track two fluid boluses. Our second aim was to assess the feasibility of the clinical implementation of MSFP in the ICU. In exploratory, as a third aim, a potential difference in response between colloids and crystalloids was assessed. As the intravascular half-life for crystalloids is around 20–40 min and for colloids 2–3 h, we hypothesized a difference in the delta MSFP after a fluid bolus (Hahn and Lyons, 2016). If present, this would question the use of crystalloids for Csys and Vs determinations.

Materials and methods

Participants

This was a prospective cohort study in post-surgical patients after coronary artery bypass grafting (CABG). The study is

written according to the Strobe guidelines for cohort studies and was conducted in accordance with the Declaration of Helsinki (von Elm et al., 2007; Cuschieri, 2019). The study took place at the ICU of the Amsterdam University Medical Centers, located at the Academic Medical Center (AMC). The study was approved by the Medical Ethics Committee (NL5531.018.15) and was registered at clinicaltrials.nl before the start of the study (NCT03139929). Written informed consent was obtained prior to surgery. Patients were included between 2017 and 2019. Adult patients (>18 years old) scheduled to undergo elective CABG surgery were included. The exclusion criteria before surgery were morbid obesity (BMI > 40), right- or left-sided heart failure, significant valvular regurgitation or stenosis, arrhythmias, intra-cardiac shunts, symptomatic peripheral vascular disease, and symptomatic pulmonary disease.

During surgery, anesthesia was provided as per routine care. At the end of surgery, noradrenalin, propofol, and/or sufentanil were continued for transport to the ICU. The exclusion criteria at the ICU were a contraindication for fluid loading and persistent hemodynamic instability. Hemodynamic instability was defined as a persistent mean arterial pressure (MAP) below 55 mmHg, a cardiac index below 1.5 L/min/m² or patients in which the MAP remained highly fluctuating (delta 40 mmHg in 10 min) after optimizing initial treatment. A maximum of 1 hour was allowed for patients to fulfil the hemodynamic stability criteria after arrival in the ICU. During study measurements, no alterations in the respiratory rate, positive end expiratory pressure (PEEP), fraction of inspired oxygen (FiO₂), and position of the patient were allowed. Also, the rates of anesthetic, analgesic, and vasoactive drugs were set before the start of the study and could not be altered during study measurements.

Study measurements

Study measurements were performed by a dedicated study team consisting of one member to control the ventilator (inspiratory holds), one member for circulation (administering fluid bolus), one member as the annotator, and one as the supervising ICU consultant.

Arterial blood pressure (ABP) was monitored *via* a catheter in the radial artery, and CVP was monitored *via* a catheter inserted into the right internal jugular vein. Both were connected to a pressure transducer, and both pressure transducers were referenced to the intersection of the anterior axillary line and the fifth intercostal space. Beat-to-beat CO was obtained by Modelflow pulse contour analysis (de Wilde et al., 2007). Measurements were recorded at a sample frequency of 100 Hz and 0.2 mmHg of resolution.




FIGURE 1
Visual study protocol. ABG, arterial blood gas.

MSFP was measured employing successive inspiratory holds as previously published (Maas et al., 2009; Maas et al., 2012a). In short, four inspiratory holds were executed at different pressure levels, namely, 5, 15, 25, and 35 cmH₂O above PEEP. The CVP and CO data at those inspiratory holds were fitted by linear regression. A previous animal study demonstrated three holds sufficient to reliably assess MSFP (Maas et al., 2011). Therefore, an MSFP measurement was judged successful if at least three holds were performed. If the third inspiratory hold (25 cmH₂O above PEEP) resulted in a significant decrease in MAP (defined as a MAP below 50 mmHg), the fourth inspiratory hold (35 cmH₂O above PEEP) could be omitted as decided by the supervising ICU consultant.

For this study, MSFP was measured at three timepoints: at baseline (T = 0), after 100 ml of fluid loading (T = 1), and after a second bolus of 500 ml (T = 2), Figure 1. Both fluid bolus sizes were chosen based on previous studies (Maas et al., 2012a; Smorenberg et al., 2018) and were given at the same infusion speed of 50 ml/min. The supervising ICU consultant could terminate fluid infusion if the ABP increased excessively. No blood pressure cut-off values were defined as the allowed maximum systolic blood pressure (SBP) could differ per patient.

To assess the effects of the type of fluid on MSFP, 50% of the patients who fulfilled hemodynamic stability criteria prior to start of the study were randomized between crystalloid (Sterofundin, BBraun) or colloid infusion (Tetraspan, BBraun) (Hahn and Lyons, 2016)

Outcomes

MSFP was determined as explained previously, by extrapolating CO versus CVP at different inspiratory plateau pressures to CVP is zero. CVP was used as a surrogate for right

atrial pressure. The driving pressure for venous return (VR_{dp}) was defined as MSFP-CVP. Venous return (VR) was defined as VR_{dp} divided by resistance to venous return (RVR). RVR is the reciprocal of the slope of the VR curve, or $RVR = (MSFP - CVP) / CO$. The total systemic vascular resistance (R_{sys}) was calculated as the ratio between the pressure difference of MAP and CVP with CO (Jansen et al., 2010; Maas et al., 2012a; Maas et al., 2012b).

With MSFP measured before and after fluid administration, a pressure-volume relationship could be constructed. C_{sys} is the slope of this relation, or $\Delta \text{volume} / \Delta \text{MSFP}$ (Maas et al., 2012a). $C_{sys} = \text{fluid bolus} / (\text{MSFP after bolus} - \text{MSFP before bolus})$. $V_s = C_{sys} \times \text{MSFP}$.

Since V_s and C_{sys} may vary widely if sympathetic tone or blood flow distribution varies, hemodynamic stability was required during the two volume challenges. Hemodynamic instability during the study was defined as a change in heart rate exceeding 10 beats per minute between two time points (T = 0, T = 1, and T = 2), a decrease in MAP despite fluid administration, or a change in the respiratory rate.

Protocol deviations, to assess feasibility, were defined as any deviation from the study protocol. The supervising ICU consultant was asked to clarify the rationale for the protocol deviation.

Statistical analysis

Continuous data are presented as means with standard deviations when normally distributed or as medians with interquartile ranges (IQR 25–75th) when the data were not normally distributed. The distribution was assessed visually based on Q–Q plots and histograms. Categorical data are presented as frequencies with percentages. The paired t-tests

TABLE 1 Baseline characteristics.

	<i>n</i> = 20
Age	66 ± 8.7
Men	20 (100%)
Height (in cm)	178.7 ± 6.5
Weight (in kg)	88.6 ± 11.2
BMI	27.8 ± 3.7
ASA I	0
ASA II	0
ASA III	16 (80%)
ASA IV	4 (20%)
Medical history	
Hypertension	9 (45%)
Heart failure	0
COPD	0
OSAS	1 (5%)
Obesity	4 (20%)
Diabetes	6 (30%)
Renal insufficiency	1 (5%)
Hypothyroidism	2 (10%)
Relevant medication	
Beta-blocker	16 (80%)
Type of surgery	
CABG on pump	20 (100%)
Surgery duration (min)	252.5 (237.8–324.5)
CPB duration (min)	87.5 (72.0–132.3)
Aortic clamp time (min)	61.5 (45.0–73.5)
TEE after cardiac bypass by cardiac anesthesiologists	
Good LVF and RVF	18 (90%)
Moderate LVF	2 (10%)

ASA, American Society of Anesthesiologists; BMI, body mass index; COPD, chronic obstructive pulmonary disease; OSAS, obstructive sleep apnea syndrome; CABG, coronary artery bypass grafting; CPB, cardiopulmonary bypass; TEE, transesophageal echocardiography; LVF, left ventricular function; RVF, right ventricular function; Min, minutes. Continuous data are presented as a mean with standard deviation (\pm) or as a median with interquartile ranges (IQR 25th–75th). Categorical data are presented as numbers with percentages (%). The ASA classifications were as follows: 1) a healthy person, 2) a patient with mild systemic disease, 3) a patient with severe systemic disease, and 4) a patient with severe systemic disease that is a constant threat to life.

or the non-parametric Wilcoxon signed-rank test were used for comparison of hemodynamic variables on T = 0, T = 1, and T = 2. The independent t-test or the non-parametric Mann–Whitney *U* test were used for differences between colloid and crystalloid groups. A post hoc analysis was performed to correct for patients who did not receive the total amount of 500 ml during the second fluid bolus. The post hoc analysis was performed by dividing the planned fluid administration (=500 ml) by the actual administered fluids (in ml) and multiplying with the delta MSFP.

A *p* value of <0.05 was considered to indicate significance. All analyses were performed using MATLAB version 14 and SPSS version 28.

Sample size analysis

To detect the 500 ml bolus, a sample size of seven patients was calculated to have 90% power to detect a difference in *V*_s means of 500 ml, assuming a standard deviation of 400 ml, using a paired *t*-test with a 0.05 two-sided significance level.

To detect the 100 ml bolus, a sample size of 38 patients was calculated to have 90% power to detect a difference in *V*_s means of 100 ml, assuming a standard deviation of 210 ml, using a paired *t*-test with a 0.05 two-sided significance level.

Assuming a drop-out rate of 10%, a sample size of 42 patients was calculated based on the first 100cc-fluid bolus, between T = 0 and T = 1. As this was the smallest fluid bolus, it required the largest number of patients.

Sample sizes were calculated using nQuery Advanced, version 8.5.1.

Results

Study population

For this prospective cohort study, 121 patients were assessed for eligibility, of these 44 patients had undergone solely CABG surgery and were enrolled in the ICU. A total of 20 patients completed the trial. The exclusion of 24 patients before the start of study measurements at ICU was because of hemodynamic instability (*n* = 18) or because of logistic reasons (*n* = 6), e.g., night-time or incomplete study team, [Supplementary Material S1](#). The median age was 65 years and 100% were men. [Table 1](#) and [Supplementary Material S2](#) show the baseline characteristics. A total of four out of 20 (20%) patients were judged fluid loading responsive (FLR), defined as a 12% increase in CO after the second fluid bolus ([Cherpanath et al., 2013](#)). No serious adverse events occurred in both the colloid and crystalloid groups.

Mean systemic filling pressure

Providing a fluid bolus increased MSFP as expected, with mean MSFP at T = 0 of 20.08 mmHg \pm 3.77, at T = 1 of 21.88 mmHg \pm 4.72 and at T = 2 of 26.82 mmHg \pm 5.58, [Table 2](#).

Stressed volume and compliance

Sixteen patients (80%) fit the inclusion criteria for hemodynamic stability (i.e., stable heart rate and no change in

TABLE 2 Hemodynamic changes after two fluid boluses.

	T = 0	T = 1	T = 2	p1	p2
MSFP	20.08±3.77	21.88 ± 4.72	26.82 ± 5.58	0.005	<0.001
Fluid bolus		100 ± 0	465 ± 103		
		100 (100–100)	500 (425–500)		
HR	69 ± 11	68 ± 10	66 ± 9	0.028	0.049
MAP	72 ± 6	79 ± 7	88 ± 11	<0.001	<0.001
CO	5.25 ± 1.53	5.12 ± 1.26	5.40 ± 1.35	0.397	0.003
CVP	6.86 ± 2.62	7.24 ± 2.70	8.63 ± 3.35	0.004	<0.001
SVR	1158 (934–1450)	1322 (1067–1607)	1393 (1169–1617)	0.025	0.478
VRdp	13.22 ± 2.38	14.97 ± 3.50	18.19 ± 3.63	0.012	<0.001
RVR	2.53 (1.89–3.27)	2.78 (2.37–3.66)	3.38 (2.94–4.11)	0.044	0.004
Rsys	13.15 (9.45–16.62)	14.65 (11.33–17.90)	15.44 (11.35–19.46)	0.004	0.204
PPV	8.94 (7.72–11.48)	7.99 (5.69–10.51)	5.29 (3.26–8.33)	0.351	<0.001
SVV	7.96 (5.92–9.19)	5.98 (4.87–8.00)	3.35 (2.58–6.28)	0.030	<0.001

MSFP, mean systemic filling pressure. HR, heart rate. MAP, mean arterial pressure. CO, cardiac output. CVP, central venous pressure. SVR, systemic vascular resistance [$80 \times (\text{MAP} - \text{CVP}) / \text{CO}$]. VRdp: driving pressure for venous return (MSFP-CVP). RVR: resistance to venous return $[(\text{MSFP} - \text{CVP}) / \text{CO}]$. Rsys: total systemic vascular resistance (MAP-CVP)/CO. Data are presented as a mean with standard deviation (\pm) or as a median with IQR (25th–75th) depending on the normality. *p* values 1 and 2 are determined with paired *t*-tests or the non-parametric related-samples Wilcoxon signed-rank test, depending on normality of the data. *p*1 demonstrates timepoint 0 versus timepoint 1. *p*2 demonstrates timepoint 1 versus timepoint 2. The bold values represent statistically significant differences

the respiratory rate during the study period) and were used for V_s and C_{sys} calculations, [Table 3](#) and [Supplementary Material S3](#).

Since V_s after 100 ml of crystalloid did not consistently result in an increase in MSFP, it was judged not reliable to present mean/median V_s and C_{sys} at T = 1, [Supplementary Material S3](#). Following, T = 2 had a median of 2028.95 ml (IQR 1605.08–3163.51) and C_{sys} at T = 2 had a median of 72.74 ml mmHg⁻¹ (IQR 55.77–132.58). Corrected for body weight, this translates to a median V_s of 24.17 ml kg⁻¹ (IQR 15.68–38.48) and a median C_{sys} of 0.87 ml mmHg⁻¹ kg⁻¹ (IQR 0.54–1.49).

Protocol deviations and clinical feasibility

[Table 3](#) summarizes the protocol deviations and reasons. In all 20 patients (100%), the predefined minimum of three holds could be performed, thus in all patients, MSFP determination was possible. In 10 out of 20 patients (50%), a total of four holds could be executed (at 5, 15, 25, and 35 cmH₂O above PEEP). The reason for not performing a fourth hold was a significant temporary decrease in MAP (mean lowest MAP 49 mmHg \pm 5.81, for less than 20 s) after the third hold.

In 5 out of 20 patients, the second fluid bolus (500 cc) was terminated before the total volume was infused because of a considerable increase in the systolic blood pressure (mean highest systolic blood pressure of 169 mmHg \pm 11.97).

[Supplementary Material S4](#) summarizes this single-center experience concerning the feasibility of MSFP, V_s , and C_{sys} calculations in the ICU.

Exploratory analyses: colloid vs. crystalloid

In the dissecting type of fluids, the choice to cease the infusion of fluids between T = 1 and T = 2 was 3 out of 15 (20%) in the crystalloid group and 2 out of 5 (40%) in the colloid group.

The independent *t*-test demonstrated a significant difference in the response on the first fluid bolus (100 ml) between crystalloids and colloids ($p = 0.019$), [Figure 2](#). The paired *t*-test demonstrated the first colloid bolus resulted in a significant increase in MSFP ($p = 0.038$), whereas the first crystalloid bolus infusion did not ($p = 0.110$). For the second fluid bolus, no significant difference in delta MSFP between the types of fluids was found ($p = 0.122$), [Figure 2](#). However, as the administered amount of fluid during the second bolus of colloid was lower than the administered bolus of crystalloid ([Table 3](#)), this was not a fair comparison. A post hoc analysis demonstrated that when the ceased fluid infusions were extrapolated to the planned 500 ml bolus, there was a significant difference in delta MSFP between crystalloid and colloid infusions, $p = 0.01$, [Figure 2](#) and [Supplementary Material S5](#).

Discussion

This prospective study demonstrated the expected increase in MSFP, derived from 3–4 successive inspiratory holds after fluid loading. The clinical feasibility for MSFP determination was deemed sufficient, although labor intensive, but the clinical feasibility for V_s and C_{sys} was judged ambiguously. The results of this study demonstrate the potential of MSFP and

TABLE 3 Protocol deviations + reasons.

Fluid administration	
Complete first bolus (100 ml)	20/20 (100%)
Complete second bolus (500 ml)	15/20 (75%)
Crystalloid complete second bolus	12/15 (80%)
Crystalloid infused	Mean 483.67 ml ± 92.78 Median 500 IQR 500-500
Colloid complete second bolus	3/5 (60%)
Colloid infused	Mean 410.00 ml ± 124.50 Median 500 IQR 275-500
Reason ceasing infusion	Considerable increase in blood pressure
ABP at which infusion was ceased	
Maximum SBP	169 mmhg ± 11.97
Δ SBP	47 mmhg ± 6.58
MAP	97 mmhg ± 4.10
Δ MAP	24 mmhg ± 4.24
Inspiratory holds	
5, 15, and 25 cmH ₂ O above PEEP	20/20 (100%)
5, 15, 25, and 35 cmH ₂ O above PEEP	10/20 (50%)
Reasons not performing the fourth hold	Considerable decrease in MAP during the third hold
Lowest MAP during the third hold	49 mmhg ± 5.81
Δ MAP during third hold	25 mmhg ± 5.78
Lowest CO during the third hold	2.27 l ± 0.95
Vs and Csys calculations	
Haemodynamic instability during the study	4/20 (20%)
Reasons	
Δ HR more than 10 bpm between two timepoints (T = 0, T = 1, and T = 2)	2/20 (10%)
Starting to trigger ventilator after the 35-mmHg hold	1/20 (5%)
Decrease in MAP after fluid administration	1/20 (5%)


Continuous data are presented as a mean with standard deviation (±) or a median with inter quartile ranges (IQR 25th–75th). Categorical data are presented as frequencies with percentages. SBP, systolic blood pressure. MAP, mean arterial pressure. PEEP, positive end expiratory pressure. CO, cardiac output. HR, heart rate. Vs, stressed volume. Csys, compliance. Δ = delta.

might partly explain why MSFP, Vs, and Csys, derived from inspiratory holds, are not yet widely implemented in clinical care.

Previous studies

Our results are in line with previous studies demonstrating the effect of a fluid bolus on MSFP (Keller et al., 2011; Guerin et al., 2015). Although MSFP is thought central for the characterization of the circulation, the subsequent derived values such as Vs and Csys are subject to physiologic variability, though if accurate, of potentially greater clinical value (Rothe, 1993; Vos et al., 2020; Persichini et al., 2022).

In 1990, Vs was calculated to represent 30% ± 17% of total predicted blood in patients on the cardiac bypass for major vascular surgery (Magder and De Varennes, 1998). During hypothermic cardiac arrest, the cardiac bypass pump was turned off, and the blood that drained passively into a reservoir represented a mean Vs of 1,290 ml ± 296, which equaled 20.2 ml kg⁻¹ ± 1.0 (Magder and De Varennes, 1998). This is close to the 19.5 ml kg⁻¹ ± 12.1 previously found in intact patients with the inspiratory hold technique (Maas et al., 2009), where Vs was a mean of 1,677 ml at baseline. In the present study, the median Vs was 2,028.95 ml at T = 2 (median 24.17 ml kg⁻¹) after the total 600 ml fluid loading.



Previous clinical studies have found C_{sys} to be 80 and 64.3 ml mmHg⁻¹ (Maas et al., 2009; Maas et al., 2012a). In the present study, the median C_{sys} was 72.74 ml mmHg⁻¹. Thus, the present V_s and C_{sys} values are in line with previous studies.

Clinical feasibility

An MSFP measurement takes around 4–5 min. Estimating V_s and C_{sys} thus requires at least 10 min (including fluid loading time) and it assumes the administered fluid is added to the stressed volume compartment (Hahn and Lyons, 2016) (Hahn et al., 2016). Interestingly, 100 ml of colloids did significantly increase MSFP, whereas 100 ml of crystalloids did not. A previous study also found the response on crystalloid variable in post-CABG patients (Sondergaard et al., 2019). Perhaps a capillary leak syndrome with endothelial glycocalyx shedding can partly explain our results, or that the expected transudation of crystalloid into the interstitium occurred rapidly in patients after cardiac surgery (Dekker et al., 2019). Furthermore, V_s and C_{sys} calculations are based upon the assumption that the fluid administered adds directly to the stressed compartment (V_s) without alterations in the unstressed compartment. This assumption might not always be true if fluid administration also results in a shift of blood flow distribution across vascular beds with differing proportions of unstressed and stressed vascular volumes. We conclude a 100-ml bolus of crystalloids to be insufficient to reliably calculate C_{sys} and V_s in this specific ICU population (Aya et al., 2016) (Aya et al., 2017).

Further scrutiny of the MSFP measurement, in the present study, shows that supervising ICU physicians were less inclined to allow the fourth hold (35 cmH₂O) compared to previous

studies because of (transient) hypotension (Maas et al., 2009; Maas et al., 2012a). Excluding the final 35 cm H₂O inspiratory hold step in patients in whom a total of four holds could be executed, no significant change in the MSFP estimate was found; *p* = 0.696. This demonstrates that using high inspiratory hold levels may not be necessary. We noted that for executing the inspiratory holds, a deep level of sedation is necessary, exceeding the common level of sedation in ICU patients. Finally, although inspiratory holds have been shown to prevent postoperative pulmonary complications in non-ARDS patients (Hartland et al., 2015), they should not be performed in ARDS patients (Cavalcanti et al., 2017).

Limitations

In only 20 patients, instead of the planned 42, study measurements could be performed. For the 20 patients, we were sufficiently powered (>90%) to detect a difference in V_s means of 500 ml, but a post hoc sample size analysis demonstrated this to reduce the power for detection of the 100 ml bolus to 56%. The majority of patients were excluded because they did not meet the hemodynamic stability criteria to start the study in the ICU. Our criteria could be too strict, or our cardiac surgery population more severely ill. Comparison with previous studies was not possible, as these numbers were not reported.

The colloid versus crystalloid analysis should be regarded as exploratory, but it could be used as a stepping stone for new trials. The post hoc analysis requires MSFP to increase linearly with infused fluids, as suggested by previous data (Maas et al., 2012a). Still, we cannot prove this linear association to be true for the presented data, so future studies should confirm our results.

The high number of protocol deviations in the study protocol might not solely describe clinical feasibility but can also illustrate that the supervising ICU consultants in the study hospital were more conservative.

In this study, we used Modelflow pulse contour to calculate CO. For MSFP, Vs and Csys absolute CO values are not necessary; trends are sufficient, [Supplementary Material S6](#). However, for RVR, absolute values become relevant. Modelflow can be calibrated with thermodilution and echocardiography (Swan et al., 1970; de Vaal et al., 2005; Harvey et al., 2005; Tuggle, 2009).

An unintended but important finding is that all the patients studied were men. All the initially included women were deemed hemodynamically unstable in the ICU. Aiming to include women in order to obtain a study population that is reflective of the clinical population and in order to translate findings across genders remains important (Bybee and Stevens, 2013; Garcia et al., 2016; Gulati et al., 2021).

What needs to happen to bring MSFP to clinical care, and what for?

MSFP determined with inspiratory holds is of great interest for research purposes, but based on this study, not yet ready for clinical use. Yet, less invasive alternatives for determining MSFP do exist (Wijnberge et al., 2018). The MSFP analogue is based on a model of the circulation, it is a calculation with CO, CVP, and MAP as input data. The MSFP analogue is much simpler to measure but was thought to suffer from greater inaccuracies, probably because of the assumptions in the calculation (Maas et al., 2012c; Meijs et al., 2022). The calculation uses standard arterial and venous compliances and resistances that might be inaccurate during acute disease states. However, a recent animal study concluded MSFP analogue to be the most reliable method to indirectly measure MSFP (Werner-Moller et al., 2019). The jury is still out, and this contradiction in results invites future research. A third method to estimate MSFP is based on the stop-flow principle, determined with a rapidly inflating cuff (halting blood flow) around the upper arm. A previous study demonstrated all three methods to track a fluid bolus (Maas et al., 2012c).

If MSFP determined with inspiratory holds is to become more commonly used, studies need to define the minimum number of inspiratory holds for accurate MSFP determination and should assess whether holds with lower plateau pressures also result in accurate MSFP values (Persichini et al., 2012; Guerin et al., 2015). Furthermore, if knowing accurate Vs and Csys is required, then defining the optimal fluid type (colloid vs. crystalloid) and the minimal volume challenge needed to determine Vs and Csys needs to

be assessed. Being able to quickly and reliably calculate MSFP, Csys and Vs could be beneficial in guiding hemodynamic care in various types of patients (Rothe, 1993; Vos et al., 2020; Persichini et al., 2022). For example, in current sepsis resuscitation, first fluids are administered and subsequently (after >2 L of fluids is added in a normal sized adult patient), a vasopressor is started. Based on a Guytonian approach to the circulation, however, it would make more sense to start a vasopressor earlier in the treatment to recruit unstressed to the stressed volume (Adda et al., 2021; Persichini et al., 2022). Recruiting unstressed to stressed volume is an important survival mechanism of the human body. Measuring MSFP, Vs, and Csys might lead to a reduction in the total amount of fluids administered for resuscitation (Ospina-Tascón et al., 2020; Persichini et al., 2022). Despite the use of invasive hemodynamic monitoring options available in the ICU, we still lack direct and repetitive estimation of the effective circulating volume (Vs). Working with MSFP might enable us to go beyond fluid loading responsiveness and help us better understand the physiology during various clinical scenarios (Crozier et al., 2015; Moller et al., 2019; Persichini et al., 2022). Future studies should study whether adding MSFP, Vs, and Csys to our clinical arsenal actually result in improved patient outcomes.

Conclusion

Mean systemic filling pressure, estimated with inspiratory holds, behaves predictably conform known physiological principles. Clinical feasibility for Csys and Vs calculations was judged ambiguously based on the lack of required hemodynamic stability and on the assumption of administered fluids to stay intravascular. Future studies should address the clinical obstacles found in this study, and less-invasive alternatives to determine MSFP should be further explored.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary Material](#); further inquiries can be directed to the corresponding authors.

Ethics statement

The studies involving human participants were reviewed and approved by the Medical Ethics Committee of the Amsterdam UMC location AMC (NL5531.018.15). The patients/participants provided their written informed consent to participate in this study.

Author contributions

Study conception: BG. Data collection: MW, RK, LT, JB, NL, AV, DV, and BG. Data analyses: MW and JJ. Statistical analyses: MW. Interpretation of the data: all authors. Drafting the manuscript: MW, JJ, MP, and BG. Revision of the manuscript: all authors.

Conflict of interest

Author BG was employed by Healthplus.ai.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Adda, I., Lai, C., Teboul, J. L., Guerin, L., Gavelli, F., and Monnet, X. (2021). Norepinephrine potentiates the efficacy of volume expansion on mean systemic pressure in septic shock. *Crit. Care* 25, 302. doi:10.1186/s13054-021-03711-5
- Aya, H. D., Rhodes, A., Chis Ster, I., Fletcher, N., Grounds, R. M., and Cecconi, M. (2017). Hemodynamic effect of different doses of fluids for a fluid challenge: A Quasi-randomized controlled study. *Crit. Care Med.* 45, e161–e168. doi:10.1097/CCM.0000000000002067
- Aya, H. D., Ster, I. C., Fletcher, N., Grounds, R. M., Rhodes, A., and Cecconi, M. (2016). Pharmacodynamic analysis of a fluid challenge. *Crit. Care Med.* 44, 880–891. doi:10.1097/CCM.0000000000001517
- Bybee, K. A., and Stevens, T. L. (2013). Matters of the heart: Cardiovascular disease in U.S. Women. *Mo. Med.* 110, 65–70.
- Cavalcanti, A. B., Suzumura É, A., Laranjeira, L. N., Paisani, D. M., Damiani, L. P., Guimarães, H. P., et al. (2017). Effect of Lung recruitment and Titrated positive end-expiratory pressure (PEEP) vs low PEEP on Mortality in patients with acute respiratory distress syndrome: A randomized clinical trial. *Jama* 318, 1335–1345. doi:10.1001/jama.2017.14171
- Cherpanath, T. G., Geerts, B. F., Lagrand, W. K., Schultz, M. J., and Groeneveld, A. B. (2013). Basic concepts of fluid responsiveness. *Neth. Heart J.* 21, 530–536. doi:10.1007/s12471-013-0487-7
- Cooke, K., Sharvill, R., Sondergaard, S., and Aneman, A. (2018). Volume responsiveness assessed by passive leg raising and a fluid challenge: A critical review focused on mean systemic filling pressure. *Anaesthesia* 73, 313–322. doi:10.1111/anae.14162
- Crozier, T. M., Wallace, E. M., and Parkin, W. G. (2015). Haemodynamic assessment in pregnancy and pre-eclampsia: A Guytonian approach. *Pregnancy Hypertens.* 5, 177–181. doi:10.1016/j.preghy.2015.01.005
- Cuschieri, S. (2019). The STROBE guidelines. *Saudi J. Anaesth.* 13, S31–S34. doi:10.4103/sja.SJA_543_18
- de Vaal, J. B., de Wilde, R. B., van den Berg, P. C., Schreuder, J. J., and Jansen, J. R. (2005). Less invasive determination of cardiac output from the arterial pressure by aortic diameter-calibrated pulse contour. *Br. J. Anaesth.* 95, 326–331. doi:10.1093/bja/aei189
- de Wilde, R. B., Schreuder, J. J., van den Berg, P. C., and Jansen, J. R. (2007). An evaluation of cardiac output by five arterial pulse contour techniques during cardiac surgery. *Anaesthesia* 62, 760–768. doi:10.1111/j.1365-2044.2007.05135.x
- de Wit, F., van Vliet, A. L., de Wilde, R. B., Jansen, J. R., Vuyk, J., Aarts, L. P., et al. (2016). The effect of propofol on haemodynamics: Cardiac output, venous return, mean systemic filling pressure, and vascular resistances. *Br. J. Anaesth.* 116, 784–789. doi:10.1093/bja/aew126
- Dekker, N. A. M., Veerhoek, D., Koning, N. J., van Leeuwen, A. L. I., Elbers, P. W. G., van den Brom, C. E., et al. (2019). Postoperative microcirculatory perfusion and endothelial glycocalyx shedding following cardiac surgery with cardiopulmonary bypass. *Anaesthesia* 74, 609–618. doi:10.1111/anae.14577
- Garcia, M., Mulvagh, S. L., Merz, C. N., Buring, J. E., and Manson, J. E. (2016). Cardiovascular disease in women: Clinical perspectives. *Circ. Res.* 118, 1273–1293. doi:10.1161/CIRCRESAHA.116.307547
- Guerin, L., Teboul, J. L., Persichini, R., Dres, M., Richard, C., and Monnet, X. (2015). Effects of passive leg raising and volume expansion on mean systemic pressure and venous return in shock in humans. *Crit. Care* 19, 411. doi:10.1186/s13054-015-1115-2
- Gulati, M., Hendry, C., Parapid, B., and Mulvagh, S. L. (2021). Why We need Specialised Centres for Women's hearts: Changing the Face of cardiovascular care for women. *Eur. Cardiol.* 16, e52. doi:10.15420/scr.2021.49
- Hahn, R. G., Drobin, D., and Zdolsek, J. (2016). Distribution of crystalloid fluid changes with the rate of infusion: A population-based study. *Acta Anaesthesiol. Scand.* 60, 569–578. doi:10.1111/aas.12686
- Hahn, R. G., and Lyons, G. (2016). The half-life of infusion fluids: An educational review. *Eur. J. Anaesthesiol.* 33, 475–482. doi:10.1097/EJA.0000000000000436
- Hartland, B. L., Newell, T. J., and Damico, N. (2015). Alveolar recruitment maneuvers under general anesthesia: A systematic review of the literature. *Respir. Care* 60, 609–620. doi:10.4187/respcare.03488
- Harvey, S., Harrison, D. A., Singer, M., Ashcroft, J., Jones, C. M., Elbourne, D., et al. (2005). Assessment of the clinical effectiveness of pulmonary artery catheters in management of patients in intensive care (PAC-Man): A randomised controlled trial. *Lancet (London, Engl.)* 366, 472–477. doi:10.1016/S0140-6736(05)67061-4
- Helmerhorst, H. J., de Wilde, R. B., Lee, D. H., Palmén, M., Jansen, J. R., van Westerloo, D. J., et al. (2017). Hemodynamic effects of short-term hyperoxia after coronary artery bypass grafting. *Ann. Intensive Care* 7, 20. doi:10.1186/s13613-017-0246-9
- Jansen, J. R., Maas, J. J., and Pinsky, M. R. (2010). Bedside assessment of mean systemic filling pressure. *Curr. Opin. Crit. Care* 16, 231–236. doi:10.1097/MCC.0b013e3283378185
- Keller, G., Desebbe, O., Benard, M., Bouchet, J. B., and Lehot, J. J. (2011). Bedside assessment of passive leg raising effects on venous return. *J. Clin. Monit. Comput.* 25, 257–263. doi:10.1007/s10877-011-9303-3
- Maas, J. J., de Wilde, R. B., Aarts, L. P., Pinsky, M. R., and Jansen, J. R. (2012b). Determination of vascular waterfall phenomenon by bedside measurement of mean systemic filling pressure and critical closing pressure in the intensive care unit. *Anesth. Analg.* 114, 803–810. doi:10.1213/ANE.0b013e318247fa44
- Maas, J. J., Geerts, B. F., and Jansen, J. R. C. (2011). Evaluation of mean systemic filling pressure from pulse contour cardiac output and central venous pressure. *J. Clin. Monit. Comput.* 25, 193–201. doi:10.1007/s10877-011-9294-0
- Maas, J. J., Geerts, B. F., van den Berg, P. C., Pinsky, M. R., and Jansen, J. R. (2009). Assessment of venous return curve and mean systemic filling pressure in postoperative cardiac surgery patients. *Crit. Care Med.* 37, 912–918. doi:10.1097/CCM.0b013e3181961481
- Maas, J. J., Pinsky, M. R., Aarts, L. P., and Jansen, J. R. (2012a). Bedside assessment of total systemic vascular compliance, stressed volume, and cardiac function curves

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2022.1041730/full#supplementary-material>

in intensive care unit patients. *Anesth. Analg.* 115, 880–887. doi:10.1213/ANE.0b013e31825fb01d

Maas, J. J., Pinsky, M. R., de Wilde, R. B., de Jonge, E., and Jansen, J. R. (2013). Cardiac output response to norepinephrine in postoperative cardiac surgery patients: Interpretation with venous return and cardiac function curves. *Crit. Care Med.* 41, 143–150. doi:10.1097/CCM.0b013e318265ea64

Maas, J. J., Pinsky, M. R., Geerts, B. F., de Wilde, R. B., and Jansen, J. R. (2012c). Estimation of mean systemic filling pressure in postoperative cardiac surgery patients with three methods. *Intensive Care Med.* 38, 1452–1460. doi:10.1007/s00134-012-2586-0

Magder, S., and De Varennes, B. (1998). Clinical death and the measurement of stressed vascular volume. *Crit. Care Med.* 26, 1061–1064. doi:10.1097/00003246-199806000-00028

Meijs, L. P. B., van Houte, J., Conjaerts, B. C. M., Bindels, A., Bouwman, A., Houterman, S., et al. (2022). Clinical validation of a computerized algorithm to determine mean systemic filling pressure. *J. Clin. Monit. Comput.* 36, 191–198. doi:10.1007/s10877-020-00636-2

Moller, P. W., Hana, A., Heinisch, P. P., Liu, S., Djafarzadeh, S., Haenggi, M., et al. (2019). The effects of Vasoconstriction and volume expansion on Veno-arterial ECMO flow. *Shock* 51, 650–658. doi:10.1097/SHK.0000000000001197

Ospina-Tascón, G. A., Hernandez, G., Alvarez, I., Calderón-Tapia, L. E., Manzano-Nunez, R., Sánchez-Ortiz, A. I., et al. (2020). Effects of very early start of norepinephrine in patients with septic shock: A propensity score-based analysis. *Crit. Care* 24 (52), 52. doi:10.1186/s13054-020-2756-3

Persichini, R., Lai, C., Teboul, J. L., Adda, I., Guérin, L., and Monnet, X. (2022). Venous return and mean systemic filling pressure: Physiology and clinical applications. *Crit. Care* 26, 411. doi:10.1186/s13054-015-1115-2

Persichini, R., Silva, S., Teboul, J. L., Jozwiak, M., Chemla, D., Richard, C., et al. (2012). Effects of norepinephrine on mean systemic pressure and venous return in human septic shock. *Crit. Care Med.* 40, 3146–3153. doi:10.1097/CCM.0b013e318260c6c3

Rothe, C. F. (1993). Mean circulatory filling pressure: Its meaning and measurement. *J. Appl. Physiol.* 74, 499–509. doi:10.1152/jappl.1993.74.2.499

Smorenberg, A., Cherpanath, T. G. V., Geerts, B. F., de Wilde, R. B. P., Jansen, J. R. C., Maas, J. J., et al. (2018). A mini-fluid challenge of 150mL predicts fluid responsiveness using Modelflow(R) pulse contour cardiac output directly after cardiac surgery. *J. Clin. Anesth.* 46, 17–22. doi:10.1016/j.jclinane.2017.12.022

Sondergaard, S., Larsson, J. S., and Möller, P. W. (2019). The haemodynamic effects of crystalloid and colloid volume resuscitation on primary, derived and efficiency variables in post-CABG patients. *Intensive Care Med. Exp.* 7 (13), 13. doi:10.1186/s40635-019-0224-7

Swan, H. J., Ganz, W., Forrester, J., Marcus, H., Diamond, G., and Chonette, D. (1970). Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. *N. Engl. J. Med.* 283, 447–451. doi:10.1056/NEJM197008272830902

Tuggle, D. (2009). Obtaining pulmonary artery catheter data is too labor intense to be reliable. *Crit. Care Med.* 37, 1833. doi:10.1097/CCM.0b013e3181a097ba

Vincent, J. L., and Pinsky, M. R. (2018). We should avoid the term "fluid overload. *Crit. Care* 22, 214. doi:10.1186/s13054-018-2141-7

von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., Vandenbroucke, J. P., et al. (2007). The Strengthening the reporting of Observational studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Lancet (London, Engl.)* 370, 1453–1457. doi:10.1016/S0140-6736(07)61602-X

Vos, J. J., Kalmar, A. F., and Scheeren, T. W. L. (2020). Bedside assessment and clinical utility of mean systemic filling pressure in acute care. *J. Emerg. Crit. Care Med.* 4, 25. doi:10.21037/jeccm.2020.03.04

Werner-Moller, P., Sondergaard, S., Jakob, S. M., Takala, J., and Berger, D. (2019). Effect of volume status on the estimation of mean systemic filling pressure. *J. Appl. Physiol.* (1985) 126, 1503–1513. doi:10.1152/japplphysiol.00897.2018

Wijnberge, M., Sindhunata, D. P., Pinsky, M. R., Vlaar, A. P., Ouweneel, E., Jansen, J. R., et al. (2018). Estimating mean circulatory filling pressure in clinical practice: A systematic review comparing three bedside methods in the critically ill. *Ann. Intensive Care* 8, 73. doi:10.1186/s13613-018-0418-2