



Universiteit  
Leiden  
The Netherlands

## **Thrombocytopenia limits the feasibility of salvage lomustine chemotherapy in recurrent glioblastoma: a secondary analysis of EORTC 26101**

Rhun, E. le; Oppong, F.B.; Bent, M. van den; Wick, W.; Brandes, A.A.; Taphoorn, M.J.B.; ... ; Weller, M.

### **Citation**

Rhun, E. le, Oppong, F. B., Bent, M. van den, Wick, W., Brandes, A. A., Taphoorn, M. J. B., ... Weller, M. (2022). Thrombocytopenia limits the feasibility of salvage lomustine chemotherapy in recurrent glioblastoma: a secondary analysis of EORTC 26101. *European Journal Of Cancer*, 178, 13-22. doi:10.1016/j.ejca.2022.10.006

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3748085>

**Note:** To cite this publication please use the final published version (if applicable).



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.ejancer.com](http://www.ejancer.com)



## Original Research

# Thrombocytopenia limits the feasibility of salvage lomustine chemotherapy in recurrent glioblastoma: a secondary analysis of EORTC 26101



Emilie Le Rhun<sup>a,b,\*</sup>, Felix Boakye Oppong<sup>c</sup>, Martin van den Bent<sup>d</sup>, Wolfgang Wick<sup>e</sup>, Alba A. Brandes<sup>f</sup>, Martin JB. Taphoorn<sup>g</sup>, Michael Platten<sup>h</sup>, Ahmed Idbaih<sup>i</sup>, Paul M. Clement<sup>j</sup>, Matthias Preusser<sup>k</sup>, Vassilis Gelfinopoulos<sup>c</sup>, Thierry Gorlia<sup>c</sup>, Michael Weller<sup>a</sup>

<sup>a</sup> Departments of Neurology and Clinical Neuroscience Center, University Hospital and University of Zurich, Zurich, Switzerland

<sup>b</sup> Departments of Neurosurgery, Clinical Neuroscience Center, University Hospital and University of Zurich, Zurich, Switzerland

<sup>c</sup> EORTC Headquarters, Brussels, Belgium

<sup>d</sup> Erasmus MC Cancer Center, Rotterdam, the Netherlands

<sup>e</sup> Department of Neurology and Neurooncology Program at the National Center for Tumor Diseases, University Hospital Heidelberg and German Cancer Research Center, Heidelberg, Germany

<sup>f</sup> Medical Oncology Department, Azienda Unità Sanitaria Locale di Bologna-IRCCS Istituto Delle Scienze Neurologiche, Bologna, Italy

<sup>g</sup> Leiden University Medical Center, Departments of Neurology, Leiden, And Haaglanden Medical Center, The Hague, the Netherlands

<sup>h</sup> Medical Faculty Mannheim, MCTN, Department of Neurology, Mannheim, Germany

<sup>i</sup> Sorbonne Université, Institut Du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS, AP-HP, Hôpital Universitaire La Pitié Salpêtrière, DMU Neurosciences, Paris, France

<sup>j</sup> Department of Oncology, KU Leuven, Leuven Cancer Institute-KU Leuven, Leuven, Belgium

<sup>k</sup> Department of Medicine I, Division of Oncology, Medical University of Vienna, Vienna, Austria

Received 6 August 2022; received in revised form 12 October 2022; accepted 13 October 2022

Available online 20 October 2022

## KEYWORDS

CCNU;  
Chemotherapy;  
Myelosuppression;  
Platelets;

**Abstract Background:** Thrombocytopenia represents the main cause of stopping alkylating chemotherapy for toxicity. Here, we explored the incidence, and the consequences for treatment exposure and survival, of thrombocytopenia induced by lomustine in recurrent glioblastoma. **Methods:** We performed a retrospective analysis of the associations of thrombocytopenia with treatment delivery and outcome in EORTC 26101, a randomised trial designed to define the role

\* Corresponding author: Department of Neurosurgery, University Hospital Zurich, Frauenklinikstrasse 10, 8091, Zurich, Switzerland.  
E-mail address: [emilie.lerhun@usz.ch](mailto:emilie.lerhun@usz.ch) (E. Le Rhun).

of lomustine versus bevacizumab versus their combination in recurrent glioblastoma.

**Results:** A total of 225 patients were treated with lomustine alone (median 1 cycle) (group 1) and 283 patients were treated with lomustine plus bevacizumab (median 3 lomustine cycles) (group 2). Among cycle delays and dose reductions of lomustine for toxicity, thrombocytopenia was the leading cause. Among 129 patients (57%) of group 1 and 187 patients (66%) of group 2 experiencing at least one episode of thrombocytopenia, 36 patients (16%) in group 1 and 93 (33%) in group 2 had their treatment modified because of thrombocytopenia. Lomustine was discontinued for thrombocytopenia in 16 patients (7.1%) in group 1 and in 38 patients (13.4%) in group 2. On adjusted analysis accounting for major prognostic factors, dose modification induced by thrombocytopenia was associated with inferior progression-free survival in patients with *MGMT* promoter-methylated tumours in groups 1 and 2. This effect was noted for overall survival, too, but only for group 2 patients. Conclusion: Drug-induced thrombocytopenia is a major limitation to adequate exposure to lomustine chemotherapy in recurrent glioblastoma. Mitigating thrombocytopenia to enhance lomustine exposure might improve outcome in patients with *MGMT* promoter-methylated tumours.

© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The current standard of care treatment for patients with newly diagnosed glioblastoma is surgery as safely feasible followed by radiotherapy with concomitant temozolomide and six cycles of maintenance temozolomide chemotherapy [1]. Recurrence is inevitable, but standards of care for recurrent glioblastoma are less well defined. A minority of patients in the range of 20% may be candidates for second surgery or re-irradiation, but neither of these interventions has been shown to prolong survival in a randomised clinical trial [2]. The majority of patients who are eligible for salvage treatment receive systemic therapy, mostly with lomustine, a nitrosourea compound, or, depending on availability, bevacizumab, an antibody to vascular endothelial growth factor. Lomustine alone has been increasingly considered a standard of care regimen for clinical trials in recurrent glioblastoma [3–5], including the AGILE platform trial [6].

Lomustine, like probably all nitrosoureas, is more active in patients with tumours with promoter methylation of the *O*<sup>6</sup>-methylguanine DNA methyltransferase (*MGMT*) gene [7–9]. However, notably in patients pre-exposed to temozolomide, lomustine chemotherapy carries a high risk of myelosuppression, mostly thrombocytopenia, which may necessitate dose reductions, dose delays, discontinuation of chemotherapy, platelet transfusions and may cause haemorrhages. In the REGAL trial, a phase III randomised study evaluating the efficacy of the vascular endothelial growth factor receptor inhibitor cediranib alone or in combination with lomustine versus lomustine alone [4], Common Terminology Criteria for Adverse Events (CTCAE) grade 3 and 4 thrombocytopenias were observed in 38% and 20%, respectively, in patients treated with cediranib plus

lomustine, compared with 22% and 3% in the lomustine alone arm, and 2% and 1% in the cediranib alone arm.

The present secondary analysis of EORTC 26101 [8] sought to explore the extent to which thrombocytopenia interferes with adequate delivery of treatment in patients with recurrent glioblastoma. The rationale for this study was the availability of novel agents like romiplostim that may counteract chemotherapy-induced thrombocytopenia, including glioblastoma [10].

## 2. Patients and methods

### 2.1. Patients

To assess the incidence and clinical significance of lomustine-induced thrombocytopenia at first recurrence of glioblastoma, we analysed data from the phase II and III parts of EORTC 26101 (NCT01290939) [8] (Note S1).

### 2.2. Statistical analysis

We analysed coded individual patient data including date of randomisation, WHO performance status, tumour volume at study entry, steroid use, *MGMT* promoter methylation status, number of treatment cycles, haematological and non-haematological toxicity according to CTCAE v4.0 during study treatment, lomustine dose delays and modifications, lomustine discontinuation, progression-free survival (PFS) and overall survival (Note S2).

SAS version 9.4 (© 2002–2012 per SAS Institute Inc., Cary, NC, USA.) was used for the analyses. For the primary analyses, statistical significance was established at a level of 5%. In descriptive comparison of subgroups,

percentage difference of 10% or more was considered clinically relevant.

### 3. Results

#### 3.1. Incidence of thrombocytopenia during the study

Analyses were performed on 508 patients: 225 patients treated with lomustine alone (group 1) and 283 patients treated with lomustine plus bevacizumab (group 2) (Fig. S1). Patient characteristics in both groups were similar (Table S1). The median number of lomustine cycles was 1 (range 1–17) in group 1 and 3 (range 1–9) in group 2 [8]. The baseline platelet counts were similar between groups (Table S1). The frequencies with grade of thrombocytopenia per all cycles per patient are detailed in Table S2: 96 patients (43%) in group 1 and 96 patients (34%) in group 2 experienced no single cycle with thrombocytopenia. Conversely, 129 patients (57%) in group 1 and 187 patients (66%) in group 2 experienced at least one cycle with thrombocytopenia. Among the patients who developed at

least one episode of thrombocytopenia, at least one more episode was noted in 44 of 129 patients (34%) in group 1 and 107 of 187 patients (57%) in group 2. The likelihood of experiencing no thrombocytopenia remained relatively stable over sequential cycles of lomustine in a range of 50–70% and was similar in both groups (Table S3).

#### 3.2. Impact of thrombocytopenia on lomustine exposure

The overall reasons for lomustine dose delays, dose reductions and discontinuation in both patient groups are shown in Table 1. Lomustine treatment delay, dose reduction and treatment discontinuation due to thrombocytopenia per cycle are shown in Table S4. Progression was the major reason for stopping lomustine, but thrombocytopenia was the major toxicity causing dose delays, dose reductions and discontinuation. Groups did not differ for dose delays and dose-reduced cycles, but relatively more group 1 patients stopped lomustine for

Table 1  
Overall causes of lomustine dose delays, dose reductions and discontinuation.

	All lomustine alone cycles (n = 464) Group 1	All lomustine-plus bevacizumab cycles (n = 1071) Group 2
<b>Delayed cycles</b>		
Number of delayed cycles (n, %)		
for any reason	41 (8.8)	111 (10.4)
for toxicity	33 (7.1)	97 (9.1)
for non-haematological toxicity	3 (0.6)	23 (2.1)
for haematological toxicity	30 (6.5)	76 (7.1)
for thrombocytopenia	23 (5.0)	53 (4.9)
for other reasons	8 (1.7)	12 (1.1)
<b>Dose-reduced cycles</b>		
Number of dose-reduced cycles (n, %)		
for any reason	82 (17.7)	143 (13.4)
for toxicity	53 (11.4)	111 (10.4)
for non-haematological toxicity	4 (0.9)	16 (1.5)
for haematological toxicity	50 (10.8)	97 (9.1)
for thrombocytopenia	29 (6.3)	76 (7.1)
for other reasons	29 (6.3)	32 (3.0)
	Lomustine alone patients (n = 225) Group 1	Lomustine plus bevacizumab patients (n = 283) Group 2
<b>Lomustine discontinuation<sup>a</sup></b>		
for toxicity	25 (11.1)	58 (20.5)
for non-haematological toxicity	4 (1.8)	16 (5.7)
for haematological toxicity	21 (9.3)	42 (14.8)
for thrombocytopenia	16 (7.1)	38 (13.4) <sup>a</sup>
for progressive disease	188 (83.6)	184 (65.0) <sup>a</sup>
for other reasons	12 (5.3)	42 (14.8)
<b>Any reason for lomustine modification</b>		
for toxicity	59 (26.2)	141 (49.8)
for non-haematological toxicity	7 (3.1)	42 (14.8)
for haematological toxicity	54 (24.0)	116 (41.0)
for thrombocytopenia	36 (16.0)	93 (32.9)
for progressive disease	188 (83.6)	184 (65.0)
for other reasons	12 (5.3)	44 (15.5)

<sup>a</sup> One patient discontinued treatment for both thrombocytopenia and progressive disease.

progression whereas more group 2 patients stopped lomustine for toxicity (Table 1).

A total of 42 patients (23%) in group 1 and of 64 patients (26%) in group 2 experienced grade 3 or 4 thrombocytopenia. The frequencies of

thrombocytopenia by CTCAE grade, associated delays of the next cycle, dose reductions and discontinuation of lomustine are summarised in Table 2. In group 1, 36 patients experienced treatment modification due to thrombocytopenia, corresponding to 16% (36/225) of

Table 2

Lomustine exposure, thrombocytopenia and haemorrhage.

	Lomustine alone (n = 225) Group 1	Lomustine plus bevacizumab (n = 283) Group 2
<b>Number of lomustine cycles (n, range)</b>		
Median number of lomustine cycles	1 (1–17)	3 (1–9)
<b>Worst grade thrombocytopenia during treatment</b>		
Grade 0	95 (42.2)	94 (33.2)
Grade 1	36 (16.0)	49 (17.3)
Grade 2	41 (18.2)	64 (22.6)
Grade 3	33 (14.7)	59 (20.8)
Grade 4	19 (8.4)	15 (5.3)
Not reported	1 (0.4)	2 (0.7)
<b>Lomustine dose delay</b>		
<b>CTCAE grade 1 thrombocytopenia</b>		
1 delayed cycle (n, %)	4 (1.8)	12 (4.2)
2 delayed cycles (n, %)	0	2 (0.7)
Median duration of delay in days (median,range)	17.5 (14–32)	14.5 (8–28)
<b>CTCAE grade 2 thrombocytopenia</b>		
1 delayed cycle (n, %)	8 (3.6)	17 (6.0)
2 delayed cycles (n, %)	0	1 (0.4)
Median duration of delay in days (median,range)	14.5 (13–19)	14.0 (8–28)
<b>CTCAE grade 3 thrombocytopenia</b>		
1 delayed cycle for grade 3 (n, %)	9 (4.0)	12 (4.2)
2 delayed cycles for grade 3 (n, %)	0	1 (0.4)
Median duration of delay in days (median,range)	14.0 (8–28)	15.0 (8–35)
<b>CTCAE grade 4 thrombocytopenia</b>		
1 delayed cycle (n, %)	2 (0.9)	4 (1.4)
2 delayed cycles (n, %)	0	0
Median duration of delay in days (median, range)	13.5 (13–14)	11 (8–14)
<b>CTCAE any grade thrombocytopenia</b>		
1 delayed cycle (n, %)	23 (10.3)	45 (15.8)
2 delayed cycles (n, %)	0	4 (1.5)
Median duration of delay in days (median,range)	14.0 (8–32)	14.0 (8–35)
<b>Lomustine dose reduction</b>		
for CTCAE grade 1 (not foreseen per protocol) (n, %)	2 (0.9)	11 (3.9)
for CTCAE grade 2 (not foreseen per protocol) (n, %)	6 (2.7)	19 (6.7)
for CTCAE grade 3 (n, %)	16 (7.1)	32 (11.3)
for CTCAE grade 4 (n, %)	4 (1.8)	9 (3.2)
for any CTCAE grade (n, %)	28 (12.4)	71 (25.1)
<b>Lomustine dose discontinuation</b>		
for CTCAE grade 1 (n, %)	2 (0.9)	11 (3.9)
for CTCAE grade 2 (n, %)	4 (1.8)	7 (2.5)
for CTCAE grade 3 (n, %)	4 (1.8)	17 (6.0)
for CTCAE grade 4 (n, %)	6 (2.7)	3 (1.1)
for any CTCAE grade (n, %)	16 (7.1)	38 (13.5)
<b>Platelet transfusion (n, %)</b>		
Patients with at least one platelet transfusion	8 (3.6)	5 (1.7)
Patients with more than one platelet transfusion	6 (2.7)	1 (0.04)
<b>Bleeding</b>		
Intracranial haemorrhage (n, %)		
CTCAE Grade 1	0	3 (1.1)
CTCAE Grade 2	2 (0.9)	2 (0.7)
CTCAE Grade 3	1 (0.4)	0
CTCAE Grade 4	0	0
CTCAE Grade 5	0	1 (0.3)
Any CTCAE grade	3 (1.3)	6 (2.1)
Extracranial haemorrhage	0	0

Abbreviations: n: number of patients, %: percentage.

all patients in group 1 and to 28% (36/129) of patients experiencing at least one episode of thrombocytopenia. In group 2, 93 patients experienced treatment modification due to thrombocytopenia, corresponding to 33% (93/283) of all patients in group 2 and to 50% (93/187) of patients experiencing at least one episode of thrombocytopenia (Table S2, Table S4). Among patients experiencing treatment modification induced by thrombocytopenia, 14 of 36 patients (39%) in group 1 and 27 of 93 patients (29%) in group 2 had their lomustine treatment modified due to thrombocytopenia more than once (Table S5).

### 3.3. Dose delays

Twenty-three patients (10%) in group 1 and 49 patients (17%) in group 2 had a dose delay for any grade of thrombocytopenia. The median duration of delay was 14 days (Table 2). A similar number of patients with delayed cycles was observed for grade 1/2 versus grade 3/4 thrombocytopenia in group 1 (5% vs. 5%) and group 2 (11% vs. 6%).

### 3.4. Dose reductions

Twenty-eight patients (12%) in group 1 versus 71 patients (25%) in group 2 had a lomustine dose reduction for any grade of thrombocytopenia (Table 2). As foreseen per protocol, most dose reductions were due to CTCAE grade 3/4 thrombocytopenia: 20 of 28 patients in group 1 and 41 of 71 patients in group 2. Thrombocytopenia was responsible for 29 of 53 cycles (55%) dose-reduced for toxicity in group 1 and 76 of 111 cycles (68%) dose-reduced for toxicity in group 2 (Table 1).

### 3.5. Discontinuation

Lomustine was discontinued for thrombocytopenia in 16 patients (7%) in group 1 and in 38 patients (13%) in group 2 (Tables 1 and 2), mainly for CTCAE grade 3/4 thrombocytopenia: 10 of 16 patients in group 1 and 20 of 38 patients in group 2. Sixteen of 25 discontinuations for toxicity (64%) in group 1 and 38 of 58 discontinuations for toxicity (66%) in group 2 were due to thrombocytopenia (Table 1).

### 3.6. Platelet transfusions and haemorrhages

The numbers of documented platelet transfusions (<4%) and haemorrhages were low (<3%). Three patients in group 1 had an intracranial haemorrhage, 2 in the context of grade 2 and 1 in the context of grade 3 thrombocytopenia. Six patients in group 2 had an intracranial haemorrhage, 3 with grade 1, 2 with grade 2 and 1 with grade 5 intracranial haemorrhage. No extracranial haemorrhage was reported in either group (Table 2).

### 3.7. Association of MGMT promoter methylation status with treatment modification induced by lomustine

Table S6 shows that, patients with MGMT promoter-methylated glioblastoma experienced more interference with study treatment than patients with MGMT promoter-unmethylated tumours. This was expected since these patients remain longer on lomustine owing to the better activity of lomustine in this patient population [8]. Lomustine was discontinued for thrombocytopenia in 2 (3%) versus 7 (11%) patients in group 1, and in 12 (12%) versus 16 (20%) patients in group 2, with tumours without versus with MGMT promoter methylation.

### 3.8. Risk factors for treatment-induced thrombocytopenia

To explore the association between the number of the lomustine cycle or other baseline characteristics and the odds of lomustine treatment modification due to thrombocytopenia, lomustine treatment cycle was dichotomised as first, second and third cycle, versus later cycles. In adjusted analysis, the number of cycles of lomustine treatment was not significantly associated with the odds of lomustine treatment modification for thrombocytopenia, presumably owing to the fact that dose reduction aims at preventing further episodes of thrombocytopenia and does so. Lower baseline platelet counts assessed as a continuous variable ( $p < 0.001$ ), female sex ( $p < 0.001$ ) and WHO performance status above 0 ( $p = 0.025$ ) were associated with higher odds of lomustine treatment modification by thrombocytopenia (Table S7), whereas interval to last prior chemotherapy cycle was not (Note S3).

### 3.9. Association of treatment modification by thrombocytopenia with PFS

In a risk-adjusted analysis, treatment modification by thrombocytopenia was not significantly associated with PFS. Maximal diameter of the tumour at baseline, steroid use at baseline and MGMT promoter methylation status were significantly associated with PFS. PFS was better in patients with smaller ( $\leq 40$  mm) initial tumour diameter ( $p = 0.005$ ), who did not take steroids at baseline ( $p = 0.035$ ), and who had tumours with MGMT promoter methylation ( $p < 0.001$ ) (Table 3). However, a more detailed analysis revealed markable differences by MGMT promoter methylation status and treatment (Table 4, Fig. 1): there was an association of treatment modification with superior PFS in patients with MGMT promoter unmethylated tumours, but this effect was largely driven by group 2 patients because only 5 patients in group 1 had a dose modification, owing to the low number of cycles of lomustine in this group. In contrast, patients with MGMT promoter methylated tumours experienced inferior PFS when treatment was modified for thrombocytopenia (Table 4, Note S4).



Table 3

Association between treatment modification by thrombocytopenia and outcome.

Variable	Progression-free survival				Overall survival			
	n = 429, n (%)	HR	95% CI	p-value	n = 505, n (%)	HR	95% CI	p-value
Unadjusted analysis								
Treatment modification by thrombocytopenia <sup>a</sup>								
No	310 (72.3)	1			381 (75.4)	1		
At least one episode	119 (27.7)	0.99	0.86–1.15	0.942	124 (24.6)	0.82	0.73–0.92	<0.001
Adjusted analysis								
Treatment modification by thrombocytopenia <sup>a</sup>								
No	310 (72.3)	1			381 (75.4)	1		
At least one episode	119 (27.7)	0.90	0.76–1.06	0.223	124 (24.6)	0.77	0.68–0.88	<0.001
Age category at baseline								
60 years or below	248 (57.8)	1			291 (57.6)	1		
More than 60 years	181 (42.2)	1.06	0.93–1.22	0.364	214 (42.4)	1.10	0.99–1.21	0.070
Sex								
Male	264 (61.5)	1			312 (61.8)	1		
Female	165 (38.5)	1.00	0.87–1.15	0.995	193 (38.2)	0.98	0.89–1.08	0.679
WHO performance status at baseline								
PS 0	137 (31.9)	1			171 (33.9)	1		
PS 1 or 2	292 (68.1)	1.00	0.86–1.16	0.969	334 (66.1)	0.98	0.88–1.09	0.675
Largest tumour diameter at baseline								
≤40 mm	242 (56.4)	1			275 (54.5)	1		
>40 mm	187 (43.6)	1.25	1.07–1.45	0.005	230 (45.5)	1.50	1.34–1.68	<0.001
Steroid use at baseline								
No	220 (51.3)	1			254 (50.3)	1		
Yes	209 (48.7)	1.18	1.01–1.37	0.035	251 (49.7)	1.69	1.50–1.89	<0.001
MGMT promoter								
Unmethylated	136 (31.7)	1			166 (32.9)	1		
Methylated	126 (29.4)	0.40	0.34–0.48	<0.001	143 (28.3)	0.60	0.52–0.67	<0.001
Undetermined/Not done	167 (38.9)	0.58	0.49–0.69	<0.001	196 (38.8)	0.82	0.73–0.92	<0.001
Surgery for recurrence								
No	338 (78.8)	1			405 (80.2)	1		
Yes	91 (21.2)	1.01	0.86–1.18	0.940	100 (19.8)	1.09	0.97–1.22	0.145

Abbreviations: MGMT O<sup>6</sup>-methylguanine DNA methyltransferase, N number of patients, PS performance status, WHO World Health Organization.<sup>a</sup> Lomustine treatment modification for thrombocytopenia.

### 3.10. Association of treatment modification due to thrombocytopenia with OS

Lomustine treatment modification by thrombocytopenia ( $p < 0.001$ ), smaller ( $\leq 40$  mm) initial tumour diameter ( $p < 0.001$ ), no baseline steroid use ( $p < 0.001$ )

and MGMT promoter methylation ( $p < 0.001$ ) were associated with superior OS (Table 3). Again, a more detailed analysis revealed notable differential associations by MGMT promoter methylation status and treatment: the association of treatment modification with superior OS in patients with MGMT promoter

Table 4

Association of treatment modification by thrombocytopenia with PFS by MGMT promoter methylation status and treatment, adjusted for baseline covariates.

Treatment (lomustine) modification by thrombocytopenia	Pooled			MGMT promoter unmethylated			MGMT promoter methylated		
	N (%)	HR (95% CI)	p-value	N (%)	HR (95% CI)	p-value	N (%)	HR (95% CI)	p-value
Pooled arms (LOM alone + LOM/BEV)									
No	N = 429			N = 136			N = 126		
Yes	310 (72.3)	1		107 (78.7)	1		79 (62.7)	1	
	119 (27.7)	0.90 (0.76–1.06)	0.223	29 (21.3)	0.39 (0.27–0.57)	<0.001	47 (37.3)	1.84 (1.44–2.35)	<0.001
LOM alone arm (group 1)									
No	N = 159			N = 43			N = 50		
Yes	125 (78.6)	1		38 (88.4)	1		33 (66.0)	1	
	34 (21.4)	0.62 (0.46–0.84)	0.002	5 (11.6)	0.27 (0.10–0.70)	0.007	17 (34.0)	1.73 (1.17–2.57)	0.006
LOM/BEV arm (group 2)									
No	N = 270			N = 93			N = 76		
Yes	185 (68.5)	1		69 (74.2)	1		46 (60.5)	1	
	85 (31.5)	1.09 (0.89–1.33)	0.431	24 (25.8)	0.41 (0.27–0.64)	<0.001	30 (39.5)	1.90 (1.34–2.68)	<0.001

Abbreviations: MGMT O<sup>6</sup>-methylguanine DNA methyltransferase, N number of patients.




Fig. 1. Progression-free survival and overall survival by MGMT status and treatment modification induced by thrombocytopenia per landmark/cycle. A: PFS in patients with unmethylated MGMT tumour, B: PFS in patients with methylated MGMT tumour, C: OS in patients with unmethylated MGMT tumour, D: OS in patients with methylated MGMT tumour. Landmark analysis performed at each cycle for PFS showed a trend for longer PFS in patients with an MGMT unmethylated glioblastoma when lomustine administration was modified by thrombocytopenia and a longer PFS in patients with a methylated glioblastoma when lomustine could be administered as planned. Landmark analysis performed at each cycle for overall survival showed a trend for longer overall survival in patients with an unmethylated glioblastoma when lomustine administration was modified by thrombocytopenia. The results of the stacked analysis performed pooling the six datasets are presented in Table S8.

unmethylated tumours remained, yet divergent associations became apparent for patients with *MGMT* promoter methylated tumours. The association of inferior OS with treatment modification for thrombocytopenia

resembled the observation made for PFS in group 2, but the opposite association was seen for OS in group 1 in that patients with treatment modification had superior OS despite inferior PFS (Table 5, Fig. 1, Note S5).

Table 5

Association of treatment modification by thrombocytopenia with OS by *MGMT* promoter methylation status and treatment, adjusted for baseline covariates.

Treatment (lomustine) modification by thrombocytopenia	Pooled			<i>MGMT</i> promoter unmethylated			<i>MGMT</i> promoter methylated		
	N (%)	HR (95% CI)	p-value	N (%)	HR (95% CI)	p-value	N (%)	HR (95% CI)	p-value
Pooled arms (LOM alone + LOM/BEV)									
No	N = 505			N = 166			N = 143		
Yes	381 (75.4)	1		136 (81.9)	1		94 (65.7)	1	
	124 (24.6)	0.77 (0.68–0.88)	<0.001	30 (18.1)	0.54 (0.41–0.70)	<0.001	49 (34.3)	1.05 (0.82–1.36)	0.697
LOM alone arm (group 1)									
No	N = 223			N = 67			N = 65		
Yes	188 (84.3)	1		62 (92.5)	1		48 (73.8)	1	
	35 (15.7)	0.69 (0.56–0.85)	<0.001	5 (7.5)	0.56 (0.33–0.95)	0.030	17 (26.2)	0.55 (0.35–0.85)	0.008
LOM/BEV arm (group 2)									
No	N = 282			N = 99			N = 78		
Yes	193 (68.4)	1		74 (74.7)	1		46 (59.0)	1	
	89 (31.6)	0.80 (0.68–0.95)	0.009	25 (25.3)	0.55 (0.40–0.76)	<0.001	32 (41.0)	1.71 (1.27–2.31)	<0.001

Abbreviations: MGMT O<sup>6</sup>-methylguanine DNA methyltransferase, N number of patients.



#### 4. Discussion

Chemotherapies used for the treatment of primary brain tumours frequently induce myelotoxicity, and thrombocytopenia represents the main cause of stopping alkylating agent chemotherapy for toxicity [1,11,12]. Consequences of thrombocytopenia include postponed chemotherapy courses, dose reductions, discontinuation of chemotherapy, as well as haemorrhages, including cerebral intratumoural bleeding, with increased risk in patients treated with corticosteroids [13,14]. Until recently, platelet transfusions were the only treatment option for severe chemotherapy-induced thrombocytopenia. The efficacy of a new thrombopoietin mimetic agent romiplostim (AMG 531, Nplate®, Amgen), to counteract chemotherapy-induced thrombocytopenia, has been shown in a phase II open label multicenter single arm phase II trial (NCT 02227576) in newly diagnosed glioblastoma [10].

Here, we used the clinical trial database of the EORTC trial 26101 [8] to estimate the impact of thrombocytopenia on lomustine exposure and outcome of patients with recurrent glioblastoma treated with lomustine alone or in combination with bevacizumab (Note S6).

The apparent association between drug-induced thrombocytopenia and improved outcome in patients with tumours lacking *MGMT* promoter methylation, mainly driven by patients in group 2, remains incompletely understood. Of note, lomustine is considered essentially inactive in this group of tumours [9]. Moreover, the overall short exposure to lomustine (Table S6) may render this analysis sensitive to chance findings (Figs. S3 and S4). The alternative explanation that lomustine may be detrimental in this group of patients requires further study although the outcome data of the BELOB trial that allow to compare bevacizumab alone with the combination of bevacizumab and lomustine do not support this notion [7]. Importantly, the positive association of treatment modification with outcome was largely driven by group 2 patients (Tables 4 and 5).

The occurrence of dose delay, dose reduction or treatment interruption was associated with inferior PFS in patients with *MGMT* promoter-methylated tumours. The association of lomustine modification induced by thrombocytopenia with inferior PFS in patients with *MGMT* promoter-methylated tumours appears to confirm the hypothesis that thrombocytopenia prevents adequate lomustine exposure. Although group 1 patients started with a higher dose of lomustine than group 2 patients, there was no difference in overall survival, potentially confounded by the cotreatment with bevacizumab in group 2 [8]. That patients with *MGMT* promoter-methylated tumours in group 1 who had treatment modification for thrombocytopenia experienced inferior PFS, but still better OS is a puzzling observation that revives the idea that myelosuppression may predict overall outcome [15,16], a hypothesis that we did not confirm at least in the

newly diagnosed setting [17]. Furthermore, temozolomide dose intensification in the newly diagnosed setting did not improve outcome [18]. Overall, our study may lend support to the hypothesis that individualised dosing may be superior to flat dosing based on body surface area and indicates that individual dose escalation schemes may make sense. We also provide a clear-cut post hoc rationale for the RIGOLETTO EORTC 1926 trial (NCT04933942) that explores romiplostim salvage in the setting of lomustine-induced thrombocytopenia.

Since bevacizumab is still used in combination with lomustine in various countries and centres, based on the clear superiority for PFS in the 26101 trial over lomustine alone, we included an analysis of this combination with some important observations. Lomustine was discontinued for thrombocytopenia more often with the combination, which may confirm that bevacizumab contributes to thrombocytopenia although patients also merely received more lomustine in this cohort (Table S6). It may also reflect the assumption that, at least in patients with tumours lacking *MGMT* promoter methylation, lomustine is perceived as not contributing to the treatment effect.

Our study has limitations. Although the data were assembled prospectively, the present analysis is a post hoc analysis, and items such as platelet transfusions and haemorrhages may have been underreported. Confounding effects of comedications could not be modelled. Data on number of cycles of temozolomide prior to treatment with lomustine were not available to explore a potential role as a risk factor. Further, the observations regarding the comparison of groups 1 and 2 need to be interpreted with caution. Group 1 started with a higher dose of lomustine, but longer PFS mediated by bevacizumab allowed patients in group 2 to receive a median of 3 rather than 1 lomustine cycles as in the control arm [8]. We cannot distinguish whether thrombocytopenia *per se* or the introduction of changes to the treatment regimen were prognostic in our adjusted analyses.

More individualised treatment regimens may have to be explored in a disease setting where standards of care have not been improved over decades. These findings encourage the evaluation of therapeutic approaches allowing to maintain the recommended doses of lomustine in patients with recurrent glioblastoma with *MGMT* promoter methylation, as previously proposed for allowing adequate temozolomide exposure in the newly diagnosed setting [10].

#### Funding

This publication was supported by the European Organisation for Research and Treatment of Cancer (EORTC) Brain Tumor Group. We are grateful to F. Hoffmann - La Roche from Switzerland for supporting

the independent EORTC Study 26101. Felix B. Oppong's work as Fellow at EORTC Headquarters was supported by a grant from the EORTC Brain Tumor Group.

### Author contributions

Experimental design and its implementation: ELR, FBO, TG, MW.

Acquisition, analysis, or interpretation of data: all authors.

Statistical analysis: FBO, TG.

Writing of the manuscript: all authors.

### Conflict of interest statement

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: ELR has received grant research from Bristol Meyer Squibb; honoraria for lectures or advisory board from Adastral, Bayer, Janssen, Leo Pharma, Pierre Fabre, and Seattle Genetics. FBO declares no conflict of interest. MvdB received honoraria for advisory boards from Genenta, Carthera, Boehringer, Astra Zeneca, Chimerix, and Agios. WW has received research support from Apogenix GmbH, Merck, Sharp & Dohme (MSD), Pfizer and Roche and honoraria for lectures from MSD. AAB declares no conflict of interest. MT declares no conflict of interest. MP has received research grants from Pfizer and Roche and honoraria for advisory boards from Bayer. AI declares Research grants from Carthera, Transgene, Sanofi, Servier, Nutritheragene, advisory board for Leo Pharma, Novocure and Boehringer Ingelheim Int, travel funding from Novocure, Carthera and Leo Pharma outside the submitted work. PC has received grant research from Astra Zeneca; honoraria for lectures or advisory board from Bayer, BMS, Leo Pharma, Merck, MSD, Rakuten Medical and Takeda. MP has received honoraria for lectures, consultation or advisory board participation from the following for-profit companies: Bayer, Bristol-Myers Squibb, Novartis, Gerson Lehrman Group (GLG), CMC Contrast, GlaxoSmithKline, Mundipharma, Roche, BMJ Journals, MedMedia, Astra Zeneca, AbbVie, Lilly, Medahead, Daiichi Sankyo, Sanofi, Merck Sharp & Dome, Tocagen, Adastral, Gan & Lee Pharmaceuticals. VG declares no conflict of interest. TG declares no conflict of interest. MW has received research grants from Philogen and Quercis, and honoraria for lectures or advisory board participation or consulting from Adastral, Medac, Nerviano Medical Sciences, Novartis, Orbus, Philogen and y-Mabs.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejca.2022.10.006>.

### References

- [1] Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 2005;352:987–96. <https://doi.org/10.1056/NEJMoa043330>.
- [2] Weller M, van den Bent M, Preusser M, Le Rhun E, Tonn JC, Minniti G, et al. EANO guidelines on the diagnosis and treatment of diffuse gliomas of adulthood. *Nat Rev Clin Oncol* 2021;18:170–86. <https://doi.org/10.1038/s41571-020-00447-z>.
- [3] Wick W, Puduvalli VK, Chamberlain MC, van den Bent MJ, Carpentier AF, Cher LM, et al. Phase III study of enzastaurin compared with lomustine in the treatment of recurrent intracranial glioblastoma. *J Clin Oncol* 2010;28:1168–74. <https://doi.org/10.1200/JCO.2009.23.2595>.
- [4] Batchelor TT, Mulholland P, Neyns B, Nabors LB, Campone M, Wick A, et al. Phase III randomized trial comparing the efficacy of cediranib as monotherapy, and in combination with lomustine, versus lomustine alone in patients with recurrent glioblastoma. *J Clin Oncol* 2013;31:3212–8. <https://doi.org/10.1200/JCO.2012.47.2464>.
- [5] Lombardi G, De Salvo GL, Brandes AA, Eoli M, Rudà R, Faedi M, et al. Regorafenib compared with lomustine in patients with relapsed glioblastoma (REGOMA): a multicentre, open-label, randomised, controlled, phase 2 trial. *Lancet Oncol* 2019;20:110–9. [https://doi.org/10.1016/S1470-2045\(18\)30675-2](https://doi.org/10.1016/S1470-2045(18)30675-2).
- [6] Wen PY, Mellinger IK, Buxton MB, Cavenee WK, Colman H, Groot JFD, et al. GBM AGILE: a global, phase 2/3 adaptive platform trial to evaluate multiple regimens in newly diagnosed and recurrent glioblastoma. [n.d].
- [7] Taal W, Oosterkamp HM, Walenkamp AME, Dubbink HJ, Beerepoot LV, Hanse MCJ, et al. Single-agent bevacizumab or lomustine versus a combination of bevacizumab plus lomustine in patients with recurrent glioblastoma (BELOB trial): a randomised controlled phase 2 trial. *Lancet Oncol* 2014;15:943–53. [https://doi.org/10.1016/S1470-2045\(14\)70314-6](https://doi.org/10.1016/S1470-2045(14)70314-6).
- [8] Wick W, Gorlia T, Bendszus M, Taphoorn M, Sahm F, Harting I, et al. Lomustine and bevacizumab in progressive glioblastoma. *N Engl J Med* 2017;377:1954–63. <https://doi.org/10.1056/NEJMoa1707358>.
- [9] Weller M, Le Rhun E. How did lomustine become standard of care in recurrent glioblastoma? *Cancer Treat Rev* 2020;87:102029. <https://doi.org/10.1016/j.ctrv.2020.102029>.
- [10] Le Rhun E, Devos P, Houillier C, Cartalat S, Chinot O, Di Stefano AL, et al. Romiplostim for temozolomide-induced thrombocytopenia in glioblastoma: the PLATUM trial. *Neurology* 2019;93:e1799–806. <https://doi.org/10.1212/WNL.0000000000008440>.
- [11] Stupp R, Hegi ME, Gorlia T, Erridge SC, Perry J, Hong Y-K, et al. Cilengitide combined with standard treatment for patients with newly diagnosed glioblastoma with methylated MGMT promoter (CENTRIC EORTC 26071-22072 study): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2014;15:1100–8. [https://doi.org/10.1016/S1470-2045\(14\)70379-1](https://doi.org/10.1016/S1470-2045(14)70379-1).
- [12] Weller M, Butowski N, Tran DD, Recht LD, Lim M, Hirte H, et al. Rindopimut with temozolomide for patients with newly diagnosed, EGFRvIII-expressing glioblastoma (ACT IV): a randomised, double-blind, international phase 3 trial. *Lancet Oncol* 2017;18:1373–85. [https://doi.org/10.1016/S1470-2045\(17\)30517-X](https://doi.org/10.1016/S1470-2045(17)30517-X).
- [13] Gerber DE, Grossman SA, Zeltzman M, Parisi MA, Kleinberg L. The impact of thrombocytopenia from temozolomide and radiation in newly diagnosed adults with high-grade gliomas. *Neuro Oncol* 2007;9:47–52. <https://doi.org/10.1215/15228517-2006-024>.
- [14] Garcia Rodríguez LA, Hernández-Díaz S. The risk of upper gastrointestinal complications associated with nonsteroidal anti-

- inflammatory drugs, glucocorticoids, acetaminophen, and combinations of these agents. *Arthritis Res* 2001;3:98–101. <https://doi.org/10.1186/ar146>.
- [15] Grossman SA, Ellsworth S, Campian J, Wild AT, Herman JM, Laheru D, et al. Survival in patients with severe lymphopenia following treatment with radiation and chemotherapy for newly diagnosed solid tumors. *J Natl Compr Cancer Netw* 2015;13:1225–31. <https://doi.org/10.6004/jnccn.2015.0151>.
- [16] Saito T, Sugiyama K, Hama S, Yamasaki F, Takayasu T, Nosaka R, et al. Prognostic importance of temozolomide-induced neutropenia in glioblastoma, IDH-wildtype patients. *Neurosurg Rev* 2018;41:621–8. <https://doi.org/10.1007/s10143-017-0903-3>.
- [17] Le Rhun E, Oppong FB, Vanlancker M, Stupp R, Nabors B, Chinot O, et al. Prognostic significance of therapy-induced myelosuppression in newly diagnosed glioblastoma. *Neuro Oncol* 2022;noac070. <https://doi.org/10.1093/neuonc/noac070>.
- [18] Gilbert MR, Wang M, Aldape KD, Stupp R, Hegi ME, Jaeckle KA, et al. Dose-dense temozolomide for newly diagnosed glioblastoma: a randomized phase III clinical trial. *J Clin Oncol* 2013;31:4085–91. <https://doi.org/10.1200/JCO.2013.49.6968>.