



Universiteit
Leiden
The Netherlands

Specific activation of ERK pathways by chitin oligosaccharides in embryonic zebrafish cell lines

Snaar, B.E.; Krens, S.F.G.; Robina, I.; Wang, L.X.; Spaink, H.P.

Citation

Snaar, B. E., Krens, S. F. G., Robina, I., Wang, L. X., & Spaink, H. P. (2003). Specific activation of ERK pathways by chitin oligosaccharides in embryonic zebrafish cell lines. *Glycobiology*, 13(10), 725-732. doi:10.1093/glycob/cwg103

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3748443>

Note: To cite this publication please use the final published version (if applicable).

Specific activation of ERK pathways by chitin oligosaccharides in embryonic zebrafish cell lines

Bogusława E. Snaar-Jagalska^{1,2}, Simon F. G. Krens²,
Inmaculada Robina³, Lai-Xi Wang⁴, and Herman
P. Spaink²

²Institute of Biology, Leiden University, Clusius Laboratory, Wassenaarseweg 64, 2333 AL Leiden, The Netherlands; ³Departamento de Química Orgánica, Facultad de Química, Universidad de Sevilla, Apartado 553, E-41071 Sevilla, Spain; ⁴Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

Received on May 8, 2003; revised on July 8, 2003; accepted on July 15, 2003

Chitin oligosaccharides (COs) play a role in plant development and are presumed to affect body plan formation during vertebrate embryogenesis. The mechanisms of COs recognition and cellular processes underlying embryonic development are still not understood. We analyze the possible link with the mitogen-activated protein kinase pathway that is conserved in evolution through the plant and animal kingdom and has been implicated in diverse cellular processes, including cell growth, proliferation, differentiation, survival, and vertebrate development. We show that *in vivo* stimulation of embryonic zebrafish cells ZF13 and ZF29 with chitin tetrasaccharides at 10^{-9} M concentration transiently induced activation/phosphorylation of extracellular regulated kinases (ERKs), with a maximum after 15 min. Furthermore the biological specificity of chitin tetrasaccharides and various derivatives was examined. The replacement of one or two GlcNAc residues of the chitin backbone by glucose and fucosylation of chitin tetrasaccharides at the reducing terminus caused a complete loss of their activity. We also tested a chitin tetrasaccharide analogue in which the oxygen atoms in glycosidic linkages were replaced by sulfur atoms. This analog, which could not be enzymatically hydrolyzed, was as potent an inducer as chitin tetrasaccharide. These results suggest that the observed activation of ERKs is chitin tetrasaccharide-specific and does not require further enzymatic processing. We examined possible signaling pathways leading to ERK activation by COs by use of phosphospecific antibodies and inhibitors. We conclude that a high-affinity CO receptor system exists that links to the Raf, MEK, and ERK pathway in zebrafish cells.

Key words: chitin oligosaccharides/ERK/zebrafish cells

Introduction

Chitin, a repeating $\beta(1 \rightarrow 4)$ -linked homopolymer of the monosaccharide *N*-acetylglucosamine (GlcNAc), is one of

the most widespread and abundant molecules in the biosphere (Varki, 1996). Chitin oligosaccharide (CO) derivatives play a role in the leguminous plant development and possibly also in vertebrate embryogenesis (Bakkers *et al.*, 1997). Rhizobial NodC protein, responsible for CO synthesis, is homologous to the developmentally regulated DG42/HAS protein of vertebrates (Bulawa and Wasco, 1991; Semino and Robbins, 1995). Members of the DG42/HAS family have been shown to be involved in the biosynthesis of both hyaluronan and COs (DeAngelis and Acbyuthan, 1996; Meyer and Kreil, 1996; Semino *et al.*, 1996; Yoshida *et al.*, 2000; Van der Holst *et al.*, 2001). A possible biological function for COs was elucidated by microinjection studies of fertilized zebrafish eggs with anti-DG42 antiserum and the rhizobial NodZ fucosyltransferase (Bakkers *et al.*, 1997). These treatments led to severe defects in trunk and tail development, suggesting that COs may function as signaling molecules in cell growth, differentiation, and development of vertebrates (Semino *et al.*, 1998; Semino and Allende, 2000).

Mammalian cells respond to a variety of extracellular stimuli via activation of specific mitogen-activated protein kinases (MAPKs) that orchestrate the transduction of the signal from receptors at the cell surface to the nucleus and play a major role in the integration of multiple biological responses. Three main classes of MAPK are recognized: the classic MAPK, also known as extracellular-regulated kinases, ERK1 and ERK2, and c-Jun N-terminal kinase (JNK) and p38 MAPK, which are activated by dual phosphorylation at neighboring threonine and tyrosine residues in the activation loop. Dephosphorylation of either residue results in MAPK inactivation (Johnson and Lapadat, 2002). The general scheme of ERK activation involves a cascade of phosphorylation events initiated by stimulation of the Ras proto-oncogene following activation of growth factor receptors. The cascade starts with the activation of one or more Raf family kinases, which phosphorylate and activate the MAPK kinases (MEK1/2). MEK in turn catalyzes dual phosphorylation and activation of ERKs (Chang and Karin, 2001). Once activated, ERKs accumulate in the nucleus by as-yet undefined mechanisms that seem to involve the dissociation of ERK from MEK (Fukuda *et al.*, 1997) and/or a blockade of its export from the nucleus by neosynthesized nuclear anchors (Lenormand *et al.*, 1998).

The ERK pathway has been implicated in diverse cellular processes, including cell growth, proliferation, differentiation, and survival (Marshall, 1995; Ballif and Blenis, 2001). Hence, finely tuned regulation of ERK activation is essential in conveying appropriate signals. The intensity, duration, and subcellular localization of ERK activation are

¹To whom correspondence should be addressed; e-mail: jagalska@rulbim.leidenuniv.nl

well regulated. Scaffolding proteins and docking sites provide the means to avoid cross-activation between MAPK signaling pathways and permit precise and even cell-specific subcellular localization of ERKs (Pouysségur *et al.*, 2002). The variable responses elicited by this cascade in different cell types are also presumably determined by the cell-specific combination of downstream substrates.

Activated ERK phosphorylates numerous substrates in all cellular compartments (Lewis *et al.*, 1998). More than 50 different ERK substrates have been identified so far. These include ubiquitous or lineage-restricted transcribed factors, the kinases RSK and MNK and proteins involved in nucleotide biosynthesis, cytoskeleton organization, ribosomal transcription, and membrane traffic (Sturgill *et al.*, 1988; Marais *et al.*, 1993; Treisman, 1996; Fukunaga and Hunter, 1997; Lewis *et al.*, 2000; Stefanovsky *et al.*, 2001). RSKs, 90-kDa ribosomal S6 serine/threonine kinases family (also known as MAPK-activated protein kinase-1, MAPKAP-K1) are activated by ERKs *in vitro* and *in vivo* via phosphorylation (Sturgill *et al.*, 1988; Dalby *et al.*, 1998).

The presence and function of MAPK pathways in zebrafish has been recently described. The insulin-like growth factor stimulates zebrafish cell proliferation by activating MAPK and PI3-kinase-signaling pathways (Pozios *et al.*, 2001), while Fgf/MAPK signaling is required for development of the subpallial telencephalon and somite boundary formation in zebrafish embryos (Shinya *et al.*, 2001; Sawada *et al.*, 2001). Hence components upstream and downstream of the specific MAPK cascades in zebrafish are not known yet.

Here we show that chitin tetrasaccharides rapidly and specifically activate the ERK pathway in embryonic zebrafish cells. These findings provide strong evidence that CO signaling is initiated on the plasma membrane via activation of high-affinity oligosaccharide receptor system, which transmits the signal to nucleus probably using the Ras, Raf, MEK, ERK cascade.

Results

Activation of ERK by chitin tetrasaccharides

To investigate a possible role of COs as signaling molecules in vertebrate development, the embryonic zebrafish cell lines ZF13 and ZF29 were used as a model (Peppelenbosch *et al.*, 1995) and the canonical MAPK pathway was selected as a well-characterized junction between extracellular signals and integrated biological responses. EGF was used as a control of functionality in the system.

Zebrafish cells were serum starved for 24 h, stimulated with 5×10^{-6} M chitin tetrasaccharides and EGF for the indicated times, lysed, and processed for western blotting. The activated/phosphorylated form of MAPKs, ERKs was monitored by specific dual phospho-p44/42 MAP kinase antibodies (dp ERK), which detect p44 and p42 MAP kinase (ERK1 and ERK2) only when catalytically activated by phosphorylation at Thr202 and Tyr204. The antibody does not appreciably cross-react with the corresponding phosphorylated threonine and tyrosine residues of either JNK/SAPK or p38 MAPK homologs. It does not




Fig. 1. Time-dependent and specific activation of ERKs by COs and EGF. Serum-starved zebrafish cells (ZF13) were stimulated with 5×10^{-6} M COs (A) and EGF (B) for increasing periods of time (0–60 min). ERK activation were monitored by immunoblotting of total cell lysates with specific phospho-MAPK (Thr202/Tyr204) antibody (dp ERK). The total amount of ERK was visualized using an ERK1 antibody. These results are representative examples of five experiments and were identical for ZF29 cells (data not shown). The lysates obtained from not stimulated (0) and 15 min stimulated (15') cells with 5×10^{-6} M CO and protein marker (PM) were subjected to western blotting. The specificity of ERK phosphorylation was determined with anti-phospho-ERK1/2, anti-phospho-p38 MAPK, and phospho-SAPK/JNK antibodies (C).

cross-react with up to 2 μ g nonphosphorylated p44/42 MAPK. The total protein level in all performed experiments was control by ERK1 antibody, which reacts with ERK1 (p44) and, to a lesser extent ERK2 (p42) by western blotting. Bands intensities were quantified by densitometry, and ERK activity was compared to the basal level at $t = 0$.

In vivo stimulation of ZF13 and ZF29 with chitin tetrasaccharide and EGF transiently induced the 8–15-fold activation/phosphorylation of ERKs, with a maximum after 15 min (Figure 1). The half maximal response was obtained at 10^{-9} M concentration (Figure 2). Although dp ERK antibodies are able to recognize both phosphorylated




Fig. 2. Structures of CO derivatives. CO: chitin oligosaccharide; thioCO: CO in which the oxygen atoms of the $\beta(1 \rightarrow 4)$ linkages are substituted by sulfur atoms; CO-PNP: CO with PNP modification at C1 position; CO(Glc)-MP and CO(2Glc)-MP: one or two GlcNAc residues are replaced by glucose, respectively, and with MP modification at C1 position; CO-fuc: fucosylated CO.

ERK1 and ERK2 in all experiments, one band of 44 kDa was preferentially detected, suggesting that ERK1 is mainly activated by COs and EGF. The anti-phospho-p38 MAPK and anti-phospho-SAPK/JNK antibodies showed absence of cross-reaction in the CO activation process observed with phospho-ERK1/2 antibody, ensuring the specificity in the western blots (Figure 1C).

ERK activation is specific for chitin tetrasaccharides, CO, and thio-CO

We examined the biological specificity of chitin tetrasaccharides and various derivatives (Figure 2). The replacement of one or two GlcNAc residues of the chitin

backbone by glucose and fucosylation of chitin tetrasaccharides by NodZ protein abolished the biological activity of CO(2glc)-MP and CO-fuc in the tested concentration of 10^{-7} M (Figure 3A). This loss of activity was apparently not due to modifications at C1 position of the chitin tetrasaccharide molecule because *para*-nitrophenyl chitin (CO-PNP) that is structurally very similar to *para*-methoxyphenyl (MP) modified at this position showed normal activation of ERK (Figure 2 and 3A). The C2 and C3 position of CO molecule seem to be essential for its biological property. We also tested a chitin tetrasaccharide analog, thioCO, in which the oxygen atoms in glycosidic linkages were replaced by sulfur atoms. This analog, which cannot be enzymatically hydrolyzed (Wang and Lee, 1995)




Fig. 3. Dose dependence and specificity of ERK activation by CO and derivatives. Serum-starved zebrafish cells (ZF13) were stimulated with various concentrations of CO, CO(2glc)-MP or CO-fuc (A) and thioCO (B) for 15 min. Phosphorylated and nonphosphorylated forms of ERK were detected by dp ERK and ERK1 antibodies. These results are representative of three experiments. 10^{-7} M EGF was used as a control in A and B.

was as potent an inducer as chitin tetrasaccharide. The half maximal responses, induced by 15 min treatment with CO and thioCO, were obtained at 10^{-9} M concentration for both compounds (Figure 3A, 3B). These results suggest that the observed activation of ERKs is chitin tetrasaccharide-specific and does not require further enzymatic processing.

Signaling pathway to ERK activation by chitin tetrasaccharides

In general, ERK activation involves a cascade of phosphorylation events initiated by stimulation of the Ras proto-oncogene following activation of growth factors receptors. The cascade starts with the activation of Raf, which phosphorylates MAPK kinase (MEK) on two serine residues, and then ERK is dually phosphorylated on a tyrosine and threonine residues by MEK. Activated ERK phosphorylates specific serines and threonines of target protein substrates. Several downstream targets of ERK1/2 have been identified, for instance, p90RSK is activated by ERK1/2 via phosphorylation at Ser380 (Dalby *et al.*, 1998). To examine a possible signaling pathway to ERKs activation by chitin tetrasaccharides, we made use of phospho-Raf (Ser259), phospho-MEK1/2 (ser217/221), and phospho-p90RSK (Ser380) antibodies to follow expected phosphorylation events in ERK activation cascade. The activation of MEK1/2 and subsequently ERK were blocked by a highly selective inhibitor of MEK1 and MEK2, UO126, at 10^{-5} M concentration (Crews *et al.*, 1992). The result in Figure 4A show that CO, thioCO, and EGF induced phosphorylation/activation of Raf, MEK1/2, and p90RSK with the same kinetics and specificity observed for ERK activation.




Fig. 4. Signaling pathway involved in ERK activation by CO and CO derivatives. Serum-starved zebrafish cells (ZF13) were stimulated with 10^{-7} M of CO, CO derivatives, and EGF for 15 min. Lysates were subjected to western blotting with phosphospecific p-Raf, p-MEK1/2, and p-p90RSK antibodies (A). Inhibition of CO and EGF induced activation of ERK by MEK inhibitor. Serum-starved zebrafish cells (ZF13) were pretreated for 1 h with 10^{-5} M of a highly selective MEK1/2 inhibitor (UO126) prior to 15 min stimulation of ERK phosphorylation by 10^{-7} M CO and EGF (B). Phosphorylated and nonphosphorylated forms of ERKs were detected by dp ERK and ERK1 antibodies.

Again CO-fuc and CO(2glc)-MP were found to be not active (Figure 3A, 4A). Preincubation of cells for 1 h with MEK1/2 inhibitor before stimulation prevented stimulation of ERK phosphorylation by CO and EGF (Figure 4B). From this we conclude that chitin oligosaccharides are recognized by a high-affinity receptor system that activates Raf, MEK, and ERK pathway in these zebrafish cell lines.

Discussion

Many human diseases, including the formation of tumors and their subsequent metastatic spread through the body, are caused by defects in carbohydrates recognition (Schuster and Nelson, 2000). Furthermore, all human pathogens also naturally produce a great variety of carbohydrates and use an unknown number of these molecules to modulate the innate immune response of the host (Aderem and Ulevitch, 2000). Most of these carbohydrates are polymers either in free form or linked to proteins. However, evidence exists that also small oligosaccharides, such as COs, play a crucial role in animal development. The genetic and cellular basis of the CO recognition mechanisms underlying their function in embryonic development, tumorigenesis, or infectious diseases is still not understood. The zebrafish model system offers novel opportunities to study the interplay between signal perception, transmission, and

development. In this study we address whether COs are able to function as signaling molecules to activate the canonical ERK pathway in embryonic zebrafish cell cultures. The MAPK/ERK pathway was chosen as biological tool in this project because it plays a major role in the integration of multiple biological responses controlling development (Pouyssegur *et al.*, 2002). The activation of ERKs and downstream components was monitored by phospho-specific antibodies because the general scheme of activation involves a cascade of phosphorylation events following activation of specific receptors by appropriate signal.

First, we found that chitin tetrasaccharides transiently induced phosphorylation of ERKs, probably ERK1, with a maximum after 15 min stimulation, at 10^{-9} M concentration. This response was very specific for the structure of the glycan backbone. The replacement of one or two GlcNAc residues by glucose and fucosylation by the NodZ protein resulted in a loss of their biological activity. In contrast, an analog of CO, thioCO, which could not be enzymatically hydrolyzed, was as potent an inducer as COs, excluding a role of possible degradation products. Furthermore, we also examined the signaling pathway to ERK activation by COs and concluded that a high-affinity oligosaccharide receptor system exists that transduces the signal to Raf, MEK, and ERK. Once activated/phosphorylated ERK phosphorylates p90RSK kinase, known as MAPK-activated protein kinase-1, MAPKAP-K1. Phosphorylated RSKs in turn activate transcription factors, such as CREB, and are involved in cell cycle control, memory formation, and suppression of apoptotic cell death (Nebreda and Gavin, 1999). Moreover, inactivation mutations in the gene encoding human RSK2 are associated with Coffin-Lowry syndrome, a disease that results in severe mental retardation as well as progressive skeletal deformation (Treisman, 1996). The recently discovered RSK4 is deleted in patients with X-linked mental retardation (XLMR) and may be a candidate XLMR gene (Yntema *et al.*, 1999).

Our results strongly support the model shown in Figure 5, but final genetic identification of all components of CO signaling remain to be determined. Several lines of evidence in different cell types indicate that p90RSKs are activated by MAPKs *in vivo* via a Ras-dependent protein kinase cascade that is triggered by growth factors or tumor-promoting phorbol esters (Alessie *et al.*, 1995). Moreover, other physiological substrates of p90RSK have been identified with different effects in development, cell cycle, suppression of apoptotic cell death, and progressive skeletal deformation (Treisman, 1996; Nebreda and Gavin, 1999). These findings promote future studies of p90RSK and its physiological substrates in zebrafish cells and embryos.

We recently cloned two zebrafish ERK genes in which the phosphorylation consensus is identical to the mammalian species and forms the expected epitope recognized by the dp ERK antibody. With a morpholino knock-down approach we plan to find specific CO targets and study their function in zebrafish cells and embryo development. Recent studies in plants have identified leucine-rich repeat (LRR) receptors involved in recognizing carbohydrate microbial factors, such as the Nod factors (Asai *et al.*, 2002; Stracke *et al.*, 2002; Endre *et al.*, 2002). LRR domains are also found in plants as part of disease resistance




Fig. 5. Model for specific activation of ERK pathway by COs in embryonic zebrafish cells. Chitin tetrasaccharides probably activate a high-affinity receptor system (Rx) that transduces the signal to phosphorylation of Raf, MEK1/2, ERK1/2, and p90RSK. An arrow indicates an alternative path through the membrane.

casades, where they function as receptors important for innate immunity. In animals, the relatives of the plant receptors, so called Toll-like receptors, are also involved in the recognition of microbes, meaning that a similar carbohydrate-recognition mechanism could underlie the plant and animal innate immune system (Beutler, 2002; Spalink, 2002). Recently we identified ~20 Toll-like receptors in the zebrafish database from Sanger Centre (unpublished data). Currently we study their expression during embryogenesis and in response to treatment of cell lines with different carbohydrates to elucidate their role in CO signaling cascade.

In future research we aim to identify CO receptors and link them to downstream pathways, such as the ERK pathway, to fully understand CO function in vertebrate development.

Materials and methods

Materials

Polyclonal phospho-p44/42 MAP kinase antibody, phospho-ERK1/2 pathway sampler kit, phospho-p38 MAPK antibody, phospho-SAPK/JNK antibody, and MEK1/2 inhibitor (U0126) were purchased from Cell Signaling Technology (Leusden, The Netherlands). Control ERK

antibody K-23 was from Santa Cruz Biotechnology (Santa Cruz, CA). Secondary antibody, enhanced chemiluminescence kit, and rainbow molecular weight markers were from Amersham Life Science (Roosendaal, The Netherlands). Murine EGF was obtained from Invitrogen (Breda, The Netherlands). Leibovitz L-15 medium (L-15) and fetal calf serum were purchased from Invitrogen.

Chitin tetrasaccharides were purchased from Seikagaku (Tokyo). Replacement of one or two central GlcNac residues of the chitin tetrasaccharide by Glc unit was received as described (Wang and Lee, 1995; Robina *et al.*, 2002; Southwick *et al.*, 2002). Fucosylation of COs by NodZ protein was performed as described previously (Bakkers *et al.*, 1997).

The syntheses of thioCO

The syntheses of di-*N*-acetyl-4-thiochitobiose (TG₂), α -methyl di-*N*-acetyl-4-thiochitobioside (MTG₂), α -methyl tri-*N*-acetyl-4,4'-dithiochitotrioside (MTG₃), α -methyl tetra-*N*-acetyl-4,4',4''-trithiochitotetraoside (MTG₄), and β -methyl tri-*N*-acetyl-4,4'-dithiochitotrioside (β -MTG₃) were reported in Wang and Lee (1995). The reducing thiochito-oligomers, tri-*N*-acetyl-4,4'-dithiochitotriose (TG₃) and tetra-*N*-acetyl-4,4',4''-trithiochitotetraose (TG₄), were prepared as follows.

TG₃. The peracetylated 4,4'-dithiochitotriose (20 mg) (Wang and Lee, 1995) was de-*O*-acetylated with sodium methoxide in methanol (10 mM, 5 ml) at room temperature for 6 h. Water (1 ml) was added and the solution was neutralized with Dowex 50w-x8 resin (H⁺ form). After removal of the resin by filtration, the filtrate was concentrated. The trisaccharide was dissolved in a small amount of water and applied to a Sephadex G-10 column (1.5 × 90 cm), which was preequilibrated and eluted with 50 mM acetic acid. Fractions containing the trisaccharide were pooled and lyophilized to give 15.5 mg TG₃ (yield: 81%). ¹H-nuclear magnetic resonance (NMR) of TG₃ (500 MHz, D₂O, 60°C): δ 5.248 (d, J = 3.4 Hz, 0.35 H, H-1 α), 4.776–4.732 (two pairs of d, J = 10.2 Hz, 2 H, H-1',1''), 4.695 (d, J = 8.4 Hz, 0.65 H, H-1 β), 4.057–3.470 (m, 16 H, sugar protons), 2.957–2.856 (two pairs of t, 2 H, J = 10.4 Hz, H-4,4'), 2.059–2.049 (singlets, 9 H, 3 *N*-acetyl).

TG₄. The α -methyl 4,4',4''-trithiochitotetraoside (4.5 mg) (Wang and Lee, 1995) was per-*O*-acetylated with acetic anhydride-pyridine (1:1, 1.5 ml) at room temperature for 5 h. Ice water (2 ml) was added, and the solution was evaporated to dryness. Trace of volatiles was removed by coevaporation of the residue with toluene (3 × 2 ml) to give 7.5 mg of the peracetylated α -methyl 4,4',4''-trithiochitotetraoside. The compound obtained was subjected to acetylation with Ac₂O–AcOH–H₂SO₄ (8:2:0.1, 2 ml) to selectively remove the 1-*O*-methyl group. After acetylation at room temperature for 12 h, the solution was cooled to 0°C, and a cold solution of aqueous sodium acetate (0.2 M, 3 ml) was added. The mixture was evaporated to dryness and the residue was partitioned between CHCl₃ (10 ml) and water (2 ml). The organic layer was separated and successively washed with 5% hydrochloric acid (1 ml), saturated sodium

bicarbonate (1 ml), and water (2 × 1 ml), dried with anhydrous sodium sulfate, and filtered. The filtrate was evaporated to afford the peracetylated 4,4',4''-trithiochitotetraose as a white solid, which was subsequently de-*O*-acetylated with sodium methoxide in methanol (10⁻² M, 5 ml). The product was purified by gel filtration with a Sephadex G-10 column the same way as described for the preparation of TG₃ to give 3.5 mg TG₄ (overall yield in three steps: 79%). ¹H-NMR of TG₄ (500 MHz, D₂O, 60°C): δ 5.246 (d, J = 3.4 Hz, 0.39 H, H-1 α), 4.769–4.725 (3 pairs of d, J = 10.3 Hz, 3 H, H-1',1'',1'''), 4.693 (d, J = 8.0 Hz, 0.61 H, H-1 β), 4.059–3.467 (m, 21 H, sugar protons), 2.945–2.872 (three pairs of t, 3 H, J = 10.3 Hz, H-4,4',4''), 2.050–2.047 (singlets, 12 H, 4 *N*-acetyl).

Cell lines

Cell lines ZF13 and ZF29c-1 were obtained from Hubrecht laboratory. These cells have been derived from dechorionated, disaggregated, 20-h-old zebrafish embryos. They have a fibroblast-like morphology (Peppelenbosch *et al.*, 1995). For our experiments cells were grown at 25°C in 2 ml 67% L-15 medium supplemented with 10% fetal calf serum in 24-well plates (Greiner, Alphen aan de Rijn, The Netherlands) until the plates became 50% confluent. Before stimulation experiments, the medium was changed to serum-free medium (SFM) for 18–24 h to reduce the basal level of phosphorylation. This SFM was then replaced with 1 ml SFM plus indicated stimulus for various times. Stimulation was terminated by quick replacement of SFM for 600 μ l, 65°C sodium dodecyl sulfate (SDS) sample buffer (62.5 mM Tris–HCl, 3% w/v SDS, 10% glycerol, 50 mM dithiothreitol, 0.01% w/v bromophenol blue). The cells were scraped off the dish, and the cell lysates were transferred to a new tube, boiled for 3 min, and separated on 10% polyacrylamide slab gels (10 μ g protein per lane).

Western immunoblotting

SDS-polyacrylamide gel electrophoresis was performed as described by Laemmli (1970). After electrophoresis, proteins were transferred to nitrocellulose membrane (Schleicher & Schuell, Den Bosch, The Netherlands) by western blotting. The membranes were blocked in 5% w/v nonfat dry milk in Tris-buffered saline-Tween 20 (TBST). The blots were incubated with a 1:1000 to 1:2000 dilution of the indicated antibody in TBST with 3% bovine serum albumin (Sigma, St. Louis, MO) for 1 h at room temperature or overnight at 4°C. Signal was detected using a 1:5000 to 1:10,000 dilution of horseradish peroxidase-conjugated anti-rabbit antibodies and the enhanced chemiluminescence method (Amersham).

Acknowledgments

We are grateful to Dr. D. Zivkovic for ZF 13 and ZF 29 cells. We thank Dr. N. Stuurman for stimulating discussion and S. Wijting for maintenance of the cell cultures.

Abbreviations

CO, chitin oligosaccharides derivatives; dp ERK, phospho-MAPK antibody; EGF, epidermal growth factor; ERK,

extracellular-regulated kinase; JNK, c-Jun N-terminal kinase; LRR, leucine-rich repeat; MAPK, mitogen-activated protein kinase; MEK, MAPK or ERK kinase; MP, *para*-methoxy-phenyl; NMR, nuclear magnetic resonance; p90RSK, MAPK-activated protein kinase; PNP, *para*-nitrophenyl; SDS, sodium dodecyl sulfate; SFM, serum-free medium, TBST, Tris-buffered saline-Tween 20.

References

- Aderem, A. and Ulevitch, R.J. (2000) Toll-like receptors in the induction of the innate immune response. *Nature*, **406**, 782–787.
- Alessie, D.R., Cuenda, A., Cohen, P., Dudley, D.T., and Saltiel, A.R. (1995) PD 098059 is a specific inhibitor of the activation of mitogen-activated protein kinase kinase *in vitro* and *in vivo*. *J. Biol. Chem.*, **270**, 27489–27494.
- Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W., Gomez-Gomez, L., Boller, T., Ausubel, F.M., and Sheen, J. (2002) MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature*, **415**, 977–983.
- Bakkers, J., Semino, C.E., Stroband, H., Kijne, J.W., Robbins, P.W., and Spaink, H.P. (1997) An important role for oligosaccharides during early embryo development of carp and zebrafish. *Proc. Natl. Sci. USA*, **94**, 7982–7986.
- Ballif, B.A., and Blenis, J. (2001) Molecular mechanisms mediating mammalian mitogen-activated protein kinase (MAPK) kinase (MEK)-MAPK cell survival signals. *Cell Growth Diff.*, **12**, 397–408.
- Beutler, B. (2002) Toll-like receptors: how they work and what they do. *Curr. Opin. Hematol.*, **9**, 2–10.
- Bulawa, C.E. and Wasco, W. (1991) Chitin and nodulation. *Nature*, **353**, 710.
- Chang, L. and Karin, M. (2001) Mammalian MAP kinase signalling cascades. *Nature*, **410**, 37–40.
- Crews, C.M., Alessandrini, A., and Erikson, R.L. (1992) The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. *Science*, **258**, 478–480.
- Dalby, K.N., Morrice, N., Caudwell, F.B., Avruch, J., and Cohen, P. (1998) Identification of regulatory phosphorylation sites in mitogen-activated protein kinase (MAPK)-activated protein kinase-1a/p90^{sk}. *J. Biol. Chem.*, **273**, 1496–1505.
- DeAngelis, P.L. and Acbyuthan, A.M. (1996) Yeast-derived recombinant DG42 protein of *Xenopus* can synthesize hyaluronan *in vitro*. *J. Biol. Chem.*, **271**, 23657–23660.
- Endre, G., Kereszt, A., Kevei, Z., Mihacea, S., Kaló, P., and Kiss, G.B. (2002) A receptor kinase gene regulating symbiotic nodule development. *Nature*, **417**, 962–966.
- Fukuda, M., Gotoh, Y., and Nishida, E. (1997) Interaction of MAP kinase with MAP kinase kinase: its possible role in the control of nucleocytoplasmic transport of MAP kinase. *EMBO J.*, **16**, 1901–1908.
- Fukunaga, R. and Hunter, T. (1997) MNK1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates. *EMBO J.*, **16**, 1921–1933.
- Johnson, G.L. and Lapadat, R. (2002) Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science*, **298**, 1911–1912.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680–685.
- Lenormand, P., Brondello, J., Brunet, A., and Pouyssegur, J. (1998) Growth factor-induced p42/p44 MAPK nuclear translocation and retention requires both MAPK activation and neosynthesis of nuclear anchoring proteins. *J. Cell Biol.*, **142**, 625–633.
- Lewis, T.S., Shapiro, P.S., and Ahn, N.G. (1998) Signal transduction through MAP kinase cascades. *Adv. Cancer Res.*, **74**, 49–139.
- Lewis, T.S., Hunt, J.B., Aveline, L.D., Jonscher, K.R., Louie, D.F., Yeh, J.M., Nehreini, T.S., Resing, K.A., and Ahn, N.G. (2000) Identification of novel MAP kinase pathway signaling targets by functional proteomics and mass spectrometry. *Mol. Cell*, **6**, 1343–1354.
- Marais, R., Wynne, J., and Treisman, R. (1993) The SRF accessory protein ELK-1 contains a growth factor-regulated transcriptional activation domain. *Cell*, **73**, 381–393.
- Marshall, C.J. (1995) Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. *Cell*, **80**, 179–185.
- Meyer, M.F. and Kreil, G. (1996) Cells expressing the dg42 gene from early *Xenopus* embryos synthesize hyaluronan. *Proc. Natl. Acad. Sci. USA*, **93**, 4543–4547.
- Nebreda, A.R. and Gavin, A.C. (1999) Perspectives: signal transduction. Cell survival demands some Rsk. *Science*, **286**, 1309–1310.
- Peppelenbosch, M.P., Tertoolen, L.G.J., De Laat, S.W., and Zivkovic, D. (1995) Ionic responses to epidermal growth factor in zebrafish cells. *Exp. Cell Res.*, **218**, 183–188.
- Pouyssegur, J., Volmat, V., and Lenormand, P. (2002) Fidelity and spatio-temporal control in MAP kinase (ERKs) signaling. *Biochem. Pharmacol.*, **64**, 755–763.
- Pozios, K.C., Ding, J., Degger, B., Upton, Z., and Duan, C. (2001) IGFs stimulate zebrafish cell proliferation by activating MAP kinase and PI3-kinase-signaling pathways. *Am. J. Physiol. Reg. Int. Comp. Physiol.*, **280**, 1230–1239.
- Robina, I., Gómez-Bujedo, S., Fernández-Bolaños, J.G., Fuentes, J., and Spaink, H.P. (2002) Synthesis and biological evaluation of oligosaccharides related to the molecule signals in plant defence and *Rhizobium*-legume symbiosis. *Tetrahedron*, **58**, 521–530.
- Sawada, A., Shinya, M., Jiang, Y., Kawakami, A., Kuroiwa, A., and Takeda, H. (2001) Fgf/MAPK signalling is a crucial positional cue in somite boundary formation. *Development*, **128**, 4873–4880.
- Schuster, J.M. and Nelson, P.S. (2000) Toll receptors: an expanding role in our understanding of human disease. *J. Leukoc. Biol.*, **67**, 767–773.
- Semino, C.E. and Allende, M.L. (2000) Chitin oligosaccharides as candidate patterning agents in zebrafish embryogenesis. *Int. J. Dev. Biol.*, **44**, 183–193.
- Semino, C.E. and Robbins, P.W. (1995) Synthesis of “Nod”-like chitin oligosaccharides by the *Xenopus* developmental protein DG42. *Proc. Natl. Acad. Sci. USA*, **92**, 3498–3501.
- Semino, C.E., Specht, C.A., Raimondi, A., and Robbins, P.W. (1996) Homologs of the *Xenopus* developmental gene *DG42* are present in zebrafish and mouse and are involved in the synthesis of nod-like chitin oligosaccharides during early embryogenesis. *Proc. Natl. Acad. Sci. USA*, **93**, 4548–4553.
- Semino, C.E., Allende, M.L., Bakkers, J., Spaink, H.P., and Robbins, P.W. (1998) Expression of *Rhizobium* oligosaccharide fucosyltransferase in zebrafish embryos disrupts normal development. *Ann. NY Acad. Sci.*, **842**, 49–54.
- Shinya, M., Koshida, S., Sawada, A., Kuroiwa, A., and Takeda, H. (2001) Fgf signalling through MAPK cascade is required for development of the subpallial telencephalon in zebrafish embryos. *Development*, **128**, 4153–4164.
- Southwick, A.M., Wang, L.-X., Long, S.R., and Lee, Y.C. (2002) Activity of *Sinorhizobium meliloti* NodAB and NodH enzymes on thiochito-oligosaccharides. *J. Bact.*, **184**, 4039–4043.
- Spaink, H.P. (2002) A receptor in symbiotic dialogue. *Nature*, **417**, 910–911.
- Stefanovsky, V.Y., Pelletier, G., Hannan, R., Gagnon-Kugler, T., Rothblum, L.I., and Moss, T. (2001) An immediate response of ribosomal transcription to growth factor stimulation in mammals is mediated by ERK phosphorylation of UBF. *Mol. Cell*, **8**, 1063–1073.
- Stracke, S., Kistner, C., Yoshida, S., Mulder, L., Sato, S., Kaneko, T., Tabata, S., Sandal, N., Stougaard, J., Szczyglowski, K., and Parniske, M. (2002) A plant receptor-like kinase required for both bacterial and fungal symbiosis. *Nature*, **417**, 959–962.
- Sturgill, T.W., Ray, L.B., Erikson, E., and Maller, J.L. (1988) Insulin-stimulated MAP-2 kinase phosphorylates and activates ribosomal protein S6 kinase II. *Nature*, **334**, 715–718.
- Treisman, R. (1996) Regulation of transcription by MAP kinase cascades. *Curr. Opin. Cell Biol.*, **8**, 205–215.
- Van der Holst, P.P.G., Schlaman, H.R.M., and Spaink, H.P. (2001) Proteins involved in the production and perception of oligosaccharides in relation to plant and animal development. *Curr. Opin. Struct. Biol.*, **11**, 608–616.
- Varki, A. (1996) Does DG42 synthesize hyaluronan or chitin: a controversy about oligosaccharides in vertebrate development. *Proc. Natl. Acad. Sci. USA*, **93**, 4523–4525.

- Wang, L.-X. and Lee, Y.C. (1995) Stereoselective synthesis of *N*-acetyl thiochitoooligosaccharides. Different behaviours of methyl *N*-acetyl- α -, and- β -thiochitobiosides during acetolysis. *J. Chem. Soc. Perkin Trans.*, 581–591.
- Yntema, H.G., Van den Helm, B., Kissing, J., Van Duijnhoven, G., Poppelaars, F., Chelly, J., Moraine, C., Fryns, J.P., Hamel, B.C., Heillbronner, H., and others. (1999) A novel ribosomal S6-kinase (RSK4; RPS6KA6) is commonly deleted in patients with complex X-linked mental retardation. *Genomics*, **62**, 332–343.
- Yoshida, M., Itano, N., Yamada, Y., and Kimata, K. (2000) *In vitro* synthesis of hyaluronan by a single protein derived from mouse HAS1 gene and characterization of amino acid residues essential for the activity. *J. Biol. Chem.*, **275**, 497–506.