



Universiteit
Leiden
The Netherlands

Systematic investigations into the role of ceramide subclass composition on lipid organization and skin barrier

Nadaban, A.

Citation

Nadaban, A. (2024, May 16). *Systematic investigations into the role of ceramide subclass composition on lipid organization and skin barrier*. Retrieved from <https://hdl.handle.net/1887/3754008>

Version: Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

License: <https://hdl.handle.net/1887/3754008>

Note: To cite this publication please use the final published version (if applicable).

CURRICULUM VITAE

Andreea Nădăban was born on the 19th of November 1993 in Arad, Romania. After graduating from the National College Elena Ghiba Birta Arad in 2012, she started her Chemistry BSc degree at the West University of Timișoara. She moved to The Netherlands in September 2015 to follow the Analytical Sciences MSc degree at the Vrije Universiteit Amsterdam and the University of Amsterdam (joint degree). During her study, she performed a 10 months internship at the Clinical Chemistry laboratory of VUmc Amsterdam. In September 2017 she started working as a LC-MS technician at the division of BioTherapeutics of the Leiden Academic Centre for Drug Research (LACDR), Leiden University. In April 2019 she started her PhD project at the LACDR under the supervision of Prof. dr. Joke Bouwstra and Prof. dr. Robert Rissmann, which resulted in this thesis.

LIST OF PUBLICATIONS

From this thesis:

- Nădăban, A., Frame, C.O., El Yachioui, D., Gooris, G.S., Dalgliesh, R.M., Malfois, M., Iacovella, C.R., Bunge, A.L., McCabe, C.M., Bouwstra, J.A. (2024). The sphingosine and phytosphingosine ceramide ratio in lipid models forming the short periodicity phase: an experimental and molecular simulation study, *Langmuir*, submitted.
- Nădăban, A., Gooris, G. S., Beddoes, C. M., Dalgliesh, R. M., Malfois, M., Deme, B., Bouwstra, J. A. (2023). The molecular arrangement of ceramides in the unit cell of the long periodicity phase of stratum corneum models shows a high adaptability to different ceramide head group structures. *Biochim. Biophys. Acta Biomembr.*, submitted.
- Rousel, J., Nădăban, A., Saghari, M., Pagan, L., Zhuparris, A. et al. (2023), Lesional skin of seborheic dermatitis patients is characterized by skin barrier dysfunction and correlating alterations in the stratum corneum ceramide composition, *Exp. Dermatol.* 33(1): 14952.
- Nădăban, A., Rousel, J., El Yachioui, D., Gooris, G. S., Beddoes, C. M., Dalgliesh, R. M., Malfois, M., Rissmann, R., Bouwstra, J.A. (2023). Effect of sphingosine and phytosphingosine ceramide ratio on lipid arrangement and barrier function in skin lipid models. *J. Lipid Res.* 64(8), 100400.
- Nădăban, A., Gooris, G. S., Beddoes, C. M., Dalgliesh, R. M., Bouwstra, J. A. (2022). Phytosphingosine ceramide mainly localizes in the central layer of the unique lamellar phase of skin lipid model systems. *J Lipid Res.* 63(9), 100258.

Author:

- Shamaprasad, P., Nădăban, A., Iacovella, C.R., Gooris, G.S., Bunge, A.L. et al. (2024), The phase behavior of skin-barrier lipids: a combined approach of experiments and simulations, *Biophys. J.*, submitted.
- Bouwstra, J.A., Nădăban, A., Bras, W., McCabe, C., Bunge, A. et al. (2023), The skin barrier: an extraordinary interface with an exceptional lipid organization, *Prog. Lipid Res.* 92: 101252.
- Rousel, J., Saghari, M., Pagan, L., Nădăban, A., Gambrah, T. et al. (2023), Treatment with the Topical Antimicrobial Peptide Omiganan in Mild-to-Moderate Facial Seborheic Dermatitis versus Ketoconazole and Placebo: Results of a Randomized Controlled Proof-of-Concept Trial, *Int. J. Mol. Sci.* 24(18): 14315.
- Helder, R.W.J., Rousel, J., Boiten, W.A., Gooris, G.S., Nădăban, A. et al. (2023), The effect of PPAR isoform (de)activation on the lipid composition in full-thickness skin models, *Exp. Dermatol.* 32(4): 469-478.
- Petracca, B., Nădăban, A., Eeman, M., Gooris, G.S., Bouwstra, J.A. (2021), Effects of ozone on stratum corneum lipid integrity and assembly, *Chem. Phys. Lipids.* 240: 105121.
- Mieremet, A., Helder, R.W.J., Nădăban, A., Boiten, W.A., Gooris, G.S. et al. (2021), Multitargeted approach for the optimization of morphogenesis and barrier formation in human skin equivalents, *Int. J. Mol. Sci.* 22(11): 5790.
- Helder, R.W.J., Rousel, J., Boiten, W.A., Gooris, G.S., Nădăban, A. et al. (2021), Improved organotypic skin model with reduced quantity of monounsaturated ceramides by inhibiting stearoyl-CoA desaturase-1, *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1866(4): 158885.
- Mieremet, A., Helder, R.W.J., Nădăban, A., Gooris, G.S., Boiten, W.A. et al. (2019), Contribution of Palmitic Acid to Epidermal Morphogenesis and Lipid Barrier Formation in Human Skin Equivalents, *Int. J. Mol. Sci.* 20(23): 6069.

A

