



**Universiteit
Leiden**
The Netherlands

Quantification of unmethylated insulin DNA using methylation sensitive restriction enzyme digital polymerase chain reaction

Leemkolk, F.E.M. van de; Nell, R.J.; Versluis, M.; Koning, E.J.P. de; Hurman, V.A.L.; Alwayn, I.P.J.; ... ; Engelse, M.A.

Citation

Leemkolk, F. E. M. van de, Nell, R. J., Versluis, M., Koning, E. J. P. de, Hurman, V. A. L., Alwayn, I. P. J., ... Engelse, M. A. (2022). Quantification of unmethylated insulin DNA using methylation sensitive restriction enzyme digital polymerase chain reaction. *Transplant International*, 35. doi:10.3389/ti.2022.10167

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3563126>

Note: To cite this publication please use the final published version (if applicable).



Quantification of Unmethylated Insulin DNA Using Methylation Sensitive Restriction Enzyme Digital Polymerase Chain Reaction

Fenna E. M. van de Leemkolk^{1,2*}, Rogier J. Nell³, Mieke Versluis³, Eelco J. P. de Koning^{1,4}, Volkert A. L. Huurman^{1,2}, Ian P. J. Always^{1,2}, Rutger J. Ploeg^{1,5}, Pieter A. van der Velden³ and Marten A. Engelse^{1,4*}

¹LUMC Transplant Center, Leiden University Medical Center, Leiden, Netherlands, ²Department of Surgery, Leiden University Medical Center, Leiden, Netherlands, ³Department of Ophthalmology, Leiden University Medical Center, Leiden, Netherlands, ⁴Department of Internal Medicine, Leiden University Medical Center, Leiden, Netherlands, ⁵Nuffield Department of Surgical Sciences, University of Oxford, Oxford, United Kingdom

Assessment of specific β -cell death can be used to determine the quality and viability of pancreatic islets prior to transplantation and hence predict the suitability of the pancreas for isolation. Recently, several groups have demonstrated that unmethylated insulin (*INS*)-DNA is correlated to β -cell death in type 1 diabetes patients and during clinical islet isolation and subsequent transplantation. Here, we present a step-by-step protocol of our novel developed method for quantification of the relative amount of unmethylated *INS*-DNA using methylation sensitive restriction enzyme digital polymerase chain reaction. This method provides a novel and sensitive way to quantify the relative amount of β -cell derived unmethylated *INS*-DNA in cellular lysate. We therefore suggest that this technique can be of value to reliably determine the purity of an islet preparation and may also serve as a measure of the quality of islets prior to transplantation measuring unmethylated *INS*-DNA as a reflection of the relative amount of lysed β -cells.

OPEN ACCESS

*Correspondence:

Fenna E. M. van de Leemkolk
f.e.m.van_de_leemkolk@lumc.nl

Marten A. Engelse
m.a.engelse@lumc.nl

Received: 31 October 2021

Accepted: 15 March 2022

Published: 07 April 2022

Citation:

van de Leemkolk FEM, Nell RJ, Versluis M, de Koning EJP, Huurman VAL, Always IPJ, Ploeg RJ, van der Velden PA and Engelse MA (2022) Quantification of Unmethylated Insulin DNA Using Methylation Sensitive Restriction Enzyme Digital Polymerase Chain Reaction. *Transpl Int* 35:10167. doi: 10.3389/ti.2022.10167

Keywords: transplantation, biomarker, unmethylated insulin DNA, methylation sensitive restriction enzyme, digital PCR, β -cell, islets

INTRODUCTION

β -cell replacement therapy has been established as a therapy for patients with complex Type 1 Diabetes (T1D) not amenable to optimal conventional diabetes management (1). One example of β -cell replacement therapy is the transplantation of deceased donor derived pancreatic islets that has proven its long-term efficacy during the past 20 years (2, 3). In order to aim for optimal post-transplant outcomes, the use of high-quality pancreatic islets is essential. Reliable assays are needed to assess the quality and viability of islets prior to transplantation. Soluble β -cell specific biomarkers may serve as a relevant diagnostic target to determine the quality and viability of islets at an early

Abbreviations: ddPCR, Digital Droplet polymerase Chain Reaction; DNA, DeoxyriboNucleic Acid; DTZ, Dithizone Staining; IEQ, Islet Equivalent; INS, Insulin; MSRE, Methylation Sensitive Restriction Enzyme; PCR polymerase Chain Reaction; T1D, Type 1 Diabetes; TTC5, Tetratricopeptide Repeat Domain 5.

stage as they can be used to assess the amount of β -cell loss during islet isolation and subsequent transplantation.

Recently, several groups have reported unmethylated Insulin (*INS*)-DNA as a specific β -cell death marker during the early development of T1D. During the progression of the disease, autoimmune destruction of β -cells occurs and unmethylated *INS*-DNA is released in the bloodstream that can be identified (4-11). As the concentration of this marker is extremely low, digital polymerase chain reaction (PCR) is often used to detect the amount of β -cell death in a quantitative manner. Recent studies using digital PCR to analyze unmethylated *INS*-DNA were based on a sodium-bisulfite conversion method that chemically converts unmethylated cytosine into uracil (6, 8-10, 12). However, this method comprises an insurmountable problem as regards heterogeneity since it depends on the completeness of the chemical conversion. Overshooting or incomplete bisulfite conversion can lead to reduced sensitivity and may hamper quantitative and qualitative interpretation (13).

To avoid bisulfite conversion whilst still allowing the possibility to specifically quantify the methylation fraction of a specific allele, we recently published a methylation sensitive restriction enzyme (MSRE) digital PCR assay (14). MSREs are used to differentiate between methylated and unmethylated alleles and in combination with digital PCR it provides the opportunity to determine specific allele quantification.

Based on this methodology we now describe here the step-by-step approach how to quantify the unmethylated *INS*-DNA fraction using a MSRE and digital PCR assay. In this proof-of-concept study, we aim to demonstrate that this novel assay can be used as a helpful method to determine the purity of an islet preparation by measuring the amount of β -cells specific genomic DNA in an islet suspension. The subsequent step to then test this particular assay as a clinically quality marker of islets prior to transplantation by measuring the relative amount of lysed β -cells was beyond the scope of this proof-of-concept study. .

METHOD

Sample Collection and DNA Isolation

Human insulinoma EndoC- β H1 cells (Univercell-Biosolutions (15), Toulouse, France) and human monocytic THP-1 cells (Invivogen, Toulouse, France) were used as a positive and negative control, respectively. Isolated human pancreatic islets with different purities were obtained from seven individual pancreases (Leiden University Medical Center, Netherlands). Human donor pancreases were used that were declined for clinical purposes according to national criteria. Written informed consent for research of pancreatic tissue from donors was present, according to local guidelines of the medical ethical committee (Leiden University Medical Center, Netherlands) and of the Dutch Transplantation Foundation as the competent authority for organ donation in Netherlands. Regarding the culture of the EndoC- β H1 and THP-1 cells and isolation and maintenance of human islets, please find further details in the Supplemental document.

- 1) Stored pellets of 2.5×10^6 EndoC- β H1 cells, 2.5×10^6 THP-1 cells and 10 μ L tissue of different purities from human islets were resuspended with phosphate buffer up to a final volume of 200 μ L.

From these samples genomic DNA was extracted using a QIAamp DNA Mini Kit (Qiagen, Venlo, Netherlands) according to the manufacturer's instructions.

- 2) DNA concentrations were measured using NanoDrop TM 1000 Spectrophotometer (Thermo Fisher Scientific, Landsmeer, Netherlands).

Treatment With Methylation Sensitive Restriction Enzyme


The restriction enzyme, HpaII (Thermo Fisher Scientific), was used according to manufacturer's instructions. The restriction enzyme was added for the *INS target DNA* (Figure 1A) as it cleaves the unmethylated *INS*-DNA and leaves the methylated *INS*-DNA intact. Each sample was either left untreated or treated with HpaII.

- 1) Take two separate units of 100 ng genomic DNA from each sample and add each of these units to a separate PCR tube (8-strip PCR tubes). Mark the first strip as "with MSRE" and the second strip as "without MSRE". Include at least one sample in each strip containing only nuclease-free H₂O (negative control).
- 2) Add 2 units/reaction of HpaII, 1.0 μ L CutSmart Buffer (Bioké, Leiden, Netherlands), and nuclease-free H₂O up to a total volume of 10 μ L to the strip marked as "with MSRE".
- 3) Add 1.0 μ L CutSmart Buffer (Bioké, Leiden, Netherlands), and nuclease-free H₂O up to a total volume of 10 μ L to the strip marked as "without MSRE".
- 4) Incubate both strips at 37°C for 1 hour.

Duplex Analysis Using Digital PCR

Primers and FAM-labelled hydrolysis probes (both Sigma-Aldrich) were designed to be 1) gene specific, 2) to contain an MSRE specific CpG site and 3) to possess optimal melting temperature ($\pm 55^\circ\text{C}$) based on the region identified previously (Supplementary Figure S1) (11, 16). Probes directed to the *INS target DNA* were labelled with FAM (Supplementary Table S1). The probe directed to the reference TTC5 (tetratricopeptide repeat domain 5) gene was labelled with HEX (BioRad, Veenendaal, Netherlands).

- 1) To prepare the PCR mastermix, add 11 μ L per reaction of Droplet PCR Supermix™ (No dUTP) (BioRad) (e.g., 110 μ L per 10 samples), 0.5 μ L per reaction 36uM forward *INS* primer (e.g., 5 μ L per 10 samples), 0.5 μ L per reaction 36uM *INS* reverse primer (e.g., 5 μ L per 10 samples), 0.5 μ L per reaction 10uM *INS* FAM probe (e.g., 5 μ L per 10 samples), 1 μ L per reaction 20x TTC5 HEX assay (e.g., 10 μ L per 10 samples) and 6.5 μ L per reaction nuclease-free H₂O (e.g., 65 μ L per 10 samples).



2) In order to set up a PCR reaction in a 96-well plate, first, add 20 μL mastermix to each well. Add 2 μL of cleaved un methylated *INS*-DNA (from the “with MSRE” PCR-strip) or un cleaved un methylated *INS*-DNA (from the “without MSRE” PCR-strip) to each appropriate well. Mix wells by pipetting up-and-down several times.

All eight wells in a column must contain cleaved un methylated *INS*-DNA (from the “with MSRE” PCR-strip) or un cleaved un methylated *INS*-DNA (from the “without MSRE” PCR-strip).


- 3) Seal the 96-well PCR plate with foil and centrifuge shortly to remove liquid from the sides of the wells.
- 4) Digital PCR is performed using the digital droplet PCR (ddPCR) method described below (Figure 1B).
 - 4.1) Use the Automated Droplet Generator (BioRad) to generate droplets according to manufacturer’s instructions.

4.2) In order to prevent evaporation of the newly formed droplets, the droplets should be collected in a second 96-well PCR plate placed into a properly frozen cooling block.

4.3) When finished, remove the 96-well PCR plate including the newly formed droplets and use a Plate Sealer (BioRad) in order to cover the 96-well PCR plate with a heat-sealed foil.

NOTE: Careful handling is strongly advised as the newly formed droplets are fragile in this stage.

- 5) Perform a PCR reaction in a T100 Thermal Cycler (BioRad) using the following protocol:
 - 10 min of activation at 95°C
 - 30s at 94°C denaturation and 60s at 60°C for 40 cycles
 - 10 min inactivation at 98°C
 - Cooling at 12°C until droplet reading



6) Analyze the DNA content of the droplets using the QuantaSoft™ software with the QX200 Droplet Reader (BioRad) according to manufacturer’s protocol.

7) Calculate for each sample the unmethylated *INS*-DNA fraction as follows:


- $$\text{Unmethylation fraction} = 1 - \frac{\frac{[INS]_{with MSRE}}{[TTC5]_{with MSRE}}}{\frac{[INS]_{without MSRE}}{[TTC5]_{without MSRE}}} * 100\%$$

RESULTS

With attention to previous studies (11, 16) on target areas of DNA methylation in the human *INS* gene, we designed a methylation sensitive restriction enzyme (MSRE) duplex digital PCR assay to determine the relative amount of

unmethylated *INS*-DNA fraction in our DNA samples of interest.


First, the assay was validated in cell line models. DNA was isolated from EndoC-βH1 cells, a cell line that was derived from human β-cells(17). The MSRE duplex digital PCR assay was performed. This results in two-dimensional plots that demonstrate four different clusters each of them representing different DNA containing droplets (Figure 2). The green cluster contains no *INS target DNA* but only *TTC5* copies; the blue cluster contains only *INS target DNA* but no *TTC5* copies; the orange cluster contains both *INS target DNA* and *TTC5* copies; the gray cluster includes the empty droplets. Without treatment of the MSRE (Figure 2A), the *INS target DNA* reflects the quantification of both unmethylated and



methylated *INS target DNA*. After treatment with the MSRE HpaII (**Figure 2B**), the un methylated *INS target DNA* is digested, resulting in less blue and orange droplets. For both, with and without treatment of MSRE, a stable independent reference, TTC5, was used to correct for input differences as it is not digested by the MSRE. When using both ratios from *INS target DNA* and reference TTC5 in the samples with and without treatment with MSRE, an un methylated *INS-DNA* fraction of 98.1% (95% CI 97.3–98.8) was determined (**Figure 2C**). With regards to DNA isolated from THP-1 cells, both ratios from *INS target DNA* and reference TTC5, when treated with (**Figure 2E**) or without (**Figure 2D**) the MSRE HpaII, were calculated and this resulted in an un methylated *INS-DNA* fraction of 3.5% (95% CI -5.2–11.5) (**Figure 2F**).

As isolated DNA from EndoC-βH1 cells was essentially un methylated for the *INS target DNA* whilst isolated DNA from THP-1 cells was mainly methylated for the *INS target DNA*, a 7-points standard curve was generated to technically validate the quantitative experimental setup. Isolated DNA from EndoC-βH1 cells diluted in the background of isolated DNA from THP-1 cells resulted in a strong linear correlation ($r^2 = 0.9953$, $Y = 0.8862 * X + 7.019$, $p < 0.0001$) (**Figure 3**).

Next, the un methylated *INS-DNA* fraction was determined in 24 human islets preparations which were isolated from seven different human donor pancreases obtained for research. For each sample, islet purity was determined, varying from <5 to 99%, via dithizone staining which is currently used by most centers to estimate the fraction of pancreatic islets in an isolated islet preparation (18, 19). In the case of a sample with <5% purity, the sample was categorized as islet depleted tissue (i.e., pancreatic tissue left over from islet isolation). After using this MSRE duplex digital PCR assay on DNA isolated from all the different purities



of the islets, the un methylated *INS-DNA* fraction was quantified (**Figure 4**). When comparing the purity of the pancreatic islets a significant linear correlation was observed (R squared = 0.8318, $p < 0.0001$). In the samples containing islet depleted tissue an un methylated *INS-DNA* fraction of 29.4%–34.5% was observed.

DISCUSSION

Previous studies have demonstrated that the human *INS* gene is controlled epigenetically by methylation as it is un methylated in β-cells and methylated in most other cell types (4, 20–22). When cells are dying or lysed - either *in vivo* or for experimentation purposes - their genomic DNA is released into the milieu. This makes un methylated-*INS* DNA a highly interesting marker to detect the death of β-cells. Several research groups have developed assays to measure the circulating fraction of un methylated *INS-DNA* in humans, often aiming to be used in the context of early detection of β-cell death in type 1 Diabetes. In 2020 Speake et al. (23) assessed the performance of three different methodologies (5, 9 11) to quantify circulating levels of un methylated *INS-DNA* in patients undergoing total pancreatectomy and subsequent islet auto-transplantation. This was considered a reliable model as damage or cell death of β-cells is known to occur during transplantation. Not only did the group measure a different CpG site or sites in the human *INS* gene in these three assays, they also applied different sample collection methods and measurement techniques (e.g., next generation sequencing or digital PCR). We agree with Speake's group that to further develop these assays, optimization of the three different techniques might be beneficial. A similarity between all three assays was that DNA was treated with sodium bisulfite. This technique, which was first described by Frommer et al. (24), is still regarded as the gold standard to analyze DNA methylation. To prevent partial conversion and subsequent misinterpretation, the chemical conversion is performed at high concentrations. As a result, however, fragmentation and degradation of DNA will occur that may lead to an incorrect quantitative interpretation

(13, 25). In addition, with regard to the bisulfite conversion kits used in these studies focusing on unmethylated-*INS* DNA, it remains a relatively time consuming technique e.g., as approximately 12–16 h are needed for the incubation period.

To circumvent or even avoid these limitations, we report in this protocol a proof-of-concept study where we have combined the MSRE with digital PCR techniques to measure unmethylated-*INS* DNA. As an MSRE can differentiate between methylated and unmethylated alleles, MSRE treatment for only 1 hour results in digestion of unmethylated DNA, with the methylated DNA remaining intact. This allows for the rapid calculation of the fraction of unmethylated alleles in our target of interest (*INS target DNA*). When using two different cell lines, a strong correlation was observed (Figure 3) demonstrating a high sensitivity and specificity of this assay.

Next, we extended the use of this assay to measure the unmethylated *INS*-DNA fraction in different purities of islets obtained after pancreas isolation (Figure 4). Interestingly, the purity of the samples was not directly proportional to the quantified unmethylated *INS*-DNA fraction as was found in the standard curve obtained from the 2 cell lines (Figure 3). When using the MSRE duplex digital PCR in islet depleted tissue (i.e. containing <5% islets) an unmethylated *INS*-DNA fraction of 29.4–34.5% was observed. Of note is that this observed fraction is likely not a limitation of the assay itself but an indication that the biological variability in methylation of the human *INS* gene promotor in non- β -cells may play an important role. Our result is in line with the study by Kuroda et al. (22) who investigated nine CpG sequences in the human *INS* gene promotor and compared the methylation pattern in this region in the ‘islet cell fraction’ and in the ‘non-islet cell fraction’. In their study they demonstrated that the human *INS* gene promotor was mainly unmethylated in the islet cell fraction and predominantly methylated in the non-islet cell fraction (i.e., 13 of 15 clones (86%) in the non-islet cell fraction exhibited at least one unmethylated CpG out of the nine CpG sequences investigated).

With regard to the samples including high purity of islets, the quantified unmethylated *INS*-DNA fraction did not reach 100% which could be explained as the ratio of β -cells versus non- β -cells (e.g., alpha and delta) in human islets is generally assumed to be 50–70% (26). This is in line with the $\pm 70\%$ unmethylated *INS*-DNA fraction we have found (Figure 4).

A limitation of this proof-of-concept study is that our protocol was performed in cell lines and in different purities of human islet preparations obtained after isolation. Further validation experiments of this assay during islet isolation, islet culture and subsequent islet transplantation are necessary. During these next steps of the process an unknown amount of β -cell destruction occurs. To be able to specifically quantify the amount of β -cell loss using this promising assay could be helpful to differentiate between low or high quality and viability of islet preparations (12, 27). In clinical islet transplantation the accurate determination of the number of (viable) β -cells in a pancreatic islet preparation is essential. Not only assessment of the islet depleted tissue fraction, but more important the total number of isolated islets in the preparation is key for a successful transplant (28). In islet transplantation, the islet yield has previously been determined using various methods such as size-dependent islet counting by visualizing islets under a microscope and subsequent measurement

of their volume (19), calculating both islet purity and graft volume or specific β -cell counting (28–31). To date, in most centers the estimation of the fraction of pancreatic islets in an isolated islet preparation is based on a method that uses dithizone staining (DTZ) (18, 19). Dithizone is a zinc chelating agent that, when added to an islet prep, results in a rapidly and reversibly red staining of islets which can therefore be distinguished from exocrine tissue. Importantly, this method cannot be used to determine the total number of β -cells in an isolated islet preparation. In addition, in case of β -cell degranulation, the red staining will not take place. Therefore, due to the human error that is intrinsic to this subjective method, an over- or under-estimation of the islet equivalent (IEQ) may easily occur. As such, determination of IEQ by eye or by digital image analysis has proven difficult within and between different centers (32).

Based on these notions, we suggest in this preliminary study that our newly developed MSRE duplex digital PCR assay using unmethylated *INS*-DNA may be a fast and easy method to specifically quantify β -cells. As shown previously, the combination of MSRE with digital PCR provides both specificity and sensitivity by quantitative assessment of target alleles (14). By measuring the concentration of the targeted unmethylated *INS*-DNA and therefore the number of lysed β -cells, this combined technique may be a promising tool to determine the fraction of β -cells immediately after islet isolation, during culture and immediately prior to islet transplantation. Pending further validation trials, the MSRE duplex digital PCR assay using unmethylated *INS*-DNA may therefore help decision making on islet quality (through the measurement of β -cell death) and islet quantity in islet transplantation centers.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Leiden University Medical Center, Leiden, Netherlands. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

FL, RN, PV and ME participated in conception and research design. The acquisition of data was performed by FL and MV. FL, RN, MV, Ed, VH, IA, RP, PV and ME participated in data analysis and interpretation. FL and ME drafted the manuscript. FL, RN, MV, Ed, VH, IA, RP, PV and ME participated in critical revision and final approval of the manuscript. PV and ME participated in study supervision. FL and ME are the guarantors of this work and, as such, had full access to all of

the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

FUNDING

RN is supported by the European Union's Horizon 2020 research and innovation program under grant agreement No 667787 (UM Cure 2020 project).

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

REFERENCES

- Shapiro AMJ, Pokrywczynska M, Ricordi C. Clinical Pancreatic Islet Transplantation. *Nat Rev Endocrinol* (2017) 13(5):268–77. doi:10.1038/nrendo.2016.178
- Farney AC, Sutherland DER, Opara EC. Evolution of Islet Transplantation for the Last 30 Years. *Pancreas* (2016) 45(1):8–20. doi:10.1097/mpa.0000000000000391
- Vantghem M-C, de Koning EJP, Pattou F, Rickels MR. Advances in β -cell Replacement Therapy for the Treatment of Type 1 Diabetes. *The Lancet* (2019) 394(10205):1274–85. doi:10.1016/s0140-6736(19)31334-0
- Akirav EM, Lebastchi J, Galvan EM, Henegariu O, Akirav M, Ablamunits V, et al. Detection of β Cell Death in Diabetes Using Differentially Methylated Circulating DNA. *Proc Natl Acad Sci U.S.A.* (2011) 108(47):19018–23. doi:10.1073/pnas.1111008108
- Lehmann-Werman R, Neiman D, Zemmour H, Moss J, Magenheimer J, Vaknin-Dembinsky A, et al. Identification of Tissue-specific Cell Death Using Methylation Patterns of Circulating DNA. *Proc Natl Acad Sci U S A* (2016) 113(13):E1826–34. doi:10.1073/pnas.1519286113
- Tersey SA, Nelson JB, Fisher MM, Mirmira RG. Measurement of Differentially Methylated INS DNA Species in Human Serum Samples as a Biomarker of Islet β Cell Death. *J Vis Exp* (2016) (118) 54838. doi:10.3791/54838
- Zhang K, Lin G, Han Y, Xie J, Li J. Circulating Unmethylated Insulin DNA as a Potential Non-invasive Biomarker of Beta Cell Death in Type 1 Diabetes: a Review and Future prospect. *Clin Epigenet* (2017) 9:44. doi:10.1186/s13148-017-0343-5
- Usmani-Brown S, Lebastchi J, Steck AK, Beam C, Herold KC, Ledizet M. Analysis of β -Cell Death in Type 1 Diabetes by Droplet Digital PCR. *Endocrinology* (2014) 155(9):3694–8. doi:10.1210/en.2014-1150
- Herold KC, Usmani-Brown S, Ghazi T, Lebastchi J, Beam CA, Bellin MD, et al. β Cell Death and Dysfunction during Type 1 Diabetes Development in At-Risk Individuals. *J Clin Invest* (2015) 125(3):1163–73. doi:10.1172/jci78142
- Simmons KM, Fouts A, Pyle L, Clark P, Dong F, Yu L, et al. Unmethylated Insulin as an Adjunctive Marker of Beta Cell Death and Progression to Type 1 Diabetes in Participants at Risk for Diabetes. *Int J Mol Sci* (2019) 20(16):3857. doi:10.3390/ijms20163857
- Fisher MM, Watkins RA, Blum J, Evans-Molina C, Chalasani N, DiMeglio LA, et al. Elevations in Circulating Methylated and Unmethylated Preproinsulin DNA in New-Onset Type 1 Diabetes. *Diabetes* (2015) 64(11):3867–72. doi:10.2337/db15-0430
- Bellin MD, Clark P, Usmani-Brown S, Dunn TB, Beilman GJ, Chinnakotla S, et al. Unmethylated Insulin DNA Is Elevated after Total Pancreatectomy with Islet Autotransplantation: Assessment of a Novel Beta Cell Marker. *Am J Transpl* (2017) 17(4):1112–8. doi:10.1111/ajt.14054
- Warnecke PM, Storzaker C, Melki JR, Millar DS, Paul CL, Clark SJ. Detection and Measurement of PCR Bias in Quantitative Methylation Analysis of Bisulphite-Treated DNA. *Nucleic Acids Res* (1997) 25(21):4422–6. doi:10.1093/nar/25.21.4422
- Nell RJ, Steenderen D, Menger NV, Weiting TJ, Versluis M, Velden PA. Quantification of DNA Methylation Independent of Sodium Bisulfite Conversion Using Methylation-sensitive Restriction Enzymes and Digital PCR. *Hum Mutat* (2020) 41(12):2205–16. doi:10.1002/humu.24111
- Ravassard P, Hazhouz Y, Pechberty S, Bricout-Neveu E, Armanet M, Czernichow P, et al. A Genetically Engineered Human Pancreatic β Cell Line Exhibiting Glucose-Inducible Insulin Secretion. *J Clin Invest* (2011) 121(9):3589–97. doi:10.1172/jci58447
- Yang BT, Dayeh TA, Kirkpatrick CL, Taneera J, Kumar R, Groop L, et al. Insulin Promoter DNA Methylation Correlates Negatively with Insulin Gene Expression and Positively with HbA1c Levels in Human Pancreatic Islets. *Diabetologia* (2011) 54(2):360–7. doi:10.1007/s00125-010-1967-6
- Tsonkova VG, Sand FW, Wolf XA, Grunnet LG, Kirstine Ringgaard A, Ingvorsen C, et al. The EndoC-Bh1 Cell Line Is a Valid Model of Human Beta Cells and Applicable for Screenings to Identify Novel Drug Target Candidates. *Mol Metab* (2018) 8:144–57. doi:10.1016/j.molmet.2017.12.007
- Latif ZA, Noel J, Alejandro R. A Simple Method of Staining Fresh and Cultured Islets. *Transplantation* (1988) 45(4):827–30.
- Ricordi C, Lacy PE, Finke EH, Olack BJ, Scharp DW. Automated Method for Isolation of Human Pancreatic Islets. *Diabetes* (1988) 37(4):413–20. doi:10.2337/diabetes.37.4.413
- Cate RL, Chick W, Gilbert W. Comparison of the Methylation Patterns of the Two Rat Insulin Genes. *J Biol Chem* (1983) 258(10):6645–52. doi:10.1016/s0021-9258(18)32461-x
- Husseiny MI, Kaye A, Zebadua E, Kandeel F, Ferreri K. Tissue-specific Methylation of Human Insulin Gene and PCR Assay for Monitoring Beta Cell Death. *PLoS one* (2014) 9(4):e94591. doi:10.1371/journal.pone.0094591
- Kuroda A, Rauch TA, Todorov I, Ku HT, Al-Abdullah IH, Kandeel F, et al. Insulin Gene Expression Is Regulated by DNA Methylation. *PLoS one* (2009) 4(9):e6953. doi:10.1371/journal.pone.0006953
- Speake C, Ylescupidez A, Neiman D, Shemer R, Glaser B, Tersey SA, et al. Circulating Unmethylated Insulin DNA as a Biomarker of Human Beta Cell Death: A Multi-Laboratory Assay Comparison. *J Clin Endocrinol Metab* (2020) 105(3):781–91. doi:10.1210/clinem/dgaa008
- Frommer M, McDonald LE, Millar DS, Collis CM, Watt F, Grigg GW, et al. A Genomic Sequencing Protocol that Yields a Positive Display of 5-methylcytosine Residues in Individual DNA Strands. *Proc Natl Acad Sci U.S.A.* (1992) 89(5):1827–31. doi:10.1073/pnas.89.5.1827
- Tanaka K, Okamoto A. Degradation of DNA by Bisulfite Treatment. *Bioorg Med Chem Lett* (2007) 17(7):1912–5. doi:10.1016/j.bmcl.2007.01.040
- Dolenssek J, Rupnik MS, Stozer A. Structural Similarities and Differences between the Human and the Mouse Pancreas. *Islets* (2015) 7(1):e1024405. doi:10.1080/19382014.2015.1024405

ACKNOWLEDGMENTS

We would like to thank our colleagues from the islet lab, for their help in islet isolation and providing sample collection and our colleagues from the digital droplet PCR group, for their expertise regarding digital droplet PCR and helpful discussions. We also would like to thank M.S. Zuurmond for the graphic design of **Figure 1**.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontierspartnerships.org/articles/10.3389/ti.2022.10167/full#supplementary-material>

27. Bellin MD, Abu-El-Haija M, Morgan K, Adams D, Beilman GJ, Chinnakotla S, et al. A Multicenter Study of Total Pancreatectomy with Islet Autotransplantation (TPIAT): POST (Prospective Observational Study of TPIAT). *Pancreatology* (2018) 18(3):286–90. doi:10.1016/j.pan.2018.02.001
28. Papas KK, Colton CK, Qipo A, Wu H, Nelson RA, Hering BJ, et al. Prediction of Marginal Mass Required for Successful Islet Transplantation. *J Invest Surg* (2010) 23(1):28–34. doi:10.3109/08941930903410825
29. Pisania A, Weir GC, O'Neil JJ, Omer A, Tchipashvili V, Lei J, et al. Quantitative Analysis of Cell Composition and Purity of Human Pancreatic Islet Preparations. *Lab Invest* (2010) 90(11):1661–75. doi:10.1038/labinvest.2010.124
30. Lehmann R, Fernandez LA, Bottino R, Szabo S, Ricordi C, Alejandro R, et al. Evaluation of Islet Isolation by a New Automated Method (Coulter Multisizer IIe) and Manual Counting. *Transplant Proc* (1998) 30(2):373–4. doi:10.1016/s0041-1345(97)01314-6
31. Huang H-H, Harrington S, Stehno-Bittel L. The Flaws and Future of Islet Volume Measurements. *Cel Transpl* (2018) 27(7):1017–26. doi:10.1177/0963689718779898
32. Kissler HJ, Niland JC, Olack B, Ricordi C, Hering BJ, Naji A, et al. Validation of Methodologies for Quantifying Isolated Human Islets: an Islet Cell Resources Study. *Clin Transplant* (2010) 24(2):236–42. doi:10.1111/j.1399-0012.2009.01052.x

Copyright © 2022 van de Leemkolk, Nell, Versluis, de Koning, Hurman, Alwayn, Ploeg, van der Velden and Engelse. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.