



Universiteit
Leiden
The Netherlands

Low-temperature spectroscopic studies of single molecules in 3-D and on 2-D hosts

Smit, R.

Citation

Smit, R. (2024, June 12). *Low-temperature spectroscopic studies of single molecules in 3-D and on 2-D hosts*. Retrieved from <https://hdl.handle.net/1887/3762935>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3762935>

Note: To cite this publication please use the final published version (if applicable).

A

APPENDIX 1

The coincidence of photons on two detectors can be related to a steady-state solution of the following optical Bloch equations, where we ignore bunching due to the triplet state:¹

$$\dot{\rho}_{11} = \gamma_{21}\rho_{22} - \frac{i\Omega}{2}\rho_{12} + \frac{i\Omega}{2}\rho_{21} \quad (\text{A.1})$$

$$\dot{\rho}_{22} = -\gamma_{21}\rho_{22} + \frac{i\Omega}{2}\rho_{12} - \frac{i\Omega}{2}\rho_{21} \quad (\text{A.2})$$

$$\dot{\rho}_{12} = -\frac{i\Omega}{2}\rho_{11} + \frac{i\Omega}{2}\rho_{22} - \pi\Gamma_0\rho_{12} \quad (\text{A.3})$$

$$\dot{\rho}_{21} = \frac{i\Omega}{2}\rho_{11} - \frac{i\Omega}{2}\rho_{22} - \pi\Gamma_0\rho_{21}. \quad (\text{A.4})$$

The term Ω is the Rabi frequency, $\pi\Gamma_0$ is the decoherence rate and γ_{21} is the depopulation rate of the excited state. The series of coupled differential equations can be solved using the Laplace transform with the condition that the population initiates from the ground state. To relate the solution for the population of the excited state ρ_{22} to an experimental photon-coincidence histogram, we need to derive the steady-state solution of ρ_{22} :

$$g^{(2)}(\tau) = \rho_{22} / \lim_{\tau \rightarrow \infty} \rho_{22}. \quad (\text{A.5})$$

Without approximations and solved with MATLAB, the theoretical solution for $g^{(2)}(\tau)$ is of the following form:

$$g^{(2)}(\tau) = 1 - e^{-\frac{|\tau|}{2}(\pi\Gamma_0 + \gamma_{21})} [\cosh(\tilde{\Omega}|\tau|) + (\frac{\pi\Gamma_0 + \gamma_{21}}{2\tilde{\Omega}}) \sinh(\tilde{\Omega}|\tau|)], \quad (\text{A.6})$$

where $\tilde{\Omega}$ is given by:

$$\tilde{\Omega} = \frac{1}{2} \sqrt{(\pi\Gamma_0)^2 - 2\pi\Gamma_0\gamma_{21} - 4\Omega^2 + \gamma_{21}^2}. \quad (\text{A.7})$$

A

Rewriting $\tilde{\Omega} = \frac{1}{2}i\sqrt{4\Omega^2 - \gamma_{21}^2 - (\pi\Gamma_0)^2 + 2\pi\Gamma_0\gamma_{21}}$ and using the identities $\sinh(\tilde{\Omega}) = -i\sin(i\tilde{\Omega})$ and $\cosh(\tilde{\Omega}) = \cos(i\tilde{\Omega})$ we obtain the more common solution:²

$$g^{(2)}(\tau) = 1 - e^{-\frac{|\tau|}{2}(\pi\Gamma_0 + \gamma_{21})} [\cos(\tilde{\Omega}|\tau|) + (\frac{\pi\Gamma_0 + \gamma_{21}}{2\tilde{\Omega}}) \sin(\tilde{\Omega}|\tau|)], \quad (\text{A.8})$$

where the new $\tilde{\Omega}$ is $\frac{1}{2}\sqrt{4\Omega^2 - \gamma_{21}^2 - (\pi\Gamma_0)^2 + 2\pi\Gamma_0\gamma_{21}}$. When the two-level system is strongly excited, the magnitude of the Rabi frequency exceeds all other terms inside the square root and the term $\tilde{\Omega}$ in the cosine and sine term reduces to the Rabi frequency Ω itself.

REFERENCES

- (1) De Vries, H.; Wiersma, D. A. *The Journal of Chemical Physics* **1980**, 72, 1851–1863.
- (2) Kummer, S.; Basche, T. *The Journal of Physical Chemistry* **1995**, 99, 17078–17081.

ACKNOWLEDGEMENTS

In the four years that I have spent in the research group Molecular Nano-Optics and Spins (MoNOS) I have been well supported by my mentor prof. dr. Michel Orrit. His vast knowledge and insights on molecular physics and optics has been a great source of inspiration for me. I look back with gratitude for all the in-depth discussions that helped me during my PhD. I would like to specially thank him for his patience to explain everything in detail to me. On the experimental side, I would like to thank dr. Amin Moradi and dr. Zoran Ristanović for the extensive training they gave me for managing the lab and the experimental setup. They prepared me well for continuing on my own in the years after they left the group. Furthermore, I am thankful to have had the pleasure of working together fruitfully with students Maialen Ortego Larrazabal and Jil Hameury and the postdoc Arash Tebyani.

For the usage of fabrication facilities and the training for handling the equipment I want to thank Douwe Scholma, Marcel Hesselberth and Luc Wigbout. I'm also indebted to engineer and strongman Harmen van der Meer, who helped me lift up the heavy cryostat to repair one of the windows that had broken. Also, the electronics department has been of great help with solving various problems encountered with electronics. For achieving low temperatures I am grateful for Wilfred van der Geest's steady supply of liquid nitrogen and helium, which fueled my experiments.

For the nice lunch breaks, support and social meetings I would like to thank all of the (former) group members of MoNOS, which includes Arash, Amin, Christopher, Deep, Eduardo, Jacco, Jacqueline, Martin, Nasrin, Patrick, Priyanshi, Leonardo, Sergii, Subhasis, Theo, Xuxing, Zoran and Zohre. For all of the challenging administrative tasks, such as exporting a laser to the US for repair, I want to thank the group's secretary Henriëtte van Leeuwen for her support. Beyond the MoNOS group I would like to thank dr. Wolfgang Löffler and Matteo Fisicaro for kindly lending me their controller box for a laser and helping me use it. Within the institute I would like to thank dr. Semonti Bhattacharyya, prof. Sense Jan van der Molen, prof. Francesco Buda, Titus de Haas and prof. Jan van Ruitenbeek for participating in fruitful scientific discussions and collaborations. Outside of the institute I would like to thank prof. Bolesław Kozankiewicz, dr. Irena Deperasińska and prof. Jean-Sébastien Lauret for collaborations and scientific discussions.

Most of all I want to thank my partner and tower of strength Rolinde Hupkes. She has been my mental support throughout my PhD. Lastly, I also thank Limon for the many creative breaks!

CURRICULUM VITAE

Robert Smit

18-08-1990

Born in Velsen, the Netherlands

EDUCATION

2009–2013

Bachelor of Science in Physics
Leiden University

2013–2015

Master of Science in Physics and Science Based Business
Leiden University

February 2020 - March 2024

PhD in Physics
Leiden University

WORK EXPERIENCE

March 2015 –August 2015

Energy market analyst internship
Northpool B.V.

September 2015 – January 2020

Energy market trader and analyst
Brainchild Capital Investments B.V

March 2024 – Present

Postdoctoral researcher
Leiden University

LIST OF PUBLICATIONS

4. Smit, R.; Ristanović, Z.; Kozankiewicz, B.; Orrit, M. *ChemPhysChem* **2022**, *23*, e202100679
3. Smit, R.; Tebyani, A.; Hameury, J.; van der Molen, S. J.; Orrit, M. *Nature Communications* **2023**, *14*, 7960
2. Adhikari, S.; Smit, R.; Orrit, M. *The Journal of Physical Chemistry C* **2024**, *128*, 3–18
1. Smit, R.; Ristanović, Z.; Deperasińska, I.; Kozankiewicz, B.; Orrit, M. *ChemPhysChem*, e202300881