

Localization of the VirA domain involved in acetosyringone-mediatedvir gene induction in Agrobacterium tumefaciens

Turk, S.C.H.J.; Lange, R.P. van; Regensburg-Tuïnk, T.J.G.; Hooykaas, P.J.J.

Citation

Turk, S. C. H. J., Lange, R. P. van, Regensburg-Tuïnk, T. J. G., & Hooykaas, P. J. J. (1994). Localization of the VirA domain involved in acetosyringone-mediatedvir gene induction in Agrobacterium tumefaciens. *Plant Molecular Biology*, *25*(5), 899-907. doi:10.1007/BF00028884

Version:Publisher's VersionLicense:Licensed under Article 25fa Copyright Act/Law (Amendment Taverne)Downloaded from:https://hdl.handle.net/1887/3765685

Note: To cite this publication please use the final published version (if applicable).

Localization of the VirA domain involved in acetosyringone-mediated vir gene induction in Agrobacterium tumefaciens

Stefan C.H.J. Turk^{1,2}, Richard P. van Lange¹, Tonny J.G. Regensburg-Tuïnk¹ and Paul J.J. Hooykaas^{1,*}

¹Institute of Molecular Plant Sciences, Clusius Laboratory, Leiden University, Wassenaarseweg 64, 2333 AL Leiden, Netherlands (* author for correspondence); ²Present address: Department of Molecular Cell Biology, University of Utrecht, Padualaan 8, 3584 CH Utrecht, Netherlands

Received 16 February 1994; accepted in revised form 20 May 1994

Key words: Agrobacterium, gene regulation, sensor protein, signal transduction, VirA protein, acetosyringone

Abstract

The VirA protein of *Agrobacterium tumefaciens* is thought to be a receptor for plant phenolic compounds such as acetosyringone. Although it is not known whether the interaction between VirA and the phenolics is direct or requires other phenolic-binding proteins, it is shown in this study that the first 280 amino acids of the VirA protein are not essential for the acetosyringone mediated *vir* gene induction response. Considering the fact that the cytoplasmic region between the amino acids 283 and 304 is highly conserved between the different VirA proteins, and that deletion of this region abolishes VirA activity, we suggest that the acetosyringone receptor domain is located in this cytoplasmic domain of the VirA protein.

Introduction

The products of the Agrobacterium tumefaciens virulence genes mediate a process that resembles bacterial conjugation [2] and by which part of the Ti plasmid (T-DNA) is transferred to the plant cell nucleus. The expression of specific T-DNA genes within the plant cell results in the production of plant hormones, such as auxin and cytokinin, which cause cell proliferation and the formation of a tumour. The expression of the virulence genes is controlled by the products of the virA and virG genes and dependent on the presence of plant phenolic compounds such as acetosyringone. The VirA and VirG proteins are members of a two-component regulatory family

and possess homology to other members such as EnvZ-OmpR and DctB-DctD. In two-component regulatory systems, one component senses environmental signals and activates the second component to carry out a function (for a recent review see [24]). In A. tumefaciens the VirA protein is thought to act as the receptor for plant phenolic compounds like acetosyringone. Whether this interaction between the phenolics and the VirA protein is direct or requires other phenolic-binding proteins is not known yet [14]. In vitro experiments showed that the cytoplasmic portion of the VirA protein has autokinase activity and is able to phosphorylate the VirG protein [8, 10, 11]. The VirG protein binds to specific sequences in the vir promoters ('vir box'), and it

is thought that upon phosphorylation the VirG protein becomes an efficient activator of vir gene transcription [12, 25, 27]. Sequence analysis of the virA gene predicted two hydrophobic regions within the VirA protein which might function as transmembrane regions [15, 18]. Indeed, topology studies with VirA-PhoA fusion proteins revealed that the VirA protein is located in the innermembrane of A. tumefaciens [19]. The topology of the VirA protein resembles that of other well-known receptor proteins such as the chemoreceptor Tar, a member of the class of methyl-accepting chemotaxis proteins that mediates chemotaxis towards attractants such as aspartate, maltose and phenol and away from repellents such as Ni^{2+} and glycerol [6, 9, 19]. For the chemoreceptor Tar it is known that the periplasmic domain plays an important role in ligand sensing, while the two transmembrane regions are thought to play an important role in the maintenance of protein conformation [23]. Studies with VirA-Tar chimeric proteins revealed that the periplasmic domain of the VirA protein is not essential for induction by phenolic compounds [19]. The periplasmic domain, however, is essential for the enhancement of vir gene induction by monosaccharides such as D-glucose [3, 29]. It is thought that the ChvE protein, which is homologous to the Escherichia coli galactose-binding protein, is able to bind monosaccharides and to enhance vir gene induction by binding to the periplasmic domain of VirA, thereby changing its conformation. This is confirmed by the observation that exchanging the periplasmic domain of VirA for that of Tar results in a protein which is non-responsive to sugars but locked in a highly responsive state which mimics the ChvE-occupied receptor conformation [31]. Melchers et al. [19] suggested that the receptor domain for the phenolic compounds might be located within, or in the vicinity of the second transmembrane domain. Hybrid proteins in which a region, containing the second transmembrane domain and the neighbouring cytoplasmic portion, was altered were not able to respond to the phenolic inducers. Recently, it was shown that a VirA deletion mutant which lacked the entire N-terminal part of the protein, including the two transmembrane regions, still had the ability to induce a virB-lacZreporter construct in the presence of $100 \,\mu M$ acetosyringone [4]. This indicated that neither of the two transmembrane domains is essential for the induction by phenolic compounds. Pazour et al. [26], however, reported that changing a leucine residue, at position 24 in the first transmembrane region, into a phenylalanine residue resulted in elevated vir gene expression in the absence of phenolic compounds. This indicated that the first transmembrane region plays an important role in determining the protein conformation which is essential for optimal VirA activity. In order to localize the phenolic receptor domain and to elucidate the role of the different regions in the activation of the VirA protein by phenolic compounds, several mutant proteins were constructed. In this report we confirm that the periplasmic and the two transmembrane regions are dispensable for the induction by phenolic compounds.

Materials and methods

Materials

Restriction endonucleases were purchased from either Promega Biotec. or New England Biolabs and used according to the suppliers' recommendations. The oligonucleotides used are listed in Table 1. *O*-Nitro- β -*D*-galactopyranoside and uridine were purchased from Sigma Chemical Co. and acetosyringone (AS) from EGA-chemie.

Bacterial strains and plasmids

Bacterial strains used were: Agrobacterium tumefaciens LBA2524 (pTiB6: Δ virA,virB-lacZ) [19] and E. coli JM101 [34] and PP1674 (dut,ung) [33].

In order to generate VirA-Tar chimeric proteins, specific restriction sites were introduced into the coding region of the *virA* gene (Fig. 1). Plasmid pST7111 was constructed by cloning a 4.5 kb *Eco* RI-*Bam* HI fragment of pRAL3254 [19], containing the complete *virA* gene of

Table 1. Oligonucleotides used in this study. Nucleotide substitutions are in bold type and restriction sites are underlined.

Mutagenesis	
22-mer (Bss HII)	5'-GGCTTC <u>GCGCGC</u> GTCTTAAAAT-3'
30-mer (Bss HII)	5'-CCCAAGAAAGATGCGCGCGCCTCCGCTCCTC-3'
33-mer (Sal I)	5'-TAACCAATCGGTTC <u>GTCGAC</u> GTAGCCTATAGAC-3'
21-mer (Bst BI)	5'-CTCACC <u>TTCGAA</u> ACATACTCC-3'
PCR	
26-mer (Bss HII)	5'-ATTAA <u>GCGCGC</u> CCGCGTAGTCACGCT-3'
23-mer	5'-CATCATCCGTACCGCTGAACGAC-3'
30-mer (Bss HII)	5'-CAAA <u>GCGCGC</u> GTATCTTTGCCCAGTGGCAA-3'
32-mer (Sal I)	5'-GTTC <u>GTCGAC</u> GTAGCCTAATGCCGTACCACGC-3'

pTi15955, into *Eco* RI-*Bam* HI-digested pTZ18R (USB corporation, Cleveland). The *virA* gene of pRAL7003 [19] was mutagenized by the method

of Kunkel *et al.* [13]. A 22-mer oligonucleotide was used to introduce a *Bss* HII restriction site in the *virA* gene. Due to the introduction of the



Fig. 1. Induction characteristics of *virA-tar* chimeric genes in *Agrobacterium*. The *vir* gene induction response of LBA2524 harbouring a wild-type *virA* gene (\circ), a *virA* gene containing several restriction sites (\triangle) or the *virA-tar* chimeric genes (\square) was tested in induction medium with 10 mM glucose and various concentrations of acetosyringone. Flasks were shaken for 18 h at 28 °C and units β -galactosidase were determined as described [20]. Open bars represent coding regions of *virA*, black and stippled bars represent coding regions of *tar*. The restriction sites *Bss* HII (B); *Xba* I (X); *Hpa* I (H) and *Sal* I (S) shown were used for the construction of the different *virA-tar* gene fusions.

Bss HII restriction site the amino acid glycine at position 15 in the VirA protein was changed into arginine. Subsequently a 4.5 kb Eco RI-Bam HI fragment, containing the mutagenized virA gene, was cloned into Eco RI-Bam HI-digested pTZ18R yielding pST7116. Using a 26-mer (Bss HII) oligonucleotide and a 23-mer oligonucleotide, a 180 bp fragment was amplified by PCR from pRBB6 [19]. This PCR fragment, coding for the first transmembrane region of the protein Tar, was digested with Bss HII and Nhe I and cloned into the Bss HII-Xba I digested pST7116, yielding pST7122. Plasmid pST7133 was constructed by mutagenizing plasmid pST7111. Using a 30-mer (Bss HII) oligonucleotide and a 33-mer (Sal I) oligonucleotide, a Bss HII and Sal I restriction site were introduced into the virA gene of pST7111. Due to the introduction of the Sal I restriction site the amino acids Lys-282 and Lys-283 were changed into Arg-282 and Arg-283. For the construction of plasmid pST7132 a 80 bp PCR fragment, coding for the second transmembrane region of the protein Tar, was amplified from pRBB6 using the 30-mer (Bss HII) oligonucleotide and the 32-mer (Sal I) oligonucleotide. This fragment was digested with Bss HII and Sal I and cloned into the Bss HII-Sal I-digested pST7133 to yield pST7132. Plasmid pST7134 was constructed by cloning a 1.2 kb Nhe I-Sna BI fragment of pST7137 into Nhe I-Sna BI-digested pST7116. For the construction of pST7136 a 600 bp PCR fragment, coding for the N-terminal part of the protein Tar, was amplified from pRBB6 using the 26-mer (Bss HII) oligonucleotide and the 32-mer (Sal I) oligonucleotide. This PCR fragment was digested with Bss HII-Sal I and cloned into Bss HII-Sal I-digested pST7134 to yield pST7136. Plasmid pST7137 was constructed by introducing a Sal I and Bst BI restriction site into the virA gene of pST7111 using a 33-mer (Sal I) and a 21-mer (Bst BI) oligonucleotide. Deletion constructs were made by digesting pST7137 with the following restriction enzymes: pST7138, ΔSal I-Bst BI; pST7139, ΔBst BI-*Eco* 47III; pST7140, ΔSal I-*Eco* 47III. The restriction sites were filled in by the Klenow fragment of DNA polymerase, followed by ligation. To verify the DNA sequence, all DNA fragments containing mutations and PCR generated DNA fragments were sequenced by the chain termination method [28].

Plasmids pST7111 to pST7140 were linearized with *Eco* RI and cloned into the unique *Eco* RI restriction site of the IncP class vector pRL750 [19], resulting in pRAL7111 to pRAL7140, respectively. Cointegrates were transferred to *A. tumefaciens* by electroporation [17].

Media and antibiotics

E. coli strains were grown at 37 °C in LC medium [7] and *A. tumefaciens* strains at 29 °C in minimal medium (MM) [7]. Induction medium for *A. tumefaciens vir* induction (IM) contained MM salts, 62.5 mM potassium phosphate pH 5.3 and 0.5% glycerol as a carbon source. Antibiotic concentrations (μ g/ml) for *E. coli* strains were: carbenicillin, 100; kanamycin, 25; tetracyclin, 10; and for *A. tumefaciens*: carbenicillin, 75; kanamycin, 100; tetracyclin, 2; rifampicin, 20; spectinomycin, 250.

Induction of Agrobacterium tumefaciens

Bacteria were grown overnight in MM to an optical density (OD_{600}) of 0.7–0.8. Cells were pelleted by centrifugation and diluted 1:20 in IM with various concentrations of acetosyringone (stock solution in DMSO). 10 mM glucose was supplemented as indicated. Flasks were shaken for 18 h at 28 °C and β -galactosidase activity was measured as described [20].

Assays were repeated at least three times in which similar data were obtained.

Results

Effects of introduced restriction sites on VirA signalling

In order to find out whether the two transmembrane regions play a role in acetosyringone recognition and/or signal transduction, the properties of several *virA-tar* hybrid genes were studied. Therefore, genes were constructed coding for chimeric proteins of which (1) the first transmembrane region, (2) the second transmembrane region, or (3) the entire N-terminus of the VirA protein were precisely exchanged for the corresponding region of the *E. coli* chemoreceptor Tar.

To be able to construct the virA-tar hybrid genes, several unique restriction sites were first introduced into the coding region of the virA gene [19, 31]. It had been determined previously that the introduction of the Xba I restriction site resulting in the VirA(Ser-44, Arg-45) mutant affected the functioning of VirA, i.e. the mutant protein no longer responded to sugars and the ChvE protein and was locked in a low responsive state [31]. Strains containing the VirA(Arg-15) protein, encoded by the virA(Bss HII) gene or the VirA(Val-251) protein, encoded by the virA(Hpa I) gene, however, showed the same induction pattern as a strain containing a wild-type VirA protein [31]. Plasmid pRAL7133 contains a virA gene in which a Sal I and Bss HII restriction site had been introduced and codes for the VirA(Arg-282, Arg-283) protein. Introduction of the Bss HII site in the virA gene did not lead to alterations in the VirA protein sequence. To determine the effects of the amino acid alterations caused by the introduction of the Sal I restriction site, plasmid pRAL7133 was introduced into A. tumefaciens strain LBA2524 and vir gene induction was measured. Figure 1b shows that LBA2524(pRAL-7133) had a similar vir gene induction response as LBA2524(pRAL7111), which harbours a wildtype VirA protein, indicating that the amino acid alterations caused by the introduction of the Sal I restriction site have no negative effect on the virgene induction response.

Role of the membrane-spanning regions

To determine the role of the different VirA domains in signal perception and transduction pRAL7122, pRAL7132 and pRAL7136 were, together with several controls, introduced into *A. tumefaciens* strain LBA2524. In this *A. tumefaciens* strain the entire *virA* gene is deleted and a *virB-lacZ* reporter gene is present in order to monitor vir gene expression [19]. The vir gene induction response of the resulting strains was determined in the presence of 10 mM D-glucose and various concentrations of acetosyringone. Figure 1 shows that in the presence of $100 \,\mu M$ acetosyringone all strains showed high vir gene induction. This indicates that the first 280 amino acids of the VirA protein are not essential for the recognition of acetosyringone. At lower concentrations of acetosyringone, however, differences in the induction response patterns of the different strains were observed. In the presence of $5 \,\mu\text{M}$ acetosyringone LBA2524(pRAL7122) and LBA2524(pRAL7132), which harbour chimeric proteins of which respectively the first or the second transmembrane region of VirA was exchanged for that of Tar, showed only 6% vir gene induction activity as compared to a strain harbouring a wild-type VirA protein. Exchanging the entire N-terminus of the VirA protein by that of Tar resulted in a chimeric protein which was nonresponsive to the presence of $5 \,\mu M$ acetosyringone. To obtain a weak vir gene induction response at least 25 μ M acetosyringone needed to be present in the induction medium in the case of LBA2524(pRAL7136).

Sugars like *D*-glucose are able to enhance the *vir* gene induction response in wild-type strains [3, 29]. Determining the responsiveness of the three strains, containing the VirA-Tar chimeric proteins, to ChvE inducing sugars revealed that LBA2524(pRAL7122), LBA2524(pRAL7132) and LBA2524(pRAL7136) are no longer responsive to sugars and the ChvE protein (data not shown). Testing the responsiveness of the three strains towards the natural inducers of the Tar protein, revealed that neither aspartate (1 mM) nor phenol (200 μ M) was able to induce the virulence genes in LBA2524 harbouring pRAL7122, pRAL7132 or pRAL7136 (results not shown).

Tumour response of the strains harbouring the chimeric proteins

Figure 1 shows that exchanging different domains of the VirA protein by those of the chemorecep-



Fig. 2. Tumour response of the virA-tar chimeric genes in Agrobacterium tumefaciens strain LBA2524 upon infection on stems of Kalanchoë tubiflora. Tumour formation was evaluated after 3 weeks.

tor Tar resulted in chimeric proteins which are still able to respond to acetosyringone, although the amount of acetosyringone needed to provoke a vir gene induction response was higher than observed with a wild-type strain. To find out whether strains harbouring these proteins were still able to provoke tumours LBA2524(pRAL-LBA2524(pRAL7132) 7122), and LBA-2524(pRAL7136) were inoculated on Kalanchoë tubiflora. Figure 2 shows that all strains were able to provoke tumours as expected, although the tumours induced by LBA2524(pRAL7136) were smaller in size than those formed by the wild-type strain LBA2524(pRAL7111).

Deletion analysis of the VirA protein

Melchers *et al.* [19] speculated that the receptor domain for the phenolic compounds might be located within or in the vicinity of the second transmembrane region. Figure 1 shows that the Nterminal part of the VirA protein, including the second transmembrane domain, is not essential for the induction by phenolics. In order to determine the role of the cytoplasmic linker domain in the acetosyringone mediated *vir* gene induction response and in an attempt to localize the receptor domain for the phenolic compounds, several deletion constructs were created (Fig. 3). Testing the responsiveness of these mutant strains to acetosyringone revealed that none of the mutants harbouring the plasmids pRAL7138, pRAL7139 or pRAL7140 was able to induce the *virB-lacZ* reporter construct in the presence of acetosyringone (Table 2). This could indicate either that the cytoplasmic region close to the second transmembrane region is essential for the induction by acetosyringone or that the deletions lead to an altered VirA protein conformation, resulting in



Fig. 3. Schematic representation of the VirA deletion mutants. The restriction sites *Sal* I (S), *Bst* BI (B) and *Eco* 47III (E) used to create the deletions are indicated.

Ta	ble 2.	Effect of	deletions	on	the	acetosyringone-mediated
vir	gene	induction	response.			

Strain/plasmid	- AS	100 µM AS	
LBA2524(pRAL7137)	31	1928	
LBA2524(pRAL7138)	24	33	
LBA2524(pRAL7139)	33	35	
LBA2524(pRAL7140)	25	25	

mutant VirA proteins which are unable to activate the VirG protein.

Discussion

The VirA protein is an inner-membrane protein which acts as a receptor for plant phenolic compounds such as acetosyringone. To determine the role of the N-terminal part of the protein in the induction response, several hybrid proteins between VirA and the chemoreceptor Tar were constructed. For the protein Tar it is known that it exists as a homodimer and that the two transmembrane regions play an important role in dimerization and the maintenance of protein conformation [6, 16]. The periplasmic domain of the Tar protein is essential for ligand recognition and it is thought that binding of aspartate locks the dimer into a tighter complex, thereby decreasing the rate of dissociation and exchange of subunits in intact receptor dimers [21]. In this paper it is shown that the first 280 amino acids of the VirA protein are dispensable for the acetosyringonemediated vir gene induction response, although exchanging this region by that of the Tar protein results in a hybrid protein which is weakly responsive to acetosyringone. This indicates that the conformation of the N-terminal part plays an important role in determining the efficiency of the vir gene induction response. Exchanging either one of the two transmembrane regions of VirA for those of Tar resulted in hybrid proteins which are inducible by acetosyringone, but non-responsive to monosaccharides such as D-glucose (Fig. 1A/ B). This indicates that the two transmembrane regions are not merely hydrophobic membranespanning regions, but rather play an important role in establishing the conformation of the periplasmic domain of the protein which is essential for the enhancement of the *vir* gene induction response by monosaccharides such as *D*-glucose. Whether the VirA protein also exists as a (homo)dimer, and whether amino acids located in the two transmembrane regions play a role in the process of dimerization (e.g. by forming a leucine zipper) is not known at the moment.

The region of the VirA protein acting as receptor domain for the phenolic compounds has not been determined yet. Turk et al. [30, 32] showed that the VirA proteins of different Agrobacterium strains, including the limited-host-range strains Ag57 and Ag162, are activated by acetosyringone. This indicates that the domain, acting as receptor for phenolic compounds must be conserved between the different VirA proteins. In an attempt to locate the region which is essential for the acetosyringone mediated vir gene induction response, Melchers et al. [19] showed that an active VirA protein which has an internal deletion of the periplasmic domain, the second transmembrane domain and part of the cytoplasmic linker region up to amino acid Arg-323 (pRAL7045) is still tumorigenic on Nicotiana glauca but nonresponsive to acetosyringone. This suggests that a domain, important for the induction by the phenolic compounds is located within the deleted part of the protein. Results shown in Fig. 1 point out that the receptor domain for the phenolic compounds must be located in the cytoplasmic portion of the VirA protein since the N-terminal part of the VirA protein, including the second transmembrane domain, is not essential for the acetosyringone mediated vir gene induction. In order to determine the role of the cytoplasmic linker domain, near the second transmembrane domain, in the acetosyringone mediated vir gene induction response, several virA deletion mutants were constructed which encode mutant VirA proteins with deletions within this cytoplasmic region up to amino acid Arg-323 (Fig. 3). Table 2 shows that all deletions within the cytoplasmic linker region near the second transmembrane region of the VirA protein knock out the induction

(pTiA6)	FLGSASVGLC	LYIITLVYRL	RKKTDWLARR
(pTi15955)	• • • • • • • • • • •	• • • • • • • • • • •	
(pRiA4)	s.v	IS	.R
(pTiC58)	· V	IS	.RAT
(pTiAg162)	AVFF.	FG.VIHK.	.RRR
pRAL7138	$\rightarrow \leftarrow$	pRAL7139	\rightarrow
LDYEELIKEI	GVCFEGEAAT	TSS	-AQAALRIIQR
••••	• • • • • • • • • •		
• • • • • • • • • • •	GG	A	G
····v	GG	LNS	sv
FVK.	DSTE.	KQSLK	EGTIEN

←---

Fig. 4. Comparison of the amino acid sequences of the linker region near the second transmembrane region from various VirA proteins. The sequence of pTiA6 is shown from amino acid 261 with the one-letter symbol. Identical amino acids are idicated by dots, missing amino acids are indicated by hyphens. The amino acids located in the second transmembrane region are indicated in bold. The regions deleted in pRAL7138 and pRAL7139, respectively, are indicated with arrows.

response. Since these deletions do not affect the kinase domain, it is likely that they affect the phenol-binding domain. However, it may equally well be that the deletions destroyed protein topology, thereby leading to mutants which are inactive or not targeted to the inner membrane of Agrobacterium. Acetosyringone is a very hydrophobic compound which accumulates in bacterial membranes (S. Turk, unpublished observations). Deleting part of the cytoplasmic linker region, between amino acids 283 and 323, affects the induction response by acetosyringone. Considering the fact that the cytoplasmic linker region between amino acids 283 and 304 is highly conserved between the different VirA proteins (Fig. 4) [5, 15, 18, 22], we suggest that the acetosyringone receptor domain is located in this cytoplasmic linker domain of the VirA protein.

References

- Ankenbauer RG, Best EA, Palanca CA, Nester EW: Mutants of the *Agrobacterium tumefaciens virA* gene exhibiting acetosyringone-independent expression of the vir regulon. Mol Plant-Microbe Interact 4: 400–406 (1991).
- 2. Beijersbergen A, Den Dulk-Ras A, Schilperoort RA,

Hooykaas PJJ: Conjugative transfer by the virulence system of *Agrobacterium tumefaciens*. Science 256: 1324–1327 (1992).

- Cangelosi GA, Ankenbauer RG, Nester EW: Sugars induce the *Agrobacterium tumefaciens* virulence genes through a periplasmic binding protein and a transmembrane signal protein. Proc Natl Acad Sci USA 87: 6708– 6712 (1990).
- Chang C, Winans SC: Functional roles assigned to the periplasmic, linker, and receiver domains of the Agrobacterium tumefaciens VirA protein. J Bact 174: 7033– 7039 (1992).
- Endoh H, Hirayama T, Aoyama T, Oka A: Characterization of the virA gene of the agropine-type plasmid pRiA4 of Agrobacterium rhizogenes. FEBS Lett 271: 28-32 (1990).
- Gardina P, Conway C, Kossman M, Manson M: Aspartate and maltose-binding protein interact with adjacent sites in the Tar chemotactic signal transducer of *Escherichia coli*. J Bact 174: 1528–1536 (1992).
- Hooykaas PJJ, Roobol C, Schilperoort RA: Regulation of the transfer of Ti plasmids of *Agrobacterium tumefaciens*. J Gen Microbiol 110: 99–109 (1979).
- Huang Y, Morel P, Powell B, Kado CI: VirA, a coregulator of Ti-specified virulence genes, is phosphorylated *in vitro*. J Bact 172: 1142–1144 (1990).
- Imae Y, Oosawa K, Mizuno T, Kihara M, Macnab R: Phenol: a complex chemoeffector in bacterial chemotaxis. J Bact 169: 371–379 (1987).
- Jin S, Prusti RK, Roitsch T, Ankenbauer RG, Nester EW: Phosphorylation of the VirG protein of *Agrobacterium tumefaciens* by the autophosphorylated VirA protein: essential role in biological activity of VirG. J Bact 172: 4945–4950 (1990).
- Jin S, Roitsch T, Ankenbauer RG, Gordon MP, Nester EW: The VirA protein of *Agrobacterium tumefaciens* is autophosphorylated and is essential for vir gene regulation. J Bact 172: 525-530 (1990).
- Jin S, Roitsch T, Christie PJ, Nester EW: The regulatory VirG protein specifically binds to a *cis*-acting regulatory sequence involved in transcriptional activation of *Agrobacterium tumefaciens* virulence genes. J Bact 172: 531– 537 (1990).
- Kunkel TA, Roberts JD, Zakour RA: Rapid and efficient site-specific mutagenesis without phenotypic selection. Meth Enzymol 154: 367–382 (1987).
- Lee K, Dudley MW, Hess KM, Lynn DG, Joerger RD, Binns AN: Mechanism of activation of *Agrobacterium* virulence genes: identification of phenol-binding proteins. Proc Natl Acad Sci USA 89: 8666–8670 (1992).
- Leroux B, Yanofsky MF, Winans SC, Ward JE, Ziegler SF, Nester EW: Characterization of the *virA* locus of *Agrobacterium tumefaciens*: a transcriptional regulator and host range determinant. EMBO J 6: 849–856 (1987).
- 16. Lynch BA, Koshland DE: disulfide cross-linking studies of the transmembrane regions of the aspartate sensory

receptor of *Escherichia coli*. Proc Natl Acad Sci USA 88: 10402–10406 (1991).

- Mattanovich D, Rüker F, da Câmara Machado A, Laimer M, Regner F, Steinkellner H, Himmler G, Katinger H: Efficient transformation of *Agrobacterium* spp. by electroporation. Nucl Acids Res 17: 6747 (1989).
- Melchers LS, Thompson DV, Idler KB, Neuteboom STC, de Maagd RA, Schilperoort RA, Hooykaas PJJ: Molecular characterization of the virulence gene virA of the Agrobacterium tumefaciens octopine Ti plasmid. Plant Mol Biol 9: 635–645 (1987).
- Melchers LS, Regensburg-Tuïnk TJG, Bourret RB, Sedee NJA, Schilperoort RA, Hooykaas PJJ: Membrane topology and functional analysis of the sensory protein VirA of Agrobacterium tumefaciens. EMBO J 8: 1919–1925 (1989).
- Miller JH: Experiments in Molecular Genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1972).
- Milligan DL, Koshland DE: Site-directed cross-linking: establishing the dimeric structure of the aspartate receptor of bacterial chemotaxis. J Biol Chem 263: 6268–6275 (1988).
- Morel P, Powell BS, Rogowsky PM, Kado CI: Characterization of the *virA* virulence gene of the nopaline plasmid, pTiC58, of *Agrobacterium tumefaciens*. Mol Microbiol 3: 1237–1246 (1989).
- Oosawa K, Simon M: Analysis of mutations in the transmembrane region of the aspartate chemoreceptor in *Escherichia coli*. Proc Natl Acad Sci USA 83: 6930–6934 (1986).
- Parkinson JS, Kofoid EC: Communication modules in bacterial signaling proteins. Annu Rev Genet 26: 71–112 (1992).
- Pazour GJ, Das A: Characterization of the VirG binding site of *Agrobacterium tumefaciens*. Nucl Acids Res 18: 6909-6913 (1990).

- Pazour GJ, Ta CN, Das A: Mutants of Agrobacterium tumefaciens with elevated vir gene expression. Proc Natl Acad Sci USA 88: 6941-6945 (1991).
- Powell BS, Kado CI: Specific binding of VirG to the vir-box requires a C-terminal domain and exhibits a minimum concentration threshold. Mol Microbiol 4: 2159– 2166 (1990).
- Sanger F, Nicklen S, Coulson A: DNA sequencing with chain terminating inhibitors. Proc Natl Acad Sci USA 74: 5463–5467 (1977).
- Shimoda N, Toyoda-Yamamoto A, Nagamine J, Usami S, Katayama M, Sakagami Y, Machida Y: Control of expression of *Agrobacterium vir* genes by synergistic actions of phenolic signal molecules and monosaccharides. Proc Natl Acad Sci USA 87: 6684–6688 (1990).
- Turk SCHJ, Melchers LS, den Dulk-Ras H, Regensburg-Tuïnk AJG, Hooykaas PJJ: Environmental conditions differentially affect vir gene induction in different Agrobacterium strains. Role of the VirA sensor protein. Plant Mol Biol 16: 1051–1059 (1991).
- Turk SCHJ, van Lange RP, Sonneveld E, Hooykaas PJJ: The chimeric VirA-Tar receptor protein is locked into a highly responsive state. J Bact 175: 5706–5709 (1993).
- Turk SCHJ, Nester EW, Hooykaas PJJ: The VirA promoter is a host-range determinant in *Agrobacterium tumefaciens*. Mol Microbiol 7: 719–724 (1993).
- 33. van Rijn PA, Goosen N, Turk SCHJ, van de Putte P: Regulation of phage Mu repressor transcription by IHF depends on the level of the early transcription. Nucl Acids Res 17: 10203–10212 (1989).
- 34. Yanisch-Perron C, Vieira J, Messing J: Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13 mp18 and pUC19 vectors. Gene 33: 103–119 (1985).