

Tangent fermions: massless fermions on a lattice Donís Vela, A.

Citation

Donís Vela, A. (2024, July 3). *Tangent fermions: massless fermions on a lattice*. Retrieved from https://hdl.handle.net/1887/3765898

Version:	Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral</u> <u>thesis in the Institutional Repository of the University</u> <u>of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/3765898

Note: To cite this publication please use the final published version (if applicable).

Stellingen

behorende bij het proefschrift

Tangent fermions: massless fermions on a lattice

- 1. The topological protection of a single Dirac cone on a space-time lattice requires not only chiral symmetry, but also the continuity of the evolution operator in the reciprocal space. [Chapter 2]
- 2. It is impossible to have a gauge invariant lattice formulation of massless fermions with a single topologically protected zeroth Landau level. [Chapter 4]
- 3. In a topological superconductor, a supercurrent cannot produce an inversion of velocity of the Majorana edge modes if the pairing is *p*-wave. [Chapter 5]
- 4. The injection of edge-vortices in a topological superconductor can be spoiled by the entrapment of excitations in the Josephson junction of the injector. [Chapter 6]
- 5. A zigzag-edge boundary condition for Dirac fermions can be replaced by a large vector potential parallel to the boundary.
- 6. The thermal-metal-insulator transition in a chiral p-wave superconductor is a percolation transition for Majorana fermions.
- 7. The exceptional points of a non-Hermitian Hamiltonian do not produce a singularity in the Josephson effect. [arXiv:2404.13976]
- 8. Although the Hamiltonian for tangent fermions is non-local, it can be written as a matrix product operator of finite bond dimension.
- 9. Within a given research area, there is a negative correlation between the length of the title of a scientific article and its quality.

Álvaro Donís Vela Leiden, 3 juli 2024