



Universiteit
Leiden
The Netherlands

The *Aspergillus nidulans* amdS gene as a marker for the identification of multicopy T-DNA integration events in *Agrobacterium*-mediated transformation of *Aspergillus awamori*

Michielse, C.B.; Ram, A.F.J.; Hondel, C.A.M.J.J. van den

Citation

Michielse, C. B., Ram, A. F. J., & Hondel, C. A. M. J. J. van den. (2004). The *Aspergillus nidulans* amdS gene as a marker for the identification of multicopy T-DNA integration events in *Agrobacterium*-mediated transformation of *Aspergillus awamori*. *Current Genetics*, 45, 399-403. doi:10.1007/s00294-004-0500-1

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3768699>

Note: To cite this publication please use the final published version (if applicable).

C. B. Michielse · A. F. J. Ram
C. A. M. J. J. van den Hondel

The *Aspergillus nidulans amdS* gene as a marker for the identification of multicopy T-DNA integration events in *Agrobacterium*-mediated transformation of *Aspergillus awamori*

Received: 28 January 2004 / Revised: 2 March 2004 / Accepted: 8 March 2004 / Published online: 26 March 2004
© Springer-Verlag 2004

Abstract The *Aspergillus nidulans amdS* selection marker was used for the identification of multicopy T-DNA insertions in *Agrobacterium*-mediated transformation of *Asp. awamori*. The selection of transformants on agar plates containing acetamide as sole nitrogen source and hygromycin resulted in a six-fold decrease in the transformation frequency, compared with the transformation frequency obtained after hygromycin selection alone. However, it was found that 47% of the transformants obtained after hygromycin and acetamide double selection contained multiple T-DNA integrations. Furthermore, it was found that the multicopy transformants could easily be identified based on their growth rate on agar plates containing acetamide medium. Based on these data, it can be concluded that the *amdS* marker can also be used as a selection marker in *Agrobacterium*-mediated transformation of *Asp. awamori* and that it is a very useful marker to identify those transformants containing multiple T-DNA integrations.

Keywords *Agrobacterium tumefaciens* · Transformation · Multicopy T-DNA integration · *amdS* selection marker · *Aspergillus awamori* · Filamentous fungi

Introduction

Agrobacterium tumefaciens, a plant pathogen, is widely used for the transformation of plants. *Agr. tumefaciens*

has the unique ability to transfer DNA (so-called transferred DNA or T-DNA), which is located between two inverted repeats, known as the left border (LB) and right border (RB), to its host. The transferred DNA is transported into the host as a single-stranded DNA molecule. Once inside the host, the DNA is targeted to the nucleus where it randomly integrates into the host genome (for reviews, see Gelvin 2000; Zupan and Zambryski 1995). *Agr. tumefaciens* is able to transfer its DNA not only to plants, but also to other organisms, including yeast, filamentous fungi and human cells (Bundock et al. 1995; de Groot et al. 1998; Kunik et al. 2001). The number of fungi that has been shown to be transformed by *Agr. tumefaciens* is rapidly increasing and *Agrobacterium*-mediated transformation (AMT) has been shown to be a good alternative to protoplast-based transformation methods (Amey et al. 2002; de Groot et al. 1998; Fitzgerald et al. 2003; Meyer et al. 2003). To date, only a limited number of selection markers have been used, including the auxotrophic *pyrG* marker and dominant selection markers based on resistance to the antibiotics hygromycin, geneticin, or phleomycin (de Groot et al. 1998; Gouka et al. 1999; Pardo et al. 2002; Vijn and Govers 2003). Analysis of the T-DNA integration patterns of transformants obtained from various fungi has revealed that the T-DNA integrates at random and predominantly as a single copy (Abuodeh et al. 2000; Bundock et al. 2002; Covert et al. 2001; Degefu and Hanif 2003; Leclerque et al. 2003; Vijn and Govers 2003), indicating that fungal AMT can be used for insertional mutagenesis and for the generation of DNA-insertion mutant banks.

To optimize AMT for various fungi, several variables [e.g. the influence of co-cultivation period and temperature, the *Agrobacterium*/spore ratio, the addition or omission of the *vir* gene inducer acetosyringone (AS) in the *Agrobacterium* pre-culture] have been found to influence transformation frequency (Abuodeh et al. 2000; Combier et al. 2003; Leclerque et al. 2003; Malonek and Meinhardt 2001; Michielse et al. 2004b; Mullins et al. 2001; Rho et al. 2001). Furthermore, it has been shown that these environmental conditions also

Communicated by U. Kück

C. B. Michielse (✉) · A. F. J. Ram · C. A. M. J. J. van den Hondel
Institute of Biology, Clusius Laboratory, Leiden University,
Wassenaarseweg 64, 2333 AL Leiden, The Netherlands
E-mail: michielse@rulbim.leidenuniv.nl
Tel.: +31-71-5274861
Fax: +31-71-5274999

A. F. J. Ram · C. A. M. J. J. van den Hondel
Department of Microbiology, TNO Nutrition,
Utrechtseweg 48, 3700 AJ Zeist, The Netherlands

have an influence on T-DNA copy number. For example, the addition of AS to the *Agrobacterium* pre-culture can result in a decrease (or an increase) in single-copy T-DNA integration (Combiér et al. 2003; Mullins et al. 2001; Rho et al. 2001). Prolongation of the co-cultivation period was shown to lead to multiple T-DNA integrations in *Magnaporthe grisea* (Rho et al. 2001). However, the length of the co-cultivation period had no influence on T-DNA copy number in *Fusarium oxysporum* (Mullins et al. 2001). Applying AMT for the generation of mutant banks, single-copy T-DNA integration is preferred as it enables linkage of an observed phenotype with a single alteration in the genome. For heterologous and homologous protein production in filamentous fungi, it has been shown that the introduction of multiple copies of the gene of interest results in an increase in protein production (Verdoes et al. 1995). Therefore, it would be beneficial to be able to generate and identify multicopy transformants obtained with AMT. Gouka et al. (1999) demonstrated the use of AMT in *Aspergillus awamori* for the generation of transformants carrying multiple copies of the gene of interest, using a binary vector carrying these multiple copies on its T-DNA.

A selection marker which is frequently used in protoplast-based transformation methods to generate transformants carrying multiple copies of the introduced DNA is the *Asp. nidulans amdS* selection marker (Hanegraaf et al. 1991; Kelly and Hynes 1985; Penttilä et al. 1987; Rodriguez and Yoder 1987). In this study, we demonstrate the use of the *Asp. nidulans amdS* selection marker and AMT in *Asp. awamori* to generate and identify multicopy T-DNA transformants.

Materials and methods

Strains, plasmids and growth conditions

Asp. awamori CBS115.52 (CBS, The Netherlands) was used as a recipient strain for transformation. *Agr. tumefaciens* LBA1100 (pAL1100ΔT-DNA, Δtra, Δocc; Beijersbergen et al. 1992), carrying pUR5750AmdS (this study, Fig. 2a) or pSDMAmdSΔBB (Michielse, Arentshorst, Ram and van den Hondel, submitted for publication; Fig. 2b) was used for AMT and was grown in LB medium (Sambrook et al. 1989) containing spectinomycin (250 μg/ml) and kanamycin (100 μg/ml) at 28°C. Introduction of plasmids into LBA1100 was performed as described by Mattanovich et al. (1989). *Escherichia coli* XL1-Blue (Stratagene) was used for construction of pUR5750AmdS and was grown in LB medium containing

kanamycin (25 μg/ml) at 37°C. The binary vector pUR5750AmdS, containing both the hygromycin and *amdS* selection markers, was constructed by inserting an *Xba*I fragment of p3SR2 (Corrick et al. 1987), corresponding to the *amdS* gene and its promoter and terminator sequences, into pUR5750 (de Groot et al. 1998) previously digested with *Xba*I.

AMT in *Asp. awamori*

AMT was performed as described by de Groot et al. (1998), with minor adjustments (Michielse et al. 2004b). Transformants were either selected on agar plates containing MM (Punt and van den Hondel 1992) supplemented with 100 μg/ml hygromycin or on agar plates containing acetamide as a sole nitrogen source (Kelly and Hynes 1985) with or without 100 μg/ml hygromycin, as indicated. All plates were supplemented with 200 μM cefotaxim to inhibit the growth of *Agr. tumefaciens*.

DNA isolation and Southern analysis

Fungal chromosomal DNA isolation and Southern analysis were performed as described by Kolar et al. (1988) and Michielse et al. (2004a), respectively. Chromosomal DNA was digested with *Bgl*III and probed with a 3.1-kb *Xho*I–*Hind*III fragment of pAN7.1 (Punt et al. 1987), corresponding to the hygromycin cassette, or with a 2.6-kb *Xba*I fragment of p3SR2 (Corrick et al. 1987), corresponding to the *amdS* expression cassette.

Results and discussion

The use of the dominant *Asp. nidulans amdS* selection marker (Hynes and Pateman 1970) to identify multicopy T-DNA transformants of *Asp. awamori* obtained with AMT was assessed. *Asp. awamori* was transformed with *Agrobacterium* strain LBA1100 carrying the binary vector pUR5750AmdS, which contains both the hygromycin and *amdS* expression cassettes on its T-DNA. *Asp. awamori* was also transformed with *Agrobacterium* strain LBA1100 carrying the binary vector pSDMAmdSΔBB, which contains only the *amdS* expression cassette on its T-DNA. Depending upon the binary vector used, transformants were selected on agar plates containing medium supplemented with hygromycin, hygromycin and acetamide, or only acetamide.

Selection of transformants obtained with *Agr. tumefaciens* carrying pUR5750AmdS on medium containing only hygromycin yielded the highest transformation efficiency (Table 1). Double selection of the transformants obtained with the same binary vector on

Table 1 Transformation frequency and T-DNA copy number obtained after different selection conditions. *amdS* *Aspergillus nidulans* acetamidase gene, *hph* hygromycin resistance gene

Binary vector	Selection	Number of transformants/10 ⁶ spores	Number of transformants analyzed	Number of transformants containing more than one T-DNA
pUR5750AmdS	<i>hph</i>	183 ± 29 ^a	10	0
pUR5750AmdS	<i>hph</i> and <i>amdS</i>	29 ± 17 ^a	15	7
pSDMAmdSΔBB	<i>amdS</i>	0.2 ± 0.4 ^b	3	1

^aThe value represents the mean of three independent of experiments

^bThe value represents the mean of four independent experiments

acetamide medium containing hygromycin resulted in a six-fold decrease in the transformation frequency (Table 1). The lower transformation efficiency is most likely due to more stringent selection conditions to obtain transformants. It is known that the use of the *amdS* marker requires high expression levels of the *amdS* gene to allow growth on medium containing acetamide. High expression levels can be the result of either multicopy integration or integration at a certain locus, which allows efficient transcription (Hanegraaf et al. 1991; Kelly and Hynes 1985; Penttila et al. 1987; Rodriguez and Yoder 1987). The selection of transformants obtained with the binary vector which carried only the *amdS* expression cassette between the T-DNA borders (pSDMAmdS Δ BB), using acetamide-containing agar plates, resulted in a low transformation frequency (Table 1). About 20 co-cultivations were performed, yielding only three transformants. Some background growth of *Asp. awamori* under these selection conditions was observed. It is likely that *Asp. awamori* contains a low level of endogenous acetamidase activity, which caused this background growth and complicated the identification of transformants. These putative transformants grew very slowly and each remained a fluffy colony of white mycelium with very few spores. It is possible that the identification of putative *amdS* positive transformants may be facilitated by performing the co-cultivation on black filter paper rather than on white nitrocellulose membranes, as both are equally efficient (Covert et al. 2001). Alternatively, the background growth might be reduced if fungal and bacterial cells are recovered from the filters after co-cultivation and subsequently spread onto selection media. This method has been shown to reduce the background growth of *Beauveria bassiana* in AMT (Leclerque et al. 2003). Since variation in co-cultivation conditions has been shown to influence transformation frequency and T-DNA copy number (Abuodeh et al. 2000; Combier et al. 2003; Leclerque et al. 2003; Malonek and Meinhardt 2001; Michielse et al. 2004b; Mullins et al. 2001; Rho et al. 2001), it might be possible to increase the transformation frequency obtained by AMT in *Asp. awamori* based on the *amdS* selection marker by using different co-cultivation conditions.

It should be noted that the addition of AS to induce expression of the virulence genes of *Agr. tumefaciens* during co-cultivation is an absolute requirement to obtain transformants. Furthermore, all transformants obtained were mitotically stable (data not shown).

All transformants obtained were tested for their ability to grow on agar plates containing acetamide or hygromycin. A total of 25 [pUR5750AmdS] transformants, of which ten were obtained after selection on minimal medium with hygromycin (designated pUR5750AmdS-H) and 15 obtained after selection on acetamide medium supplemented with hygromycin (designated pUR5750AmdS-HA), were replica-plated onto agar plates containing either minimal medium supplemented with hygromycin or acetamide medium.

All 25 transformants were able to grow on minimal medium containing hygromycin and no growth differences between the transformants were observed under these conditions (Fig. 1a,c). However, growth differences between the transformants were observed on acetamide medium (Fig. 1b,d). The ten pUR5750AmdS-H transformants remained small on acetamide medium and, even after prolonged incubation, these colonies did not grow further and did not sporulate. Seven of the 15 pUR5750AmdS-HA transformants (Fig. 1d, numbers 1, 2, 3, 6, 7, 9, 15; identified in caption) grew much faster on acetamide medium compared with the pUR5750AmdS-H transformants, suggesting that these transformants may contain more than one T-DNA copy. Also, the three [pSDMAmdS Δ BB] transformants were able to grow on agar plates containing acetamide and no major growth differences between the three transformants were observed (data not shown).

Southern analysis was performed to determine the T-DNA copy number. Chromosomal DNA was digested with *Bgl*III and probed with hygromycin (*hph*) and *amdS* genes to determine single and multicopy T-DNA integration events. A single T-DNA integration event was expected to result in the detection of one fragment when probed with the *hph* gene ($\geq 4,674$ bp) and in two fragments ($\geq 4,563$ bp, ≥ 820 bp) when probed with the *amdS* gene (Fig. 2a). Based on previous studies, we know that 0–30% of the transformants contain more than one T-DNA copy when selected on medium containing hygromycin (Michielse et al. 2004a; Michielse, unpublished data). In this study, we found that all ten pUR5750AmdS-H transformants contained a single T-DNA integration. The Southern analysis result of a representative pUR5750AmdS-H transformant is shown in Fig. 2c,d (lane 1).




Fig. 1a–d Growth phenotype of transformants grown on hygromycin and acetamide selection plates. **a, b** Ten pUR5750AmdS transformants were obtained after selection on hygromycin-containing agar plates (*top row*, from left to right: numbers 1–5; *bottom row*: numbers 6–10) and replica-plated onto hygromycin-containing agar plates (**a**) or acetamide-containing agar plates (**b**). **c, d** pUR5750AmdS transformants obtained after selection on acetamide- and hygromycin-containing agar plates (*top row*, from left to right: numbers 1–5; *middle row*: numbers 6–10; *bottom row*: numbers 11–16) and replica-plated onto hygromycin-containing agar plates (**c**) or acetamide-containing agar plates (**d**)




Fig. 2 **a** Schematic representation of pUR5750AmdS T-DNA. *LB* Left border, *RB* right border. *B* *Bg*II restriction site, *Pgpd* *Asp. nidulans* glyceraldehyde-3-phosphate dehydrogenase promoter, *hph* hygromycin resistance gene, *T trpC* *Asp. nidulans trpC* terminator, *amdS* *Asp. nidulans* acetamidase enzyme, *nptII* neomycin/kanamycin resistance gene. **b** Schematic representation of pSDMAmdSΔBB T-DNA. **c, d** Southern analysis of [pUR5750AmdS] and [pSDMAmdSΔBB] transformants. Chromosomal DNA was digested with *Bg*II and hybridized with a hygromycin (*hph*) probe (**c**) and *amdS* probe (**d**). Lane 1 pUR5750AmdS H-1, lane 2 pUR5750AmdS HA-1, lane 3 pUR5750AmdS HA-3, lane 4 pUR5750AmdS HA-2, lane 5 pSDMAmdSΔBB 6

Of the 15 pUR5750AmdS-HA transformants, eight single, four double and three triple T-DNA integrations were found. The double T-DNA integration occurred as inverted repeats at either the LB repeat (see Fig. 2c,d; 9,348-bp fragment in c, two times in d, lane 2, fragments of $\geq 4,563$ bp and ≥ 820 bp) or the RB repeat (see Fig. 2c,d; two times in c, $\geq 4,674$ -bp fragments, two times in d, lane 3, 9,126-bp and 820-bp fragments). In three of the 15 transformants, multiple fragments were found for the *hph* and the *amdS* probes, indicating a triple T-DNA integration as either tandem or inverted repeats (Fig. 2c,d, lane 4).

The transformants obtained using [pSDMAmdSΔBB] were also analyzed by Southern analysis to determine the T-DNA copy number (Fig. 2d, lane 5). With this vector, a single T-DNA integration would be expected to result in the detection of two fragments (Fig. 2a; $\geq 1,940$ bp, ≥ 875 bp). This pattern was observed in two of the three transformants obtained (data not shown). Southern analysis of the third transformant revealed the presence of four hybridizing bands. Fragments of 1,750 bp and 2,815 bp were observed (Fig. 2d, lane 5), indicating inverted and tandem integration, respectively.

Furthermore, two fragments of 8.5 kb and 8 kb (two times $\geq 1,940$ bp) were observed. This hybridization pattern indicates that this transformant contains three T-DNA copies which are linked.

Based on the Southern analysis, the pUR5750AmdS-HA1, HA2, HA3, HA6, HA7, HA14 and HA15 transformants were identified as multicopy transformants. Six out of these seven transformants grew well on acetamide plates, indicating a good correlation between the Southern data and the growth phenotype. pUR5750AmdS-HA transformant 9 grew and sporulated well on agar plates containing acetamide medium. However, its sporulating phenotype was different from the other transformants (Fig. 1d). Southern analysis revealed that this transformant contained a single T-DNA copy (data not shown). It could be that, in this transformant, the location of the T-DNA integration into the genome resulted in this phenotype. Conversely, pUR5750AmdS-HA transformant 14 was identified as a multicopy transformant, based on Southern analysis (data not shown). However, on agar plates containing acetamide, this transformant did not grow better than the single copy pUR5750AmdS-H transformants (Fig. 1). This could be due to the location of the T-DNA integration into the genome, such that the *amdS* gene is not expressed highly enough to result in better growth of this transformant than the wild-type *Asp. awamori* strain on agar plates containing acetamide.

Based on our results, it can be concluded that the *amdS* marker can be used as a selection marker for AMT in *Asp. awamori* and that with this marker growth on agar plates containing acetamide correlates with the T-DNA copy number enabling the identification of multicopy T-DNA transformants.

Acknowledgements We would like to thank Simon Flitter and Patricia vanKuyk for critically reading this manuscript. This work was supported by Unilever Research, The Netherlands.

References

- Abuodeh RO, Orbach MJ, Mandel MA, Das A, Galgiani JN (2000) Genetic transformation of *Coccidioides immitis* facilitated by *Agrobacterium tumefaciens*. *J Infect Dis* 181:2106–2110
- Amey RC, Athey-Pollard A, Burns C, Mills PR, Bailey A, Foster GD (2002) PEG-mediated and *Agrobacterium*-mediated transformation in the mycopathogen *Verticillium fungicola*. *Mycol Res* 106:4–11
- Beijersbergen AG, Den Dulk A, Schilperoort RA, Hooykaas PJ (1992) Conjugative transfer by the virulence system of *Agrobacterium tumefaciens*. *Science* 256:1324–1327
- Bundock P, Dulk-Ras A, Beijersbergen A, Hooykaas PJ (1995) Trans-kingdom T-DNA transfer from *Agrobacterium tumefaciens* to *Saccharomyces cerevisiae*. *EMBO J* 14:3206–3214
- Bundock P, Attikum H van, Dulk-Ras A, Hooykaas PJ (2002) Insertional mutagenesis in yeasts using T-DNA from *Agrobacterium tumefaciens*. *Yeast* 19:529–536
- Combiér JP, Melayah D, Raffier C, Gay G, Marmeisse R (2003) *Agrobacterium tumefaciens*-mediated transformation as a tool for insertional mutagenesis in the symbiotic ectomycorrhizal fungus *Hebeloma cylindrosporium*. *FEMS Microbiol Lett* 220:141–148

- Corrick CM, Twomey AP, Hynes MJ (1987) The nucleotide sequence of the *amdS* gene of *Aspergillus nidulans* and the molecular characterization of 5' mutations. *Gene* 53:63–71
- Covert SF, Kapoor P, Lee M, Briley A, Nairn CJ (2001) *Agrobacterium*-mediated transformation of *Fusarium circinatum*. *Mycol Res* 105:259–264
- Degefu Y, Hanif M (2003) *Agrobacterium tumefaciens*-mediated transformation of *Helminthosporium turcicum*, the maize leaf-blight fungus. *Arch Microbiol* 108:279–284
- Fitzgerald AM, Mudge AM, Gleave AP, Plummer KM (2003) *Agrobacterium* and PEG-mediated transformation of the phytopathogen *Venturia inaequalis*. *Mycol Res* 107:803–810
- Gelvin SB (2000) *Agrobacterium* and plant genes involved in T-DNA transfer and integration. *Annu Rev Plant Physiol Plant Mol Biol* 51:223–256
- Gouka RJ, Gerk C, Hooykaas PJ, Bundock P, Musters W, Verrips CT, Groot MJ de (1999) Transformation of *Aspergillus awamori* by *Agrobacterium tumefaciens*-mediated homologous recombination. *Nat Biotechnol* 17:598–601
- Groot MJ de, Bundock P, Hooykaas PJ, Beijersbergen AG (1998) *Agrobacterium tumefaciens*-mediated transformation of filamentous fungi. *Nat Biotechnol* 16:839–842
- Hanegraaf PP, Punt PJ, Hondel CA van den, Dekker J, Yap W, Verseveld HW van, Stouthamer AH (1991) Construction and physiological characterization of glyceraldehyde-3-phosphate dehydrogenase overproducing transformants of *Aspergillus nidulans*. *Appl Microbiol Biotechnol* 34:765–771
- Hynes MJ, Pateman JA (1970) The genetic analysis of regulation of amidase synthesis in *Aspergillus nidulans*. I. Mutants able to utilize acrylamide. *Mol Gen Genet* 108:97–106
- Kelly JM, Hynes MJ (1985) Transformation of *Aspergillus niger* by the *amdS* gene of *Aspergillus nidulans*. *EMBO J* 4:475–479
- Kolar M, Punt PJ, Hondel CA van den, Schwab H (1988) Transformation of *Penicillium chrysogenum* using dominant selection markers and expression of an *Escherichia coli lacZ* fusion gene. *Gene* 62:127–134
- Kunik T, Tzfira T, Kapulnik Y, Gafni Y, Dingwall C, Citovsky V (2001) Genetic transformation of HeLa cells by *Agrobacterium*. *Proc Natl Acad Sci USA* 98:1871–1876
- Leclercq A, Wan H, Abschutz A, Chen S, Mitina GV, Zimmermann G, Schairer HU (2003) *Agrobacterium*-mediated insertional mutagenesis (AIM) of the entomopathogenic fungus *Beauveria bassiana*. *Curr Genet* (in press)
- Malonek S, Meinhardt F (2001) *Agrobacterium tumefaciens*-mediated genetic transformation of the phytopathogenic ascomycete *Calonectria morganii*. *Curr Genet* 40:152–155
- Mattanovich D, Ruker F, Machado AC, Laimer M, Regner F, Steinkellner H, Himmler G, Katinger H (1989) Efficient transformation of *Agrobacterium* spp. by electroporation. *Nucleic Acids Res* 17:6747
- Meyer V, Mueller D, Strowig T, Stahl U (2003) Comparison of different transformation methods for *Aspergillus giganteus*. *Curr Genet* 43:371–377
- Michiels CB, Ram AFJ, Hooykaas PJJ, Hondel CAMJJ van den (2004a) *Agrobacterium*-mediated transformation of *Aspergillus awamori* in the absence of full length VirD2, VirC2 and VirE2 leads to the insertion of aberrant T-DNA structures. *J Bacteriol* (in press)
- Michiels CB, Ram AFJ, Hooykaas PJJ, Hondel CAMJJ van den (2004b) Role of bacterial virulence proteins in *Agrobacterium*-mediated transformation of *Aspergillus awamori*. *Fungal Genet Biol* (in press)
- Mullins ED, Chen X, Romaine P, Raina R, Geiser DM, Kang S (2001) *Agrobacterium*-mediated transformation of *Fusarium oxysporum*: an efficient tool for insertional mutagenesis and gene transfer. *Phytopathology* 91:173–180
- Pardo AG, Hanif M, Raudaskoski M, Gorfer M (2002) Genetic transformation of ectomycorrhizal fungi mediated by *Agrobacterium tumefaciens*. *Mycol Res* 106:132–137
- Penttila M, Nevalainen H, Ratto M, Salminen E, Knowles J (1987) A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*. *Gene* 61:155–164
- Punt PJ, Hondel CA van den (1992) Transformation of filamentous fungi based on hygromycin B and phleomycin resistance markers. *Methods Enzymol* 216:447–457
- Punt PJ, Oliver RP, Dingemans MA, Pouwels PH, Hondel CA van den (1987) Transformation of *Aspergillus* based on the hygromycin B resistance marker from *Escherichia coli*. *Gene* 56:117–124
- Rho HS, Kang S, Lee YH (2001) *Agrobacterium tumefaciens*-mediated transformation of the plant pathogenic fungus, *Magnaporthe grisea*. *Mol Cells* 12:407–411
- Rodriguez RJ, Yoder OC (1987) Selectable genes for transformation of the fungal plant pathogen *Glomerella cingulata* f.sp. *phaseoli* (*Colletotrichum lindemuthianum*). *Gene* 54:73–81
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*, 2nd edn. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Verdoes JC, Punt PJ, Hondel CAM van den (1995) Molecular genetic strain improvement for the overproduction of fungal proteins by filamentous fungi. *Appl Microbiol Biotechnol* 43:195–205
- Vijn I, Govers F (2003) *Agrobacterium tumefaciens* mediated transformation of the oomycete plant pathogen *Phytophthora infestans*. *Mol Plant Pathol* 4:459–467
- Zupan JR, Zambryski P (1995) Transfer of T-DNA from *Agrobacterium* to the plant cell. *Plant Physiol* 107:1041–1047