



Universiteit
Leiden
The Netherlands

Cholecystectomy following EUS-guided gallbladder drainage in patients with acute cholecystitis at high surgical risk: friend or foe?

Larghi, A.; Wanrooij, R.L.J. van; Bronswijk, M.; Vanella, G.; Kunda, R.; Pérez-Miranda, M.; ... ; Merwe, S.W. van der

Citation

Larghi, A., Wanrooij, R. L. J. van, Bronswijk, M., Vanella, G., Kunda, R., Pérez-Miranda, M., ... Merwe, S. W. van der. (2024). Cholecystectomy following EUS-guided gallbladder drainage in patients with acute cholecystitis at high surgical risk: friend or foe? *Gut*. doi:10.1136/gutjnl-2024-332273

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3947629>

Note: To cite this publication please use the final published version (if applicable).

Cholecystectomy following EUS-guided gallbladder drainage in patients with acute cholecystitis at high surgical risk: friend or foe?

We read with great interest the paper by Bang *et al.*¹ reporting their single-centre retrospective experience in 25 patients with acute cholecystitis, who were deemed at increased surgical risk and treated by EUS-guided gallbladder drainage (EUS-GBD) using LAMS. Three patients underwent surgery because of persistent biliary-type symptoms, but the presence of LAMS precluded successful laparoscopic cholecystectomy (LC) and open or subtotal cholecystectomy was required. Diverging from the recent guidelines of the European Society of Gastrointestinal Endoscopy and the American Gastroenterological Association,^{2,3} the authors concluded that EUS-GBD should only be considered in patients for whom surgery would never be an option. We thank Bang *et al.* for sharing their experience with these unfortunate surgical outcomes, which encourages further discussion on how to use this technique. However, some of the points raised in their study deserve further considerations.

First, a clear-cut definition on how patients should be defined as high surgical risk versus a 'never surgery' candidates is completely lacking. This is a critical issue that has not been adequately addressed even in the literature and needs further clarification. This semantic difference seems mostly dependent on the different probability to survive up to a reasonable recovery after surgical intervention, implying that all patients can potentially undergo surgery with different preprocedural mortality's risk.^{4,5}

Second, in contrast with Bang *et al.*¹ larger series demonstrated no significant difference in conversion rate from laparoscopic to open cholecystectomy following EUS-GBD compared with percutaneous drainage (PT-GBD).⁶ Of note, in a comparative multicentre study, surgery following EUS-GBD was associated with a significant decrease in operative time, time to symptom resolution and length of stay compared with PT-GBD.⁷ In line with this study, high technical success rates have been reported in a large cohort of patients who underwent EUS-GBD with LAMS and second-stage LC, provided LAMS removal before surgery.⁸ These pilot findings suggest that surgery can still

be offered safely to patients judged at high surgical risk, who eventually need surgery.

Third, another important point neglected in the study by Bang *et al.*¹ is the potential harm associated with PT-GBD, which can be associated with adverse events (AEs) in up to 40% of cases, thus denying patients a safer and more effective approach. Indeed, according to the pivotal randomised trial by Teoh *et al.*⁹ the number needed to harm for generating 1-year AEs is close to two when preferring PT-GBD over EUS-GBD.

At last, persistence of biliary type was observed in 47.1% of the survivors. These long-term issues might simply reflect the protocol used in the study, where LAMS was kept in situ indefinitely. Conversely, in a prospective multicentre observational study conducted in Spain where LAMS was left in situ for at least 1 year in 45 patients (55% of the entire cohort), recurrent biliary events were observed in only 6.1% of patients, with no cases of biliary-type pain-related AEs.¹⁰ Biliary events were more frequently associated with pancreaticobiliary malignancies (PBM) and were all successfully treated endoscopically. It would be helpful to know if the association with PBM was also found in the study by Bang *et al.*¹ and why no endoscopic salvage was attempted. Nevertheless, we believe that unless patients are too frail to undergo a second procedure, LAMS removal should be preferred.

In conclusion, the study from Bang *et al.*¹ underlines that the jury is still out whether surgery is safe and feasible following EUS-GBD. However, we believe that EUS-GBD can substantially improve patient care in selected cases with a standardised yet tailored LAMS management. More high-quality data are eagerly awaited to further guide our strategy, but at this moment, it is too early to abandon EUS-GBD in high surgical patients.

Alberto Larghi ,¹ Roy L J van Wanrooij,² Michiel Bronswijk,³ Giuseppe Vanella ,⁴ Rastislav Kunda,¹ Manuel Pérez-Miranda,⁵ Jeanin E Van-Hooft ,⁶ Marc A Barthet ,⁷ Paolo Giorgio Arcidiacono,⁴ Schalk Willem Van der Merwe ⁸

¹Digestive Endoscopy Unit, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Roma, Italy

²Gastroenterology and Hepatology, Amsterdam UMC, Locatie VUmc, Amsterdam, Noord-Holland, The Netherlands

³Gastroenterology-Hepatology, Department of Gastroenterology and Hepatology, Imelda General Hospital, Belgium, Bonheiden, Belgium

⁴Pancreatobiliary Endoscopy and Endosonography Division, IRCCS San Raffaele Scientific Institute, Milan, Italy

⁵Gastroenterology and Hepatology, University Hospital Rio Hortega, Valladolid, Spain

⁶Gastroenterology & Hepatology, Leiden University Medical Centre, Leiden, The Netherlands

⁷Gastroenterology, Hôpital Nord, Marseille Cedex 20, France

⁸CHROMETA, KU Leuven, Leuven, Belgium

Correspondence to Dr Alberto Larghi, Digestive Endoscopy Unit, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Roma, Italy; alberto.larghi@yahoo.it

Twitter Giuseppe Vanella @GVanell5

Contributors Study concept and design and drafting of the manuscript: AL. Revision of the manuscript for important intellectual content: all authors. Approval of the final manuscript: all the authors. Guarantor of the article: AL.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Larghi A, van Wanrooij RLJ, Bronswijk M, *et al.* Gut Epub ahead of print: [please include Day Month Year]. doi:10.1136/gutjnl-2024-332273

Received 23 February 2024

Accepted 27 February 2024

Gut 2024;0:1–2. doi:10.1136/gutjnl-2024-332273

ORCID iDs

Alberto Larghi <http://orcid.org/0000-0001-7131-3686>
Giuseppe Vanella <http://orcid.org/0000-0001-7280-1761>

Jeanin E Van-Hooft <http://orcid.org/0000-0002-4424-0079>

Marc A Barthet <http://orcid.org/0000-0001-6589-7160>
Schalk Willem Van der Merwe <http://orcid.org/0000-0002-9891-2686>

REFERENCES

- Bang JY, Arnoletti JP, Wagner A, *et al.* EUS-guided gallbladder drainage in acute cholecystitis: long-term problems with surgical approach. *Gut* 2023;gutjnl-2023
- van der Merwe SW, van Wanrooij RLJ, Bronswijk M, *et al.* Therapeutic endoscopic ultrasound: European society of gastrointestinal Endoscopy (ESGE) guideline. *Endoscopy* 2022;54:185–205.
- Irani SS, Sharzei K, Siddiqui UD. AGA clinical practice update on role of EUS-guided gallbladder drainage in acute cholecystitis: commentary. *Clin Gastroenterol Hepatol* 2023;21:1141–7.
- Parmar KL, Law J, Carter B, *et al.* Frailty in older patients undergoing emergency Laparotomy: results from the UK observational emergency laparotomy and frailty (ELF) study. *Ann Surg* 2021;273:709–18.
- Mastalerz K, Kenig J, Olszewska U, *et al.* The surgical Appar score and frailty as outcome predictors in Short- and long-term evaluation of fit and frail older patients undergoing elective laparoscopic cholecystectomy - a prospective cohort study. *Wideochir Inne Tech Maloinwazyjne* 2018;13:350–7.
- Tyberg A, Saumoy M, Sequeiros EV, *et al.* EUS-guided versus percutaneous gallbladder drainage: isn't it time to convert? *J Clin Gastroenterol* 2018;52:79–84.

- 7 Tyberg A, Duarte-Chavez R, Shahid HM, *et al.* Endoscopic ultrasound-guided gallbladder drainage versus percutaneous drainage in patients with acute cholecystitis undergoing elective cholecystectomy. *Clin Transl Gastroenterol* 2023;14:e00593.
- 8 Kunda R, Messaoudi N, Chan S, *et al.* Feasibility and safety of interval cholecystectomy after endoscopic ultrasound-guided gallbladder drainage with lumen Apposing metal Stents – time to explore broader indications. *Gastrointestinal Endoscopy* 2023;97:AB864–5.
- 9 Teoh AYB, Kitano M, Itoi T, *et al.* Endosonography-guided gallbladder drainage versus percutaneous cholecystostomy in very high-risk surgical patients with acute cholecystitis: an international randomised multicentre controlled superiority trial (DRAC 1). *Gut* 2020;69:1085–91.
- 10 Bazaga S, García-Alonso FJ, Tormo A, *et al.* RNPAL (Registro Nacional de Prótesis de Aposición Luminal, national lumen-Apposing metal Stent Registry) study group. endoscopic ultrasound-guided gallbladder drainage with long-term lumen-Apposing metal Stent Indwell: 1-year results from a prospective nationwide observational study. *J Gastroenterol Hepatol* 2024;39:360–8.