



Universiteit
Leiden
The Netherlands

Vascular disease and apathy symptoms in the very old: a cross-sectional and longitudinal meta-analysis of individual participant data

Klei, V.M.G.T.H. van der; Poortvliet, R.K.E.; Bogaerts, J.M.K.; Blom, J.W.; Mooijaart, S.P.; Teh, R.; ... ; Gussekloo, J.

Citation

Klei, V. M. G. T. H. van der, Poortvliet, R. K. E., Bogaerts, J. M. K., Blom, J. W., Mooijaart, S. P., Teh, R., ... Gussekloo, J. (2022). Vascular disease and apathy symptoms in the very old: a cross-sectional and longitudinal meta-analysis of individual participant data. *International Journal Of Geriatric Psychiatry*, 37(12). doi:10.1002/gps.5831



Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3503755>

Note: To cite this publication please use the final published version (if applicable).

Vascular disease and apathy symptoms in the very old: A cross-sectional and longitudinal meta-analysis of individual participant data

Veerle M. G. T. H. van der Klei¹  | Rosalinde K. E. Poortvliet² |
Jonathan M. K. Bogaerts² | Jeanet W. Blom² | Simon P. Mooijaart¹ | Ruth Teh³ |
Marama Muru-Lanning^{3,4} | Leah Palapar³ | Andrew Kingston⁵ |
Louise Robinson⁵  | Ngaire Kerse³ | Jacobijn Gussekloo^{1,2}

¹Department of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, The Netherlands

²Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands

³School of Population Health, University of Auckland, Auckland, New Zealand

⁴James Henare Māori Research Centre, University of Auckland, Auckland, New Zealand

⁵Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University, Campus for Ageing and Vitality, Newcastle Upon Tyne, UK

Correspondence

Veerle M. G. T. H. van der Klei, Department of Gerontology and Geriatrics, Leiden University Medical Center, Postzone C7-Q, PO Box 9600, 2300RC, Leiden, The Netherlands. Email: v.m.g.t.h.van_der_klei@lumc.nl

Funding information

Oakley Mental Health Foundation New Zealand; Medical Research Council United Kingdom; New Zealand National Centre for Research Excellence for Māori research; National Heart Foundation of New Zealand; NIHR Biomedical Research Centre at Newcastle upon Tyne Hospitals NHS Foundation Trust; Newcastle Hospitals Healthcare Charity; Biotechnology and Biological Sciences Research Council; Dunhill Medical Trust; Health Research Council of New Zealand; Ministry of Health, New

Abstract

Objectives: Previous findings suggest a vascular foundation underlying apathy, but transdiagnostic and prospective evidence on vascular apathy is scarce. This study examines the association between vascular disease and the presence and development of apathy symptoms in the very old.

Methods: Four cohorts of the *Towards Understanding Longitudinal International older People Studies* (TULIPS)-consortium were included in a two-staged, individual participant data meta-analysis using generalized linear mixed models. Vascular disease was defined as a history of any clinical atherosclerotic pathology (angina pectoris, myocardial infarction, intermittent claudication, transient ischemic attack, stroke or related surgeries) and was related to apathy symptoms as repeatedly measured by the Geriatric Depression Scale (GDS-3A ≥ 2) over a maximum of 5 years.

Results: Of all 1868 participants (median age 85 years old), 53.9% had vascular disease and 44.3% experienced apathy symptoms. Participants with vascular disease had a 76% higher risk of apathy symptoms at baseline (odds ratio (OR) 1.76, 95% confidence interval (CI) 1.32–2.35), irrespective of depressive symptoms and only partially explained by stroke. Conversely, there was no association of vascular disease with the occurrence of apathy symptoms longitudinally, both in those with apathy at baseline (OR 1.00, 95% CI 0.84–1.20) and without (OR 0.96, 95% CI 0.84–1.09).

Conclusions: Vascular disease in the very old is associated with apathy symptoms cross-sectionally, but not proven longitudinally, independent of depressive symptoms. These findings query a vascular cause underlying apathy symptoms. However, the consistency of our cross-sectional findings in direction and magnitude across the

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. International Journal of Geriatric Psychiatry published by John Wiley & Sons Ltd.

Zealand; Ministerie van Volksgezondheid, Welzijn en Sport; Medical Research Council

TULIPS-consortium do emphasize international relevance of the interplay of vascular factors and apathy in advanced age, which meaning needs further unravelling.

KEYWORDS

atherosclerosis, meta-analysis, older people, prospective, vascular apathy

Key points

- Apathy symptoms are a common burden in the very old population of which the complex pathophysiology is yet to be unraveled.
- Vascular disease might predispose older adults to developing apathy, but prospective evidence on a vascular origin for apathy in old age remains scarce.
- Vascular disease is consistently associated with increased risk of concurrent apathy symptoms, but there is no evidence for influence on future apathy symptoms in the very old.
- Optimizing cardiovascular disease (CVD) management to attempt to prevent apathy may not be warranted, at least in the very old.

1 | INTRODUCTION

Reduced motivation, or apathy, is defined as a persistent lack of goal-directed behaviour which impairs daily functioning.^{1,2} Apathy symptoms do not merely occur with different medical conditions like dementia or depression,³⁻⁷ but are also common in the general population at old age. These symptoms affect community-dwelling older adults in approximately 20%–40% and occur in the absence of depressive symptoms (i.e. isolated apathy) in 13.8% on average.⁸ Due to this widespread occurrence and little evidence about apathy prevention or recovery,^{7,9,10} there is a strong need for a transdiagnostic approach to unravel the complex pathophysiology underlying these symptoms.⁹

Vascular pathologies have been proposed to specifically precede apathy symptoms and not depressive symptoms, and to predispose older adults to developing apathy in old age: the so-called vascular apathy hypothesis.¹¹ An increasing body of literature has since emerged on the interplay between apathy and several proxies of cardiovascular disease (CVD), as comprehensively depicted in a recent systematic review.¹² Various mechanisms could drive the vascular foundation of apathy, such as atherosclerosis or structural brain damage following stroke.^{4,9,11,13} Conversely, apathy symptoms also seem to contribute to incident CVD,¹⁴ which could for example, be mediated by adverse health behaviour if motivation for health promotion declines.¹⁵ To our knowledge, only the latter perspective has been examined over time in a thorough meta-analysis,⁸ whereas prospective evidence on the vascular apathy hypothesis remains scarce¹² as well as replication of previous cross-sectional findings in the very old population (i.e. those over 80 years of age),¹¹ hampering further understanding about the direction and interpretability of these associations in old age.

Therefore, we aim to investigate the association between vascular disease and the presence as well as development of apathy symptoms in the very old, irrespective of depressive symptoms.

2 | MATERIALS AND METHODS

2.1 | Study design and participants

Our two-stage meta-analysis incorporated individual participant data (IPD) from the pre-existing *Towards Understanding Longitudinal International older People Studies* (TULIPS)-consortium.¹⁶⁻¹⁹ This is a worldwide collection of prospective, population-based studies on ageing in the very old, which was at the time established to increase power and to examine international variation in associations in old age. We included four of its cohorts from three different studies: (1) the Leiden 85-plus Study from the Netherlands (2a) the Māori (indigenous population of New Zealand) and (2b) the non-Māori parallel cohorts of the Life and Living in Advanced Age: a Cohort Study in New Zealand (LiLACS NZ) and (3) the Newcastle 85+ Study originating from the United Kingdom. In contrary to previous IPD meta-analyses in the TULIPS-consortium, the Tokyo Oldest Old Survey on Total Health from Japan had to be entirely excluded from this analysis due to unavailability of measurements on the outcome.

The Leiden 85-plus Study is an observational study established from September 1997 to September 1999, which included 599 people out of the 705 eligible older adults that lived in the town of Leiden and turned 85 years old during the baseline period (i.e. a response rate of 87%).^{11,20} LiLACS NZ was initiated between March 2010 and April 2011 and successfully recruited 421 Māori of 80–90 years old along with 516 non-Māori of 85 years old residing in the Bay of Plenty and Rotorua area (56% and 59% respectively of all eligible older adults).^{21,22} The Newcastle 85+ Study was set up in 2006 to 2007 by including 1040 85-year-old adults out of the sampling frame of 1470 people, which were born in 1921 and were registered with a participating general practice in Newcastle or the North Tyneside (i.e. a 71% response rate).²³ All the above final samples were shown to be by and large representative of the several ageing source populations, as outlined in detail,²⁰⁻²³ and within these

samples individual participants were solely excluded from this study due to data unavailability on the exposure and/or outcome.

Repeated data collection was annual over a maximum of 5 years in the Leiden 85-plus Study as well as LiLACS NZ, while follow-up was clustered at 1.5, 3 and 5 years from baseline in the Newcastle 85+ Study. The individual study protocols were approved by Medical Ethical Committees at their respective sites and all very old or their proxies had granted either oral or written informed consent.

2.2 | Vascular disease

Several manifestations of vascular disease have been assessed at baseline by combining different types of data collection per study (Supplementary Table 1). In the Leiden 85-plus Study certified and trained research staff conducted standardized interviews with physicians.¹¹ In LiLACS NZ such interviews were performed with participants and were seconded by primary health care and hospitalization record checks.²⁴ In the Newcastle 85+ Study primary health care records were thoroughly examined.²³ Electrocardiograms were recorded in all studies to objectify and to complement findings on previous myocardial infarction.^{11,23,24} Vascular disease (yes/no) was defined as a history of any clinical atherosclerotic pathology: angina pectoris, myocardial infarction, intermittent claudication, transient ischemic attack, stroke or any related coronary or peripheral artery surgery.

2.3 | Apathy and depressive symptoms

All studies repeatedly administered the 15-item Geriatric Depression Scale (GDS) to their participants,²⁵ which is appropriate to assess symptoms over time.²⁶ In line with other population-based studies and prior factor analyses,^{11,13,27} a 3-item apathy subdimension has been used to screen for apathy symptoms, which included the following questions: (1) Have you dropped many of your activities and interests? (2) Do you prefer to stay at home rather than to go out and do new things? (3) Do you feel full of energy? (GDS-3A; range 0–3 points, with greater scores implying more apathy). Since scores ≥ 2 points have shown to be indicative of a clinical diagnosis of apathy, with a sensitivity of 69% and a specificity of 85%,^{11,28} this cut-off has been applied to dichotomize the presence of apathy symptoms. Likewise, the 12 remaining items of the GDS were summed up (i.e. those not reflecting apathy, the GDS-12D; range 0–12 points, with greater scores indicating more depression) and a ≥ 2 point cut-off was applied to reflect the presence of depressive symptoms.

As apathy is increasingly regarded as an independent syndrome across disease boundaries²⁹ and vascular apathy is preferably studied independently of depression,¹² we classified apathy symptoms without these depressive symptoms as isolated apathy (concurrent GDS-3A ≥ 2 and GDS-12D < 2).⁸

Because the 15-item GDS is self-reported and its validity is decreased in older adults with dementia,²⁹ the GDS was either not

administered to participants with a Mini-Mental State Examination (MMSE) score < 19 points^{11,30} or we excluded their GDS-scores to align data availability between the cohorts (Supplementary Table 1). Exclusion of these GDS-scores was carried out by assigning all GDS-measurements in which the concurrent MMSE-score was below 19 points to missing.

2.4 | Other measures

Data on demographic variables such as age, gender, educational attainment (up to and including secondary school vs. higher education) and living situation (institutionalized or not) was gathered during the baseline interviews. LiLACS NZ participants also reported their ethnicity, whether they identified themselves as Māori, which was prioritized over other ethnicities if more than one was listed.²¹ According to overall data availability and previously applied conditions¹¹ participants were considered to have a chronic disease when any of the following diseases was present: arthritis, cancer, diabetes mellitus, chronic lung disease or Parkinson's disease. This was evaluated by combining reported histories, related medication use and blood glucose levels, which are specified in the supplement (Supplementary Table 1).

2.5 | Statistical analysis

A two-stage IPD approach was adopted to acknowledge the heterogeneity between the cohorts and to honour the preferences and practices in NZ Māori research,³¹ which means the analyses were initially performed per cohort and their distinct results were afterwards pooled for meta-analysis.

2.5.1 | Cohort-level analyses

Generalized linear mixed models were applied as these mixed models optimize the use of available data during follow-up and account for dependency of repeated measures per person.³² Two models were assessed in accordance with existing literature^{11,13}: (1) adjusted for age (only in Māori), sex and depressive symptoms (GDS-12D ≥ 2 , as a time-varying covariate) and employing a random intercept, and (2) additionally adjusted for educational attainment and chronic disease. Besides the aforementioned exclusion of total GDS-scores based on MMSE-scores, no adjustment was made for cognitive status as it was thought to possibly be part of the causal pathway of interest. The selection of the final model is described in more detail in Supplementary Table 2. The cross-sectional (main effect of vascular disease) as well as the longitudinal estimates (interaction term: vascular disease * time since baseline) were generated by use of the Lme4 package in R (version 4.0.2).^{33,34} These estimates were respectively presented as baseline and annual odds ratio (OR) of apathy symptoms (GDS-3A ≥ 2) with a 95% confidence interval (CI) (95% CI). The

latter represents the additional OR with every year increase in time since baseline when having vascular disease and was stratified by baseline apathy status. All analyses were repeated while excluding participants with a history of stroke as a proxy for subsequent structural brain damage.^{11,13}

2.5.2 | Pooled analyses

The individual ORs were thereafter pooled by random-effects models with inverse-variance weighting and summarized in forest plots by using Review Manager 5.4.³⁵ Consistency between the estimates was defined as a statistical heterogeneity (I^2) below 40%.³⁶

2.5.3 | Sensitivity analyses

To examine whether our findings were dependent on the cut-offs that were applied to define apathy and depressive symptoms, we conducted sensitivity analyses by use of the ordinal package.³⁷ We entered the continuous instead of the dichotomized GDS-3A and GDS-12D scores in ordinal logistic regression models, which were apart from that equal to the main analyses.

3 | RESULTS

3.1 | Demographic and clinical characteristics

Out of the 2576 participants of the TULIPS-consortium we excluded 706 older adults due to data unavailability (27.4% in total; exclusion on MMSE accounted for 1.5%). This meta-analysis therefore consisted of 1868 participants across the four cohorts and their baseline characteristics are shown in Table 1. The median age was 82 years old in the NZ Māori population and 85 years in the other cohorts. The majority of participants was female (60.1%), community-dwelling (i.e. 5.6% was institutionalized) and maximally attained primary or secondary school (83.8%). Most of them were diagnosed with one or more chronic disease (77.7%) and their median MMSE-score was 28 out of 30 (interquartile range (IQR) 26–29). Vascular disease was present in 53.9% of participants, explained by at least a stroke in 11.6% of all participants. Apathy symptoms were experienced by 44.3% of older adults, depressive symptoms by 34.3% and isolated apathy occurred in 22.8% of participants.

Figure 1 shows the percentage of participants experiencing apathy symptoms during follow-up, stratified by baseline apathy status. The median number of follow-up waves on the GDS varied from 2 years in the Māori population (IQR 0.0–4.0) to 3 years in the combined cohort (IQR 1.0–5.0). Overall, apathy symptoms occurred at least once again in 65.1% of those experiencing apathy symptoms at baseline, while 42.8% of those without apathy symptoms at baseline experienced apathy symptoms during follow-up. The

majority of both regarded isolated apathy symptoms, respectively 37.3% and 29.9%.

3.2 | Cross-sectional and longitudinal associations on vascular apathy

Adjusted for depressive symptoms, age (only in Māori), sex, education and chronic disease, vascular disease increased the risk of experiencing apathy symptoms at baseline by 76% (pooled OR 1.76, 95% CI 1.32–2.35), as depicted in Figure 2. The association was minimally attenuated and remained significant after exclusion of stroke patients (OR 1.60, 95% CI 1.23–2.09).

To the contrary, there was no significant association between vascular disease at baseline and the course of apathy symptoms over time, neither in those with apathy symptoms at baseline (OR 1.00, 95% CI 0.84–1.20) nor in those without apathy symptoms at baseline (OR 0.96, 95% CI 0.84–1.09), irrespective of a history of stroke (Figure 3). All minimally adjusted models showed similar results (not shown) and there was overall consistency between the individual estimates ($I^2 < 40\%$).

3.3 | Sensitivity analyses

Sensitivity analyses applying the continuous GDS-3A and GDS-12D scores showed comparable results for both the cross-sectional and the longitudinal associations on vascular apathy (not shown).

4 | DISCUSSION

In this IPD meta-analysis of four population-based cohorts of the very old, we found that vascular disease is cross-sectionally associated with the presence of apathy symptoms, independent of concurrent depressive symptoms and only partially explained by a history of stroke. In contrast, vascular disease was not associated with the occurrence of apathy symptoms over time, which was also irrespective of whether apathy symptoms were experienced at baseline.

In relationship to previous transdiagnostic literature on the vascular apathy hypothesis, our significant cross-sectional associations are not only consistent in relating vascular disease to apathy in older populations, independent of depression and/or structural brain damage following stroke, but rather add to this body of literature by extending this finding to those of advanced age (i.e. in the very old vs. a mean age of 57 and 74 years old).^{13,38} Our findings were also consistent in direction and magnitude throughout the four cohorts examined suggesting international relevance. This is surprising as the levels of apathetic and depressive symptomatology differed between the cohorts and performance of self-rated health tools can be culturally or socioeconomically mediated.^{39,40}

TABLE 1 Baseline characteristics of study participants and median follow-up time

Cohort	Leiden 85-plus study	LiLACS NZ		Newcastle 85+ study	Overall
		Māori	Non-Māori		
N =	493	243	376	756	1868
Study population (starting year)	Cohort of 85-year old inhabitants of Leiden, the Netherlands (1997)	Cohort of 80–90 year old Māori, indigenous to New Zealand (2010)	Cohort of 85-year old non-Māori inhabitants of New Zealand (2010)	Cohort of 85-year old inhabitants of Newcastle upon Tyne, the UK (2006)	
Demographics					
Age, median [IQR], years	85 [85, 85]	82 [81, 84]	85 [85, 85]	85 [85, 86]	85 [85, 85]
Female, n (%)	312 (63.3)	148 (60.9)	199 (52.9)	463 (61.2)	1122 (60.1)
≤ Secondary school, n (%)	421 (85.9)	204 (84.3)	268 (72.4)	663 (87.8)	1556 (83.8)
Institutionalized, n (%) ^a	56 (12.1)	3 (1.3)	13 (3.5)	31 (4.1)	103 (5.6)
Clinical characteristics					
Chronic disease, n (%) ^b	296 (61.7)	187 (77.6)	322 (86.1)	614 (82.3)	1419 (77.1)
MMSE-score, median [IQR]	27 [24, 28]	28 [26, 29]	28 [27, 29]	28 [26, 29]	28 [26, 29]
Depressive symptoms, n (%) ^c	144 (28.8)	73 (30.0)	86 (22.9)	337 (44.6)	640 (34.3)
Apathy, n (%)					
Apathy symptoms ^d	136 (27.6)	94 (38.7)	151 (40.2)	447 (59.1)	828 (44.3)
Isolated apathy ^e	68 (13.8)	56 (23.0)	99 (26.3)	203 (26.9)	426 (22.8)
Vascular disease, n (%)					
Vascular disease ^f	221 (44.8)	149 (61.3)	226 (60.1)	411 (54.4)	1007 (53.9)
Stroke	37 (7.5)	26 (10.7)	61 (16.2)	91 (12.0)	215 (11.6)
Follow-up, median [IQR], years	3.0 [1.0, 5.0]	2.0 [0.0, 4.0]	3.0 [1.0, 5.0]	3.0 [1.5, 5.0]	3.0 [1.0, 5.0]

Note: Missing data per cohort (Leiden 85-plus Study = L85+, LiLACS NZ–Māori = NZ-M, LiLACS NZ–non-Māori = NZ-NM, Newcastle 85+ Study = N85+): age—N85+ 1; education—L85+ 3, NZM 1, NZ-NM 6, N85+ 1; institutionalized—L85+ 30, NZM 4, NZ-NM 1; chronic diseases—L85+ 13, NZM 2, NZ-NM 2, N85+ 10; MMSE—L85+ 23, NZM 2, NZ-NM 12; stroke—NZM 5, NZ-NM 2.

Abbreviations: IQR, interquartile range; LiLACS NZ, Life and Living in Advanced Age: a Cohort Study in New Zealand; MMSE, Mini-Mental State Exam; UK, United Kingdom.

^aParticipants were regarded as institutionalized if living in care- or nursing homes in the Leiden 85-plus Study, in rest homes or private hospitals in LiLACS NZ and in residential care homes, nursing homes or long stay hospitals in the Newcastle 85+ Study.

^bChronic disease was defined as the presence of arthritis, cancer, chronic lung disease, diabetes mellitus or Parkinson's disease.

^cDepressive symptoms were defined as a score of ≥ 2 on the GDS-12D, the depression subdimension of the GDS.

^dApathy symptoms were defined as a score of ≥ 2 on the GDS-3A, the apathy subdimension of the GDS.

^eIsolated apathy was defined as having apathy symptoms in the absence of above depressive symptoms (i.e. GDS-3 A ≥ 2 and GDS-12D < 2).

^fVascular disease was defined as a history of any clinical atherosclerotic pathology (including angina pectoris, myocardial infarction, intermittent claudication, transient ischemic attack, stroke and any related coronary or peripheral artery surgery).

Conversely, our longitudinal findings in the Leiden 85-plus Study did not correspond to prior prospective findings by van der Mast et al. on vascular apathy in the same cohort,¹¹ presumably due to methodological insights (e.g. different modelling: performing

generalized and ordinal regressions on the GDS-3A instead of linear regressions). In addition, our longitudinal meta-analysis shows heterogeneity between the cohorts in being more or less likely to display apathy symptoms over time when having vascular disease (not




FIGURE 1 Percentage of participants with apathy symptoms over time, stratified by baseline apathy status. ^aBaseline apathy status was defined as the presence or absence of apathy symptoms at baseline (i.e. a score of ≥ 2 or < 2 on the GDS-3A, the apathy subdimension of the GDS). GDS, Geriatric Depression Scale; LiLACS NZ, Life and Living in Advanced Age: a Cohort Study in New Zealand.

significant). This is not unexpected as NZ Māori have been exposed to a lifetime of discrimination resulting in health disparities,⁴¹ but suggests nuance in its interpretation. As there are no other prospective, population-based studies to substantiate the vascular apathy hypothesis,¹² unequivocal understanding of a potential vascular ground for apathy symptoms remains limited. Optimizing CVD-management to attempt to prevent apathy in older adults may not be warranted, at least in advanced age.

There could be several pathophysiological mechanisms underlying our findings. One explanation is reversed causality, which has been evidently substantiated in the previously mentioned meta-analysis.⁸ Apathy symptoms increased the risk of incident myocardial infarction and stroke by 21% and 37% respectively over a median follow-up of 8.8 years in approximately 48,000 older adults with a median age of 74 years old (IQR 72–78). To elucidate whether our cross-sectional results reflect this reversed causality or whether our lack of longitudinal evidence is, for instance, explained by characteristics of the TULIPS-consortium (such as competing risks at advanced age), future research could assess this reversed association for the very old and study vascular apathy in relatively younger, more stable populations of older adults over time (e.g. over 65 years old). In addition, apathy might be a marker of subclinical (cerebral) small vessel disease which could then result in overt CVD.^{8,12} Support for

this mechanism is strengthened by associations between more white matter hyperintensities on brain imaging and higher risk of having apathy,^{12,42} which should also be studied prospectively in the general older population to elucidate the direction of their interplay.¹² Although we did not find support for vascular disease to predispose older adults to apathy symptoms, if the origin were to be vascular, mediation by cognitive decline could explain our lacking prospective associations as older people with severe cognitive impairment could not partake by design. To that extent, upcoming studies on apathy could be complemented by informant-rated scales or clinical interviews. Lastly, apathy could also be a marker or consequence of (multi)morbidity instead of specifically having a vascular origin; especially the apathy symptoms we studied, as they have been shown to closely connect to functional disability.⁴³ This calls for future apathy research to employ the diagnostic criteria² or instruments specifically designed to assess apathy in otherwise healthy, community-dwelling samples (e.g. the Apathy-Motivation Index)⁴⁴ to further unravel apathy as an independent construct.

This meta-analysis has various strengths. It addresses the need for both transdiagnostic and prospective analyses on (vascular) apathy during substantial follow-up time. The utilized data includes a globally unique collection of population-based cohort studies of very old adults, originating from diverse countries and cultures, which was




FIGURE 2 Cross-sectional association between vascular disease and apathy symptoms. Results from two-staged, generalized linear mixed model (GLMM) analyses, presented as baseline odds ratios of apathy symptoms (GDS-3A ≥ 2) by vascular disease status (including and excluding participants with a history of stroke). Fully adjusted models are shown (i.e. adjusted for depressive symptoms, age (only in Māori), sex, education and chronic diseases). Individual study estimates were pooled by random-effects models with inverse-variance weighting. CI, confidence interval; GDS, Geriatric Depression Scale; LiLACS NZ, Life and Living in Advanced Age: a Cohort Study in New Zealand; OR, odds ratio.




FIGURE 3 Longitudinal association between vascular disease at baseline and apathy symptoms during follow-up, stratified by baseline apathy status. Results from two-staged, generalized linear mixed model (GLMM) analyses, presented as annual odds ratios of apathy symptoms (GDS-3A ≥ 2) by vascular disease status (i.e. analyses including and excluding participants with a history of stroke). Fully adjusted models are shown (i.e. adjusted for depressive symptoms, age (only in Māori), sex, education and chronic diseases), stratified by baseline apathy status. Individual study estimates were pooled by random-effects models with inverse-variance weighting. ^aBaseline apathy status was defined as the presence or absence of apathy symptoms at baseline (i.e. a score of ≥ 2 or < 2 on the GDS-3A, the apathy subdimension of the GDS). CI, confidence interval; GDS, Geriatric Depression Scale; LiLACS NZ, Life and Living in Advanced Age: a Cohort Study in New Zealand; OR, odds ratio.

honoured by our two-stage IPD approach. Participants were solely excluded due to data unavailability and by using this IPD-method we verified previously presented results,¹¹ as well as provided consistency in the data and analyses across cohorts.

Although there was not much statistical heterogeneity between the pooled estimates, clinical heterogeneity between the cohorts is a potential limitation, which existed throughout time (i.e. initiation in different time periods) and at baseline (e.g. ethnicity and living situation). Furthermore, a disadvantage inherent to longitudinal research in the very old is the loss-to-follow-up due to death or ill health.²¹ Subgroup analyses were hampered (e.g. minor vs. major vascular disease) and potential other confounders could not be included (e.g. social factors). The associations studied in the TULIPS-consortium could be even more unraveled by performing a systematic review to find other comparable cohorts. Another limitation is that our conclusions are confined to apathy symptoms of the three included questions in the GDS-3A and deficits in self-awareness, commonly associated with apathy,⁷ could have impeded the reliability of this self-reported assessment. Despite these characteristics the GDS-3A has however been deemed appropriate for etiological research purposes, particularly in older populations over time.^{11,45}

In conclusion, vascular disease in the very old population is associated with apathy symptoms cross-sectionally, but there is no evidence of influence on apathy occurrence over time, independent of concomitant depressive symptoms. While these findings query the vascular apathy hypothesis and may discourage prevention of apathy by CVD-management, they emphasize the international relevance of their interplay in advanced age.

ACKNOWLEDGEMENTS

The LiLACS NZ study wishes to acknowledge the study participants, their families and whanau for supporting the study. We are extremely grateful Te RōpuKaitiaki o ngā tikanga Māori for their guidance and we acknowledge the community partners (Western Bay of Plenty Primary Health Organisation, Ngā Matāpuna Oranga Kaupapa Māori Primary Health Organisation, Te Korowai Aroha Trust, Te Rūnanga o Ngati Pikiāo, Rotorua Area Primary Health Services, Ngati Awa Research & Archives Trust, Te Rūnanga o Ngati Irapuaia and Te Whanau a Apanui Community Health Centre).

The Leiden 85-plus Study was partly funded by an unrestricted grant from the Dutch Ministry of Health, Welfare and Sports (1997–2001). The Life and LiLACS NZ - Te Puāwaitanga o Nga Tapuwae Kia Ora Tonu (LiLACS NZ), was funded by the Health Research Council of New Zealand programme grant [HRC 09/068B]; and Ministry of Health New Zealand [MOH ref: 345426/00]; and Ngā Pae o te Māramatanga (the New Zealand National Centre for Research Excellence for Māori research; funded Māori engagement and project management) project grant; National Heart Foundation project grant for investigating cardiac markers; and Oakley Mental Health Foundation project grant for investigating dementia. The Newcastle 85+ study was funded by the Medical Research Council [grant no. G0500997], the Biotechnology and Biological Sciences Research Council, the Dunhill Medical Trust, the

NIHR Biomedical Research Centre at Newcastle upon Tyne Hospitals NHS Foundation Trust and Newcastle Hospitals Health-care Charity.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Veerle M. G. T. H. van der Klei  <https://orcid.org/0000-0002-2782-1517>

Louise Robinson  <https://orcid.org/0000-0003-0209-2503>

REFERENCES

1. Robert P, Onyike CU, Leentjens AF, et al. Proposed diagnostic criteria for apathy in Alzheimer's disease and other neuropsychiatric disorders. *Eur Psychiatr*. 2009;24(2):98-104. <https://doi.org/10.1016/j.eurpsy.2008.09.001>
2. Robert P, Lanctôt KL, Agüera-Ortiz L, et al. Is it time to revise the diagnostic criteria for apathy in brain disorders? The 2018 international consensus group. *Eur Psychiatr*. 2018;54:71-76. <https://doi.org/10.1016/j.eurpsy.2018.07.008>
3. Groeneweg-Koolhoven I, Ploeg M, Comijs HC, et al. Apathy in early and late-life depression. *J Affect Disord*. 2017;223:76-81. <https://doi.org/10.1016/j.jad.2017.07.022>
4. Caeiro L, Ferro JM, Costa J. Apathy secondary to stroke: a systematic review and meta-analysis. *Cerebrovasc Dis*. 2013;35(1):23-39. <https://doi.org/10.1159/000346076>
5. Lyketsos CG, Lopez O, Jones B, Fitzpatrick AL, Breitner J, DeKosky S. Prevalence of neuropsychiatric symptoms in dementia and mild cognitive impairment: results from the cardiovascular health study. *JAMA*. 2002;288(12):1475-1483. <https://doi.org/10.1001/jama.288.12.1475>
6. Onyike CU, Sheppard JM, Tschanz JT, et al. Epidemiology of apathy in older adults: the cache county study. *Am J Geriatr Psychiatr*. 2007;15(5):365-375. <https://doi.org/10.1097/01.JGP.0000235689.42910.0d>
7. van Reekum R, Stuss DT, Ostrander L. Apathy: why care? *Focus (Am Psychiatr Publ)*. 2016;14(4):523-532. <https://doi.org/10.1176/appi.focus.140402>
8. Eurelings LS, van Dalen JW, Ter Riet G, Moll van Charante E, Richard E, van Gool W. Apathy and depressive symptoms in older people and incident myocardial infarction, stroke, and mortality: a systematic review and meta-analysis of individual participant data. *Clin Epidemiol*. 2018;10:363-379. <https://doi.org/10.2147/clep.S150915>
9. Husain M, Roiser JP. Neuroscience of apathy and anhedonia: a transdiagnostic approach. *Nat Rev Neurosci*. 2018;19(8):470-484. <https://doi.org/10.1038/s41583-018-0029-9>
10. Fahed M, Steffens DC. Apathy: neurobiology, assessment and treatment. *Clin Psychopharmacol Neurosci*. 2021;19(2):181-189. <https://doi.org/10.9758/cpn.2021.19.2.181>
11. van der Mast RC, Vinkers DJ, Stek ML, et al. Vascular disease and apathy in old age. The Leiden 85-Plus Study. *Int J Geriatr Psychiatr*. 2008;23(3):266-271. <https://doi.org/10.1002/gps.1872>
12. Wouts L, van Kessel M, Beekman ATF, Marijnissen RM, Oude Voshaar RC. Empirical support for the vascular apathy hypothesis: a structured review. *Int J Geriatr Psychiatr*. 2020;35(1):3-11. <https://doi.org/10.1002/gps.5217>
13. Ligthart SA, Richard E, Fransen NL, et al. Association of vascular factors with apathy in community-dwelling elderly individuals. *Arch Gen Psychiatr*. 2012;69(6):636-642. <https://doi.org/10.1001/archgenpsychiatry.2011.1858>

14. Eurelings LS, Ligthart SA, van Dalen JW, Moll van Charante EP, van Gool WA, Richard E. Apathy is an independent risk factor for incident cardiovascular disease in the older individual: a population-based cohort study. *Int J Geriatr Psychiatr.* 2014;29(5):454-463. <https://doi.org/10.1002/gps.4026>
15. Eurelings LS, Jaccard J, Moll van Charante EP, et al. The mediating role of cardiovascular risk factors in the relationship between symptoms of apathy and incident cardiovascular disease in community-dwelling older individuals. *Int Psychogeriatr.* 2016;28(4):669-679. <https://doi.org/10.1017/s1041610215001751>
16. Du Puy RS, Poortvliet RKE, Mooijaart SP, et al. Outcomes of thyroid dysfunction in people aged eighty years and older: an individual patient data meta-analysis of four prospective studies (towards understanding longitudinal international older people studies consortium). *Thyroid.* 2020;31(4):552-562. <https://doi.org/10.1089/thy.2020.0567>
17. Palapar L, Kerse N, Rolleston A, et al. Anaemia and physical and mental health in the very old: an individual participant data meta-analysis of four longitudinal studies of ageing. *Age Ageing.* 2021;50(1):113-119. <https://doi.org/10.1093/ageing/afaa178>
18. Bogaerts JMK, Poortvliet RKE, van der Klei V, et al. Disentangling the varying associations between systolic blood pressure and health outcomes in the very old: an individual patient data meta-analysis. *J Hypertens.* 2022;40(9):1786-1794. <https://doi.org/10.1097/hjh.0000000000003219>
19. Ravensbergen WM, Blom JW, Kingston A, et al. Declining daily functioning as a prelude to a hip fracture in older persons-an individual patient data meta-analysis. *Age Ageing.* 2022;51(1). <https://doi.org/10.1093/ageing/afab253>
20. der Wiel AB, van Exel E, de Craen AJ, et al. A high response is not essential to prevent selection bias: results from the Leiden 85-plus study. *J Clin Epidemiol.* 2002;55(11):1119-1125. [https://doi.org/10.1016/s0895-4356\(02\)00505-x](https://doi.org/10.1016/s0895-4356(02)00505-x)
21. Kerse N, Teh R, Moyes SA, et al. Cohort profile: Te Puawaitanga o Nga Tapuwae Kia Ora Tonu, life and living in advanced age: a cohort study in New Zealand (LiLACS NZ). *Int J Epidemiol.* 2015;44(6):1823-1832. <https://doi.org/10.1093/ije/dyv103>
22. Dyall L, Kepa M, Hayman K, et al. Engagement and recruitment of Māori and non-Māori people of advanced age to LiLACS NZ. *Aust N Z J Publ Health.* 2013;37(2):124-131. <https://doi.org/10.1111/1753-6405.12029>
23. Collerton J, Davies K, Jagger C, et al. Health and disease in 85 year olds: baseline findings from the Newcastle 85+ cohort study. *BMJ.* 2009;339:b4904. <https://doi.org/10.1136/bmj.b4904>
24. Hayman KJ, Kerse N, Dyall L, et al. Life and living in advanced age: a cohort study in New Zealand--e Puawaitanga o Nga Tapuwae Kia Ora Tonu, LiLACS NZ: study protocol. *BMC Geriatr.* 2012;12(1):33. <https://doi.org/10.1186/1471-2318-12-33>
25. Yesavage JA, Sheikh JI. Geriatric Depression Scale (GDS): recent evidence and development of a shorter version. *Clin Gerontol.* 1986;5(1-2):165-173. https://doi.org/10.1300/J018v05n01_09
26. Vinkers DJ, Gussekloo J, Stek ML, Westendorp RGJ, van der Mast RC. The 15-item Geriatric Depression Scale (GDS-15) detects changes in depressive symptoms after a major negative life event. The Leiden 85-plus Study. *Int J Geriatr Psychiatr.* 2004;19(1):80-84. <https://doi.org/10.1002/gps.1043>
27. Grool AM, Geerlings MI, Sigurdsson S, et al. Structural MRI correlates of apathy symptoms in older persons without dementia: AGES-Reykjavik Study. *Neurology.* 2014;82(18):1628-1635. <https://doi.org/10.1212/wnl.00000000000000378>
28. Starkstein SE, Mayberg HS, Preziosi TJ, et al. Reliability, validity, and clinical correlates of apathy in Parkinson's disease. *J Neuropsychiatry Clin Neurosci.* 1992;4(2):134-139. <https://doi.org/10.1176/jnp.4.2.134>
29. Kørner A, Lauritzen L, Abelskov K, et al. The geriatric depression scale and the cornell scale for depression in dementia. A validity study. *Nord J Psychiatr.* 2006;60(5):360-364. <https://doi.org/10.1080/08039480600937066>
30. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12(3):189-198. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
31. Dyall L, Kerse N, Hayman K, et al. Pinnacle of life--Māori living to advanced age. *N Z Med J.* 2011;124(1331):75-86.
32. Gueorguieva R, Krystal JH. Move over ANOVA: progress in analyzing repeated-measures data and its reflection in papers published in the Archives of General Psychiatry. *Arch Gen Psychiatr.* 2004;61(3):310-317. <https://doi.org/10.1001/archpsyc.61.3.310>
33. Team RC.R: A Language and Environment for Statistical Computing. Accessed July 22, 2021. <https://www.R-project.org/>
34. Bates DMM, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Software.* 2015;67(1):1-48. <https://doi.org/10.18637/jss.v067.i01>
35. Review Manager (RevMan) [Computer Program]. Version 5.4. The Cochrane Collaboration; 2020.
36. Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions.* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration; 2011. Accessed August 28, 2020. www.handbook.cochrane.org
37. RHB C. Ordinal—Regression Models for Ordinal Data. R Package Version 2019.12-10. Accessed July 22, 2021. <https://CRAN.R-project.org/package=ordinal>
38. Sugawara N, Yasui-Furukori N, Umeda T, et al. Ankle brachial pressure index as a marker of apathy in a community-dwelling population. *Int J Geriatr Psychiatr.* 2011;26(4):409-414. <https://doi.org/10.1002/gps.2541>
39. Scott KM, Sarfati D, Tobias MI, Haslett SJ. A challenge to the cross-cultural validity of the SF-36 health survey: factor structure in Maori, Pacific and New Zealand European ethnic groups. *Soc Sci Med.* 2000;51(11):1655-1664. [https://doi.org/10.1016/s0277-9536\(00\)00083-6](https://doi.org/10.1016/s0277-9536(00)00083-6)
40. Lima-Costa MF, Steptoe A, Cesar CC, De Oliveira C, Proietti FA, Marmot M. The influence of socioeconomic status on the predictive power of self-rated health for 6-year mortality in English and Brazilian older adults: the ELSA and Bambui cohort studies. *Ann Epidemiol.* 2012;22(9):644-648. <https://doi.org/10.1016/j.annepidem.2012.06.101>
41. Harris R, Cormack D, Tobias M, et al. Self-reported experience of racial discrimination and health care use in New Zealand: results from the 2006/07 New Zealand Health Survey. *Am J Publ Health.* 2012;102(5):1012-1019. <https://doi.org/10.2105/ajph.2011.300626>
42. Clancy U, Gilmartin D, Jochems ACC, Knox L, Doubal FN, Wardlaw JM. Neuropsychiatric symptoms associated with cerebral small vessel disease: a systematic review and meta-analysis. *Lancet Psychiatr.* 2021;8(3):225-236. [https://doi.org/10.1016/s2215-0366\(20\)30431-4](https://doi.org/10.1016/s2215-0366(20)30431-4)
43. van Wanrooij LL, Borsboom D, Moll van Charante EP, Richard E, van Gool WA. A network approach on the relation between apathy and depression symptoms with dementia and functional disability. *Int Psychogeriatr.* 2019;31(11):1655-1663. <https://doi.org/10.1017/s1041610218002387>
44. Ang YS, Lockwood P, Apps MA, Muhammed K, Husain M. Distinct subtypes of apathy revealed by the apathy motivation index. *PLoS One.* 2017;12(1):e0169938. <https://doi.org/10.1371/journal.pone.0169938>
45. Bertens AS, Moonen JE, de Waal MW, et al. Validity of the three apathy items of the Geriatric Depression Scale (GDS-3A) in

measuring apathy in older persons. *Int J Geriatr Psychiatry*. 2017;32(4):421-428. <https://doi.org/10.1002/gps.4484>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: van der Klei VMGTH, Poortvliet RKE, Bogaerts JMK, et al. Vascular disease and apathy symptoms in the very old: a cross-sectional and longitudinal meta-analysis of individual participant data. *Int J Geriatr Psychiatry*. 2022;1-10. <https://doi.org/10.1002/gps.5831>