



Universiteit  
Leiden  
The Netherlands

## Unveiling the third dimension: vertical structure as a probe of planet formation conditions

Paneque Carreño, T.

### Citation

Paneque Carreño, T. (2024, September 13). *Unveiling the third dimension: vertical structure as a probe of planet formation conditions*. Retrieved from <https://hdl.handle.net/1887/4083418>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4083418>

**Note:** To cite this publication please use the final published version (if applicable).

## Chapter 5

---

# Vertical CO surfaces as a probe for protoplanetary disk mass and carbon depletion

---

T. Paneque-Carreño, A. Miotello, E. F. van Dishoeck,  
G. Rosotti, B. Tabone

submitted to *Astronomy & Astrophysics*

## Abstract

*Context:* As the sample of mid-inclination disks with measured CO emission surfaces grows, a fundamental unanswered question is how these vertical profiles connect to their host properties.

*Aims:* This project aims to relate the vertical extent of protoplanetary disks as traced by  $^{12}\text{CO } 2 - 1$  to key stellar and physical parameters. In order to produce a result that is applicable towards an observational analysis, we benchmark our results with ALMA observations of CO emission from nineteen disks.

*Methods:* We produce a grid of disk models using the physical-chemical code DALI, for a template T Tauri and Herbig star. Our models use an iterative solver to calculate the hydrostatic equilibrium equations and determine a physically-motivated density structure. Key stellar and disk parameters such as stellar luminosity and temperature, total disk mass, carbon abundance and critical radius are varied to determine their effect on the CO emitting surface. Each vertical profile is fitted by an exponentially tapered power-law and characterized by the  $z/r$  value that represents the structure inwards of 80% of the tapering radius.

*Results:* The CO emission surface location is determined by the disk mass ( $M_d$ ) and the level of carbon depletion. T Tauri and Herbig systems show different vertical profiles, with disks around T Tauri stars being more vertically extended. We derive a  $z/r$ - $M_d$  relationship, which for each stellar type has a degeneracy with the volatile carbon abundance. In order to reconcile total disk mass estimates from the characteristic  $z/r$  and the values obtained based on dust continuum analysis, a volatile carbon depletion of 10-100 (with respect to the ISM) is needed for the majority of our sources. Our carbon depletion values are in agreement with previous literature estimates, highlighting the potential of this method to rapidly calculate key disk parameters.

## 5.1 Introduction

Planets form and grow within protoplanetary disks made of dust and gas which orbit around young stellar objects. Through high spatial and spectral resolutions of the Atacama Large Millimeter/submillimeter Array (ALMA) we have been able to trace the extent and structure of these disks in dust and gas across all three dimensions; azimuthal, radial and vertical (e.g. Huang et al. 2018b; Andrews 2020; Law et al. 2021a,b; Paneque-Carreño et al. 2023). To date, most studies have focused on the information offered by the radial and azimuthal structure (for a review see Bae et al. 2023). Radial extent compared to the stellar age is used as an indicator on the governing evolution processes (Rosotti et al. 2019; Tazzari et al. 2021a; Long et al. 2022; Trapman et al. 2023) and radial substructures characterize a number of chemical and physical scenarios (Andrews 2020; Öberg et al. 2021b, 2023). Azimuthal enhancements can signal towards the presence of planetary companions (Ragusa et al. 2017; Pérez et al. 2020; Booth et al. 2023) and some azimuthal asymmetries can be linked to inner warps or other disk processes (Pérez et al. 2018a; Paneque-Carreño et al. 2022). Vertical structure, however, has been less accessible from an observational point of view and therefore less studied in comparison to the other two dimensions.

Directly probing the vertical structure and molecular layering of disks through observations of edge-on disks has offered valuable insight on the thermal and density conditions (Dutrey et al. 2017; Ruíz-Rodríguez et al. 2021), as well as dust-settling processes (Villanave et al. 2020, 2023). In the recent years, novel methodologies (Pinte et al. 2018a) have proven that the vertical structure can also be traced from mid-inclination ( $40\text{--}60^\circ$ ) disks, opening an exciting avenue to explore this dimension in larger samples (Teague et al. 2021b; Rich et al. 2021; Law et al. 2021b, 2022; Paneque-Carreño et al. 2023; Pinte et al. 2023). From a theoretical point of view, the vertical distribution of the gaseous disk material will follow hydrostatic equilibrium between the stellar gravitational potential and thermal pressure support (Armitage 2015). This implies that a number of properties such as stellar mass and radiation field, but also disk temperature structure and density distribution are expected to affect the vertical extent (D’Alessio et al. 1998; Aikawa et al. 2002). While the bulk of the disk material is composed of molecular hydrogen ( $\text{H}_2$ ), we are not able to trace the disk structure through its emission, as it is extremely hard to detect throughout the cold outer ( $r > 50$  au) disk region. Due to this difficulty, less abundant, but bright tracers, such as carbon monoxide (CO) are preferred to characterize the disk structure and conditions (for a review see Miotello et al. 2023).

This work focuses on the protoplanetary disk properties that affect the vertical structure as traced by bright CO ( $2 - 1$ ) emission. CO is the main carrier of gas-phase carbon in the interstellar medium and its chemistry is relatively simple and reliably implemented in various physical-chemical models (e.g. Aikawa et al. 2002; Gorti & Hollenbach 2008; Bruderer 2013; Miotello et al. 2014; Voitke et al. 2016; Yu et al. 2016; Molyarova et al. 2017). For this reason CO is a preferred molecule to study the disk extent (radially and vertically, Ansdell et al. 2018; Sanchis et al. 2021; Law et al. 2021a, 2022; Rich et al. 2021), kinematical processes (Pinte et al.

2023), temperature structure (Law et al. 2021b; Paneque-Carreño et al. 2023; Stapper et al. 2023) and disk mass (which usually requires analysis of rarer CO isotopologues, Miotello et al. 2014, 2016; Booth et al. 2019; Stapper et al. 2024). Vertically, CO gas emission originates from the warm molecular layer defined by the boundary of UV photo-dissociation in the upper regions and low temperatures that cause CO freeze-out ( $\sim 20$  K) closer to the midplane (Aikawa et al. 2002). At typical temperatures (40–60 K) and CO column densities ( $\geq 5 \times 10^{16} \text{ cm}^{-2}$ ) of the molecular layer in protoplanetary disks, the CO millimeter emission will be optically thick ( $\tau \geq 1$ , van der Tak et al. 2007). Indeed, previous works have shown that the emitting surface traced by CO, as extracted from emission channel maps, follows the CO  $\tau \sim 1$  region of the disk structure (Paneque-Carreño et al. 2023). Efforts in linking the observationally derived CO surfaces to system or emission properties such as stellar mass, brightness temperature or disk radius have only provided tentative trends (Law et al. 2022, 2023b) and it is not yet well understood how the wide morphology of CO vertical profiles relates to the host disk.

A crucial assumption that most studies related to CO emission make is that its abundance with respect to  $\text{H}_2$ , as set by the volatile carbon abundance, follows the canonical interstellar medium (ISM) value of  $\sim 10^{-4}$  (e.g. Andrews et al. 2011; Ansdell et al. 2016; Flaherty et al. 2015; Stapper et al. 2024). This is in tension with observational results that show that between 1-2 orders of magnitude of volatile carbon depletion must be present to account for disk masses derived from HD observations (Favre et al. 2013; McClure et al. 2016; Schwarz et al. 2016; Calahan et al. 2021) and derivations of CO depletion, with respect to ISM abundance, from optically thin CO isotopologue emission profiles (Zhang et al. 2019, 2021). Lower-than-ISM CO abundances may be caused by chemical reprocessing of CO into other molecules (Bergin et al. 2014; Bosman et al. 2018; Diop et al. 2024) or physical processing through which carbon is removed from the disk atmosphere by icy dust grains that grow and settle towards the disk midplane (Krijt et al. 2016, 2018, 2020). These processes are in addition to photodissociation and freeze-out, which will also lower the CO abundance throughout the disk and are included in standard thermochemical models (e.g., Bruderer 2013; Woitke et al. 2016). Overall, it is expected that the CO abundance may change as the disk evolves and therefore it is not inherited from the natal cloud environment (Visser et al. 2009; Bergner et al. 2020; Zhang et al. 2021). To the best of our knowledge, there are no previous studies that model or observationally analyze the effect of volatile carbon depletion on the vertical emitting surface of disks.

The remainder of this paper is organized as follows. Section 2 details the setup of thermochemical models used in our analysis, indicating the relevance of accounting for hydrostatic equilibrium for a more realistic approach. Section 3 shows our main results, an overview of the sampled parameter space and the effect of various disk and stellar properties on the location of the  $\tau = 1$  CO emission layer. We present a  $z/r$ –Disk mass relation for both T Tauri and Herbig disk systems and apply it to a sample of archival protoplanetary disk observations. Our results showcase the necessity of considering carbon depletion in order to obtain disk masses that are in agreement with literature values. Section 4 has a detailed

discussion of our results, comparing them to previous studies and literature values of carbon depletion. Finally, Section 5 indicates the main conclusions of this work.

## 5.2 Modelling

The results shown in this paper are based on an extensive grid of models run with the thermochemical code DALI (Bruderer 2013). For details on the code, how the chemical network is setup and the calculation of the temperature structure, dust populations and flux values we reference to past works using the same code (e.g. Bruderer et al. 2012; Bruderer 2013; Miotello et al. 2016; Leemker et al. 2022; Stapper et al. 2024). Our models are run using the chemical network from Miotello et al. (2014), that accounts for CO isotope-selective effects, considering selective photodissociation, fractionation reactions, self-shielding, and freeze-out (for more details see Miotello et al. 2014, 2016).

### 5.2.1 Initial DALI setup

The initial setup of the radial structure for the gas and dust in DALI follows the self-similar solution to a viscously evolving disk (Lynden-Bell & Pringle 1974; Andrews et al. 2011).

$$\Sigma(R) = \Sigma_c \left( \frac{R}{R_c} \right)^{-\gamma} \exp \left[ - \left( \frac{R}{R_c} \right)^{2-\gamma} \right] \quad (5.1)$$

where  $R_c$  is the characteristic radius and  $\gamma$  the surface density exponent. The initial vertical structure of the disk follows a Gaussian distribution, where the gas density ( $\rho$ ) corresponds to,

$$\rho(z) = \rho_c(r) \exp \left[ - \frac{z^2}{(2H^2)} \right] \quad (5.2)$$

and  $H$  is the disk scale height. To set the scale height aspect ratio ( $H/r$ ), we have modified the usual DALI prescription (where the aspect ratio is typically noted as  $h$ , see for example Leemker et al. 2022) to match the parametrization of Zhang et al. (2021). In our model the scale height aspect ratio as a function of radius follows,

$$H/r = h_{100} \left( \frac{R}{100 \text{au}} \right)^\varphi \quad (5.3)$$

where  $h_{100}$  is the scale height aspect ratio at 100 au. The main results in this study are based on the fiducial models for a T Tauri and Herbig system following the values presented in Table 5.1. For the T Tauri, excess UV radiation due to a mass accretion rate of  $10^{-8} M_\odot \text{yr}^{-1}$  was taken into account. It was assumed that the gravitational potential energy of the accreted mass is released with 100% efficiency as blackbody emission with  $T = 10^4 \text{K}$  (Miotello et al. 2014, 2016). For our parameters, these assumptions result in  $L_{\text{FUV}}/L_{\text{bol}} = 1.46 \times 10^{-1}$ . Beyond

**Table 5.1:** Initial parameters in fiducial DALI model

Parameter	Value
<i>Chemistry</i>	
Chemical Age	1 Myr
Volatile [C]/[H]	$1.35 \times 10^{-4}, 10^{-5}, 10^{-6}$
Volatile [O]/[H]	$2.88 \times 10^{-4}, 10^{-5}, 10^{-6}$
[PAH]	$10^{-4}$ ISM abundance
<i>Physical Structure</i>	
gas-to-dust ratio	100
$f_{\text{large}}$ (small/large grains)	0.95
$\chi$ (settling of large grains)	0.2
$\varphi$ (flaring index)	0.1
$h_{100}$ (scale height aspect ratio)	0.1
$R_c$ (characteristic radii)	100 au
$M_d$ (total disk mass)	$10^{-1}, 10^{-2}, 10^{-3}, 10^{-4} M_{\odot}$
<i>Stellar Parameters</i>	
$L_x$	$1.0 \times 10^{30} \text{ erg s}^{-1}$
$T_x$	$7.0 \times 10^7 \text{ K}$
<i>T Tauri model</i>	
$M_*$	$1.0 M_{\odot}$
$\dot{M}_{\text{acc}}$	$10^{-8} M_{\odot} \text{ yr}^{-1}$
$L_{\text{bol}}$	$1.0 L_{\odot}$
$T_{\text{eff}}$	4000 K
<i>Herbig model</i>	
$M_*$	$2.0 M_{\odot}$
$L_{\text{bol}}$	$17.0 L_{\odot}$
$T_{\text{eff}}$	9000 K

**Table 5.2:** Variation of DALI model parameters. Bold numbers indicate the fiducial model values.

Parameter	Sampled values
<i>T Tauri model</i>	
$\varphi$ (flaring index)	<b>0.1</b> , 0.2, 0.5
$f_{\text{large}}$ (small/large grains)	0.9, <b>0.95</b> , 0.99
$T_{\text{eff}}$	3500, <b>4000</b> , 4500 K
$L_{\text{bol}}$	0.2, <b>1.0</b> , $3 L_{\odot}$
$\dot{M}_{\text{acc}}$	$10^{-9}$ , <b><math>10^{-8}</math></b> , $10^{-7} M_{\odot} \text{ yr}^{-1}$
$R_c$ (characteristic radii)	15, 50, <b>100 au</b>
<i>Herbig model</i>	
[PAH]	<b><math>10^{-4}</math></b> , $10^{-2}$ , $10^{-1}$ ISM abundance
$T_{\text{eff}}$	7500, <b>9000</b> , 10000 K
$L_{\text{bol}}$	10, <b>17</b> , $30 L_{\odot}$
$R_c$ (characteristic radii)	15, 50, <b>100 au</b>

the fiducial models, we study a wider parameter space to determine the effect of key disk properties on the location of the vertical structure as traced by  $^{12}\text{CO}$ , Table 5.2 indicates the sampled values.

## 5.2.2 Hydrostatic equilibrium

The initial setup presented in the previous section is a useful approximation, but is not realistic, as it is known that the disk vertical structure does not follow a gaussian distribution, rather it is set by hydrostatic equilibrium between the stellar gravity and pressure support of the disk (given by the thermal structure). To produce more realistic models, DALI has a hydrostatic equilibrium solver implemented, which self-consistently iterates over consecutive models, starting from the initial parametrization, to compute the disk structure. We use this approach for all of our analysis. To this end, the hydrostatic equilibrium equation that must be solved in the vertical direction, using cylindrical coordinates, is:

$$\frac{1}{\rho} \frac{dP}{dz} = -\frac{d\phi}{dz} \quad (5.4)$$

where  $\rho$  is the gas density,  $P$  the gas pressure and  $\phi$  the gravitational potential. Without considering self-gravity,  $\phi$  corresponds to the stellar gravitational potential and follows,

$$\phi = -\frac{GM_*}{(r^2 + z^2)^{1/2}} \quad (5.5)$$




Pressure and density can be connected such that  $P/\rho = c_s^2 = kT/(\mu m_p)$ . Disks are not isothermal in their vertical direction, therefore, the sound speed ( $c_s$ ) will vary as function of height, depending on the density and thermal conditions. The differential equation for  $P$  is

$$\frac{1}{P} \frac{dP}{dz} = -\frac{z GM_*}{c_s^2 (r^2 + z^2)^{3/2}} \quad (5.6)$$

and can be solved through


$$P(z) = c_s(z)^2 \rho(z) = C \exp\left(-\int \frac{z GM_*}{c_s^2 (r^2 + z^2)^{3/2}} dz\right) \quad (5.7)$$

which will obtain the vertical distribution for the gas density ( $\rho(z)$ ). In DALI, the disk temperature structure is first obtained from the initial model, which computes the thermal conditions of the material considering all relevant cooling and heating processes, and then used to update the gas density distribution accounting for thermal support. The updated density is consecutively used to recalculate the temperature for the next iteration and start over until reasonable convergence of the disk structure is achieved. The integration constant  $C$  is adjusted such that  $\int \rho(z) dz$  is preserved with each iteration. In the isothermal approximation  $c_s^2$  does not have a vertical dependence and we recover the gaussian distribution indicated in equation 2.



**Figure 5.1:** Comparison between the outputs of a vertically static gaussian distribution and a model after solving self-consistently for the vertical material distribution. Each row shows the 2D distribution of key parameters, displayed using the same colorbar. From top to bottom the volumetric gas density, gas temperature and CO abundance (relative to  $\text{H}_2$ ) are shown. In the bottom row the solid line traces the  $^{12}\text{CO}$   $\tau = 1$  surface and the dashed line the  $^{13}\text{CO}$   $\tau = 1$  surface.

Figure 5.1 shows the 2D maps for the gas density, temperature and CO abundance (with respect to  $\text{H}_2$ ) of a converged DALI model for a  $10^{-2}M_\odot$  disk around a  $1M_\odot$  and  $1L_\odot$  stellar system. The static gaussian distribution corresponds to the initial DALI output, computed from the setup equations outlined in section 2.1. The self-consistent model solution is the result of six iterations solving the hydrostatic equilibrium equations, considering the previously computed thermal structure. The material appears to be more vertically extended in the self-consistent model, which is expected due to the increased pressure support of the heated atmosphere calculated from the vertical temperature gradient. Our study focuses on the  $^{12}\text{CO}$  2 – 1 emission surface, which is directly computed from the models as the location where the integrated column density of material results in an optical depth ( $\tau$ ) of 1 for  $^{12}\text{CO}$  2 – 1. Indeed, the  $\tau = 1$  layer of DALI models is consistent with the emission surfaces extracted from ALMA gas emission maps of  $^{12}\text{CO}$  2 – 1 (Paneque-Carreño et al. 2023).



**Figure 5.2:** Variation of the  $^{12}\text{CO}$   $\tau = 1$  surface when considering the hydrostatic equilibrium model solutions. Left panel shows the output for a T Tauri system and Right panel the results for a Herbig system. Both models correspond to a disk mass of  $10^{-2}M_\odot$ . The dashed lines trace the surface from the parametric gaussian distribution and the solid black line the final hydrostatic equilibrium solution considered. Each red line represents a model iteration of hydrostatic equilibrium in DALI.


When accounting for hydrostatic equilibrium, the  $\tau = 1$  surface varies both in its vertical and radial extent from the surface extracted from the parametric model. Figure 5.2 indicated the variations for the  $\tau = 1$  layer in a  $10^{-2}M_\odot$  disk around a  $1M_\odot$  and  $1L_\odot$  star (T Tauri model) compared to the same disk mass around a  $2M_\odot$  and  $17L_\odot$  star (Herbig model). As the hydrostatic equilibrium solution depends on the stellar mass, temperature and density distribution, it is expected that the surfaces will be different in each case. Convergence is reached rapidly after a couple of iterations, therefore all presented models in this work will consider the output from the 6th iteration of the self-consistent hydrostatic equilibrium solver.

### 5.3 Results

As has been shown in multiple studies, CO molecular surfaces can be parametrized through an exponentially tapered power-law (Law et al. 2021b, 2023b; Paneque-Carreño et al. 2023). To model each  $^{12}\text{CO}$   $\tau = 1$  surface from the self-consistent DALI outputs after solving the hydrostatic equilibrium equations, we select the following prescription,

$$z(r) = z_0 \times \left(\frac{r}{100 \text{ au}}\right)^\phi \times \exp\left[\left(\frac{-r}{r_{\text{taper}}}\right)^\psi\right] \quad (5.8)$$


The best fit values for  $z_0$ ,  $\phi$ ,  $r_{\text{taper}}$  and  $\psi$  are obtained using a Monte Carlo Markov Chain (MCMC) sampler as implemented by emcee (Foreman-Mackey et al. 2019). Each surface is then identified based on its characteristic  $z/r$  value, which corresponds to the best-fit line of constant  $z/r$  that traces the height profile within 80% of  $r_{\text{taper}}$  (Law et al. 2023b). This assures us that we are characterizing the rising portion of the  $^{12}\text{CO}$  surface, as exemplified in Figure 5.3 where a schematic of the method is shown.



**Figure 5.3:** Schematic of the method used to measure the  $^{12}\text{CO}$  surface  $z/r$  values. Each surface traced by the grey line is the output of a T Tauri model, in order of increasing height and radial extent they have disk masses of  $10^{-4}$ ,  $10^{-3}$ ,  $10^{-2}$  and  $10^{-1} M_{\odot}$ . The shaded red curve shows the best-fit parametric model fitted to the molecular surface, from where  $r_{\text{taper}}$  is the taper radius. The measured  $z/r$  is obtained by fitting the surface region within 80% of  $r_{\text{taper}}$ , this distance is indicated by the vertical dashed line. The characteristic  $z/r$  is traced by the straight black line for each surface.

### 5.3.1 Parameters that affect vertical CO surface


Our fiducial models for each stellar type are indicated in Table 5.1. To determine which parameter variation most strongly affects the measured  $z/r$  we compute additional models with a varied stellar luminosities and temperatures as reported in Table 5.2 covering the range of reported values from the comparison observations. Disk structure values for the flaring angle ( $\varphi$ ), small-to-large dust grain fraction ( $f$ ), PAH abundance, surface density characteristic radius ( $R_c$ ), total disk mass ( $M_d$ ) and volatile carbon abundance ( $C_{\text{abu}}$ ) are also sampled. For each model the characteristic  $z/r$  is computed as described at the beginning of Section 3. Figure 5.4 presents the percentual variation from the fiducial model for disk masses of  $10^{-3} M_\odot$  and  $10^{-2} M_\odot$ . The full vertical profiles for each parameter can be found in Figures 5.A.1 and 5.A.2.



**Figure 5.4:** Percentual variation of the  $^{12}\text{CO}$  surface  $z/r$  caused by various stellar and disk parameters. Left panel indicates the results for the T Tauri system and right panel for the Herbig system. Each parameter corresponds to a specific color. The circle markers indicate the effects on a disk with mass  $10^{-3} M_\odot$  and the diamond markers the effects on a disk with mass  $10^{-2} M_\odot$ . Specific details on the parameter range and properties are shown in Table 5.2.

The largest variation in  $z/r$  is caused by varying the total disk mass, which is sampled at  $10^{-4} M_\odot$ ,  $10^{-3} M_\odot$ ,  $10^{-2} M_\odot$  and  $10^{-1} M_\odot$ . An order of magnitude change in the disk mass can vary the emission height by more than 50%. Volatile carbon abundance also presents a strong effect in the emission surfaces. Indeed, an order of magnitude decrease in the assumed volatile carbon abundance with respect to the ISM can lower the characteristic  $z/r$  by  $\sim 30\%$  and two orders of magnitude in carbon depletion produce  $z/r$  variations beyond 50%. These two parameters have a degenerate contribution to the vertical surface, as can be seen from Figure 5.5. Increasing disk mass will push the CO emission layer higher, while for the same disk mass a lower volatile carbon abundance (higher depletion) will push the emitting layer towards the midplane. This effect of carbon abundance in the vertical structure had been previously observed in Calahan et al. (2021) when constraining the disk model of TW Hya.

Across all masses, for the same degree of carbon depletion, the T Tauri systems display a larger  $z/r$  than the Herbig disks (see Fig. 5.5). This is expected due to the larger stellar mass of the Herbig model, which will translate into a higher gravitational potential (equation 5). The parameter space between our models is covered using a cubic spline which we use to extrapolate our results beyond our mass grid, to nearby values. We refer to this result as the  $z/r$ - $M_d$  relationship, noting that volatile carbon depletion will shift it and must be accounted for.



**Figure 5.5:** Relation between the  $^{12}\text{CO}$  surface  $z/r$  and the total disk mass ( $z/r - M_d$  relationship) with a dependence on the volatile carbon abundance. Left panel shows results for T Tauri systems and right panel for Herbig systems. Circles indicate ISM-like carbon abundance (no depletion), diamonds trace a carbon depletion of 10 and squares a depletion of 100, with respect to the ISM.

All other parameters, including order of magnitude variations in stellar accretion rate and PAH abundance (see Table 5.2), cause a variation of less than  $\sim 30\%$ . We stress that these results only hold for the  $^{12}\text{CO}$  2 – 1 surface  $z/r$ , other molecules, particularly those more sensitive to UV fields are likely more significantly affected by some of the sampled parameters.

### 5.3.2 Model comparison to observations

In order to benchmark our model results and the  $z/r$ - $M_d$  relationship, we estimate the gas masses and expected  $z/r$  in an observational sample of 19 disks, as indicated in Table 5.3. The studied disks all have estimates of their total disk mass from dust continuum observations, additionally some of them have disk mass estimates from modelling of CO, HD and  $\text{N}_2\text{H}^+$  fluxes and dynamical analysis of rotation curves.

Most disk mass values from dust continuum emission are extracted by converting the total continuum flux into a dust mass, assuming a dust temperature and grain opacity and then using the canonical assumption of a gas-to-dust ratio of 100 (Hildebrand 1983; Ansdell et al. 2016; Andrews et al. 2013; Barenfeld et al. 2016; Pascucci et al. 2016; Stapper et al. 2022). Excluding the sources from the MAPS

**Table 5.3:** Observational sample, stellar parameters and estimated total disk masses

Star	$M_*$	$L_*$	$T_*$	Previous disk mass estimates [ $M_\odot$ ]				References
	[ $M_\odot$ ]	[ $L_\odot$ ]	[ $T_\odot$ ]	Dust continuum	CO flux	HD emission	Dynamical	
<b>T Tauri Disks</b>								
MY Lup	1.27	0.87	5129	$1.5 \times 10^{-2}$	$8.3 \times 10^{-5}$	-	-	4, 3, 12, 5
AS 209	1.2	1.41	4266	$7.5 \times 10^{-2}$	$4.5 \times 10^{-3}$	-	$2 \times 10^{-4}$	1, 13, 16, 18
WaOph 6	1.12	2.9	4169	$1.4 \times 10^{-2}$	-	-	-	4, 3, 14
IM Lup	1.1	2.57	4266	$3.67 \times 10^{-1}$	$2 \times 10^{-1}$	-	$1.1 \times 10^{-1}$	1, 13, 16, 18
GM Aur	1.1	1.2	4350	$7.3 \times 10^{-2}$	$2 \times 10^{-1}$	-	$1.2 \times 10^{-1}$	1, 13, 16, 18
DoAr 25	1.06	0.95	4266	$4.2 \times 10^{-2}$	-	-	-	4, 3, 5
LkCa 15	0.7	1.098	4162	$2.7 \times 10^{-2}$	-	-	-	5
GW Lup	0.62	0.33	3631	$1.5 \times 10^{-2}$	$9.6 \times 10^{-5}$	-	-	4, 3, 12, 5
Sz 91	0.55	0.26	3720	$2.2 \times 10^{-3}$	-	-	-	4, 10, 5
DM Tau	0.50	0.24	3705	$1.6 \times 10^{-2}$	-	-	-	4, 9, 5
Elias 2-27	0.46	0.91	3890	$3.4 \times 10^{-2}$	-	-	$8 \times 10^{-2}$	2, 3, 5, 17
<b>Herbig Disks</b>								
HD 100546	2.10	23.4	9750	$1.2 \times 10^{-2}$	$3.2 \times 10^{-2}$	$< 8 \times 10^{-2}$	-	4, 8, 11, 15, 19
HD 97048	2.70	44.2	10500	$4.7 \times 10^{-2}$	$1 \times 10^{-1}$	$< 9.4 \times 10^{-2}$	-	4, 8, 11, 15, 19
HD 100453	1.66	6.31	7400	$5.3 \times 10^{-3}$	$3.2 \times 10^{-2}$	$< 1 \times 10^{-2}$	-	6, 11, 15, 19
HD 34282	1.59	13.64	8625	$2.6 \times 10^{-2}$	$1.2 \times 10^{-1}$	-	-	7, 11, 15
MWC 480	2.1	21.9	8250	$1.2 \times 10^{-1}$	$1 \times 10^{-1}$	-	$1.5 \times 10^{-1}$	1, 13, 15, 16
HD 163296	2.0	17.0	9332	$8.4 \times 10^{-2}$	$1 \times 10^{-1}$	$< 6.7 \times 10^{-2}$	$1.3 \times 10^{-1}$	1, 13, 15, 16, 19
HD 142666	1.73	9.1	7500	$7.6 \times 10^{-3}$	$3.2 \times 10^{-2}$	-	-	4, 8, 11, 15
Ak Sco	1.2	2.4	6250	$1.8 \times 10^{-3}$	$2.3 \times 10^{-3}$	-	-	8, 11, 15

**References.** (1) Öberg et al. 2021b (2) Veronesi et al. 2021 (3) Andrews et al. 2018 (4) Law et al. 2022 (5) Manara et al. 2023 (6) Rosotti et al. 2020 (7) Merín et al. 2004 (8) Fairlamb et al. 2015 (9) Pegues et al. 2020 (10) Tsukagoshi et al. 2019 (11) Stapper et al. 2022 (12) Miotello et al. 2017 (13) Sierra et al. 2021 (14) Brown-Sevilla et al. 2021 (15) Stapper et al. 2024 (16) Martire et al. 2024 (17) Veronesi et al. 2021 (18) Zhang et al. 2021 (19) Kama et al. 2020

program (Öberg et al. 2021b), the total disk masses estimated from CO millimeter emission for the Herbig sample are calculated in Stapper et al. (2024) based on an extensive grid of models, as is the case of MY Lup and GW Lup from the T Tauri sample (Miotello et al. 2017). For the disks in the MAPS program (AS 209, IM Lup, GM Aur, MWC 480, HD 163296 Öberg et al. 2021b) the total disk mass calculated from dust continuum accounts for dust scattering processes and has a more accurate fitting of the thermal conditions (Sierra et al. 2021), on the other hand, the CO disk masses come from detailed thermochemical modelling of Zhang et al. (2021). Upper limits of disk mass from HD emission were also calculated using thermochemical disk models to match the non-detection threshold (Kama et al. 2020). Through the combination of  $N_2H^+$  and  $C^{18}O$  emission, thermochemi-


cal models assuming a degree of disk ionization have computed the total disk mass and expected carbon depletion (Trapman et al. 2022). Dynamical masses have been obtained for some disks, through the detection of super-keplerian motion in the outer radius of the disk rotation profile using isothermal (Veronesi et al. 2021) and thermally stratified (Martire et al. 2024) vertical disk prescriptions.

This sample has been constructed from mid-inclination disks with available  $^{12}\text{CO } 2-1$  data cubes from which the surfaces have been extracted and published in previous works (Law et al. 2022, 2023b; Stapper et al. 2023; Paneque-Carreño et al. 2023). In all cases, the vertical structure from the observations is obtained following the methodology presented in Pinte et al. (2018a), where the surfaces are constrained based on the emission maxima across the molecular channel maps. All  $^{12}\text{CO}$  surfaces for the Herbig disk sample are taken from Stapper et al. (2023), where the ALFAHOR code (Paneque-Carreño et al. 2023) is used to extract the vertical profile. For the T Tauri disks, part of the sample (AS 209, WaOph 6, IM Lup, GM Aur, Elias 2-27) is presented in Paneque-Carreño et al. (2023), also using ALFAHOR. The remainder of the T Tauri disks are published in Law et al. (2023b) using an initial version of the DISKSURF code (Teague et al. 2021b) that in some cases underestimated the vertical profiles due to contamination from the back side of the disk (Paneque-Carreño et al. 2023). In order to do a uniform analysis, the disks previously studied using DISKSURF (MY Lup, DoAr 25, LkCa 15, GW Lup, Sz 91, DM Tau) are re-analysed using the ALFAHOR, yielding comparable results but with a lower scatter. All vertical profiles from the observational sample are presented in Figures 5.A.3 and 5.A.4, compared to the vertical profiles from the fiducial DALI models of varying disk mass with ISM-like carbon abundance.

As done for the DALI models, the vertical profiles of the observational sample are processed to obtain their characteristic  $z/r$ . To accurately account for the spread in the vertical profile,  $z/r$  is calculated for the mean value and also for the lower and upper limits of the profiles, which indicate the dispersion within radial bins and will give the uncertainty on the characteristic  $z/r$ .

Based on the  $z/r - M_d$  relationship shown in Figure 5.5 we estimate the expected  $z/r$  for the reported total disk mass derived from dust continuum and available HD upper limits, assuming an ISM-like carbon abundance. These values are compared against the observationally derived  $z/r$  values in Figure 5.6. For the majority of the sample, if we assume an ISM-like carbon abundance, our measured  $z/r$  is below the expected value based on the dust masses considering a gas-to-dust ratio of 100. The only exceptions are DM Tau, Elias 2-27, HD 97048 and HD 34282. Throughout the sample, the differences are more noticeable in the T Tauri systems, as Herbig systems have better agreement within the uncertainty of our method. We note that taking into consideration carbon depletion would lower the expected  $z/r$  of the literature mass measurements (see Fig. 5.5), obtaining a better agreement with the observational emission heights.

Overall, these results may indicate that the assumed gas-to-dust ratio in the systems is much lower than 100 (effectively lowering the literature estimates from the dust flux) or that our model assumptions are failing to account for a significant factor affecting the  $z/r$  value, such as the presence of carbon depletion. We favor



**Figure 5.6:** Comparison between the measured  $^{12}\text{CO}$  surface  $z/r$  for a sample of disks and their expected values, based on literature total disk mass estimates from dust continuum observations. Black dots indicate the  $z/r$  values obtained in this study for our disk sample, open red circles trace the expected  $z/r$  values estimated from the dust continuum total disk masses and downwards triangles the expected upper limits of  $z/r$  from HD measurements. Left panel shows the results for a T Tauri system and right panel for a Herbig system. Each horizontal line indicates the  $z/r$  of our DALI models with disk masses of  $10^{-4} M_{\odot}$ ,  $10^{-3} M_{\odot}$ ,  $10^{-2} M_{\odot}$  and  $10^{-1} M_{\odot}$ , in increasing  $z/r$  order respectively.

the second scenario as dust-derived masses are expected to be lower limits of the total disk mass due to the assumptions on uncertain dust properties required for their calculation and usually unaccounted for effects such as optical depth and scattering (see discussion and references in Miotello et al. 2023). Additionally, dynamical estimates of total disk mass have shown typical gas-to-dust ratios  $\gtrsim 100$  (Lodato et al. 2023; Martire et al. 2024), further indicating that the total disk mass derived from dust measurements should be a lower limit to the actual value.


### 5.3.3 Accounting for volatile carbon depletion

The initial abundance of volatile carbon in our models will determine the amount of CO formed in each case. We have shown in the previous sections that this also has a significant effect in the extracted vertical CO surfaces, with low-carbon models producing shallower emission layers. There is independent observational motivation to assume carbon depletion for our disk sample, as measurements of CO abundance in some of the sources have shown depletion factors of 10-100 from the typically assumed ISM value of  $\sim 10^{-4}$  (Zhang et al. 2019, 2021).

Figure 5.7 compares the published masses from various methods (Table 5.3) to the derived disk mass from our  $z/r$  values, calculated accounting for varying degrees of carbon depletion in the  $z/r - M_d$  relationship. For disks that show a large scatter in their literature values we focus on the total disk mass extracted from the dust mass assuming a gas-to-dust ratio of 100 as main indicator of the



expected value.



**Figure 5.7:** Estimated total disk masses from  $z/r$  analysis compared to literature values. Left panel shows the results for the T Tauri systems and right panel for the Herbig sources. For each disk the available literature values of the disk mass are shown in colored crosses and grey triangles for the HD upper limits. Black symbols correspond to the estimates from the characteristic  $z/r$ , where circles, diamonds and squares have been calculated using the  $z/r - M_d$  relationship curves for 0, 10 and 100 values of volatile carbon depletion, respectively.

In some disks it is sufficient to consider a volatile carbon depletion of 10 to match the dust masses, however many require carbon depletion of 100 to be in agreement, particularly in the T Tauri systems. We briefly discuss here the case of HD 163296 and MWC 480, which have well constrained CO depletion profiles from optically thin  $C^{18}O$  and  $C^{17}O$  emission (Zhang et al. 2021). The HD 163296 literature analysis indicates an almost constant CO abundance profile, depleted by a factor 10 with respect to the ISM value beyond  $\sim 50$  au and MWC 480 follows a similar pattern going towards even lower values of CO depletion beyond  $\sim 100$  au (Zhang et al. 2021). Our analysis indicates that for HD 163296 a global volatile carbon depletion of  $\sim 10$  is required to match the literature disk mass estimates to those derived from the CO  $z/r$ . For MWC 480 the preferred carbon depletion from our results ranges between 10-100, also in agreement with the spatially resolved values of Zhang et al. (2021).

For all disks, the inferred total mass we obtain from the characteristic  $z/r$ , for each carbon depletion value, are presented in Table 5.4 together with the preferred carbon depletion. The volatile carbon depletion value should be taken with caution, as it has been visually estimated based on comparison to the dust-derived total disk mass. In the optimal scenario it may be a lower limit of the disk integrated volatile carbon depletion with respect to the ISM.

**Table 5.4:** Studied systems, their calculated characteristic  $^{12}\text{CO}$   $z/r$  and associated total disk mass for various volatile carbon depletion scenarios.

Star	Characteristic $^{12}\text{CO}$ $z/r$	Total disk mass [ $\log(M_{\odot})$ ]			Preferred carbon depletion
		ISM C abundance	C depletion: 10	C depletion: 100	
<b>T Tauri Disks</b>					
MY Lup	$0.16^{+0.03}_{-0.05}$	$-3.52^{+0.38}_{-0.48}$	$-2.99^{+0.37}_{-0.44}$	$-2.16^{+0.33}_{-0.44}$	100
AS 209	$0.21^{+0.05}_{-0.08}$	$-2.95^{+0.53}_{-0.85}$	$-2.45^{+0.55}_{-0.81}$	$-1.67^{+0.49}_{-0.74}$	-
WaOph 6	$0.30^{+0.05}_{-0.07}$	$-2.19^{+0.34}_{-0.63}$	$-1.67^{+0.35}_{-0.64}$	$-0.97^{+0.36}_{-0.59}$	10
IM Lup	$0.40^{+0.03}_{-0.05}$	$-1.55^{+0.17}_{-0.34}$	$-1.02^{+0.18}_{-0.35}$	$-0.26^{+0.21}_{-0.39}$	10-100
GM Aur	$0.38^{+0.05}_{-0.08}$	$-1.67^{+0.27}_{-0.56}$	$-1.13^{+0.27}_{-0.57}$	$-0.40^{+0.32}_{-0.6}$	10
DoAr 25	$0.24^{+0.03}_{-0.05}$	$-2.73^{+0.32}_{-0.50}$	$-2.22^{+0.33}_{-0.49}$	$-1.48^{+0.30}_{-0.43}$	100
LkCa 15	$0.25^{+0.04}_{-0.06}$	$-2.60^{+0.34}_{-0.56}$	$-2.09^{+0.35}_{-0.56}$	$-1.36^{+0.33}_{-0.49}$	100
GW Lup	$0.25^{+0.08}_{-0.11}$	$-2.60^{+0.59}_{-1.1}$	$-2.09^{+0.6}_{-1.08}$	$-1.36^{+0.58}_{-0.96}$	10-100
Sz 91	$0.12^{+0.04}_{-0.06}$	$-3.94^{+0.47}_{-0.52}$	$-3.38^{+0.43}_{-0.46}$	$-2.54^{+0.42}_{-0.55}$	100
DM Tau	$0.45^{+0.05}_{-0.09}$	$-1.27^{+0.22}_{-0.53}$	$-0.73^{+0.23}_{-0.54}$	-	0
Elias 2-27	$0.55^{+0.05}_{-0.09}$	$-0.82^{+0.17}_{-0.49}$	$-0.27^{+0.17}_{-0.50}$	-	0
<b>Herbig Disks</b>					
HD 100546	$0.19^{+0.08}_{-0.1}$	$-2.27^{+0.74}_{-1.31}$	$-1.83^{+0.76}_{-1.09}$	$-1.14^{+0.84}_{-1.15}$	10
HD 97048	$0.32^{+0.04}_{-0.07}$	$-1.14^{+0.27}_{-0.52}$	$-0.57^{+0.41}_{-0.66}$	-	0
HD 100453	$0.14^{+0.08}_{-0.09}$	$-2.83^{+0.87}_{-1.18}$	$-2.30^{+0.76}_{-1.03}$	$-1.64^{+0.83}_{-1.06}$	10
HD 34282	$0.37^{+0.04}_{-0.08}$	$-0.83^{+0.24}_{-0.49}$	-	-	0
MWC 480	$0.24^{+0.04}_{-0.05}$	$-1.74^{+0.32}_{-0.53}$	$-1.31^{+0.37}_{-0.52}$	$-0.56^{+0.41}_{-0.57}$	10-100
HD 163296	$0.28^{+0.02}_{-0.04}$	$-1.42^{+0.18}_{-0.29}$	$-0.94^{+0.23}_{-0.34}$	$-0.14^{+0.14}_{-0.39}$	10
HD 142666	$0.15^{+0.05}_{-0.05}$	$-2.69^{+0.48}_{-0.65}$	$-2.19^{+0.42}_{-0.53}$	$-1.52^{+0.45}_{-0.56}$	10-100
Ak Sco	$0.07^{+0.03}_{-0.05}$	$-3.82^{+0.48}_{-0.46}$	$-3.14^{+0.43}_{-0.49}$	$-2.51^{+0.43}_{-0.48}$	100

## 5.4 Discussion

## 5.5 Discussion

### 5.5.1 Low gas-to-dust or high carbon depletion?

It has been previously discussed in the literature that current CO gas models compared to observations do not allow us to distinguish between a low gas-to-dust system (lower gas mass than dust-derived total disk mass) or a high volatile carbon depletion scenario (Miotello et al. 2017; Calahan et al. 2021). Our results are in agreement with this entanglement, however we favor the high carbon depletion over low gas-to-dust due to evidence for CO abundance evolution over time in young stars (Bergner et al. 2020; Zhang et al. 2020), measurements of CO depletion in sources from our sample (Zhang et al. 2019, 2021) and results from dynamical

mass estimates that indicate gas-to-dust ratios  $\gtrsim 100$  (Martire et al. 2024) rather than much less than 100.

As discussed in Miotello et al. (2023), various methods can yield orders of magnitude differences in the calculated disk mass. This is particularly apparent for the literature total disk mass values of T Tauri systems MY Lup and GW Lup. For these disks our analysis indicates that a high carbon depletion of  $\sim 100$  is needed to reach the reported dust-derived mass value, but the reported gas mass, which is several ( $\sim 2$ ) orders of magnitude lower in both cases, was extracted without accounting for carbon depletion (Miotello et al. 2017). It is therefore likely that a higher carbon depletion rate would put all values in agreement, an alternative that was suggested by Miotello et al. (2017) to account for the low gas-to-dust ratios that the measurement would otherwise entail.

We note that for the Herbig sample, the gas mass estimates used in this work were mostly obtained from DALI models of isotopologue CO emission (Stapper et al. 2024), measuring the fluxes and disk radii in these tracers. These models assumed an ISM volatile carbon abundance, with a parameter space similar to our models, however they obtain mass values higher than the dust-based masses, consistent with a gas-to-dust ratio  $\gtrsim 100$  (see right panel of Figure 5.7). We are confident that carbon depletion must happen at least for MWC 480 and HD 163296, as it has been measured independently through dedicated models and observations (Zhang et al. 2021). HD 100546 also has estimates of volatile carbon depletion between 2-10, consistent with our results (Bruderer et al. 2012; Kama et al. 2016). The overall discrepancy between our work and analysis using models of CO isotopologue fluxes is the consideration of hydrostatic equilibrium in our models. We show in Figure 5.B.1 that for high disk masses, when the gas emission becomes optically thick, not accounting for hydrostatic equilibrium can underestimate the CO flux, therefore overestimating the calculated disk masses. Due to the optical depth, the use of model fluxes in the high disk mass regime also leads to large uncertainties of up to an order in magnitude in the calculated mass, which also affects the comparison (Miotello et al. 2017; Stapper et al. 2024). Overall, this highlights the sensitivity of vertical surfaces as a probe for total disk mass and volatile carbon depletion.

## 5.5.2 Synergy with other methods

Using our  $z/r - M_d$  relationships can be useful to have an initial idea on the system properties, contrary to other methods, we are able to utilize a bright and readily detected molecule to have a first-order joint constraint of disk mass and carbon abundance. Indeed, if the calculated  $z/r$  is  $\geq 0.3$  then the total disk mass is  $\geq 10^{-2} M_\odot$ . For lower  $z/r$ , the uncertainty on the carbon abundance creates a span of several orders of magnitude in the total disk mass value. However, if gas-to-dust ratio of at least 100 is assumed, which is reasonable from accretion rate values that point towards gas-rich disks (Manara et al. 2016, 2023) and dynamical mass estimates (Lodato et al. 2019; Martire et al. 2024), this work has shown that the total mass calculated from the dust continuum flux may be used as an anchor to estimate the lower limit of carbon depletion (see section 3.3).

Combining the information from CO surfaces with dynamical mass estimates, as has been done for a small sample in this work would put strong constraints on the overall carbon depletion. Furthermore, the use of rarer CO isotopologues and C-abundance-sensitive tracers such as  $\text{N}_2\text{H}^+$  has demonstrated to be a good tool to discern between the effects of varying disk mass and volatile carbon depletion (Anderson et al. 2019, 2022; Trapman et al. 2022).  $\text{N}_2\text{H}^+$  models have the uncertainty of the disk ionization rate and complication of relying on a fainter molecular line, however in combination with measurements of the CO vertical profile it could be possible to determine all three parameters, key for understanding protoplanetary disk conditions. We highlight that our results for DM Tau and GM Aur using the CO surface and dust-derived disk masses are in great agreement with the total disk mass and carbon depletion values determined in Trapman et al. (2022) through  $\text{N}_2\text{H}^+$  and  $\text{C}^{18}\text{O}$  flux analysis.

### 5.5.3 Caveats and future perspectives

While exciting and very useful for an initial overview of the disk conditions, there are a few caveats to our results. In particular, the studied models have been developed using a full-disk prescription, a standard radial density profile and a single disk-integrated volatile carbon depletion value. However observations show disks with a plethora of substructure (e.g. Andrews et al. 2018; Andrews 2020; Long et al. 2019) and that CO abundance profiles varies radially (Zhang et al. 2019, 2021). Highly structured or transition disks such as AS 209 and LkCa 15 are likely not accurately represented by our models. Future work, beyond the scope of this paper, will test the effect of large cavities and other substructure on the vertical profile and  $z/r$ - $M_d$  relationship.

We note that emission surfaces obtained at high enough spatial resolution may give direct information on the surface density profile, allowing for a more accurate description of the material distribution without the need to use a characteristic  $z/r$  (see for example, Paneque-Carreño et al. 2023). Through this approach, the total disk mass may be more accurately determined and scaled depending on the carbon abundance in a source-by-source analysis, without the uncertainties of general models.

Most importantly, future work must focus on different molecular tracers, such as rarer CO isotopologues and bright optically thin molecules, for example CN and  $\text{C}_2\text{H}$ , that have been proposed to be sensitive tracers of the upper disk regions (Cazzoletti et al. 2018; Bergin et al. 2016; Paneque-Carreño et al. 2021, 2024). Our results in this work are only applicable to the CO emitting layer, however it is likely that tracing the vertical structure with other molecular species will give direct insight into varied ongoing disk processes.

## 5.6 Conclusions

In this work we have presented a  $z/r$ - $M_d$  relationship, linking the vertical extent of CO 2 – 1 to the disk mass and volatile carbon abundance. This was done through

the study of thermochemical DALI models, computed accounting for hydrostatic equilibrium. Our models tested the effect of various stellar and disk parameters on the vertical structure and our theoretical predictions were benchmarked on a set of observations. The main findings and conclusions are the following:

1. We determine that for T Tauri and Herbig systems, the total disk mass and volatile carbon abundance are the main parameters that set the location of the CO 2 – 1 emission surface. This surface can be characterized by a constant characteristic  $z/r$  tracing the rising portion of the vertical profile.
2. The  $z/r$ - $M_d$  relationship for T Tauri systems is different than that of disks around Herbig stars. For the same disk mass and assumed volatile carbon abundance, material around T Tauri stars extends further vertically, as expected due to the difference in the stellar masses and gravitational potential.
3. Using the total disk masses inferred from continuum dust observations (assuming a gas-to-dust ratio of 100) we are able to calibrate the  $z/r$ - $M_d$  relationship for each individual system and estimate the order of magnitude of carbon depletion.
4. Fifteen out of the nineteen studied systems (79%) show indicators of volatile carbon depletion (compared to the ISM abundance). For disks with previous constraints on their carbon abundance, our results yield comparable values. T Tauri disks seem to be more carbon depleted than Herbig systems.

Overall, this study has demonstrated the utility of a bright and readily-detected molecular tracer, CO 2–1, that may be used to probe two of the most fundamental disk parameters through its vertical extent: total disk mass and carbon abundance. As the sample of disks with high resolution data grows, studies of vertical layering in mid-inclination disks will be used to deepen our understanding not only on the thermal and density conditions, but also on the amount and composition of the materials available for planet formation.


## Appendix

### 5.A Full emission surfaces from models and observational sample


In the main text our results are expressed in  $z/r$ . For visual comparison, here we show the full vertical profiles extracted for the CO emission surfaces in models and observations. Figures 5.A.1 and 5.A.2 show the emission surfaces tracing the  $\tau = 1$  layer, as extracted from the DALI model output of the fiducial models and models varying specific parameters (see Table 5.2 for details).

Figures 5.A.3 and 5.A.4 display the emission surfaces extracted from the CO channel map emission of each disk using ALFAHOR (Paneque-Carreño et al. 2023; Stapper et al. 2023). For comparison, the location of the  $\tau = 1$  layer from the

fiducial models using  $R_c = 100$  au of each sampled disk mass are plotted in each panel.




**Figure 5.A.1:** Vertical profiles tracing  $\tau = 1$  layer of the DALI model grid for  $M_{\text{disk}} = 10^{-2} M_{\odot}$ . Top row shows the sampled parameters for the T Tauri systems and bottom row for Herbig. In each panel the black dashed line corresponds to the fiducial model and only one parameter is varied, as indicated by the legend and colors.




**Figure 5.A.2:** Same as Figure 5.A.1 but for fiducial model with  $M_{\text{disk}} = 10^{-3} M_{\odot}$ .

## 5.B CO isotopologue model fluxes using hydrostatic equilibrium solver

When using the hydrostatic equilibrium solver in DALI, the disk density converges to a physical structure given by the equilibrium between the stellar gravitational potential and the thermal structure. The disk integrated fluxes may vary when comparing the hydrostatic equilibrium output to the models with static gaussian distributions. For optically thin emission in low-mass disks, the fluxes will be larger in the static models, which means the disk masses will be underestimated in




**Figure 5.A.3:** Observational CO emission surfaces for the T Tauri disk sample. Magenta lines indicate the observational results and data vertical dispersion, black lines show the  $\tau = 1$  layer from the fiducial DALI models using  $R_c = 100$  au for different disk masses indicated by different line styles. Grey lines mark constant  $z/r$  surfaces of 0.1, 0.2, 0.3, 0.4 and 0.5.



**Figure 5.A.4:** Same as Figure 5.A.3, but for the Herbig sample and models.

comparison to those obtained from model fluxes when hydrostatic equilibrium is considered. In the case of high-mass disks the reverse happens, the static gaussian models underestimate the fluxes, therefore overestimate the disk masses.

Figure 5.B.1 shows this effect displaying the total flux difference and percentual variation. This flux variation explains the generally higher masses of Stapper et al. (2024) and the lower masses obtained by Miotello et al. (2017) for MY Lup and GW Lup compared to our mass values assuming an ISM carbon abundance (see Fig. 5.7).



**Figure 5.B.1:** Comparison of retrieved fluxes from DALI models when using hydrostatic equilibrium and not. Left panel shows the  $^{13}\text{CO}$  and  $\text{C}^{18}\text{O}$  emission fluxes, black dots corresponds to models obtained after iterating and solving the hydrostatic equilibrium equations. Colored dots show the fluxes from models with a single iteration using a static gaussian material distribution. Right two panels display the percentual variation of the model flux from the static models compared to the one which uses hydrostatic equilibrium to determine its thermal and density structure.



