



Universiteit
Leiden
The Netherlands

Metabolomics in community-acquired pneumonia: exploring metabolomics-based biomarkers for diagnosis and treatment response monitoring of community-acquired pneumonia

Hartog, I. den

Citation

Hartog, I. den. (2024, September 17). *Metabolomics in community-acquired pneumonia: exploring metabolomics-based biomarkers for diagnosis and treatment response monitoring of community-acquired pneumonia*. Retrieved from <https://hdl.handle.net/1887/4083598>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4083598>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 3

Differential metabolic host response to pathogens associated with community-acquired pneumonia

Ilona den Hartog, Naama Karu, Laura B. Zwep, G. Paul Voorn, Ewoudt M.W. van de Garde, Thomas Hankemeier, J.G.C. van Hasselt. Differential metabolic host response to pathogens associated with community-acquired pneumonia. *Metabolism Open* **18** (2023).

Abstract

Background: Metabolic changes induced by the host immune response to pathogens found in patients with community-acquired pneumonia (CAP) may provide insight into its pathogenesis. In this study, we characterized differences in the host metabolic response to common CAP-associated pathogens.

Method: Targeted metabolomic profiling was performed on serum samples obtained from hospitalized CAP patients (n=119) at admission. We quantified 347 unique metabolites across multiple biochemical classes, including amines, acylcarnitines, and signaling lipids. We evaluated if unique associations between metabolite levels and specific CAP-associated pathogens could be identified.

Results: Several acylcarnitines were found to be elevated in *C. burnetii* and herpes simplex virus (HSV), and lowered in *M. pneumoniae* as compared to other pathogens. Phenylalanine and kynurenine were found elevated in *L. pneumophila* as compared to other pathogens. S-methylcysteine was elevated in patients with *M. pneumoniae*, and these patients also showed lowered cortisol levels in comparison to almost all other pathogens. For the herpes simplex virus, we observed a unique elevation of eicosanoids and several amines. Many lysophosphatidylcholines showed an altered profile in *C. burnetii* versus *S. pneumoniae*, *L. pneumophila*, and respiratory syncytial virus. Finally, phosphatidylcholines were negatively affected by the influenza virus in comparison to *S. pneumoniae*.

Conclusions: In this exploratory analysis, metabolites from different biochemical classes were found to be altered in serum samples from patients with different CAP-associated pathogens, which may be used for hypothesis generation in studies on differences in pathogen host response and pathogenesis of CAP.

3.1 Introduction

Community-acquired pneumonia (CAP) is a common infection of the lower respiratory tract, caused by bacterial or viral pathogens [51]. Causative pathogens that are commonly found in CAP patients include *Streptococcus pneumoniae*, respiratory viruses, *Haemophilus influenzae*, and, to a lesser extent *Mycoplasma pneumoniae* and *Legionella pneumophila* [51, 29, 30]. CAP is associated with high mortality and morbidity, especially in the elderly [3, 31], and with distinct differences in clinical outcomes between different CAP-associated pathogens [52, 53].

There is an increasingly recognized role of the patient-associated host metabolic response to infection and its association with overall innate immune system activation and clinical outcomes [17, 34, 54]. Previous studies have found distinct effects of this host metabolic response in COVID-19 and sepsis [55, 56, 57, 58]. We previously evaluated the discriminatory power of host-associated metabolites to support the microbial diagnosis of CAP and identified metabolic biomarkers to support the diagnosis of atypical bacterial pathogens [59].

In this study, we aimed to further characterize the differential metabolic host response associated with distinct CAP-associated pathogens, which may support understanding the role of metabolic changes in the pathogenesis associated with specific pathogens. To this end, we quantified 347 unique metabolites in 119 serum samples taken at hospital admission from patients with CAP and studied their association with the pathogens found in these patients.

3.2 Materials and methods

3.2.1 Study population

We analyzed previously collected serum samples obtained from 119 hospitalized patients with CAP, obtained as part of two previously conducted clinical studies executed between October 2004 and September 2010 and collected within 24 hours after hospital admission [29, 30]. For all patients, microbial disease etiology was confirmed using conventional diagnostic methods such as culturing, PCR, and urinary antigen tests. Inflammatory markers such as CRP and leukocyte counts were also available. Most patients (80%) were diagnosed with bacterial infections and the sample size per pathogen varied between 3 to 48 samples. Pathogen groups, a group of patients in which the same pathogen was found, including only one patient were excluded. None of the included patients had received treatment with corticosteroids. The patient distribution per pathogen showed no significant differences in sex and pneumonia severity index (PSI) score. However, the mean age differed significantly between pathogen groups ($p < 0.05$). All patients that participated in the previous clinical studies provided informed consent for the secondary use of their materials. Data were handled in an anonymized

way. An overview of the patient characteristics is provided in Table 3.1 and additional information is available in Table 3.2.

Table 3.1 Patient characteristics per CAP-associated pathogen.

Patient characteristic	<i>S. pneumoniae</i> (n=48)	<i>L. pneumophila</i> (n=18)	<i>C. burnetii</i> (n=17)	<i>C. psittaci</i> (n=7)	<i>M. pneumoniae</i> (n=5)	Influenza virus (n=11)	HS virus (n=6)	RS virus (n=4)	Parainfluenza virus (n=3)
Age (years) *									
Mean (SD)	62.2 (18.9)	62.9 (12.5)	47.4 (10.0)	62.6 (14.5)	38.8 (10.4)	67.0 (15.9)	71.3 (18.6)	69.3 (25.9)	83.3 (5.77)
Sex									
Male	22 (45.8%)	13 (72.2%)	14 (82.4%)	5 (71.4%)	2 (40.0%)	9 (81.8%)	5 (83.3%)	2 (50.0%)	2 (66.7%)
PSI score									
<50	9 (18.8%)	0 (0%)	5 (29.4%)	1 (14.3%)	3 (60.0%)	1 (9.1%)	0 (0%)	1 (25.0%)	0 (0%)
51-70	7 (14.6%)	3 (16.7%)	9 (52.9%)	0 (0%)	1 (20.0%)	3 (27.3%)	1 (16.7%)	0 (0%)	0 (0%)
71-90	5 (10.4%)	6 (33.3%)	1 (5.9%)	2 (28.6%)	1 (20.0%)	1 (9.1%)	1 (16.7%)	1 (25.0%)	0 (0%)
91-130	23 (47.9%)	7 (38.9%)	1 (5.9%)	4 (57.1%)	0 (0%)	6 (54.5%)	2 (33.3%)	1 (25.0%)	2 (66.7%)
131>	4 (8.3%)	2 (11.1%)	1 (5.9%)	0 (0%)	0 (0%)	0 (0%)	2 (33.3%)	1 (25.0%)	1 (33.3%)
Duration of symptoms before admission *									
Mean (SD)	4.06 (3.03)	5.11 (1.23)	5.06 (1.71)	4.14 (2.04)	13.4 (16.1)	4.45 (2.34)	5.17 (4.45)	6.50 (5.07)	4.00 (2.65)
CRP upon admission									
Mean (SD)	265 (166)	306 (90.8)	233 (115)	272 (110)	145 (86.0)	207 (87.9)	271 (121)	289 (135)	220 (142)
Leukocyte count upon admission*									
Mean (SD)	16.9 (5.99)	13.0 (4.24)	9.49 (1.56)	10.7 (5.03)	10.5 (3.29)	10.1 (4.93)	15.3 (4.89)	14.2 (7.94)	13.7 (9.30)

* Significant difference between pathogens (p -value < 0.05). Abbreviations: HS: herpes simplex; RS: respiratory syncytial; PSI: pneumonia severity index; CRP: C-reactive protein; SD: standard deviation.

3.2.2 Bioanalytical procedures

Metabolomic profiling was performed by the Biomedical Metabolomics Facility of Leiden University (BMFL), Leiden, The Netherlands. Details of the bio-analytical procedures are described elsewhere [59]. Briefly, the serum samples were quantitatively analyzed using five liquid- and one gas- chromatography methods coupled to mass spectrometry detectors, with a total coverage of 596 metabolite targets from 25 biochemical metabolite classes such as organic acids, amino acids, biogenic amines, acylcarnitines, and lipids. Following data integration and quality control steps, 374 unique metabolites were reported and underwent statistical analysis.

3.2.3 Data analysis

The metabolite levels were log-transformed and standardized prior to statistical analysis. Biological and biochemical relevant sums and ratios of metabolites (Table 3.3) were calculated from the raw metabolite data and were added as variables to the dataset, undergoing the same analysis together with metabolites.

Exploratory data analysis was performed using principal component analysis (PCA). To identify differences in metabolite levels between pathogens, we performed analysis of variance (ANOVA) with a false discovery rate (FDR) multiple testing correction. P-adjusted significant metabolites ($q < 0.1$) were tested with Tukey's posthoc test to identify which pathogen groups the metabolite was able to distinguish between. In addition to the significance threshold, we applied a cutoff of $>20\%$ fold change between the medians of the raw metabolite levels of each group.

All data analysis was performed using R. The scripts used for the data analysis are available on GitHub at <http://github.com/vanhasseltlab/MetabolicHostResponseToCAP>.

3.3 Results and discussion

3.3.1 Metabolic markers differentiate between pathogens

The metabolomics analysis of the patient serum samples yielded 347 quantifiable unique metabolites from different biochemical metabolite classes. No visible separation between samples obtained from patients infected by the various pathogens was apparent using PCA (Figure 3.6). Systematic analysis of individual metabolites for a specific pathogenic species yielded 64 unique metabolites that were significantly altered (ANOVA, $q < 0.1$ and $FC > 20\%$) as compared to other pathogenic species (Figure 3.1-3.5). All significant results are shown together in Figure 3.8 and Table 3.4. Large variation in group size for specific pathogenic species ($n=3-48$) and large within pathogen species group variation was present (Figure 3.7), limiting the ability to identify associations between metabolites and specific pathogenic species. Nonetheless, although often not statistically significant, trends between groups of metabolites and specific pathogenic

species were apparent, which is why we have maintained these in visualization for each metabolic class (Figure 3.1-3.4). The discriminating metabolites belong to different biochemical classes. The results are reported by biochemical class to allow focused biochemical discussion. A comprehensive qualitative overview of the main associations between pathogen species and metabolites identified has been summarized in Figure 3.5.

3.3.2 Carnitine and acylcarnitines

We identified significant differences in carnitines between serum samples of CAP patients that carried the pathogens *C. burnetii*, herpes simplex virus (HSV), or *M. pneumoniae* compared to CAP patients with other pathogens (Figure 3.1). The most consistent and statistically significant differences were found in patients with *C. burnetii* compared to patients with *L. pneumophila* or *S. pneumoniae* infections. Patients with *C. burnetii* showed 60-80% elevated long-chain acylcarnitines (LCAC), leading to a 2-fold lower ratio of free carnitine/total acylcarnitines. Changes in acylcarnitines can be related to mitochondrial energy production shifts between glycolysis and beta-oxidation of long-chain fatty acids [60, 61]. Naturally, such shifts depend on the glucose stores and the availability of carnitine as a limiting factor.

We observed increased levels of carnitine and short-chain acylcarnitines (SCAC) for *L. pneumophila* vs. *C. burnetii* and *M. pneumoniae*. Elevated carnitine and short-chain acylcarnitines suggest successful fatty acid beta-oxidation. Conversely, we also saw elevated mid-chain acylcarnitines (MCAC), most pronounced in patients with herpes simplex virus versus *C. burnetii*. MCAC could indicate incomplete beta-oxidation of long-chain fatty acids and may be related to mitochondrial dysfunction in the diseased state [62]. In contrast, LCAC were elevated mainly in patients with *C. burnetii* infection, suggesting a low rate of beta-oxidation rather than an incomplete process. It is unknown how well the levels of circulating acylcarnitines reflect the cellular levels, however, accumulation of long-chain acylcarnitines in lung cells indicated a defective beta-oxidation process, increasing oxidative stress and potentially leading to epithelial cell death [63]. Moreover, LCAC accumulation in the air-fluid interface of the lung can reduce lung function due to the inhibition of pulmonary surfactant [64].

It has been reported that impaired utilization of acylcarnitines for energy production can be caused by various viruses, including influenza and respiratory syncytial virus [63], although the latter two did not show any significant results in our study.

Overall, for acylcarnitines high within-pathogen group variance was present (Figure 3.7), which limited identifying further metabolite-pathogen associations. Acylcarnitine levels are associated with food consumption, certain medications such as valproic acid, and carnitine supplementation [65, 66, 67].




Figure 3.1 Fold changes of carnitine and acylcarnitines as median values per pathogen group compared to CAP patients with other pathogens. Significant differences ($q < 0.1$ and $FC > 20\%$) are marked with a green star.

3.3.3 Organic acids, amino acids, and related compounds

Significant differences in serum amino acids and derivatives were found in all groups of patients infected by CAP-associated pathogens (Figure 3.2). Several significantly different metabolites were only found in infections by specific pathogens. For example, patients with *L. pneumophila* had elevated kynurenine (1.5-1.7 fold) and phenylalanine (1.4-1.6 fold), while *M. pneumoniae* showed elevated S-methylcysteine (1.3-1.7 fold). Also, glutamate was elevated in *C. burnetii* compared to *L. pneumophila* (1.8 fold). Most of the stronger findings can be related to inflammation and oxidative stress and were associated with negative health outcomes in studies of various infections.

Elevated kynurenine was observed in patients with CAP and sepsis [68, 69], and severe COVID-19 [56, 70, 71]. Kynurenine increases due to enhanced IDO (Indoleamine-2,3-dioxygenase) conversion of tryptophan during the inflammatory response. IDO is expressed in the lungs and blood cells, and its activity and expression are induced by pro-inflammatory cytokines, viral proteins, and lipopolysaccharides [72, 73, 74]. Kynurenine acts as an immune suppressor via binding to the aryl-hydrocarbon receptor (AHR) [75], and also inhibits nitric oxide synthase (NOS), thereby increasing oxidative stress. Kynurenine also contributes to vasodilation and hypotension in sepsis [58, 76] and increases endothelial activation which leads to impaired microvascular reactivity [68, 77].

Elevated phenylalanine was associated with immune activation and negative outcome in patients with COVID-19 [55, 56], post-trauma, or sepsis [57, 58]. A possible explanation for elevated phenylalanine in the inflammatory environment is ROS-induced oxidation of the cofactor tetrahydrobiopterin (BH4) which is essential for the conversion




Figure 3.2 Fold changes of organic acids, amino acids, and derivatives as median values per pathogen group compared to CAP patients with other pathogens. Significant differences ($q < 0.1$ and $FC > 20\%$) are marked with a green star.

of phenylalanine to tyrosine and the downstream catecholamines by the enzyme phenylalanine 4-hydroxylase [57, 78].

Glutamate, a neurotransmitter in the CNS, also modulates the activity of immune cells via the NMDA receptor, and excessive production was related to damaged endothelial cells [79, 80, 81].

We found elevated uracil in patients with influenza and herpes simplex virus compared to patients with *C. psittaci* (1.2-1.9 fold). Patients with *S. pneumoniae* and *L. pneumophila* showed higher 3-hydroxyisobutyric acid (1.5-1.6 fold) and lower isoleucine (1.4 fold lower) compared to *C. burnetii*.

3.3.4 Signaling lipids

The signaling lipids covered in the applied platforms included oxylipins, endocannabinoids, their precursor fatty acids, and also cortisol. Metabolites that showed significant results were included in Figure 3.3.

Although we did not find significant differences in fatty acids in this study, the oxylipin profile of patients with herpes simplex virus was unique in comparison to other infections, showing significantly higher (all $FC > 1.8$) prostaglandin F2 alpha (PGF2a), 9,10,13-TriHOME (TriHOME), and thromboxane B2 (TXB2). Oxylipins are immune modulators produced by the peroxidation of C20 or C22 polyunsaturated fatty acids (PUFA). The omega-6 PUFA oxylipins exhibit mainly pro-inflammatory action

during the acute phase of inflammation. Some of these oxylipins also aid in the transition from innate to adaptive immune response. The arachidonic acid-produced PGF₂-alpha is a direct metabolite of PGE₂ and its increase aligns with the existing evidence that prostaglandin synthesis reactivates the herpes virus and increases its spread [82]. Although fluctuating along the course of inflammation, low levels of prostanoids (PGE₂ and TXB₂) are characteristic of sepsis non-survivors [83, 84] and COVID-19 [85]. TriHOMEs were reported to be dysregulated in respiratory diseases such as asthma and COPD [86, 87].

Of the covered endocannabinoids, several fatty acyl ethanolamides (EAs) were significantly different between patients with various infections. Docosatetraenoyl ethanolamide (DEA), which is the ethanolamide of the omega-6 adrenic acid (22:4) docosatetraenoic acid, was elevated in patients with *C. psittaci* compared to patients with *S. pneumoniae* or influenza virus (1.3-1.4 fold). Patients with the influenza virus showed the lowest levels of both DEA and dihomo-gamma-linolenoyl ethanolamide (DGLEA), the ethanolamide of the omega-6 (20:3) dihomo-gamma-linolenic acid. In comparison to other pathogens, patients with herpes simplex virus showed trends of elevated palmitoleoyl ethanolamide (POEA) and alpha-linolenoyl ethanolamide (aLEA), which are the ethanolamides of the omega-7 (16:1) palmitoleic acid, and the omega-3 (18:3) alpha-linolenic acid, respectively. A 2-fold significant elevation of POEA was measured in comparison to patients with *C. burnetii* and a 1.4-fold aLEA elevation in comparison to *M. pneumoniae*. Endocannabinoids do not only act as CNS modulators but are also peripheral immune mediators. Their pro- or anti-inflammatory activity varies and depends on their acyl group, the type of cells (immune; endothelial, etc.), and the receptors they bind to. In times of increased demand, for example during an acute immune response, and depending on diet and fat reserves, increased levels of endocannabinoids may reflect accelerated catabolism of cellular lipids, to serve as precursors for free fatty acids and oxylipins. It is also possible to see a shift towards the synthesis of fatty acyl EAs from fatty acids that occurs without medication [88], or due to treatment with corticosteroids and NSAIDs, that inhibit PLA₂ and COX enzymes [89, 90]. The last significant signaling lipid, cortisol, was lower in patients with *M. pneumoniae* compared to all other pathogens (2-3.3 fold lower), and lower for *C. burnetii* patients in comparison to *S. pneumoniae* and *L. pneumophila* patients (1.6 fold lower, Figure 3.8). Reduced excretion of cortisol indicates lower activation of the hypothalamic-pituitary-adrenal (HPA) axis in response to stress and innate immune activation following any infection. Cortisol stimulates gluconeogenesis and reduces inflammation, therefore its levels also affect (directly or indirectly) other metabolites measured in this study. In multiple studies on infections of the lower respiratory system, higher cortisol levels at hospital admission were predictive of mortality [91], indicating its relation to acute inflammation and disease severity.

3.3.5 Phospholipids and derivatives

Phospholipids form cell membranes and are hydrolyzed into lysophospholipids and long-chain fatty acids by various phospholipase A₂ (PLA₂) enzymes. The host's innate




Figure 3.3 Fold changes of oxylipin, endocannabinoid, and cortisol as median values per pathogen group compared to CAP patients with other pathogens. Significant differences ($q < 0.1$ and $FC > 20\%$) are marked with a green star.

immune response to infection, as well as the pathogen itself, can induce the activity of PLA2 [92]. Patients with *S. pneumoniae* infection showed the lowest levels of any lysophospholipids (LPCs) compared to other pathogens, alongside the highest levels of intact phosphatidylcholines (PCs), suggesting reduced activity of PLA2. Such a profile was also characteristic of sepsis [91, 93], possibly due to the varying degree of expression and activities of different PLA2 enzymes. Many LPCs and the total LPC/PC ratio were significantly higher in *C. burnetii* compared to *S. pneumoniae*, *L. pneumophila*, and RS virus (1.5-4 fold, Figure 3.4). Bacteria can use host lipids as building blocks for bacterial membrane formation and as an energy source. Also, host lipids are crucial for the entry of bacteria, viruses or toxins into cells. For example, *C. burnetii* has been found to use phospholipase A to integrate lipids into its membrane. [94]. *L. pneumophila* increased its virulence by secretion of PLA2 that destructed alveolar surfactant phospholipids [95]. Conversely, many intact PCs showed significantly lower levels in patients with the influenza virus in comparison to those with *S. pneumoniae*. The M protein of viruses like the influenza virus can interact with lipids and phosphatidylcholine vesicles allowing membrane fusion [96, 97, 98, 99].

Regarding lysophosphatidyls with serine or inositol functional groups (LPS and LPI), patients with parainfluenza virus showed 3-5 fold higher LPS (22:6) compared to *S. pneumoniae*, *C. psittaci*, and herpes simplex virus. Patients with *L. pneumophila* infection showed 30-90% higher LPIs compared to *S. pneumoniae*, influenza, and herpes simplex virus.




Figure 3.4 Fold changes of phospholipids including (lyso)phosphatidylcholines median values per pathogen group as median values per pathogen group compared to CAP patients with other pathogens. Significant differences ($q < 0.1$ and $FC > 20\%$) are marked with a green star.


3.3.6 Summary of metabolic findings per pathogen

The consolidation of the obtained results associates specific bacterial and viral CAP infections with metabolic perturbations related to inflammation, oxidative stress, lipids utilization, and energy production (Figure 3.5). Patients with *S. pneumoniae* infection showed high levels of the stress hormone cortisol that can be excreted as an HPA axis response to pro-inflammatory cytokines and hypoxia. Moreover, patients with *S. pneumoniae* infections consistently showed the highest PCs in comparison to all other pathogen groups, and very low lysophospholipids, suggesting lower activation of PLA2 enzymes, as described before in studies on sepsis [58, 91]. The lowest levels of LPCs were found in patients with *L. pneumophila* infection, who also showed the highest levels of cortisol, kynurenine, and phenylalanine, all characteristic of acute inflammatory response. Possibly indicating signs of oxidative stress and potentially also mitochondrial dysfunction [100], these patients had increased production of energy via beta-oxidation of fatty acids, evidenced by the elevated end-products of free carnitine and short-chain acylcarnitines. In comparison, patients with *C. burnetii* infection showed inhibition of fatty acid beta-oxidation, suggested by high levels of long-chain acylcarnitines, which are associated with increased ROS. These patients also had elevated glutamate accompanied by the highest levels of lysophospholipids, all can be linked to either inflammation, oxidative stress, or endothelial dysfunction [80, 81, 91, 98, 101]. Patients with *C. psittaci* infection possibly suffered from metabolic defects causing incomplete beta-oxidation of long-chain fatty acids, suggested by the elevated mid-chain fatty acids. This interrupted metabolism may be related to their relatively high levels of omega-6 fatty acyl ethanolamides, and high kynurenine, indicating acute inflammation. Despite the small sample size, patients with herpes simplex infection showed unique disruption in lipid metabolism. This included accumulation of long- and medium-chain acylcarnitines, elevated non-omega-6 fatty acyl ethanolamides, and very high levels of certain oxylipins, that possibly support the spread of the infection.

3.3.7 Study limitations and future perspectives

In this study, the sample size of patient groups with different CAP-associated pathogenic species was unbalanced and in several cases low (Table 3.1). Furthermore, a large within-group variance of metabolite levels was observed, which may impact our findings, especially in small sample size pathogen groups. Finally, the mean age of patients was unbalanced across the cohort, possibly acting as an additional confounder, since many aspects of metabolism change with age [102]. Overall, these factors limit the power to detect statistically significant metabolite-pathogen associations and require careful consideration of the relationships identified and further validation in a larger and more balanced study cohort. Nonetheless, several of the identified relationships are in line with expectations and other literature reports, supporting the biological relevance of our analysis.

Our study only considered metabolite levels at a single time point, even though patients will enter the clinical at different stages of their infection. The use of longitudinal



metabolite levels measured in patients represents an important next step to help to correct for such differences, and, to elucidate the relationship between changes in metabolite levels and biochemical or cellular biomarkers for immunological and/or inflammatory response markers, e.g. C-reactive protein levels, cytokines, and leukocyte cell counts [12].

Importantly, we have made available the full raw dataset underlying this study, which could be of interest for further bioinformatics analyses of the immune-metabolic responses reported [103].

3.4 Conclusion

In this study, we identified metabolites associated with the host response to specific CAP-associated pathogens, which is of relevance to further elucidate the pathogenesis of specific pathogens. We identified 64 metabolites that were significantly different between several pathogens. Specific bacterial and viral pathogens could be associated with metabolic perturbations that can be related to inflammation, oxidative stress, lipids utilization, and energy production. Further research should be conducted to validate our results and to further unravel the mechanisms of the metabolic host response to different pathogens causing CAP.

3.5 Supplementary data




Figure 3.6 Principal component analysis (PCA) plot of all pathogens. The percentages denote the percentage of variance explained by each principal component. Data were log-transformed and standardized prior to analysis.




Figure 3.7 Box-scatter plots of selected metabolites from different biochemical classes, that showed significant differences between pathogen patient groups. The box shows the lower quartile, the median, and the upper quartile of the metabolite ratios for each pathogen, with the whiskers denoting the minimum and maximum metabolite ratios within the interquartile range. Individual samples are shown as grey dots. Different letters above each bar represent significant differences between the groups.




Figure 3.8 Fold change of metabolite levels for all comparisons and all biochemical classes. Fold change values are shown only for metabolites with a significance of $q < 0.1$ & $FC > 20\%$.

Table 3.2 Extended patient characteristics

This table is available as pdf file (Table A.1) on the website of the publisher at
<https://doi.org/10.1016/j.metop.2023.100239>

Table 3.3 Metabolite ratios and sums

Metabolite sum or ratio name in R	Metabolite sum or ratio formula
BCAA_sum	isoleucine + leucine + valine
TCA_cycle_sum	Citric acid + lactic acid + malic acid + fumaric acid
urea_cycle_sum	Citrulline + arginine + ornithine + fumaric acid
lc_Carnitines_sum	Myristoylcarnitine + Hexadecenoylcarnitine + Palmitoylcarnitine + Stearoylcarnitine + Dodecenoylcarnitine + Tetradecenoylcarnitine + Linoleylcarnitine + Oleoylcarnitine + Tetradecadienylcarnitine
mc_Carnitines_sum	Hexanoylcarnitine + Octanoylcarnitine + Octenoylcarnitine + Decanoylcarnitine + Lauroylcarnitine + Nonacylcarnitine + Pimelylcarnitine + Decenoylcarnitine
sc_Carnitines_sum	Acetylcarnitine + Propionylcarnitine + Isobutyrylcarnitine + Butyrylcarnitine + Tiglylcarnitine + Methylbutyrylcarnitine + Isovalerylcarnitine
Cer_sum	Cer(d18:1/22:1) + Cer. (d18:1/24:1. + Cer(d18:1/24:0) + Cer(d18:1/16:0) + Cer(d18:1/23:0) + Cer(d18:1/24:0)
SM_sum	Sphingomyelin (d18:1/14:0) + (d18:1/15:0) + (d18:1/16:0) + (d18:1/16:1) + (d18:1/17:0) + (d18:1/18:0) + (d18:1/18:1) + (d18:1/18:2) + (d18:1/20:0) + (d18:1/20:1) + (d18:1/21:0) + (d18:1/22:0) + (d18:1/22:1) + (d18:1/23:0) + (d18:1/ 23:1) + (d18:0/24:0) + (d18:0/24:1) + (d18:0/24:2) + (d18:0/25:0) + (d18:0/25:1)
LPC_sum	Lysophosphatidylcholine (14:0) + (16:0) + (16:1) + (18:0) + (18:1) + (18:2) + (18:3) + (20:4) + (20:5) + (22:6) + (O-16:1) + (O-18:1)
PC_sum	Diacyl-phosphatidylcholine (32:0) + (32:1) + (32:2) + (34:1) + (34:2) + (34:3) + (34:4) + (36:1) + (36:2) + (36:3) + (36:4) + (36:5) + (36:6) + (38:2) + (38:3) + (38:4) + (38:5) + (38:6) + (38:7) + (40:4) + (40:5) + (40:6) + (40:7) + (40:8) + (O-34:1) + (O-34:2) + (O-34:3) + (O-36:2) + (O-36:3) + (O-36:4) + (O-36:5) + (O-36:6) + (O-38:4) + (O-38:5) + (O-38:6) + (O-38:7) + (O-40:6) + (O-42:6) + (O-44:5)
HT5_Trp_ratio	Serotonine / Tryptophan
ADMA_Arg_ratio	ADMA / Arginine
SDMA_Arg_ratio	SDMA / Arginine
Carnitine_sum_lc_Carnitines_ratio	Carnitine / LCAC sum
Carnitine_sum_mc_Carnitines_ratio	Carnitine / MCAC sum
Carnitine_sum_sc_Carnitines_ratio	Carnitine / SCAC sum
DCA_CA_ratio	DCA / CA
FA_14.1_14.0	FA (14:1) / FA (14:0)
FA_16.1_16.0	FA (16:1) / FA(16:0)
Gln_Glu	Glutamine / Glutamic acid
Kyn_Trp	Kynurenine / Tryptophan
sum_BCAA_sum_Phe_Tyr_ratio	BCAA sum / (Phenylalanine + Tyrosine)
sum_CER_sum_SM_ratio	Cer sum / SM sum
sum_LPC_sum_PC_ratio	LPC sum / PC sum

Table 3.4 Significant metabolites for the host response to different CAP pathogens (q < 0.1, FC > 20%)

This table is available as Excel file (Table A.3) on the website of the publisher at
<https://doi.org/10.1016/j.metop.2023.100239>

