



Universiteit  
Leiden  
The Netherlands

## Attitudes of Dutch intensive care unit clinicians towards oxygen therapy

Grim, C.C.A.; Cornet, A.D.; Kroner, A.; Meinders, A.J.; Brouwers, A.J.B.W.; Reidinga, A.C.; ... ; ATONIC Study Grp

### Citation

Grim, C. C. A., Cornet, A. D., Kroner, A., Meinders, A. J., Brouwers, A. J. B. W., Reidinga, A. C., ... Ruijter, W. de. (2020). Attitudes of Dutch intensive care unit clinicians towards oxygen therapy. *Netherlands Journal Of Medicine*, 78(4), 167-174. Retrieved from <https://hdl.handle.net/1887/3182574>

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3182574>

**Note:** To cite this publication please use the final published version (if applicable).

# Attitudes of Dutch intensive care unit clinicians towards oxygen therapy

C.C.A. Grim<sup>1\*</sup>

Leiden University Medical Centre, Leiden, the Netherlands.

\*Corresponding author: C.C.A.Grim@lumc.nl

<sup>1</sup>On behalf of the members of the ATONIC study group: A.D. Cornet, Medisch Spectrum Twente, Enschede; A. Kröner, Gelre Hospital, Apeldoorn; A.-J. Meinders, St. Antonius Hospital; A.J.B.W. Brouwers, Franciscus Gasthuis & Vlietland, Rotterdam; A.C. Reidinga, Martini Hospital, Groningen; D.J. van Westerloo, Leiden University Medical Center, Leiden; D.C.J.J. Bergmans, Maastricht University Medical Center, Maastricht; D. Gommers, Erasmus University Medical Center; D. Versluis, Haaglanden Medical Center, Den Haag; D. Weller, Maasstad Hospital, Rotterdam; E. Christiaan Boerma, Medical Center Leeuwarden, Leeuwarden; E. van Driel, Alrijne Hospital, Leiderdorp; E. de Jonge, Leiden University Medical Center, Leiden; F.J. Schoonderbeek, Ikazia Hospital, Rotterdam; H.J.F. Helmerhorst, Leiden University Medical Center, Leiden; H.G. Jongasma-van Netten, Meander Medical Center, Amersfoort; J. Weenink, Spaarne Gasthuis, Haarlem; K.J. Woittiez, VieCuri Medical Center, Venlo; K.S. Simons, Jeroen Bosch Hospital, 's-Hertogenbosch; L. van Welie, Gelderland Valley Hospital; M. Petjak, Groene Hart Hospital, Gouda; M.J. Sigtermans, Diakonessenhuis, Utrecht; M. van der Woude, Zuyderland Medical Center, Geleen; O.L. Cremer, University Medical Center Utrecht, Utrecht; P. Bijlstra, Elisabeth-TweeSteden Hospital, Tilburg; P. van der Heiden, Reinier de Graaf Gasthuis, Delft; R.K.L. So, Albert Schweitzer Hospital, Dordrecht; R. Vink, Tergooi, Hilversum; T. Jansen, Haga Hospital, Den Haag; W. de Ruijter, Noordwest Ziekenhuisgroep, Alkmaar

## ABSTRACT

**Background:** Over the last decade, there has been an increasing awareness for the potential harm of the administration of too much oxygen. We aimed to describe self-reported attitudes towards oxygen therapy by clinicians from a large representative sample of intensive care units (ICUs) in the Netherlands.

**Methods:** In April 2019, 36 ICUs in the Netherlands were approached and asked to send out a questionnaire (59 questions) to their nursing and medical staff (ICU clinicians) eliciting self-reported behaviour and attitudes towards oxygen therapy in general and in specific ICU case scenarios.

**Results:** In total, 1361 ICU clinicians (71% nurses, 24% physicians) from 28 ICUs returned the questionnaire. Of responding ICU clinicians, 64% considered oxygen-induced lung injury to be a major concern. The majority of respondents considered a partial pressure

of oxygen ( $\text{PaO}_2$ ) of 6-10 kPa (45-75 mmHg) and an arterial saturation ( $\text{SaO}_2$ ) of 85-90% as acceptable for 15 minutes, and a  $\text{PaO}_2$  7-10 kPa (53-75 mmHg) and  $\text{SaO}_2$  90-95% as acceptable for 24-48 hours in an acute respiratory distress syndrome (ARDS) patient. In most case scenarios, respondents reported not to change the fraction of inspired oxygen ( $\text{FiO}_2$ ) if  $\text{SaO}_2$  was 90-95% or  $\text{PaO}_2$  was 12 kPa (90 mmHg).

**Conclusion:** A representative sample of ICU clinicians from the Netherlands were concerned about oxygen-induced lung injury, and reported that they preferred  $\text{PaO}_2$  and  $\text{SaO}_2$  targets in the lower physiological range and would adjust ventilation settings accordingly.

## KEYWORDS

Conservative oxygenation, intensive care unit, mechanical ventilation, oxygen therapy, questionnaire, survey

## INTRODUCTION

Optimal oxygen therapy in critical care remains a subject of debate. Almost all critically ill patients receive supplemental oxygen to prevent tissue hypoxia, but the appropriate oxygen dose remains unclear.<sup>1</sup> Recent guidelines recommend targeting an oxygen saturation of 94-98% and stopping oxygen therapy when saturation reaches 96%.<sup>2,3</sup> This recommendation is mainly based on observational studies suggesting harmful effects of liberal oxygen therapy.<sup>4-8</sup> The first randomised controlled trial performed in one Italian ICU found an absolute mortality reduction of 8.6% for critically ill patients when limiting oxygen supplementation titrated to conservative oxygen saturation targets.<sup>9</sup> However, a recent large multicentre randomised controlled trial found no difference in outcomes of adults undergoing conservative or usual oxygen therapy in the ICU, even though targeted oxygenation was frequently not achieved.<sup>10</sup>

In recent years, self-reported views of nurses and physicians with regards to oxygen therapy in critically ill patients have evolved towards a more restrictive approach. In the Netherlands, oxygen-induced lung injury was seen as a major concern.<sup>11</sup> However, in actual clinical practice, these concerns were not accommodated, and the majority of partial pressure of oxygen (PaO<sub>2</sub>) values recorded were higher than self-reported targets. Hereafter, conservative oxygenation targets were introduced into daily clinical practice.<sup>5</sup> A recent study showed an increase in the number of ICU clinicians concerned about oxygen-induced lung injury, which was also reflected in actual clinical practice.<sup>12</sup> However, these Dutch studies were performed in three ICUs actively involved in research focused on oxygen therapy for critically ill patients. It therefore remains unknown whether these beliefs are widely supported and whether the results of previous studies can be generalised to ICU clinicians across the Netherlands. In the present study, we aimed to describe self-reported attitudes towards oxygen therapy from a large sample of ICU clinicians across the Netherlands. We further aimed to assess if there were differences in attitudes between nurses and physicians, age categories, and type of ICU.

## MATERIALS AND METHODS

### Questionnaire

The questionnaire was a translated and comprehensive version of previously used surveys from Canada and Australia/New Zealand, and was also previously used in the Netherlands.<sup>11-14</sup> The anonymous online questionnaire consisted of 59 multiple-choice questions that were designed to elicit self-reported behaviour of ICU clinicians with respect to oxygen therapy in general

and in specific case scenarios (complete questionnaire in Dutch: Supplement 1)\*. It included questions about oxygen-induced lung injury, risks of mechanical ventilation, indices of tissue oxygenation, and arterial saturation (SaO<sub>2</sub>) and PaO<sub>2</sub> targets for short and long-time periods in an acute respiratory distress syndrome (ARDS) patient receiving mechanical ventilation. The last part of the questionnaire investigated whether the respondent would adjust a fraction of inspired oxygen (FiO<sub>2</sub>) of 50% for given SaO<sub>2</sub> and for given PaO<sub>2</sub> in the following case scenarios: ARDS, cardiac ischaemia, cerebral ischaemia, sepsis, percutaneous coronary intervention (PCI) stent, and untreatable anaemia.

### Target population

In April 2019, 36 ICUs in the Netherlands were approached with the request to send out a web-based questionnaire (LimeSurvey) to their nursing and medical staff. Subsequently, ICU clinicians were invited by email to complete the questionnaire. A reminder was sent out if deemed necessary. The invited ICUs included mixed medical and surgical adult ICUs in university and non-university hospitals across the Netherlands. The ethical reviewing board was informed and had no objection (G18.110).

### Statistical analysis

Questionnaire responses are presented as a proportion of respondents per question. Differences in questionnaire responses between nurses and physicians, university and non-university ICUs and age categories were analysed using the Chi square test of Fisher's exact test as appropriate. A p-value below 0.05 was considered statistically significant. Statistical analyses were conducted using R 1386 3.4.4.

## RESULTS

### Characteristics of questionnaire respondents

Between April and August 2019, 1361 questionnaire responses were received from ICU clinicians (61% completed all questions). ICU clinicians from 28 (78%) out of 36 invited ICUs in the Netherlands participated in the online questionnaire. These 28 participating ICUs comprised the majority of all available ICU beds in the Netherlands (428 of 831 beds, data from the 2018 Dutch National Intensive Care Evaluation, <https://stichting-nice.nl/datainbeeld/public>). Two hundred-twenty-six (19%) ICU clinicians that participated in the questionnaire were from four university hospitals. Respondents consisted of 847 (71%) nurses, 287 (24%) physicians (8% residents, 1% fellows and 15% intensivists), and 53 (5%) with another type of practice (e.g., ventilation practitioner, physician's

**Table 1.** Responses to questions regarding concerns, risks, and indices of oxygen therapy and oxygenation for critically ill patients.

Answers	Total n (%)	Nurses n (%)	Physicians n (%)	University n (%)	Non-university n (%)	> 40 years of age n (%)	< 40 years of age n (%)
<b>Question: Is oxygen-induced lung injury a concern when placing a patient on mechanical ventilation?</b>							
YES, a major concern							
due to the high incidence of injury	46 (4)	31 (4)	12 (4)	8 (4)	38 (4)	21 (3)	25 (5)
due to the severity of injury	434 (38)	335 (42)	79 (28)	77 (36)	357 (39)	232 (36)	202 (41)
due to the high incidence and severity of injury	248 (22)	181 (23)	55 (20)	56 (26)	192 (21)	135 (21)	113 (23)
YES, but not a major concern	358 (32)	224 (28)	121 (43)	69 (32)	289 (32)	226 (35)	131 (27)
NO, it is not a concern	49 (4)	34 (4)	14 (5)	7 (3)	42 (5)	29 (5)	20 (4)
Total number of respondents	1135	805	281	217	918	643	491
		p < 0.01		p = 0.51		p = 0.03	
<b>Question: In your opinion, which one of the following two situations poses a greater threat of lung injury for mechanically ventilated patients?</b>							
High FiO <sub>2</sub>	194 (17)	169 (20)	19 (7)	24 (11)	170 (18)	109 (17)	85 (17)
High tidal volumes and high ventilator pressures	953 (82)	649 (78)	261 (92)	198 (88)	755 (80)	531 (82)	420 (82)
Don't know	18 (2)	14 (2)	4 (1)	2 (1)	16 (2)	11 (2)	7 (1)
Total number of respondents	1165	832	284	224	941	651	512
		p < 0.01		p = 0.01		p = 0.90	
<b>Question: In situations when maximum SaO<sub>2</sub> achievable is low (± 85%) or when FiO<sub>2</sub> requirements are high, do you assess indices of tissue oxygenation?</b>							
NO	294 (28)	212 (30)	65 (23)	237 (28)	57 (28)	175 (29)	119 (27)
YES, lactate	629 (60)	424 (59)	185 (65)	505 (60)	124 (62)	350 (59)	279 (63)
YES, microcirculation with OPS/SDF imaging	12 (1)	9 (1)	1 (0)	11 (1)	1 (1)	7 (1)	5 (1)
YES, other	108 (10)	72 (10)	32 (11)	89 (11)	19 (10)	66 (11)	41 (9)
Total number of respondents	1043	717	283	201	842	598	444
		p = 0.09		p = 0.85		p = 0.54	
<b>Question: Independent of FiO<sub>2</sub>, after what duration would a stable SaO<sub>2</sub> of 85% begin to raise concerns?</b>							
< 2 hours	754 (75)	569 (81)	156 (61)	136 (66)	618 (78)	357 (76)	397 (75)
2-24 hours	184 (18)	108 (15)	66 (26)	42 (21)	142 (18)	82 (17)	102 (19)
24-48 hours	41 (4)	18 (3)	20 (8)	19 (9)	22 (3)	23 (5)	18 (3)
48-72 hours	11 (1)	7 (1)	4 (2)	6 (3)	5 (1)	2 (0)	9 (2)
> 72 hours	10 (1)	2 (0)	8 (3)	2 (1)	8 (1)	8 (2)	2 (0)
Total number of respondents	1000	704	254	205	795	472	528
		p < 0.01		p < 0.01		p = 0.04	
P-values represents difference in questionnaire responses in subgroups, tested with chi square or fisher exact as appropriate. Total number of respondents may differ per question because not all questions were answered by all participants. Number of respondents are shown, with percentages (n (%)). FiO <sub>2</sub> = fraction of inspired oxygen; OPS = orthogonal polarization spectral.; SaO <sub>2</sub> = arterial oxygen saturation; SDF = sidestream dark field; SvO <sub>2</sub> = mixed venous oxygen saturation.							

assistant). Of the respondents, 258 (22%) were 18-30 years of age, 374 (32%) 31-40 years of age, 239 (20%) 41-50 years of age, 259 (22%) was 51-60 years of age, and 55 (5%) 61-70 years of age.


**Questionnaire responses**

The responses to questions regarding concerns when placing a patient on mechanical ventilation are listed in table 1. A majority (64%) of respondents considered oxygen-induced lung injury a major concern when initiating mechanical ventilation; 17% of respondents reported that high FiO<sub>2</sub> posed a greater threat of lung injury than high tidal volumes and high ventilator pressures. Significantly more ICU nurses than physicians considered oxygen-induced lung injury a major concern during mechanical ventilation; 20% of the nurses reported that high FiO<sub>2</sub> posed a greater threat of lung injury than high tidal volumes and high ventilator pressures,

compared with 7% of physicians (p < 0.01). Compared to nurses, significantly fewer physicians would be concerned with a stable SaO<sub>2</sub> of 85% within two hours (61% physicians vs. 81% nurses, p < 0.01). More clinicians from university ICUs responded that high tidal volumes and high ventilator pressures posed a greater threat of lung injury compared to clinicians from non-university ICUs (p = 0.01). Compared to university ICU clinicians, more clinicians from non-university ICUs reported they would begin to raise concern after a shorter duration of time with a stable SaO<sub>2</sub> of 85% (p < 0.01). ICU clinicians of younger than 40 years of age more often reported they were majorly concerned with oxygen induced lung injury compared to older clinicians (p = 0.03).

Figure 1 shows the percentage of respondents accepting various oxygenation ranges in a young to middle-aged mechanically ventilated patient with ARDS. More

**Figure 1.** Self-reported acceptable SaO<sub>2</sub> and PaO<sub>2</sub> ranges for 15 minutes and 24-48 hours. Bars represent percentage of respondents. In the questionnaire, respondents were presented with a case of a young-to-middle-aged ARDS patient requiring mechanical ventilation. Ventilator settings are optimised with respect to PaO<sub>2</sub>/FiO<sub>2</sub> ratio and haemodynamic indices. There is no evidence to indicate end organ ischaemia, and haemodynamics are stable.



FiO<sub>2</sub> = fraction of inspired oxygen; PaO<sub>2</sub> = partial pressure of oxygen; SaO<sub>2</sub> = arterial oxygen saturation

**Table 2.** Questionnaire responses to questions regarding FiO<sub>2</sub> response for case scenarios with given SaO<sub>2</sub> and PaO<sub>2</sub>.

		FiO <sub>2</sub> response			FiO <sub>2</sub> response		
		Higher n (%)	No change n (%)	Lower n (%)	Higher n (%)	No change n (%)	Lower n (%)
		ARDS			Sepsis		
SaO <sub>2</sub>	80-85%	919 (95)	51 (5)	1 (0)	872 (98)	12 (1)	3 (0)
	85-90%	536 (56)	417 (43)	9 (1)	764 (86)	118 (13)	2 (0)
	90-95%	9 (1)	760 (79)	189 (20)	103 (12)	740 (84)	40 (5)
	95-100%	2 (0)	107 (11)	865 (89)	6 (1)	257 (29)	629 (71)
PaO <sub>2</sub>	6 kPa	791 (97)	26 (3)	0	785 (99)	7 (1)	1 (0)
	9 kPa	175 (22)	610 (75)	31 (4)	316 (40)	462 (58)	13 (2)
	12 kPa	6 (1)	281 (35)	528 (65)	16 (2)	434 (55)	340 (43)
	16 kPa	2 (0)	18 (2)	801 (98)	1 (0)	45 (6)	752 (94)
		Cardiac ischaemia			PCI stent		
SaO <sub>2</sub>	80-85%	934 (99)	7 (1)	1 (0)	869 (99)	8 (1)	0
	85-90%	874 (94)	53 (6)	0	813 (93)	61 (7)	0
	90-95%	310 (33)	601 (64)	23 (3)	139 (16)	707 (81)	26 (3)
	95-100%	15 (2)	528 (56)	396 (42)	4 (1)	299 (34)	578 (66)
PaO <sub>2</sub>	6 kPa	793 (99)	7 (1)	0	785 (99)	9 (1)	1 (0)
	9 kPa	488 (61)	307 (38)	7 (1)	417 (53)	368 (46)	9 (1)
	12 kPa	31 (4)	512 (64)	255 (32)	19 (2)	471 (60)	301 (38)
	16 kPa	3 (0)	95 (12)	706 (88)	2 (0.3)	51 (6)	746 (93)
		Cerebral ischaemia			Untreatable anaemia		
SaO <sub>2</sub>	80-85%	903 (99)	12 (1)	2 (0.2)	809 (93)	62 (7)	1 (0)
	85-90%	829 (92)	72 (8)	4 (0.4)	705 (81)	154 (18)	7 (1)
	90-95%	129 (14)	743 (82)	30 (3)	413 (48)	378 (44)	63 (7)
	95-100%	6 (1)	314 (34)	595 (65)	56 (6)	509 (58)	308 (35)
PaO <sub>2</sub>	6 kPa	775 (98)	12 (2)	2 (0.3)	759 (96)	32 (4)	1 (0.1)
	9 kPa	345 (44)	431 (55)	12 (2)	524 (66)	260 (33)	12 (2)
	12 kPa	16 (0)	452 (57)	320 (41)	125 (16)	485 (61)	185 (23)
	16 kPa	3 (0)	52 (6)	740 (93)	16 (2)	197 (25)	584 (73)

All clinical situations represent patients in the ICU, who have been invasively mechanically ventilated for at least 5 days, with FiO<sub>2</sub> set at 50%. ARDS: patient with acute respiratory distress syndrome and pneumonia; cardiac ischaemia: patient with signs of cardiac ischaemia (ST-depressions in het anterior leads (max 3 mm) and pneumonia; cerebral ischaemia: patient with recent cerebral ischaemia and one-side hemiplegia; sepsis: patient with liver abscess and sepsis; untreatable anaemia: Jehovah's Witness with stable haemoglobin of 1.8 mmol/l after gastric bleeding; Higher: increase FiO<sub>2</sub> higher than current 50%; no change maintain FiO<sub>2</sub> at current 50%; Lower: decrease FiO<sub>2</sub>, lower than current 50%. Number of respondents are shown, with percentages (n (%)).  
FiO<sub>2</sub> = fraction of inspired oxygen; PaO<sub>2</sub> = partial pressure of oxygen; SaO<sub>2</sub> = arterial oxygen saturation

physicians reported that lower SaO<sub>2</sub> and PaO<sub>2</sub> were acceptable, compared to nurses (supplementary table 1\*, p < 0.01). More university ICU clinicians chose lower SaO<sub>2</sub> ranges as lowest acceptable for under 15 minutes and 24-48 hours, compared to non-university clinicians

(p < 0.01). Older ICU clinicians reported lower PaO<sub>2</sub> as acceptable for up to 15 minutes (p = 0.01), compared to younger clinicians.

The proportions of ICU clinicians adjusting FiO<sub>2</sub> levels in different case scenarios are listed in table 2. For the case

scenarios, most respondents reported they would increase  $\text{FiO}_2$  if  $\text{SaO}_2$  was 80-85% or if  $\text{PaO}_2$  was 6 kPa (45 mmHg). If  $\text{PaO}_2$  was 16 kPa (120 mmHg), most respondents ( $\geq 73\%$  of respondents per case scenario) reported they would lower  $\text{FiO}_2$ . The majority of ICU clinicians reported in most case scenarios, that they would leave  $\text{FiO}_2$  unchanged if  $\text{SaO}_2$  was 90-95% or if  $\text{PaO}_2$  was 9 or 12 kPa (68 or 90 mmHg). Respondents favoured lower  $\text{SaO}_2$  and  $\text{PaO}_2$  levels for the ARDS cases. Overall, nurses, more frequently than physicians, would increase  $\text{FiO}_2$  if  $\text{SaO}_2$  or  $\text{PaO}_2$  was lower. Physicians more often responded that they would decrease  $\text{FiO}_2$  or leave it unchanged if  $\text{SaO}_2$  or  $\text{PaO}_2$  was lower, except for the untreatable anaemia case scenarios (supplementary table 2). Minor differences were found between university and non-university clinicians and between ICU clinicians < 40 years and > 40 years of age (supplementary tables 3 and 4).

## DISCUSSION

This national questionnaire study assessed clinicians working in the majority of ICUs across the Netherlands and showed that ICU clinicians consider oxygen-induced lung injury a major concern; high ventilator pressures and high tidal volumes were considered a greater threat than high  $\text{FiO}_2$ . For shorter periods of time, ICU clinicians accepted  $\text{SaO}_2$  levels as low as 85% and  $\text{PaO}_2$  levels as low as 6 kPa (45 mmHg), but a higher limit of 90% and 7 kPa (53 mmHg) is preferred if the situation lasts longer. It seems that ICU clinicians consider a  $\text{PaO}_2$  of 6 kPa (45 mmHg) as too low, a  $\text{PaO}_2$  of 16 kPa (120 mmHg) as too high, and a  $\text{PaO}_2$  of 12 kPa (90 mmHg) as optimal, because they reported adjusting ventilation settings accordingly. In general, compared to nurses, physicians had a more open attitude towards conservative oxygen therapy and would allow lower  $\text{SaO}_2$  and  $\text{PaO}_2$  levels without adjusting  $\text{FiO}_2$ . Older and university ICU clinicians would accept lower oxygenation, possibly due to more experience or more awareness of the potential adverse effects.

To our knowledge, this is the first nation-wide study describing self-reported attitudes towards oxygen therapy of ICU clinicians working in the 28 ICUs consisting of the majority of available beds and admissions in the Netherlands. Previous studies have focused on a selection of ICUs nationally or internationally.<sup>11,13,15,16</sup> Perhaps, these ICUs were more oxygen-focused and this could have influenced the questionnaire results. We believe our results are a better reflection of the general attitudes of ICU clinicians towards oxygen therapy in daily critical care. The first study assessing attitudes of intensivists was performed in Canada in 1999 and found that 51% of respondents considered oxygen-induced lung injury a major concern.<sup>14</sup> In our cohort of ICU clinicians, 64%

reported oxygen-induced lung injury to be important, which was similar to results of a survey in 2013 of 90 ICU nurses and physicians in Australia.<sup>15</sup> However, in 2010, 542 critical care nurses from Australia and New Zealand were surveyed and only 22% considered oxygen-induced lung injury a major concern.<sup>16</sup> In our cohort, nearly 70% of ICU nurses considered it a significant concern. Apparently, the number of clinicians considering oxygen-induced lung injury in daily practice has increased over the years. This is most likely due to increasing evidence about oxygen-induced lung injury and the effect of conservative oxygen therapy on patient outcomes. However, apart from a time-dependent effect, it may also reflect a geographical difference or difference by chance. In a recent Dutch questionnaire study, performed after the implementation of a conservative oxygenation protocol in three ICUs, 76% of respondents considered oxygen-induced lung injury to be a major concern<sup>12</sup>, which was an increase of nearly 20% compared to the assessment before the implementation,<sup>11</sup> and 10% more than what we found in this study. This supports our hypothesis that in ICUs where oxygen-related research is conducted clinicians are more concerned about oxygen-induced lung injury than in other ICUs.

Seventeen percent of the current respondents considered high  $\text{FiO}_2$  to be a greater threat of lung injury than high tidal volumes and ventilator pressures, which was similar to 13% previously found.<sup>13,14</sup> This is remarkable, as evidence for lung-injury by high tidal volumes is well accepted,<sup>17</sup> while evidence for the risks of high  $\text{FiO}_2$  in ICU patients is still controversial.<sup>9,10</sup> Compared to an earlier study,<sup>16</sup> we found that fewer nurses considered barotrauma to be a greater threat than high  $\text{FiO}_2$ , compared to physicians. Possibly, physicians may be more often convinced by evidence of ventilator-induced lung injury due to high tidal volumes and pressures.<sup>18-20</sup>

The preferred  $\text{PaO}_2$  ranges for short and long time periods in an ARDS patient reported by the current respondents were comparable to the  $\text{PaO}_2$  range previously recommended in studies in ARDS patients by the ARDS Clinical Trials Network.<sup>21</sup> These findings were similar to an earlier survey performed in critical care physicians from seven northern European countries where the majority chose a  $\text{PaO}_2$  of 10 kPa (75 mmHg) for an ARDS patient.<sup>22</sup> Our questionnaire results also suggest that physicians tolerate lower  $\text{PaO}_2$  and  $\text{SaO}_2$  values than nurses. Physicians may be more comfortable with lower oxygenation and with taking actions and accountability with the risk of hypoxia. Physicians may also be better informed about the potential downsides of supplemental oxygen therapy. In a previous survey of nurses, more experienced nurses were more likely to answer that they would never be concerned with a stable  $\text{SpO}_2$  of 90%.<sup>16</sup>

Our study has the following clinical implications. As this study reflects the beliefs and attitudes of a representative

sample of ICU clinicians from the Netherlands, the results may be useful to customise training, clinical decision making, and protocols. Furthermore, our study provides data for determining PaO<sub>2</sub> and SaO<sub>2</sub> targets in future interventional or observational studies. The current study also gives insight into the differences in attitudes of clinicians. These differences in attitudes could be explained by variances in education and training and by barriers experienced by the clinicians. When implementing new oxygenation strategies in daily critical care, it is important to acknowledge these differences of attitudes and actively engage and educate all clinicians to improve team compliance.

The primary strength of our study is its size and thereby its representativeness. The participation rate of 78% of the invited ICUs was high. In addition, participating ICUs consisted of more than half of the available ICU beds and more than half of all ICU admissions in the Netherlands in 2018. Moreover, the distribution of nurses and physicians was representative of a typical staff constitution of ICUs in the Netherlands. The questionnaire strongly resembles questionnaires previously used in other studies, allowing for comparisons and exploring trends over time and continents.<sup>11,13,14</sup> Furthermore, our results show what clinicians think about oxygen therapy in critical care and for specific pathologies, which could be helpful when reviewing the impact of guidelines for critical care or specific pathologies.

Our study has some limitations. The questionnaire data is self-reported and may not reflect actual practice and does not reflect the practice of non-responders. It has been shown that self-reported attitudes towards oxygen therapy are generally more conservative than actual practice.<sup>11</sup> The cases included in the survey do not represent the complexity of patients in daily practice. SaO<sub>2</sub> and PaO<sub>2</sub> ranges and values in the survey were chosen arbitrarily. Because this was an explorative study, we chose not to correct for multiple testing.

In conclusion, our study provides new insights into the attitudes of ICU clinicians towards oxygen therapy across the Netherlands. The majority of ICU clinicians reported concern about oxygen-induced lung injury and preferred PaO<sub>2</sub> and SaO<sub>2</sub> targets in the lower physiological range. Physicians reported being more conservative with oxygen therapy and decreased FiO<sub>2</sub> at lower SaO<sub>2</sub> and PaO<sub>2</sub> values, compared to nurses.

## DISCLOSURES

All authors declare no conflicts of interest. No funding or financial support was received.

*\*The supplementary information (Dutch questionnaire / S-tables 1-4) is available upon request; please contact the corresponding author.*

## REFERENCES

- Girardis M, Alhazzani W, Rasmussen BS. What's new in oxygen therapy? *Intensive Care Med.* 2019;45:1009-11.
- Siemieniuk RAC, Chu DK, Kim LH-Y, et al. Oxygen therapy for acutely ill medical patients: a clinical practice guideline. *BMJ.* 2018;363:k4169.
- O'Driscoll BR, Howard LS, Earis J, Mak V. British Thoracic Society Guideline for oxygen use in adults in healthcare and emergency settings. *BMJ Open Respir Res.* 2017;4:e000170.
- Helmerhorst HJ, Arts DL, Schultz MJ, et al. Metrics of Arterial Hyperoxia and Associated Outcomes in Critical Care. *Crit Care Med.* 2017;45:187-95.
- Helmerhorst HJ, Schultz MJ, van der Voort PH, et al. Effectiveness and Clinical Outcomes of a Two-Step Implementation of Conservative Oxygenation Targets in Critically Ill Patients: A Before and After Trial. *Crit Care Med.* 2016;44:554-63.
- Damiani E, Adrario E, Girardis M, et al. Arterial hyperoxia and mortality in critically ill patients: a systematic review and meta-analysis. *Crit Care.* 2014;18:711.
- de Jonge E, Peelen L, Keijzers PJ, et al. Association between administered oxygen, arterial partial oxygen pressure and mortality in mechanically ventilated intensive care unit patients. *Crit Care.* 2008;12:R156-R.
- Chu DK, Kim LHY, Young PJ, et al. Mortality and morbidity in acutely ill adults treated with liberal versus conservative oxygen therapy (IOTA): a systematic review and meta-analysis. *Lancet.* 2018;391:1693-705.
- Girardis M, Busani S, Damiani E, et al. Effect of Conservative vs Conventional Oxygen Therapy on Mortality Among Patients in an Intensive Care Unit: The Oxygen-ICU Randomized Clinical Trial. *JAMA.* 2016;316:1583-9.
- ICU-ROX Investigators and the Australian and New Zealand Intensive Care Society Clinical Trials Group, Mackle D, Bellomo R, et al. Conservative Oxygen Therapy during Mechanical Ventilation in the ICU. *N Engl J Med.* 2020;382(11):989-98.
- Helmerhorst HJ, Schultz MJ, van der Voort PH, et al. Self-reported attitudes versus actual practice of oxygen therapy by ICU physicians and nurses. *Ann Intensive Care.* 2014;4:23.
- Grim CC, Helmerhorst HJ, Schultz MJ, et al. Changes in Attitudes and Actual Practice of Oxygen Therapy in ICUs after Implementation of a Conservative Oxygenation Guideline [published online ahead of print, 2020 Mar 24]. *Respir Care.* 2020;respcare.07527.
- Eastwood GM, Reade MC, Peck L, Jones D, Bellomo R. Intensivists' opinion and self-reported practice of oxygen therapy. *Anaesth Intensive Care.* 2011;39:122-6.
- Mao C, Wong DT, Slutsky AS, Kavanagh BP. A quantitative assessment of how Canadian intensivists believe they utilize oxygen in the intensive care unit. *Crit Care Med.* 1999;27:2806-11.
- Eastwood GM, Peck L, Young H, Suzuki S, Garcia M, Bellomo R. Intensive care clinicians' opinion of conservative oxygen therapy (SpO<sub>2</sub> 90-92%) for mechanically ventilated patients. *Aust Crit Care.* 2014;27:120-5.
- Eastwood GM, Reade MC, Peck L, Baldwin I, Considine J, Bellomo R. Critical care nurses' opinion and self-reported practice of oxygen therapy: a survey. *Aust Crit Care.* 2012;25:23-30.
- Acute Respiratory Distress Syndrome N, Brower RG, Matthay MA, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med.* 2000;342:1301-8.
- Putensen C, Theuerkauf N, Zinserling J, Wrigge H, Pelosi P. Meta-analysis: ventilation strategies and outcomes of the acute respiratory distress syndrome and acute lung injury. *Ann Intern Med.* 2009;151:566-76.



19. Neto AS, Simonis FD, Barbas CS, et al. Lung-Protective Ventilation With Low Tidal Volumes and the Occurrence of Pulmonary Complications in Patients Without Acute Respiratory Distress Syndrome: A Systematic Review and Individual Patient Data Analysis. *Crit Care Med.* 2015;43:2155-63.
20. Serpa Neto A, Simonis FD, Barbas CS, et al. Association between tidal volume size, duration of ventilation, and sedation needs in patients without acute respiratory distress syndrome: an individual patient data meta-analysis. *Intens Care Med* 2014;40:950-7.
21. Brower RG, Lanken PN, MacIntyre N, et al. Higher versus lower positive end-expiratory pressures in patients with the acute respiratory distress syndrome. *N Engl J Med.* 2004;351:327-36.
22. Schjorring OL, Toft-Petersen AP, Kusk KH, et al. Intensive care doctors' preferences for arterial oxygen tension levels in mechanically ventilated patients. *Acta Anaesth Scand.* 2018;62:1443-51.