



Universiteit
Leiden

The Netherlands

Early clinical and imaging features of rheumatoid arthritis: towards a more complete picture

Dijk, B.T. van

Citation

Dijk, B. T. van. (2024, September 24). *Early clinical and imaging features of rheumatoid arthritis: towards a more complete picture*. Retrieved from <https://hdl.handle.net/1887/4092968>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4092968>

Note: To cite this publication please use the final published version (if applicable).

Part I

Clinical characteristics
and early recognition
of arthritis



Chapter 2

Development and validation of a clinical rule for recognition of early inflammatory arthritis

Robin M. ten Brinck*¹, Bastiaan T. van Dijk*¹, Hanna W. van Steenbergen¹, Saskia le Cessie², Mattijs E. Numans³, Samantha L. Hider^{4,5} Christian Mallen⁴, Annette H.M. van der Helm-van Mil¹

* both contributed equally

¹ Department of Rheumatology, Leiden University Medical Centre, Leiden, The Netherlands.

² Department of Epidemiology and Statistics, Leiden University Medical Centre, Leiden, The Netherlands.

³ Department of Public Health and Primary Care, Leiden University Medical Centre, Leiden, The Netherlands.

⁴ Research Institute for Primary Care and Health Sciences, Keele University, Newcastle, United Kingdom.

⁵ Haywood Academic Rheumatology Centre, Haywood Hospital, Stoke-on-Trent, Staffordshire, United Kingdom.

BMJ Open. 2019 Feb 22;8(11):e023552.

ABSTRACT

Objectives. National and international guidelines recommend prompt referral of patients presenting with inflammatory arthritis (IA), but general practitioners (GPs) feel uncertain in their proficiency to detect synovitis through joint examination, the method of choice to identify IA. Our objective was to develop and validate a rule composed of clinical characteristics to assist GPs and other physicians in identifying IA when in doubt.

Design. Split-sample derivation and validation study.

Setting. The Leiden Early Arthritis *Recognition* Clinic (EARC); a screening clinic for patients in whom GPs suspected but were unsure of the presence of IA.

Participants. 1,288 consecutive patients visiting the EARC.

Primary and secondary outcome measures. Associations of clinical characteristics with presence of IA were determined using logistic regression in 644 patients, after which the results were validated in the other 644 patients (split-sample validation). To facilitate application in clinical practice, a simplified rule (with scores ranging 0 to 7.5) was derived and validated.

Results. IA was identified by a rheumatologist in 41% of patients. In univariable analysis, male gender, age ≥ 60 years, symptom duration < 6 weeks, morning stiffness > 60 minutes, a low number of painful joints (1-3 joints), presence of patient-reported joint swelling, and difficulty with making a fist were associated with IA in the derivation dataset. Using multivariable analysis, a simplified rule consisting of these seven items was derived and validated, yielding an Area Under the Receiver Operator Characteristic curve (AUC) of 0.74 (95%CI 0.70-0.78) in the derivation dataset. Validation yielded an AUC of 0.71 (95%CI 0.67-0.75). Finally, the model was repeated to study predicted probabilities with a lower prevalence of inflammatory arthritis to simulate performance in primary care settings.

Conclusions. Our rule, composed of clinical parameters, had reasonable discriminative ability for IA and could assist physicians in decision-making in patients with suspected IA, increasing appropriateness of healthcare utilisation.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- A clinical rule could help to select patients to refer for additional investigations (laboratory or imaging) or to secondary care. This could promote early identification of inflammatory arthritis and increase appropriateness of healthcare utilisation.
- Data were collected prospectively in a population of patients in whom general practitioners had doubt on the presence of inflammatory arthritis.
- The main limitation is that data were not collected in primary care itself, but in a setting intermediary between primary and secondary care. Further external validation in GP settings is therefore required.

BACKGROUND

Early initiation of disease modifying anti-rheumatic drugs is strongly associated with improved outcomes of rheumatoid arthritis (RA).[1] National and international guidelines attempt to facilitate this by emphasising prompt referral of patients presenting with inflammatory arthritis (IA) to a rheumatologist. The European League Against Rheumatism (EULAR) taskforce for the management of early IA recommends referral within 6 weeks of the onset of symptoms[2], while in the United Kingdom (UK) the National Institute for Health and Care Excellence (NICE) guidelines advise referral to a rheumatologist in patients with new, persistent (>3–4 weeks) synovitis within three working days.[3] However, it was demonstrated that this referral timeline is achieved in only 17% of patients.[4] On average, RA patients are seen four (and sometimes more than eight) times by general practitioners (GPs) before they refer to secondary care[5–8], which may reflect the difficulty of differentiating patients with early IA from patients with other types of common musculoskeletal symptoms. A recent qualitative study revealed that GPs acknowledge the importance of early detection and referral, but feel uncertain in their proficiency to detect synovitis through joint examination, the method of choice to identify IA.[2,9] As a consequence, referral to a rheumatologist may be delayed, which contributes to overall treatment delay in early RA as observed in Europe.[10,11]

This is further complicated by the high incidence of consultations for various common musculoskeletal symptoms and the low incidence of early IA in primary care.[12] The consultation prevalence of any musculoskeletal symptom in primary care in the UK approximates 2405 per 10,000 per year [13], making it the most common organ system consulted for at GP practices.[12–14] Although musculoskeletal symptoms are common, GPs suspect IA (based on pattern recognition) in only a very small minority of patients.[5] In these patients, GPs often lack confidence in joint assessment for synovitis.

To support early detection, several initiatives have been developed, including triage systems. The best studied triage system (the Early Inflammatory Arthritis Questionnaire) was developed and validated for patients attending secondary and tertiary care.[15-17] Furthermore, several referral guidelines for GPs[6,18-22], and public awareness campaigns have been developed, for instance one attempting to simplify pattern recognition to the “S-Factor”: Stiffness, Swelling, Squeezing. However, none of these initiatives were designed using primary care data, and all assume that GPs can differentiate between the presence and absence of joint swelling[6,18-20], which continues to be a barrier to the early detection of IA.

Altogether there is a contradiction with the need to refer as quickly as possible while evidence who must be referred or, in line with this, in whom additional investigations are appropriate, is lacking. To solve the issue, we have developed and validated a rule composed of clinical characteristics, by taking advantage of data from a setting intermediate between primary and secondary care. This intermediate setting of the Early Arthritis *Recognition* Clinic was a local solution to promote early referrals and may not be easily implemented in other regions. The clinical rule derived from these data, however, is easy to apply and may assist in the decision-making process in patients with musculoskeletal symptoms with suspected IA at other places, in order to promote early identification of IA.

METHODS

Study population

To promote early recognition of early IA, the Early Arthritis *Recognition* Clinic (EARC) was initiated in September 2010 in Leiden, the Netherlands. The outpatient clinic of the department of Rheumatology of the Leiden University Medical Centre (LUMC) is the only referral centre in a healthcare region of ~400,000 people. GPs were instructed to refer patients to the EARC in whom they were unsure about the presence of IA (instead of a ‘wait-and-see’ approach or performing additional tests). The EARC system has reduced referral delay from 8 to 2 weeks, and improved early identification of IA.[11,23] To emphasise the importance of early identification of IA and aiming to inform on the purpose of the EARC, a region-wide educational campaign was conducted among regional GPs.

In addition to (and distinct from) the EARC, the LUMC also has an Early Arthritis Clinic (EAC). The EAC was established in 1993 to include and follow patients with early arthritis and to offer the possibility of rapid access to rheumatology care, usually

within a week of referral. To differentiate between the clinics, GPs were instructed to refer to the EAC if there was a clear synovitis or very high suspicion of IA (i.e. to continue as they had before, since there was no benefit for such patients to go the EARC first) and to refer to the EARC when in doubt about the presence of IA (i.e. to not 'wait-and-see' or order additional tests). Thus, patients included in this study represent the difficult group in whom GPs were uncertain of the presence of suspected IA; patients with a very high degree of suspicion were referred directly to the EAC.

The EARC screening clinic was held twice a week between 2010–2014 and once a week from 2014 onwards. After GP referral, patients can visit the EARC without an appointment. All patients who visited the EARC between 2010 and September 2015 were studied.

Data collection

At the EARC, patients completed a short questionnaire about their joint symptoms, after which they were seen by an experienced rheumatologist (AvdHvM or other senior rheumatologists) who performed a full 66-joint examination. If synovitis was determined by physical examination, patients were fast-tracked to visit the EAC within 1 week for further evaluation and treatment. Patients without IA were discharged to primary care. The questionnaire completed by patients, provided in Online Appendix S1, contained questions on age, gender, date of symptom onset, date of first visit to GP, presence of a (sub)acute symptom onset (versus a gradual symptom-onset), morning stiffness (duration in minutes), which part of the day symptoms were worst, and whether they had difficulty with making a fist. Patients were asked to indicate on a 52-joint mannequin which joints were painful and which joints they considered to be swollen. IA, defined as synovitis confirmed by the rheumatologist at physical examination, was used as outcome.

Collected data was anonymised and entered in a research database at chronological order of visiting the EARC. The local medical ethical committee approved this study.

Derivation and validation of the model

We used half of the dataset for derivation and the other half for validation of results (split-sample validation). To prevent bias by (unknown) effects of inclusion period, patients with odd ID-numbers (1, 3, etc) were included in the derivation dataset and those with even ID-numbers (2, 4, etc) were used for validation.

To prevent exclusion of patients with one or more missing variables, we imputed missing values using chained equations[24]; frequencies of missing variables are presented in Online Appendix S2. The variables 'difficulty with making a fist' and 'self-reported joint

swelling' were most frequently missing as these were added to the questionnaire after April 1st 2012, thus absence of these data was considered to occur completely at random.

We conducted logistic regression analysis modelling with presence of IA (defined as synovitis on physical examination by the rheumatologist) as dependent variable. Continuous variables were categorised using clinically relevant cut-offs: age: <40 / 40–59.9 / ≥60 years; duration of symptoms: <6 / 6–11 / 12–51.9 / ≥52 weeks; duration of morning stiffness: ≤60 / >60 minutes; number of painful joints: 0 / 1–3 / 4–10 / ≥11; number of swollen joints: 0 / 1–3 / 4–10 / ≥11. We performed univariable logistic regression to evaluate associations between dependent variables and presence of IA. Variables with p-values <0.05 in univariable analyses were entered in multivariable regression analyses to obtain a model with a small number of variables. If several categories within a variable had similar regression coefficients in multivariable modelling, we pooled these categories and repeated the analysis. In sub-analysis, we also performed a multivariable logistic regression model with the pooled categories using backward selection.

To obtain a simplified rule applicable in daily care, we rounded the regression coefficients of the final multivariable logistic regression model to the nearest 0.5 (irrespective of p-value). This resulted in an easily calculable risk score. For each value of the risk score, we determined test characteristics (i.e. sensitivity and specificity) and predicted probabilities of the presence of inflammatory arthritis.

We evaluated the overall discriminative ability of the models using the Area Under the Receiver Operating Characteristic curve (AUC). The model's calibration was assessed by generating a calibration plot to measure goodness of fit, where the data was partitioned in 10 equally sized groups based on the predicted probabilities using the final fitted multivariable model. In each group, the average predicted probability on current IA was compared with the observed prevalence, both in the derivation and validation dataset. Additionally, the Hosmer-Lemeshow statistic was calculated.

To estimate performance of our simplified rule in a setting with a different prevalence of IA (e.g. primary care), a simulation was performed. Accurate data on prevalence of IA in GP practices is lacking, and therefore an estimation was made based on previous literature. One study revealed that 27% assigned with the International Classification of Primary Care-1 code for suspected IA in their medical record had confirmed RA (n=38), polyarthritis (n=5), or oligoarthritis (n=8) following rheumatologist's assessment. Another study among GPs found that 18% of patients with suspected IA was referred; though data on rheumatologists' diagnoses were not provided.[25] Guided by these scarce data obtained in GP practices, performance of the model was simulated with an

estimated prevalence of 20%.^[5] The intercept of the regression model was adjusted as described by King et al.^[26,27] and we plotted average estimated predicted probabilities against the regression and simplified risk score.

Statistical analyses were performed using Statistical Package for the Social Sciences (SPSS, version 23.0). P-values <0.05 were considered significant.

Patient involvement

Patient research partners agreed with the pathway of care at the EARC. They also provided feedback on the questionnaire, which was expanded in 2012 with two questions.

RESULTS

Patients

1,288 patients in whom GPs were unsure about the presence of IA visited the EARC between 2010 and 2015; of these patients, 41% had synovitis at joint examination. The frequency of inflammatory arthritis was stable throughout the study years (Online Appendix S3). Baseline characteristics of patients in both the derivation and validation dataset are presented in Table 1.

Table 1. Characteristics of patients visiting the Early Arthritis Recognition Clinic

	Derivation (N=644)	Validation (N=644)	P-value**
Male, n (%)	190 (30)	198 (31)	0.62
Age in years, mean ± SD	52 ± 16	51 ± 17	0.27
Symptom duration in weeks, median (IQR)	10 (3–45)	12 (4–45)	0.18
Acute onset of symptoms*, n (%)	252 (39)	238 (37)	0.45
Symptoms worst in the early morning, n (%)	372 (58)	351 (55)	0.10
Morning stiffness in minutes, median (IQR)	10 (0–30)	10 (0–30)	0.33
Number of painful joints, median (IQR)	7 (2–15)	6 (3–15)	0.69
Number of patient-reported swollen joints, median (IQR)	2 (1–5)	2 (1–5)	0.19
Difficulty with making a fist, n (%)	329 (51)	301 (47)	0.06
Arthritis present at joint examination by experienced rheumatologist, n (%)	271 (42)	252 (39)	0.28

* Patients were asked to define onset of symptoms; either acute onset of symptoms or gradual onset of symptoms, see Online Appendix S1.

** Unpaired t-tests, chi-squared tests and Mann-Whitney U tests were used as appropriate. Abbreviations: IQR = interquartile range; SD = standard deviation.

Model derivation

In univariable analyses, male gender, age ≥ 60 years, symptom duration of < 6 weeks, an acute onset of symptoms, morning stiffness > 60 minutes, a low number of painful joints (1–3 joints), presence of patient-reported joint swelling (1–3 joints), and difficulty with making a fist were associated with the presence of IA in the derivation dataset (Table 2). ‘Symptoms worst in the early morning’ was not associated with IA and therefore not included in multivariable analysis. Two multivariable models were created with categorised variables; first a model with categories similar to the univariable analysis (Table 3, model 1), and secondly a model pooling categories per variable with similar regression coefficients (Table 3, model 2). Performing this second model in the derivation dataset revealed that male gender, age ≥ 60 years, symptom duration of < 6 weeks, a low number of painful joints (1–3 joints), and presence of patient-reported joint swelling were independently associated with the presence of IA (Table 3). The AUC of model 2 was 0.75 (95%CI 0.70–0.79) in the derivation dataset. In sub-analysis, model 2 was repeated with a backward selection procedure, showing similar regression coefficients (Online Appendix S4).

Generation of a simplified rule

In order to facilitate usage in routine clinical practice, a simplified model was generated (Online Appendix S5). The obtained regression coefficient of ‘acute onset of symptoms’ in multivariable modelling was -0.015 , yielding 0 points. Also after exclusion of this variable, the regression coefficients of the other seven variables in the model did not change yielding similar points. This resulted in a simplified rule consisting of seven scored items and a total score ranging from 0 to 7.5 with corresponding predicted risks (Figure 1). Risks of IA predicted by the model as a function of the regression score (i.e. the sum of the regression coefficients times the value of the corresponding covariates) are presented in Figure 2A; as shown, simplification did not majorly affect the predicted risks. The calibration plot shows that predicted probabilities correlated well with the observed proportions of patients with IA (Online Appendix S6). The Hosmer-Lemeshow test for the derivation dataset yielded a P-value of 0.36. If cut-offs are required and a highly sensitive approach is preferred ($> 90\%$ sensitivity), this is obtained by a cut-off score of ≥ 4 . When a highly specific approach is preferred ($> 90\%$ specificity), this is obtained by a cut-off score of ≥ 6 . Test characteristics for all cut-off points are presented in Online Appendix S7. The AUC of the simplified score, measuring discrimination, was 0.74 (95%CI 0.70–0.78; Online Appendix S8).

Validation

The final multivariable model (model 2) was applied in the validation dataset, revealing similar results (Table 3). The AUC was 0.72 (95%CI 0.68–0.77). Figure 2A shows the predicted probabilities of the simplified rule are almost similar to those obtained in the derivation data. The AUC of the simplified rule was 0.71 (95%CI 0.67–0.75) in the validation dataset. The calibration plot is shown in Online Appendix S6, the Hosmer-Lemeshow test for the validation dataset yielded a P-value of 0.43.

Table 2. Univariable logistic regression in the derivation dataset with presence of synovitis upon joint examination as outcome

		Arthritis (N=271)	No arthritis (N=373)	OR (95% CI)
Male, n (%)		104 (38)	86 (23)	2.1 (1.5–2.9)
Age, n (%)	<40	49 (18)	104 (28)	(ref)
	40–59.9	109 (40)	172 (46)	1.3 (0.9–2.0)
	≥60	113 (42)	97 (26)	2.5 (1.6–3.8)
Symptom duration in weeks, n (%)	<6	124 (46)	103 (28)	3.8 (2.4–5.9)
	6–11	38 (14)	62 (17)	1.9 (1.1–3.9)
	12–51.9	66 (24)	75 (20)	2.7 (1.7–4.5)
	≥52	43 (16)	132 (36)	(ref)
Acute onset of symptoms *, n (%)		122 (45)	131 (35)	1.5 (1.1–2.1)
Symptoms worst in early morning, n (%)		158 (58)	214 (57)	1.1 (0.7–1.6)
Morning stiffness >60 min, n (%)		45 (17)	40 (11)	1.7 (1.03–2.7)
Number of painful joints, n (%)	0	1 (0)	10 (3)	(ref)
	1–3	110 (41)	82 (22)	13.2 (1.7–105.5)
	4–10	76 (28)	123 (33)	6.1 (0.8–49.0)
	≥11	84 (31)	158 (42)	5.2 (0.7–41.3)
Number of patient-reported swollen joints, n (%)	0	18 (7)	71 (19)	(ref)
	1–3	115 (42)	119 (32)	3.7 (2.0–6.9)
	4–10	87 (32)	115 (31)	2.9 (1.5–5.5)
	≥11	51 (19)	68 (18)	2.9 (1.4–5.9)
Difficulty with making a fist, n (%)		156 (58)	172 (46)	1.6 (1.1–2.4)

* Patients were asked to define onset of symptoms; either acute onset of symptoms or gradual onset of symptoms, see Online Appendix S1.

Abbreviations: CI = confidence interval; OR = odds ratio.

Table 3. Multivariable logistic regression analyses with synovitis upon joint examination as outcome
 Model 1 includes categories of clinically applicable cut-offs; if within variables several categories had similar regression coefficients, categories were pooled (Model 2).


	Model 1			Model 2		
	Derivation		Derivation		Validation	
	OR (95% CI)		OR (95% CI)	B	OR (95% CI)	
Male	1.7 (1.1–2.5)		1.7 (1.1–2.5)	0.517	1.7 (1.1–2.4)	
Age (years)	<40	(ref)	0–59.9	(ref)	(ref)	(ref)
	40–59.9	1.5 (0.96–2.5)	≥60	2.1 (1.4–3.1)	0.750	2.1 (1.5–3.0)
	≥60	2.9 (1.7–4.8)				
Symptom duration (weeks)	<6	3.8 (2.3–6.4)	<6	3.6 (2.2–6.0)	1.279	3.4 (2.0–5.7)
	6–11	1.7 (0.92–3.1)	6–51.9	2.2 (1.4–3.6)	0.797	1.9 (1.2–3.0)
	12–51.9	2.9 (1.7–5.0)	≥52	(ref)	(ref)	(ref)
	≥52	(ref)				
Acute onset of symptoms*	1.0 (0.67–1.5)		0.99 (0.66–1.5)	-0.015	1.0 (0.7–1.5)	
Morning stiffness (minutes)	>60	1.6 (0.88–2.9)	>60	1.6 (0.91–2.9)	0.485	1.2 (0.6–2.3)
Number of painful joints	0	(ref)	0	(ref)	(ref)	(ref)
	1–3	9.3 (1.1–78.2)	1–3	10.0 (1.2–83.4)	2.300	7.9 (0.9–68.6)
	4–10	4.5 (0.53–37.6)	≥4	4.5 (0.54–37.1)	1.497	5.2 (0.6–45.1)
	≥11	3.3 (0.39–28.4)				
Number of patient-reported swollen joints	0	(ref)	0	(ref)	(ref)	(ref)
	1–3	3.2 (1.6–6.4)	≥1	3.5 (1.9–6.6)	1.253	3.7 (1.9–7.0)
	4–10	3.4 (1.7–7.0)				
	≥11	4.3 (1.9–10.0)				
Difficulty with making a fist	1.6 (0.97–2.5)		1.6 (0.99–2.6)	0.467	1.4 (0.9–2.2)	
Intercept				-4.6		
AUC	0.76 (0.71–0.80)		0.75 (0.70–0.79)		0.72 (0.68–0.77)	

* Patients were asked to define onset of symptoms; either acute onset of symptoms or gradual onset of symptoms, see Online Appendix S1.

Variables with p-values <0.05 in univariable analysis in the derivation set were entered in multivariable regression analyses.


Abbreviations: AUC = Area under the Receiver Operating Characteristics curve; B = beta; CI = confidence interval; OR = odds ratio.

Figure 1. The Clinical Arthritis Rule (CARE) and corresponding predicted risks of the presence of inflammatory arthritis per score



Observed risks of current inflammatory arthritis were obtained by calculating the proportion of patients with a positive outcome (rheumatologist-confirmed synovitis) for each value of the risk score in the derivation dataset.

Figure 2. The Clinical Arthritis Rule (CARE) and presentation of the predicted probabilities of the presence of current inflammatory arthritis based on the regression model, and the simplified score as observed in the derivation and validation datasets (A), and estimated predicted probabilities in a simulation with a pre-test probability (i.e. prevalence) of inflammatory arthritis of 20% (B)



Predicted probabilities of the final multivariable logistic regression model, fitted in the derivation set as function of the regression score (i.e. the sum of the regression coefficients times the value of the corresponding covariates (green line)). Furthermore, for each value of the simplified score the mean predicted probability is plotted in the derivation and validation dataset (blue and orange dots).

Figure 3. A stylised representation of the Clinical Arthritis RuleE, to be used in patients in whom GPs doubt about the presence of inflammatory arthritis

The Clinical Arthritis Rule (CARE)

Welcome to the Clinical Arthritis Rule (CARE) calculator!

This calculator estimates the risks of the presence of inflammatory arthritis based on the research of Ten Brinck *et al.* More information on this calculator can be found at the bottom of the page. The rule is developed for use in patients in whom GPs or other physicians doubt about the presence of inflammatory arthritis. The calculator estimates the risk of the presence of synovitis, detectable at joint examination by experienced rheumatologists.

Gender:

- Male
- Female

Age

Symptom duration:

- < 6 weeks
- 6-51 weeks
- ≥ 52 weeks

Morning stiffness duration:

- ≤ 60 minutes
- > 60 minutes

Number of patient-reported painful joints:


- 1-3 joints
- ≥ 4 joints

Patient-reported joint swelling:

- Yes
- No

Difficulty with making a fist:

- Yes
- No



Calculate risk!

This calculator has the option to calculate the risks of current inflammatory arthritis in settings with different pre-test probabilities (i.e. prevalences). The calculator was derived and validated in patients in whom GPs suspected inflammatory arthritis and in whom inflammatory arthritis was confirmed in 41% of cases. Data obtained in (other) GP practices suggested that the pre-test risk in patient with suspected inflammatory arthritis is approximately 20%. Therefore the calculator can also estimate the risk on inflammatory arthritis in this setting.

The web application that provides predictions on the predicted risk of inflammatory arthritis for individual patients as can be accessed at <http://caretool.eu/>.

Simulation of accuracy in a setting with a lower prevalence of IA

In contrast to test characteristics, predicted probabilities depend on the prior risk (i.e. prevalence) of IA. The frequency of IA among primary care patients with GP-determined clinical suspicion of IA may be different than that observed in the EARC. Based on observations in GP practices[5,25], a simulation was run for the regression and simplified score with a prevalence of inflammatory arthritis set at 20%. Estimated predicted probabilities for different scores of the multivariable model and simplified rule (in derivation and validation datasets) are presented in Figure 2B.

Our simplified rule was implemented in a web application that provides predictions on the presence of current IA for individual patients; a screenshot is presented in Figure 3. The web application is accessible online at <http://caretool.eu/>.

DISCUSSION

GPs play a crucial role in early identification of RA and often lack confidence in detecting joint synovitis.[9] In an attempt to solve the contradiction between the need to refer very early and absence of evidence who must be referred, we provided an evidence-based and simple method to identify the presence of IA in patients in whom IA is suspected. This clinical rule helps to select patients to refer for additional investigations (laboratory or imaging) or to secondary care. Hence, the Clinical Arthritis Rule could increase appropriateness of healthcare utilisation.

This study is different from studies that derived tools to facilitate triage of patients who have been referred to secondary or tertiary care,[15-17] as our study did not aim to prioritise patients that are already referred. In addition, we aimed to facilitate recognition of IA (as this would necessitate prompt referral to a rheumatologist) and did not perform a longitudinal study to predict development of specific diagnoses (e.g. RA) later-on. This explains why several factors were found to be associated with presence of IA that are not generally considered typical for RA (male gender, a low number of painful joints, a short symptom duration). GPs generally do well in identifying those at high risk for development of RA (i.e. women with subacute smouldering polyarticular, symmetric complaints), and therefore with this tool we aimed to assist GPs in decision-making for more atypical or non-classical presentations of IA leading to doubt (e.g. due to overlap of symptoms with other diagnoses). Indeed, many of the patients who did not have synovitis at the EARC had symptoms due to diagnoses that are characterised by longstanding or extensive joint pain (e.g. osteoarthritis, fibromyalgia), explaining higher scores for a short symptom duration or a low number of symptomatic joints.

Adding other clinical variables might increase the discriminative ability of the model. Potential examples include the squeeze test of the metacarpophalangeal joints (although the diagnostic accuracy was shown to be only moderate),[28] information on family history, or functional impairments. These items were not routinely collected before December 2015. Adding data on laboratory investigations to our rule could potentially also increase its discriminative ability. However, our data do not permit us to evaluate this, as additional investigations were done afterwards and only in patients with synovitis at joint examination.

A strength of our EARC for the purpose of this study is that GPs in our region are familiar with the need for early referral and that regional healthcare logistics make rheumatological care rapidly available for patients with arthritis, with the EARC as ultimate service for patients in whom GPs suspect (but are unsure about) IA. With the availability of the EARC every week and lack of any waiting list for the EARC, we assume a low number of patients not showing up at the EARC despite being encouraged by their GP to visit the EARC. As the EARC serves as a unique bridge between primary and secondary care, its patients closely resemble the population GPs have contact with and have doubts about. Although the EARC is successful in our region[11,23], this approach may be more difficult to implement in other centres or regions due to a shortage of rheumatologists, or long traveling distances to rheumatology outpatient clinics, and as such a different system is needed to aid GPs in identifying IA. This prompted us to derive a validated rule composed of clinical characteristics that could assist GPs in decision-making for more atypical or non-classical (but nevertheless suspect) presentations of IA, as classical presentations usually don't cause GPs' concern.

GPs were discouraged (both by our local communication with GPs and according to national guidelines for GPs) to perform autoantibody testing.[29] Autoantibody testing in primary care in this region was infrequent[5], unlike in other parts of the world. Autoantibody testing may falsely reassure doctors and patients, especially when results are negative, and as such we believe a model based on clinical presentation is more appropriate to facilitate rapid referral.

Another strength is that we studied patients in whom the GPs have indicated a lack of confidence to identify the presence of synovitis. Patients with clinically obvious IA had early access to rheumatologic care already. This may enhance the generalisability of the present data to the setting of doubt in primary care. Furthermore, the use of real-life observational data in our study may boost external validity of the results.

A disadvantage of our setting is that the data were not collected in primary care itself, but in a setting intermediary between primary and secondary care. Although


musculoskeletal symptoms are a very common reason for consulting primary care, suspected IA is relatively unusual, and the average full-time GP diagnoses only one new patient with RA each year.[30] Additionally, although the EARC is easily accessible on a weekly basis, the exact number of patients who were referred but did not visit the EARC is unknown. Validation in primary care is required. We studied the difficult group of patients in whom GPs were uncertain of the presence of suspected IA. The prevalence of such patients in primary care may be higher and, as a consequence, the actual prevalence of IA among suspected IA patients may be lower than 41% in primary care. Since the post-test probabilities strongly depend on the prevalence (i.e. pre-test probability), a simulation was performed with an estimated prevalence of IA that was half the prevalence as observed in our data (20%). The choice of 20% was based on literature from primary care. Although not much is known about suspected IA in primary care, two studies suggested a prevalence of IA among suspected patients of 18 to 27%.[5, 25] We demonstrated the predictive accuracy of the model using a simulated prevalence of 20%. Because of the limitation that no other data are available on the prevalence of IA when GPs suspect IA, this estimated prevalence could be an overestimation. However, the observed data could also be an underestimation as in our setting GPs were instructed to refer patients with high suspicion/definite arthritis to the regular outpatient clinic. Further external validation in GP settings is therefore required.

GPs in our region are well informed about the importance of early detection of IA, but the GPs in our region feel that their actual detection skills are not different from that of GPs elsewhere. However if the detection skills of our GPs are different from that of GPs in other regions, a lower prevalence of IA (and therefore lower pre-test probabilities) may be present. As a consequence, the rule may yield lower post-test probabilities. This effect may have been dealt with in the simulation analysis, but still external validation in primary care and preferably in different regions or countries is necessary.

We expect that our rule (Clinical Arthritis Rule - CARE) might support GPs and other health care professionals in the decision-making process in patients with musculoskeletal symptoms in whom they suspect IA, regardless of the region. Of course, the consequences of an increased score will likely depend on the setting and relation with secondary care: it can either influence the decision to directly refer a patient or to first ask for additional laboratory tests (e.g. acute phase reactants or autoantibodies; Figure 4). A clinical decision aid may be of value to this end as well, as for most laboratory investigations the diagnostic accuracy depends on the prior risk. Using a simple clinical decision aid first may be more cost-effective than performing additional investigations in all patients in whom there is doubt about IA. Depending on the setting and consequences of a high score, either a sensitive method or a specific method may be

preferred; for this reason cut-offs for both situations are provided. The web application, also easily assessable by phone, facilitates implementation of the Clinical Arthritis RuleE by GPs, physicians, and other health care professionals such as physiotherapists in their daily work.

Figure 4. Flowchart of decision-making in patients with suspected early IA based on clinical characteristics and the role of the Clinical Arthritis RuleE



Abbreviations: GP = general practitioner; IA = inflammatory arthritis; CARE = Clinical Arthritis RuleE.

CONCLUSION

In conclusion, this study developed a clinical rule that supports the identification of patients suspected of having IA by physicians that feel insufficiently experienced in assessment of synovitis by joint examination. We hope the current data are a prelude to a data-driven method that supports GPs, physicians, and other health care professionals in decision-making in patients with suspected early IA.

REFERENCES

1. Van Nies JA, Krabben A, Schoones JW, *et al.* What is the evidence for the presence of a therapeutic window of opportunity in rheumatoid arthritis? A systematic literature review. *Ann Rheum Dis.* 2014 May;73(5):861-70.
2. Combe B, Landewé R, Daïen CI, *et al.* 2016 update of the EULAR recommendations for the management of early arthritis. *Ann Rheum Dis.* 2017 Jun;76(6):948-959.
3. NICE quality standard: Rheumatoid arthritis in over 16s [QS33] - Quality statement 1: Referral. [Internet] Available from: <https://www.nice.org.uk/guidance/qs33/chapter/Quality-statement-1-Referral>. [accessed on 10 May 2017].
4. Ledingham JM, British Society for Rheumatology National Audit Project. Importance of early detection and referral of rheumatoid arthritis. *BMJ.* 2016 Apr 18;353:i2172.
5. Newsum EC, de Waal MWM, van Steenberg HW, *et al.* How do general practitioners identify inflammatory arthritis?—A cohort analysis of Dutch general practitioner electronic medical records. *Rheumatology.* 2016 May;55(5):848-53.
6. Emery P, Breedveld FC, Dougados M, *et al.* Early referral recommendation for newly diagnosed rheumatoid arthritis: evidence based development of a clinical guide. *Ann Rheum Dis.* 2002 Apr;61(4):290-7.
7. Mankia K, Nam J, Emery P. Identifying arthralgia suspicious for progression to rheumatoid arthritis. *Ann Rheum Dis.* 2017 Jun;76(6):e14.
8. National Audit Office. Services for people with rheumatoid arthritis. [Internet] Available from: <http://www.nao.org.uk/report/services-for-people-with-rheumatoid-arthritis/>. [accessed on 10 May 2017].
9. Meyfroidt S, Stevens J, De Lepeleire J, *et al.* A general practice perspective on early rheumatoid arthritis management: A qualitative study from Flanders. *Eur J Gen Pract.* 2015;21(4):231-7.
10. Raza K, Stack R, Kumar K, *et al.* Delays in assessment of patients with rheumatoid arthritis: variations across Europe. *Ann Rheum Dis.* 2011 Oct;70(10):1822-5.
11. Van der Linden MP, Le Cessie S, Raza K, *et al.* Long-term impact of delay in assessment of patients with early arthritis. *Arthritis Rheum.* 2010 Dec;62(12):3537-46.
12. Nielen MM, Spronk I, Davids R. [Incidentie en prevalentie van gezondheidsproblemen in de Nederlandse huisartsenpraktijk in 2014]. From: NIVEL Zorgregistratie Eerste Lijn [Internet]. 2017; Available from: <https://www.nivel.nl/nl/NZR/incidenties-en-prevalenties> [accessed on 10 May 2017].
13. Jordan KP, Kadam UT, Hayward R, *et al.* Annual consultation prevalence of regional musculoskeletal problems in primary care: an observational study. *BMC Musculoskelet Disord.* 2010 Jul 2;11:144.
14. Jordan KP, Jöud A, Bergknot C, *et al.* International comparisons of the consultation prevalence of musculoskeletal conditions using population-based healthcare data from England and Sweden. *Ann Rheum Dis.* 2014 Jan;73(1):212-8.
15. Bell MJ, Tavares R, Guillemin F, *et al.* Development of a self-administered early inflammatory arthritis detection tool. *BMC Musculoskelet Disord.* 2010 Mar 17;11:50.
16. Tavares R, Huang S, Bykerk VP, Bell MJ. A parallel group cohort to determine the measurement properties of an early inflammatory arthritis detection tool. *Rheumatology.* 2013 Nov;52(11):2077-85.
17. Villeneuve E, Nam JL, Bell MJ, *et al.* A systematic literature review of strategies promoting early referral and reducing delays in the diagnosis and management of inflammatory arthritis. *Ann Rheum Dis.* 2013 Jan;72(1):13-22.
18. Croydon Health Services NHS Foundation Trust. Early Inflammatory Arthritis Clinic Referral Form. [Internet] Available from: https://www.croydonhealthservices.nhs.uk/Downloads/GP_resources/

- Referral%20and%20service%20information/Arthritis/Primary%20Care%20Referral%20form%20for%20Early%20Inflammatory%20Arthritis%20Clinic.pdf. [accessed on 10 May 2017].
19. Hampshire Hospitals NHS Foundation Trust. New Early Arthritis Referral form. [Internet] Available from: https://www.hampshirehospitals.nhs.uk/media/419347/early_arthritis_referral_additional_information.pdf. [accessed on 10 May 2017].
 20. Musgrove Park Hospital. Early Arthritis Referral Form. [Internet] Available from: <http://www.tsft.nhs.uk/media/353049/early-arthritis-final.pdf>. [accessed on 10 May 2017].
 21. Hülsemann JL, Mattussek S, Siegmund-Schultze E, Zeidler H. [Early diagnosis of arthritis and spondyloarthritis within the framework of integrated health care in Lower Saxony]. *Z Rheumatol*. 2006 Feb;65(1):70-4.
 22. Guy's and St Thomas' NHS Foundation Trust. Early arthritis Clinic Referral Criteria. [Internet] Available from: <http://www.guysandstthomas.nhs.uk/our-services/rheumatology/specialties/early-arthritis/referrals.aspx#na>. [accessed on 10 May 2017].
 23. Van Nies JA, Brouwer E, Van Gaalen FA, *et al*. Improved early identification of arthritis: evaluating the efficacy of Early Arthritis Recognition Clinics. *Ann Rheum Dis*. 2013 Aug;72(8):1295-301.
 24. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med*. 2011 Feb 20;30(4):377-99.
 25. Knuiman CA, Schers HJ. [Het beloop van aspecifieke artritis]. *Huisarts Wet*. 2011;54(5):234-236.
 26. King G, Zeng L. Estimating risk and rate levels, ratios and differences in case-control studies. *Stat Med*. 2002;21(10):1409
 27. King G, Zeng L. Inference in Case-Control Studies. *Encyclopedia of Biopharmaceutical Statistics*. 2004. Available from: <https://gking.harvard.edu/files/gking/files/1s-enc.pdf>.
 28. Van den Bosch WB, Mangnus L, Reijnierse M, *et al*. The diagnostic accuracy of the squeeze test to identify arthritis: a cross-sectional cohort study. *Ann Rheum Dis*. 2015 Oct;74(10):1886-9.
 29. Janssens HJEM, Lagro HAHM, Van Peet PG, *et al*. [NHG-Standaard Artritis (Eerste versie)]. *Huisarts Wet*. 2009;52(9):439-53.
 30. Symmons D, Turner G, Webb R, *et al*. The prevalence of rheumatoid arthritis in the United Kingdom: new estimates for a new century. *Rheumatology*. 2002 Jul;41(7):793-800.

