



Universiteit
Leiden
The Netherlands

Acute pancreatitis: from treatment to prevention

Sissingh, N.J.

Citation

Sissingh, N. J. (2024, October 17). *Acute pancreatitis: from treatment to prevention*. Retrieved from <https://hdl.handle.net/1887/4098024>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4098024>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 7

Gallstones as a cause in presumed acute alcoholic pancreatitis: *observational multicentre study*

British Journal of Surgery 2024

Authors

Noor J. Sissingh, Fleur E.M. de Rijk, Hester C. Timmerhuis,
Devica S. Umans, Marie-Paule G.F. Anten, Stefan A.W. Bouwense,
Foke van Delft, Brechje C. van Eijck, Willemien G. Erkelens, Wouter L.
Hazen, Sjoerd D. Kuiken, Rutger Quispel, Tessa E.H. Römken, Matthijs
P. Schwartz, Tom C. Seerden, Marcel Spanier, Tessa Verlaan, Frank P.
Vleggaar, Rogier P. Voermans, Robert C. Verdonk, Jeanin E. van Hooft

for the Dutch Pancreatitis Study Group

ABSTRACT

Background: Data on the incidence and clinical relevance of gallstones in patients with suspected acute alcoholic pancreatitis are lacking and are essential to minimize the risk of recurrent acute pancreatitis. The aim of this study was to assess the incidence of gallstones and the associated rate of recurrent acute pancreatitis in patients with presumed acute alcoholic pancreatitis.

Methods: Between 2008 and 2019, 23 hospitals prospectively enrolled patients with acute pancreatitis. Those diagnosed with their first episode of presumed acute alcoholic pancreatitis were included in this study. The term gallstones was used to describe the presence of cholelithiasis or biliary sludge found during imaging. The primary outcome was pancreatitis recurrence during 3 years of follow-up.

Results: A total of 334 patients were eligible for inclusion, of whom 316 patients were included in the follow-up analysis. Gallstone evaluation, either during the index admission or during follow-up, was performed for 306 of 334 patients (91.6%). Gallstones were detected in 54 patients (17.6%) with a median time to detection of 6 (IQR 0-42) weeks. During follow-up, recurrent acute pancreatitis occurred in 121 of 316 patients (38.3%), with a significantly higher incidence rate for patients with gallstones compared with patients without gallstones (59% versus 34.2% respectively; $p<0.001$), while more patients with gallstones had stopped drinking alcohol at the time of their first recurrence (41% versus 24% respectively; $p=0.020$). Cholecystectomy was performed for 19 patients with gallstones (36%). The recurrence rate was lower for patients in the cholecystectomy group compared with patients who did not receive inadequate or no treatment (5/19 versus 19/34 respectively; $p=0.038$).

Conclusions: Gallstones were found in almost one in every five patients diagnosed with acute alcoholic pancreatitis. Gallstones were associated with a higher rate of recurrent pancreatitis, while undergoing cholecystectomy was associated with a reduction in this rate.

INTRODUCTION

The incidence of acute pancreatitis continues to rise, with an average annual increase of 3.67% in the USA (1). Biliary disease and alcohol are the most common causes (2, 3). Identification of the underlying aetiology is important to guide targeted interventions; cholecystectomy can prevent recurrent biliary events in patients with biliary pancreatitis (4), while alcohol cessation support can reduce the risk of recurrent acute pancreatitis (5). Yet, the precise alcohol threshold that defines alcoholic aetiology is unclear and likely varies between patients (6). Previous studies have used different definitions, with thresholds ranging from more than 3 or 4 units per day (7, 8) to more than 4 units in the 24h before onset (9). As a result, the diagnosis often depends on ruling out other potential causes. Current guidelines recommend a comprehensive evaluation that includes a personal and family history, laboratory tests, and a transabdominal ultrasonography (TUS) (10). If TUS shows cholelithiasis, biliary sludge, or dilated bile duct(s), biliary disease is generally considered the most likely cause (11), at least for those who are not excessive drinkers.

Distinguishing between biliary and alcoholic aetiologies can be challenging. First, the lack of clear criteria for alcoholic pancreatitis may lead clinicians to rely on subjective interpretations of excessive alcohol consumption. This approach may bypass TUS, as recommended by the guidelines (10), potentially leading to the misdiagnosis of patients with biliary aetiology as having alcoholic pancreatitis alone. Second, the diagnostic accuracy of TUS in detecting sludge is limited, especially in the acute phase (12, 13). Therefore, patients with suspected idiopathic acute pancreatitis often undergo TUS for a second time and, if necessary, endoscopic ultrasonography (EUS) (9, 14). Conversely, for patients labelled as having alcoholic pancreatitis, the diagnostic work-up may stop after a single (possibly suboptimal) TUS at the time of diagnosis. This scenario exposes patients to potential recurrent biliary events, including acute pancreatitis. Finally, even if a biliary aetiology is identified, it may go untreated because alcohol is considered the primary trigger of pancreatitis. This again raises the risk of future biliary complications.

In the absence of available literature, the aim of this study was to assess the incidence of gallstones and the associated rate of recurrent acute pancreatitis in a large prospective nationwide cohort of patients with presumed acute alcoholic pancreatitis.

METHODS

This study was performed according to the principles of the Declaration of Helsinki and the STROBE guideline (available as Supplementary material) (15). This study was not pre-registered in an independent, institutional registry.

Study design and population

This study is a post-hoc analysis of the Dutch Pancreatitis Study Group's prospective nationwide registry of acute pancreatitis (PWN-CORE). For this study, all patients from 23 hospitals between 2008 and 2019 were screened for eligibility. Acute pancreatitis was defined according to the revised Atlanta classification (16).

Eligible patients were adults with a first episode of 'presumed' alcoholic pancreatitis, diagnosed when the treating physician considered alcohol as the most likely cause, and no treatment was initiated for other aetiological factors. Patients were excluded if they had chronic pancreatitis according to the M-ANNHEIM criteria at the time of the first diagnosis of acute pancreatitis (17), if they had a previous episode of acute pancreatitis for which data could not be retrieved, or if data were incomplete. PWN-CORE was approved by a medical ethics committee (W19.088). Written informed consent was obtained from each participant.

Data collection

Clinical data, including patient characteristics and results of laboratory and imaging tests, were prospectively collected at the time of initial hospitalization using a standardized case record form. Follow-up data on imaging, and readmissions and outpatient hospital visits for recurrent pancreatitis, biliary complications, and biliary interventions were collected retrospectively from medical records and evaluated until 3 years after the initial admission. Only initial admission data were used for patients who were lost to follow-up. Data were imported into data management software by two researchers (N.J.S. and F.E.M.d.R.). Any discrepancies were resolved by discussion with the involvement of an expert (R.C.V.) until consensus was reached.

Outcomes

The primary outcome was the rate of recurrent acute pancreatitis according to the revised Atlanta classification during the 3 years after the initial admission (16). This follow-up interval was chosen because it was not considered feasible to link the initial episode of pancreatitis to subsequent gallstone detection and possible recurrence beyond this interval. Secondary outcomes included biliary events, biliary interventions, adherence to guidelines for performing the standard diagnostic work-up (10), and diagnostic yield of additional imaging after an initial negative TUS result.

Definitions

Alcohol consumption, as reported immediately upon hospital admission, was converted into standard units per week (1 standard unit equals 10 g) for regular drinkers using an online calculation tool (18). Those who occasionally consumed more than 4 (for women) or 5 (for men) standard units were classified as binge drinkers (19). The term gallstones was used to describe the presence of cholelithiasis or biliary sludge found during the following imaging tests: transabdominal TUS, EUS, magnetic resonance imaging (MRI), and magnetic resonance cholangiopancreatography (MRCP) (14, 20-22). Biliary events included acute cholecystitis, cholangitis, obstructive choledocholithiasis requiring endoscopic retrograde cholangiopancreatography, and biliary colic. Acute cholecystitis and cholangitis were defined according to the Tokyo classification (23, 24). Biliary colic was defined according to the Rome IV criteria (25). A complete standard diagnostic work-up was defined as serum calcium and triglycerides tests, and imaging with TUS according to the International Association of Pancreatology/American Pancreatic Association guidelines during the index admission (10). Personal and family histories (that is drug use, genetic mutations, etc.) were not included in this study due to the challenges of the retrospective design. All definitions are listed in Table S1.

Statistical analysis

All analyses were performed using SPSS® (IBM, Armonk, NY, USA). Categorical variables are presented as n (%) and continuous variables are presented as mean (s.d.) or median (interquartile range (IQR)). Statistical comparisons between patients with and without gallstones were made using the chi-squared test or Fisher's exact test for categorical data and Student's t-test or the Mann-Whitney U test for continuous data. Other pre-specified subgroup analyses were attempted based on history of cholecystectomy and initial TUS results. The diagnostic yield for each imaging modality is presented as % (95% CI). Missing data were not imputed. $P < 0.050$ was considered statistically significant.

RESULTS

Between 2008 and 2019, 2447 patients from 23 hospitals were prospectively registered. Of these, 334 were included in the present study (Figure 1). Clinical characteristics are shown in Table 1. A total of 10 patients (3.0%) had previously undergone a cholecystectomy.

Alcohol consumption was self-reported for 295 patients (Table 1). Median liver enzyme levels at admission are shown in Table 1.

During the index admission, serum calcium tests were performed for 290 patients (86.8%) and triglyceride tests were performed for 199 patients (59.6%). Abnormal calcium and triglyceride tests were found for 0 patients and 16 patients (8.0%), respectively. TUS was performed during the index admission for 276 patients (82.6%). The median number of standard alcohol units per week was higher for the group of patients who did not undergo TUS than in those who did undergo TUS (70 versus 35 units per week; respectively; $p<0.001$) (Table S2). A complete standard diagnostic work-up according to the guidelines was performed for less than half of all patients (156 of 334 patients (46.7%)).

During the 3-year follow-up interval, 19 patients (5.7%) died (of these, 3 patients died during the index admission). A total of 18 patients (5.4%) were lost to follow-up after the index admission (Figure 1), resulting in 316 patients included in the assessment for the follow-up interval.




Figure 1. Flowchart

Gallstone detection

During the index admission, 276 out of 334 patients (82.6%) underwent TUS, of whom 18 (6.5%) were found to have gallstones (Table 2). During follow-up, 198 out of 316 patients (62.7%) underwent imaging, including (repeat) TUS for 168 patients (53.2%), MRI/MRCP for 69 patients (21.8%), and EUS for 40 patients (12.7%). These imaging modalities identified gallstones in 47 patients (23.7%). Taking into account overlap, 306 out of 334 patients (91.6%) underwent at least one imaging test for gallstone evaluation, either during the index admission or during follow-up. Gallstones were found in 54 of these 306 patients (17.6%), of whom 1 was lost to follow-up. The median detection time was 6 (IQR 0-42) weeks after the index admission.

Table 1. Clinical characteristics

	N=334
Age in years – mean (SD)	50 (14)
Male sex – no. (%)	275 (82.3%)
BMI – median (IQR) (n=182)	25 (23-28)
Prior cholecystectomy – no. (%)	10 (3.0%)
Self-reported alcohol use per week – no. (%) (n=295)*	
≤21 units	49 (16.6%)
>21 units	188 (63.7%)
Binge drinking	58 (19.7%)
Smoking – no. (%) (n=304)	200 (65.8%)
Liver enzymes at admission – median (IQR)	
Aspartate aminotransferase in U/L (n=324)	39 (25-79)
Alanine aminotransferase in U/L (n=329)	40 (24-72)
Alkaline phosphatase in U/L (n=327)	86 (71-119)
Gamma-glutamyltransferase in U/L (n=330)	113 (52-395)
Bilirubin total in µmol/L (n=329)	14 (9-22)
Standard diagnostic work-up during index-admission	
Calcium testing – no. (%)	290 (86.8%)
Calcium in mmol/L – median (IQR)	2.29 (2.12-2.40)
Calcium >3 mmol/L – no. (%)	0
Triglycerides testing – no. (%)	199 (59.6%)
Triglycerides in mmol/L – median (IQR)	1.53 (0.92-2.80)
Triglycerides >11.2 mmol/L – no. (%)	16 (8.0%)
Transabdominal ultrasound – no. (%)^	276 (82.6%)
Complete standard diagnostic work-up – no. (%)	156 (46.7%)
Severity of acute pancreatitis – no. (%)#	
Mild	207 (62.0%)
Moderate severe	94 (28.1%)
Severe	33 (9.9%)
Mortality – no. (%)	19 (5.7%)
During index admission – no. (%)	3 (0.9%)

Data are presented as no. (%), mean (SD), or median (IQR). *The cut-off value of 21 units/week was chosen based on the definitions of the Dutch National Institute for Public Health and Environment, which defines excessive alcohol consumption as more than 21 units per week. ^Magnetic resonance cholangiopancreatography was used as the first diagnostic modality in 3 patients. #According to the revised Atlanta classification. Abbreviations: BMI body mass index.

For the subgroup of patients whose initial TUS during the index admission did not reveal gallstones, the diagnostic yields of subsequent imaging tests are detailed in Table 3. The overall gallstone detection rate was 11.0% (95% CI 7.6-14.4). The individual rates were: 11.5% for repeat TUS, 2.8% for MRI/MRCP, and 23.7% for EUS.

Table 2. Number and yield of imaging tests

Type of test	Patients with imaging test – no. (%)	Patients with gallstones based on imaging test – no. (%) [^]	Total of imaging tests performed – no	Gallstones demonstrated on first positive imaging– no*
TUS during index admission	276 (82.6%)	18 (6.5%)	276	12 cholecystolithiasis 1 choledocholithiasis 8 sludge in gallbladder 2 sludge in CBD
Imaging tests after index admission	198 (62.7%)	47 (23.7%)	434	37 cholecystolithiasis 2 choledocholithiasis 20 sludge in gallbladder
(Repeat) TUS	168 (53.2%)	40 (23.8%)	300	27 cholecystolithiasis 15 sludge in gallbladder
MRI/MRCP	69 (21.8%)	4 (5.8%)	86	3 cholecystolithiasis 2 sludge in gallbladder
EUS	40 (12.7%)	11 (27.5%)	48	7 cholecystolithiasis 2 choledocholithiasis 3 sludge in gallbladder
Total for all imaging tests	306 (91.6%)	54 (17.6%)	710	49 cholecystolithiasis 3 choledocholithiasis 28 sludge in gallbladder 2 sludge in CBD

Data are presented as no. (%) or no. [^]Multiple positive imaging modalities in one patients were scored as one outcome. *Please not that in several cases multiple biliary findings were observed in a single patient. Abbreviations: EUS endoscopic ultrasound, CBD common bile duct, MRCP magnetic resonance cholangiopancreatography, MRI magnetic resonance imaging, TUS transabdominal ultrasound.

Recurrent acute pancreatitis and biliary events

During follow-up, recurrent acute pancreatitis occurred in 121 of 316 patients (38%). Patients with gallstones (31 of 53 (59%)) were significantly more likely to develop recurrent acute pancreatitis than patients without gallstones (90 of 263 (34.2%)) ($p < 0.001$). Self-reported alcohol consumption at admission for the first recurrent episode was available for 114 of the 121 patients with recurrence, of whom 32 (28.1%) reported no longer consuming alcohol (41% versus 24% for patients with and without gallstones respectively $p = 0.020$). Subgroup analyses, based on the initial TUS results during the index admission, are presented in Table S3 and show recurrence rates of 53% for patients with an initial positive TUS result, 48% for patients who did not undergo TUS, and 35.0% for patients with an initial negative TUS result. Biliary events after the first episode of pancreatitis were observed in nine patients (3%) (cholangitis, 4 patients; acute cholecystitis, 2 patients; obstructive choledocholithiasis, 2 patients; and colic, 1 patient).

Table 3. Number and yield of additional imaging tests in patients with an initial negative US (n=243)[^]

Type of test	Patients with diagnostic test – no. (%)	Patients with gallstones based on diagnostic test – no. (%)	Total of diagnostic tests performed – no.	Total times gallstones were demonstrated – no.	Diagnostic yield – percentage (95% CI)
Repeat TUS	128 (52.7%)	21 (16.4%)	218	25	11.5% (6.8-15.2)
MRI/MRCP	56 (23.0%)	2 (3.6%)	71	2	2.8% (-1.0-6.6)
EUS	32 (13.2%)	8 (25.0%)	38	9	23.7% (10.1-37.3)
Total	154 (63.4%)	26 (16.9%)	327	36	11.0% (7.6-14.4)

Data are presented as no. (%), no., or percentage (95% CI). [^] A total of 15 patients were excluded as they were lost to follow-up. Abbreviations: EUS endoscopic ultrasound, MRCP magnetic resonance cholangiopancreatography, MRI magnetic resonance imaging, TUS transabdominal ultrasound.

Biliary treatment

During follow-up, 22 of 53 patients with gallstones (42%) underwent biliary intervention. The procedures performed were cholecystectomy with or without biliary endoscopic sphincterotomy (ERCP) (19 patients), biliary endoscopic sphincterotomy alone (1 patient), and percutaneous gallbladder drainage (2 patients). The remaining 31 patients (59%) received no biliary intervention. A single patient underwent an unsuccessful ERCP procedure in which no biliary access could be obtained, one patient's scheduled cholecystectomy was cancelled due to the development of metastatic disease, and another patient did not attend a scheduled appointment at the surgical division to discuss the possibility of elective cholecystectomy.

After receiving appropriate treatment (that is cholecystectomy), 5 of 19 patients developed recurrent acute pancreatitis compared with 19 of 34 patients who received inadequate treatment (that is ERCP or percutaneous gallbladder drainage alone) or no treatment (relative risk 0.47; 95% CI 0.21-1.06; p=0.038).

DISCUSSION

In this first nationwide cohort study, we found that 91.6% of patients with presumed acute alcoholic pancreatitis underwent gallstone evaluation, and 17.6% were found to have gallstones. Patients with gallstones had a nearly twofold increased risk of recurrent acute pancreatitis. In contrast, those patients with gallstones who underwent cholecystectomy had half the risk of recurrence. Cholecystectomy was performed for only 36% of patients.

The main findings are that the risk of pancreatitis recurrence significantly increased from 34.2% to 59% in the presence of gallstones for patients with presumed acute alcoholic pancreatitis, even if more of these patients were alcohol abstinent at the time of recurrence. Although the authors' study group has previously evaluated recurrence rates for patients with biliary pancreatitis and alcoholic pancreatitis (26) (rates of 12% and 24%, respectively), no studies have specifically targeted our study population, making direct comparisons difficult. Based on these results, one could speculate that a significant number of patients diagnosed with alcoholic pancreatitis may indeed have biliary pancreatitis, accompanied by excessive alcohol consumption habits, and may benefit from cholecystectomy. The results of the present study show that the recurrence rate after cholecystectomy is indeed two times lower than after no treatment.

A dual role for alcohol in the pathophysiology of acute pancreatitis, acting as either a trigger or a modulator, has previously been suggested (6). This is supported by the low lifetime risk of developing acute pancreatitis in the overall population of excessive alcohol consumers (27). However, the potential interaction of alcohol with biliary disease in the development of acute pancreatitis and vice versa remains far from clear. Furthermore, the authors believe that the exact aetiology cannot be determined when a patient presents with both gallstones and excessive alcohol consumption. Future studies should focus on identifying biochemical and clinical markers and combining them in a (machine learning) prediction model to adequately differentiate between these two aetiologies. Meanwhile, it is important to recognize and address both potential aetiologies, with the initial approach being that patients with excessive alcohol consumption undergo the same diagnostic work-up as non-drinkers.

At admission, current guidelines recommend testing for calcium and triglycerides and performing an TUS (10). In the present study, only half of the patients underwent this recommended work-up. Although TUS is both affordable and non-invasive, it was not performed in 17.4% of patients. Notably, a significant difference in alcohol consumption was observed, favouring patients who underwent TUS. This raises concerns about potential alcohol-related stigma in the management of acute pancreatitis, which should be further explored, for example in a qualitative study assessing the stigmatizing attitudes of pancreatologists towards their patients. Such attitudes have been well documented in other diseases often considered to be self-inflicted, such as HIV, obesity, and psychiatric disorders (28-30). The potential influence of stigma on the doctor-patient relationship, and its impact on a patient's quality of life (31) underscores the need to improve the understanding of the pathophysiology and to establish universally accepted diagnostic criteria for acute alcoholic pancreatitis. In the meantime, strict adherence to guidelines for patients suspected of having alcoholic pancreatitis remains critical, especially as the recurrence rate of pancreatitis was sig-

nificantly higher for patients who did not undergo TUS compared with patients with an initially negative TUS result.

Positive TUS findings for gallstones were present for 6.5% of patients during the index admission, suggesting that an initial diagnosis of just alcoholism would be incorrect. However, given the established prevalence of biliary disease in the general population in the USA (32), these findings must be interpreted with caution. Factors known to increase the risk of biliary disease include female sex, older age, and higher BMI (33), while the effect of alcohol remains controversial (34-36). Alcohol has been associated with gallstones because of its role in lipid metabolism, but studies also report that alcohol may increase gallbladder motility and decrease bile lithogenicity, thus protecting against gallstone formation.

Additional imaging was performed for 62.7% of patients with an initial negative TUS result. The overall yield was 11.0%, which may be an underestimate given the relatively low utilization rates of EUS and MRCP, the two modalities with the highest accuracy for detection of gallstones (37, 38). This supports the hypothesis that patients diagnosed with alcoholic pancreatitis may have undetected occult gallstones. However, the inclusion of patients with a high suspicion of gallstones, as indicated by prior diagnostic tests such as repeat TUS, may have influenced the high individual yield of EUS (23.7%). Nevertheless, the repeat TUS showed a notable yield of 11.5%, above the 10% cut-off considered sufficient for routine use of EUS for patients after the first episode of idiopathic acute pancreatitis (9). In the absence of prospective studies, the validity, futility and optimal timing of additional imaging for patients with excessive alcohol consumption and an initial negative TUS result require further investigation before reliable recommendations can be made.

In the present study, only 36% of patients with gallstones underwent a cholecystectomy, while the remaining patients received no or inadequate treatment (that is ERCP or percutaneous gallbladder drainage alone). This suggests that clinicians often consider gallstones in patients with presumed alcoholic pancreatitis as an incidental finding that does not warrant further treatment, which is concerning, as the incidence rate of recurrent acute pancreatitis after cholecystectomy was 5/19 compared with 19/37 for untreated patients. This finding, together with the consistent evidence for the effectiveness of cholecystectomy in preventing recurrence (4, 39), emphasizes that timely cholecystectomy should be considered once gallstones are identified. Nevertheless, the observed recurrence rate after cholecystectomy highlights the importance of broadening the focus beyond the consideration of cholecystectomy alone. Previous studies have shown that alcohol cessation reduces recurrence to almost 0% (40, 41). However, complete alcohol abstinence is notoriously difficult, as the authors have also

observed. In an earlier Finnish trial, additional alcohol reduction efforts were shown to reduce the risk of recurrent acute pancreatitis (5). Recently, the Dutch Pancreatitis Study Group has initiated the multicentre PANDA trial (42). This trial aims to determine the effectiveness of a structured alcohol cessation support programme in reducing the rate of recurrent acute pancreatitis for patients after their first episode of acute alcoholic pancreatitis when compared with the current standard practice.

The post-hoc design of the present study has limitations, leading to several drawbacks. First, it must be emphasized that the observed association between gallstones and recurrent pancreatitis and between cholecystectomy and recurrent pancreatitis does not imply causality. For example, it was not possible to account for the possibility of different aetiologies for different attacks. Second, data on continued alcohol use or cessation after the first episode of pancreatitis were collected post hoc and only at the time of recurrent episodes. Therefore, it was not possible to perform a multivariable analysis with alcohol use as a potential confounder to compare the primary outcome for patients with and without gallstones. In addition, the study utilized a pragmatic approach to assessing alcohol use, relying on patient self-reporting and making the following distinction: continued alcohol use? (yes or no). Third, it was not possible to reliably assess data on smoking after the first episode of pancreatitis, a factor that could also influence the primary outcome. Fourth, a diagnosis of acute alcoholic pancreatitis was made based on the discretion of the treating physician and therefore no predefined diagnostic work-up was required, which may have introduced bias. On the other hand, the present study reflects what is happening in current clinical practice. Another limitation is that all of the imaging studies performed were evaluated and not just those that were done to assess the presence of gallstones. Also, patients who did not undergo any imaging were included in the subgroup of patients thought to have no gallstones. Finally, subgroup analyses based on gallbladder status were not possible because only 10 out of the 334 patients had a history of cholecystectomy.

CONCLUSION

Our study found that 17.6% of patients diagnosed with acute alcoholic pancreatitis had gallstones, which were significantly associated with a higher rate of recurrent acute pancreatitis. In addition, we show that gallstone evaluation at initial admission was not consistently performed. The same was true for the performance of cholecystectomy once gallstones were identified. This is of concern, especially since our results also showed an almost significant reduction in recurrent pancreatitis after cholecystectomy. With the ever-increasing burden of acute pancreatitis, we strongly recommend better adherence to guidelines for all patients suspected of having acute alcoholic pancreatitis, including performing an transabdominal ultrasound, and considering cholecystectomy for those diagnosed with gallstones.

REFERENCES

1. Iannuzzi JP, King JA, Leong JH, Quan J, Windsor JW, Tanyingoh D, Coward S, Forbes N, Heitman SJ, Shaheen AA, Swain M, Buie M, Underwood FE, Kaplan GG. Global Incidence of Acute Pancreatitis Is Increasing Over Time: A Systematic Review and Meta-Analysis. *Gastroenterology*. 2022 Jan;162(1):122-134. doi: 10.1053/j.gastro.2021.09.043.
2. Yadav D, Lowenfels AB. Trends in the epidemiology of the first attack of acute pancreatitis: a systematic review. *Pancreas*. 2006 Nov;33(4):323-30. doi: 10.1097/01.mpa.0000236733.31617.52.
3. Roberts SE, Morrison-Rees S, John A, Williams JG, Brown TH, Samuel DG. The incidence and aetiology of acute pancreatitis across Europe. *Pancreatology*. 2017 Mar-Apr;17(2):155-165. doi: 10.1016/j.pan.2017.01.005.
4. da Costa DW, Bouwense SA, Schepers NJ, Besselink MG, van Santvoort HC, van Brunschot S, Bakker OJ, Bollen TL, Dejong CH, van Goor H, Boermeester MA, Bruno MJ, van Eijck CH, Timmer R, Weusten BL, Consten EC, Brink MA, Spanier BWM, Bilgen EJS, Nieuwenhuijs VB, Hofker HS, Rosman C, Voorburg AM, Bosscha K, van Duijvendijk P, Gerritsen JJ, Heisterkamp J, de Hingh IH, Witteman BJ, Kruij PM, Scheepers JJ, Molenaar IQ, Schaapherder AF, Manusama ER, van der Waaij LA, van Unen J, Dijkgraaf MG, van Ramshorst B, Gooszen HG, Boerma D; Dutch Pancreatitis Study Group. Same-admission versus interval cholecystectomy for mild gallstone pancreatitis (PONCHO): a multicentre randomised controlled trial. *Lancet*. 2015 Sep 26;386(10000):1261-1268. doi: 10.1016/S0140-6736(15)00274-3.
5. Nordback I, Pelli H, Lappalainen-Lehto R, Järvinen S, Rätty S, Sand J. The recurrence of acute alcohol-associated pancreatitis can be reduced: a randomized controlled trial. *Gastroenterology*. 2009 Mar;136(3):848-55. doi: 10.1053/j.gastro.2008.11.044.
6. Nordback I, Sand J, Andrén-Sandberg A. Criteria for alcoholic pancreatitis. Results of an international workshop in Tampere, Finland, June 2006. *Pancreatology*. 2007;7(2-3):100-4. doi: 10.1159/000104234.
7. Hallensleben ND, Umans DS, Bouwense SA, Verdonk RC, Romkens TE, Witteman BJ, Schwartz MP, Spanier MB, Laheij R, van Santvoort HC, Besselink MG, van Hooft JE, Bruno MJ; Dutch Pancreatitis Study Group. The diagnostic work-up and outcomes of 'presumed' idiopathic acute pancreatitis: A post-hoc analysis of a multicentre observational cohort. *United European Gastroenterol J*. 2020 Apr;8(3):340-350. doi: 10.1177/2050640619890462.
8. Schepers NJ, Hallensleben ND, Besselink MG, Anten MGF, Bollen TL, da Costa DW, van Delft F, van Dijk SM, van Dullemen HM, Dijkgraaf MGW, van Eijck CHJ, Erkelens GW, Erler NS, Fockens P, van Geenen EJM, van Grinsven J, Hollemans RA, van Hooft JE, van der Hulst RWM, Jansen JM, Kubben FJGM, Kuiken SD, Laheij R, Quispel R, de Ridder RJJ, Rijk MCM, Römkens TEH, Ruigrok CHM, Schoon EJ, Schwartz MP, Smeets XJNM, Spanier BWM, Tan ACITL, Thijs WJ, Timmer R, Venneman NG, Verdonk RC, Vleggaar FP, van de Vrie W, Witteman BJ, van Santvoort HC, Bakker OJ, Bruno MJ; Dutch Pancreatitis Study Group. Urgent endoscopic retrograde cholangiopancreatography with sphincterotomy versus conservative treatment in predicted severe acute gallstone pancreatitis (APEC): a multicentre randomised controlled trial. *Lancet*. 2020 Jul 18;396(10245):167-176. doi: 10.1016/S0140-6736(20)30539-0.
9. Umans DS, Timmerhuis HC, Hallensleben ND, Bouwense SA, Anten MG, Bhalla A, Bijlsma RA, Boermeester MA, Brink MA, Hol L, Bruno MJ, Curvers WL, van Dullemen HM, van Eijck BC, Erkelens GW, Fockens P, van Geenen EJM, Hazen WL, Hoge CV, Inderson A, Kager LM, Kuiken SD, Perk LE, Poley JW, Quispel R, Römkens TE, van Santvoort HC, Tan AC, Thijssen AY, Venneman NG, Vleggaar FP, Voorburg AM, van Wanrooij RL, Witteman BJ, Verdonk RC,

- Besselink MG, van Hooft JE; Dutch Pancreatitis Study Group. Role of endoscopic ultrasonography in the diagnostic work-up of idiopathic acute pancreatitis (PICUS): study protocol for a nationwide prospective cohort study. *BMJ Open*. 2020 Aug 20;10(8):e035504. doi: 10.1136/bmjopen-2019-035504.
10. Working Group IAP/APA Acute Pancreatitis Guidelines. IAP/APA evidence-based guidelines for the management of acute pancreatitis. *Pancreatology*. 2013 Jul-Aug;13(4 Suppl 2):e1-15. doi: 10.1016/j.pan.2013.07.063.
 11. Żorniak M, Sirtl S, Beyer G, Mahajan UM, Bretthauer K, Schirra J, Schulz C, Kohlmann T, Lerch MM, Mayerle J; LMU Microlithiasis Expert Survey Team. Consensus definition of sludge and microlithiasis as a possible cause of pancreatitis. *Gut*. 2023 Oct;72(10):1919-1926. doi: 10.1136/gutjnl-2022-327955.
 12. Thorbøll J, Vilmann P, Jacobsen B, Hassan H. Endoscopic ultrasonography in detection of cholelithiasis in patients with biliary pain and negative transabdominal ultrasonography. *Scand J Gastroenterol*. 2004 Mar;39(3):267-9. doi: 10.1080/00365520310008377.
 13. Mirbagheri SA, Mohamadnejad M, Nasiri J, Vahid AA, Ghadimi R, Malekzadeh R. Prospective evaluation of endoscopic ultrasonography in the diagnosis of biliary microlithiasis in patients with normal transabdominal ultrasonography. *J Gastrointest Surg*. 2005 Sep-Oct;9(7):961-4. doi: 10.1016/j.gassur.2005.03.002.
 14. Umans DS, Rangkuti CK, Sperna Weiland CJ, Timmerhuis HC, Bouwense SAW, Fockens P, Besselink MG, Verdonk RC, van Hooft JE; Dutch Pancreatitis Study Group. Endoscopic ultrasonography can detect a cause in the majority of patients with idiopathic acute pancreatitis: a systematic review and meta-analysis. *Endoscopy*. 2020 Nov;52(11):955-964. doi: 10.1055/a-1183-3370.
 15. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol*. 2008 Apr;61(4):344-9. doi: 10.1016/j.jclinepi.2007.11.008.
 16. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, Tsiotos GG, Vege SS; Acute Pancreatitis Classification Working Group. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. *Gut*. 2013 Jan;62(1):102-11. doi: 10.1136/gutjnl-2012-302779.
 17. Schneider A, Löhr JM, Singer MV. The M-ANNHEIM classification of chronic pancreatitis: introduction of a unifying classification system based on a review of previous classifications of the disease. *J Gastroenterol*. 2007 Feb;42(2):101-19. doi: 10.1007/s00535-006-1945-4.
 18. Alcoholhulp. Alcoholcalculator. <https://www.alcoholhulp.be/alcoholcalculator>. (accessed until 2 January 2024).
 19. NIAAA. Defining binge drinking. What colleges need to know now. Bethesda (MD): National Institutes of Health; 2007 Nov. https://www.collegedrinkingprevention.gov/sites/cdp/files/documents/1College_Bulletin-508_361C4E.pdf. (accessed until 2 January 2024).
 20. Umans DS, Timmerhuis HC, Anten MGF, Bhalla A, Bijlsma RA, Boxhoorn L, Brink MA, Bruno MJ, Curvers WL, van Eijck BC, Erkelens GW, van Geenen EJM, Hazen WL, Hoge CV, Hol L, Inderson A, Kager LM, Kuiken SD, Perk LE, Quispel R, Römken TEH, Sperna Weiland CJ, Thijssen AY, Venneman NG, Verdonk RC, van Wanrooij RLJ, Witteman BJ, Besselink MG, van Hooft JE. Prospective multicentre study of indications for surgery in patients with idiopathic acute pancreatitis following endoscopic ultrasonography (PICUS). *Br J Surg*. 2023 Nov 9;110(12):1877-1882. doi: 10.1093/bjs/znad318.

21. Garg PK, Tandon RK, Madan K. Is biliary microlithiasis a significant cause of idiopathic recurrent acute pancreatitis? A long-term follow-up study. *Clin Gastroenterol Hepatol*. 2007 Jan;5(1):75-9. doi: 10.1016/j.cgh.2006.06.023.
22. Lee SP, Nicholls JF, Park HZ. Biliary sludge as a cause of acute pancreatitis. *N Engl J Med*. 1992 Feb 27;326(9):589-93. doi: 10.1056/NEJM199202273260902.
23. Yokoe M, Hata J, Takada T, Strasberg SM, Asbun HJ, Wakabayashi G, Kozaka K, Endo I, Deziel DJ, Miura F, Okamoto K, Hwang TL, Huang WS, Ker CG, Chen MF, Han HS, Yoon YS, Choi IS, Yoon DS, Noguchi Y, Shikata S, Ukai T, Higuchi R, Gabata T, Mori Y, Iwashita Y, Hibi T, Jagannath P, Jonas E, Liau KH, Dervenis C, Gouma DJ, Cherqui D, Belli G, Garden OJ, Giménez ME, de Santibañes E, Suzuki K, Umezawa A, Supe AN, Pitt HA, Singh H, Chan ACW, Lau WY, Teoh AYB, Honda G, Sugioka A, Asai K, Gomi H, Itoi T, Kiriyaama S, Yoshida M, Mayumi T, Matsumura N, Tokumura H, Kitano S, Hirata K, Inui K, Sumiyama Y, Yamamoto M. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholecystitis (with videos). *J Hepatobiliary Pancreat Sci*. 2018 Jan;25(1):41-54. doi: 10.1002/jhbp.515.
24. Kiriyaama S, Kozaka K, Takada T, Strasberg SM, Pitt HA, Gabata T, Hata J, Liau KH, Miura F, Horiguchi A, Liu KH, Su CH, Wada K, Jagannath P, Itoi T, Gouma DJ, Mori Y, Mukai S, Giménez ME, Huang WS, Kim MH, Okamoto K, Belli G, Dervenis C, Chan ACW, Lau WY, Endo I, Gomi H, Yoshida M, Mayumi T, Baron TH, de Santibañes E, Teoh AYB, Hwang TL, Ker CG, Chen MF, Han HS, Yoon YS, Choi IS, Yoon DS, Higuchi R, Kitano S, Inomata M, Deziel DJ, Jonas E, Hirata K, Sumiyama Y, Inui K, Yamamoto M. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholangitis (with videos). *J Hepatobiliary Pancreat Sci*. 2018 Jan;25(1):17-30. doi: 10.1002/jhbp.512.
25. Cotton PB, Elta GH, Carter CR, Pasricha PJ, Corazziari ES. Rome IV. Gallbladder and Sphincter of Oddi Disorders. *Gastroenterology*. 2016 Feb 19:S0016-5085(16)00224-9. doi: 10.1053/j.gastro.2016.02.033.
26. Ahmed Ali U, Issa Y, Hagenaars JC, Bakker OJ, van Goor H, Nieuwenhuijs VB, Bollen TL, van Ramshorst B, Witteman BJ, Brink MA, Schaapherder AF, Dejong CH, Spanier BW, Heisterkamp J, van der Harst E, van Eijck CH, Besselink MG, Gooszen HG, van Santvoort HC, Boermeester MA; Dutch Pancreatitis Study Group. Risk of Recurrent Pancreatitis and Progression to Chronic Pancreatitis After a First Episode of Acute Pancreatitis. *Clin Gastroenterol Hepatol*. 2016 May;14(5):738-46. doi: 10.1016/j.cgh.2015.12.040.
27. Lankisch PG, Lowenfels AB, Maisonneuve P. What is the risk of alcoholic pancreatitis in heavy drinkers? *Pancreas*. 2002 Nov;25(4):411-2. doi: 10.1097/00006676-200211000-00015.
28. Kay ES, Rice WS, Crockett KB, Atkins GC, Batey DS, Turan B. Experienced HIV-Related Stigma in Health Care and Community Settings: Mediated Associations With Psychosocial and Health Outcomes. *J Acquir Immune Defic Syndr*. 2018 Mar 1;77(3):257-263. doi: 10.1097/QAI.0000000000001590.
29. Lawrence BJ, Kerr D, Pollard CM, Theophilus M, Alexander E, Haywood D, O'Connor M. Weight bias among health care professionals: A systematic review and meta-analysis. *Obesity (Silver Spring)*. 2021 Nov;29(11):1802-1812. doi: 10.1002/oby.23266.
30. Oliveira AM, Machado D, Fonseca JB, Palha F, Silva Moreira P, Sousa N, Cerqueira JJ, Morgado P. Stigmatizing Attitudes Toward Patients With Psychiatric Disorders Among Medical Students and Professionals. *Front Psychiatry*. 2020 Apr 30;11:326. doi: 10.3389/fpsy.2020.00326.
31. Clement S, Schauman O, Graham T, Maggioni F, Evans-Lacko S, Bezborodovs N, Morgan C, Rüsch N, Brown JS, Thornicroft G. What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychol Med*. 2015 Jan;45(1):11-27. doi: 10.1017/S0033291714000129.

32. Unalp-Arida A, Ruhl CE. Increasing gallstone disease prevalence and associations with gallbladder and biliary tract mortality in the US. *Hepatology*. 2023 Jun 1;77(6):1882-1895. doi: 10.1097/HEP.0000000000000264.
33. Halldestam I, Kullman E, Borch K. Incidence of and potential risk factors for gallstone disease in a general population sample. *Br J Surg*. 2009 Nov;96(11):1315-22. doi: 10.1002/bjs.6687.
34. You M, Arteel GE. Effect of ethanol on lipid metabolism. *J Hepatol*. 2019 Feb;70(2):237-248. doi: 10.1016/j.jhep.2018.10.037.
35. Thijs C, Knipschild P, Leffers P. Does alcohol protect against the formation of gallstones? A demonstration of protopathic bias. *J Clin Epidemiol*. 1991;44(9):941-6. doi: 10.1016/0895-4356(91)90057-g.
36. Cha BH, Jang MJ, Lee SH. Alcohol Consumption Can Reduce the Risk of Gallstone Disease: A Systematic Review with a Dose-Response Meta-Analysis of Case-Control and Cohort Studies. *Gut Liver*. 2019 Jan 15;13(1):114-131. doi: 10.5009/gnl18278.
37. Wan J, Ouyang Y, Yu C, Yang X, Xia L, Lu N. Comparison of EUS with MRCP in idiopathic acute pancreatitis: a systematic review and meta-analysis. *Gastrointest Endosc*. 2018 May;87(5):1180-1188.e9. doi: 10.1016/j.gie.2017.11.028.
38. Kondo S, Isayama H, Akahane M, Toda N, Sasahira N, Nakai Y, Yamamoto N, Hirano K, Komatsu Y, Tada M, Yoshida H, Kawabe T, Ohtomo K, Omata M. Detection of common bile duct stones: comparison between endoscopic ultrasonography, magnetic resonance cholangiography, and helical-computed-tomographic cholangiography. *Eur J Radiol*. 2005 May;54(2):271-5. doi: 10.1016/j.ejrad.2004.07.007.
39. Hallensleben ND, Timmerhuis HC, Hollemans RA, Pocornie S, van Grinsven J, van Brunschot S, Bakker OJ, van der Sluijs R, Schwartz MP, van Duijvendijk P, Römkens T, Stommel MWJ, Verdonk RC, Besselink MG, Bouwense SAW, Bollen TL, van Santvoort HC, Bruno MJ; Dutch Pancreatitis Study Group. Optimal timing of cholecystectomy after necrotising biliary pancreatitis. *Gut*. 2022 May;71(5):974-982. doi: 10.1136/gutjnl-2021-324239.
40. Nikkola J, Rätty S, Laukkanen J, Seppänen H, Lappalainen-Lehto R, Järvinen S, Nordback I, Sand J. Abstinence after first acute alcohol-associated pancreatitis protects against recurrent pancreatitis and minimizes the risk of pancreatic dysfunction. *Alcohol Alcohol*. 2013 Jul-Aug;48(4):483-6. doi: 10.1093/alcalc/agt019.
41. Pelli H, Lappalainen-Lehto R, Piironen A, Sand J, Nordback I. Risk factors for recurrent acute alcohol-associated pancreatitis: a prospective analysis. *Scand J Gastroenterol*. 2008;43(5):614-21. doi: 10.1080/00365520701843027.
42. Sissingh NJ, Nagelhout A, Besselink MG, Boermeester MA, Bouwense SAW, Bruno MJ, Fockens P, Goudriaan AE, Rodríguez-Gironde MDM, van Santvoort HC, Sijbom M, van Weert HCPM, van Hooft JE, Umans DS, Verdonk RC; Dutch Pancreatitis Study Group. Structured alcohol cessation support program versus current practice in acute alcoholic pancreatitis (PANDA): Study protocol for a multicentre cluster randomised controlled trial. *Pancreatol*. 2023 Dec;23(8):942-948. doi: 10.1016/j.pan.2023.10.015.

Table S1. Definitions

Baseline	
Acute pancreatitis	When two of the following criteria were met: upper abdominal pain, serum lipase concentration (or amylase) ≥ 3 times higher than the upper limit of normal or features of acute pancreatitis on imaging
(Presumed) alcoholic pancreatitis	Acute pancreatitis was classified as 'presumed' alcoholic if the treating physician considered alcohol as the most probably cause, and no treatment was initiated for other etiological factors.
Binge drinking	The term binge drinking was used for women consuming 4 or more standard units on an occasion or 5 or more standard units on an occasion for men
Outcomes	
Gallstones	Gallstones were defined as the presence of cholelithiasis or biliary sludge on the following imaging techniques: transabdominal ultrasound, endoscopic ultrasound, magnetic resonance imaging, and magnetic resonance cholangiopancreatography
Biliary events	Acute cholecystitis, biliary colic, cholangitis, or obstructive choledocholithiasis
Acute cholecystitis	An acute inflammation of the gallbladder, diagnosed when one item in A, B, and C is present (according to the Tokyo classification): A) Local signs of inflammation 1) Murphy's sign, or 2) Right upper abdominal quadrant mass, pain or tenderness B) Systemic signs of inflammation 1) Fever or hypothermia 2) Elevated C-reactive protein, or 3) Elevated white blood cell count C) Imaging findings characteristics of acute cholecystitis Note: Acute cholecystitis that occurs in the absence of gallstones (i.e. acalculous cholecystitis) did not meet our criteria for acute cholecystitis
Biliary colic	Intermittent pain located in the epigastrium and/or right upper quadrant lasting at least 30 minutes, severe enough to interrupt daily activities or lead to an emergency department visit, not related to bowel movements, and not relieved by postural change or acid suppression (according to Rome IV criteria)

Table S1. Definitions (*continued*)

Cholangitis	An inflammation of the bile duct(s), diagnosed when one item in A, B, and C is present (according to the Tokyo classification): A) Systemic inflammation 1) Fever, hypothermia and/or shaking chills 2) Laboratory data: evidence of inflammatory response (abnormal white blood cell counts, increase of serum CRP levels, and other changes indicating inflammation) B) Cholestasis 1) Jaundice 2) Laboratory data: abnormal liver function tests C) Imaging 1) Biliary dilatation 2) Evidence of aetiology on imaging Note: Cholangitis that occurs in the absence of gallstones did not meet our criteria for cholangitis
Obstructive choledocholithiasis	Presence of gallstones or biliary sludge in the common bile duct on imaging, requiring an ERCP, according to the treating physician
Chronic pancreatitis	A chronic inflammation of the pancreatic parenchyma, defined as typical clinical history or chronic pancreatitis (such as recurrent acute pancreatitis or abdominal pain, except for primary painless pancreatitis and one or more of the following (according to the M-ANNHEIM criteria): A) Pancreatic calcifications B) Moderate or marked ductal lesions C) Marked and persistent exocrine insufficiency D) Typical histology
Complete standard diagnostic work-up	Complete standard diagnostic work-up was defined as laboratory testing, including calcium and triglycerides, and imaging using transabdominal ultrasound, MRI or MRCP (according to IAP/APA guidelines). Note: The standard diagnostic work-up did not include the patient's family history of hereditary pancreatitis or associated genetic mutations, as well as their personal history, such as drug use, due to the challenges of collecting this data retrospectively
Positive imaging	Positive imaging is defined as imaging in which proof of gallstones were found. Note: Isolated dilatation of the intrahepatic bile ducts or CBD was not deemed conclusive evidence for establishing a biliary etiology. This is due to the limitations imposed by the post-hoc design of the study, which prevents reliably confirmation of the absence of other factors that could potentially contribute to duct dilatation, such as stenosis, obstruction caused by external compression, or opioid use
Recurrent acute pancreatitis	Recurrence of acute pancreatitis is defined as a new episode of acute pancreatitis after complete resolution of all symptoms associated with the previous episode (according to the Revised Atlanta criteria)

Table S2. Self-reported alcohol use in patients with and without transabdominal ultrasound at index admission

Self-reported alcohol use per week	All patients (n=295) [*]	Patients with TUS (n=246)	Patients without TUS (n=49)	P-value
Units – median (IQR) (n=237) [^]	42 (28-70)	35 (28-70)	70 (35-112)	<0.001
≤ 21 units – no. (%)	49 (16.6%)	43 (17.5%)	6 (12.2%)	0.369
> 21 units – no. (%)	188 (63.7%)	155 (63.0%)	33 (67.3%)	0.564
Binge drinking – no. (%)	58 (19.7%)	48 (19.5%)	10 (20.4%)	0.885

Data are presented as no. (%), or median (IQR). ^{*}The levels of self-reported alcohol use were available in 295/334 patients. [^]The 58 patients who met the definition for binge drinking were excluded to calculate the median self-reported alcohol use per week. Abbreviations: TUS transabdominal ultrasound.

Table S3. Subgroup analyses of recurrent acute pancreatitis

	All patients (n=316) ^A	Patients with positive first TUS (n=17) ^A	Patients with negative TUS (n=243) ^B	Patients with no TUS (n=56) ^C	P-value	
					A vs. B	B vs. C
Recurrent acute pancreatitis	121 (38.3%)	9 (52.9%)	85 (35.0%)	27 (48.2%)	0.136	0.065

Data are presented as no. (%). ^A A total of 18 patients were excluded as they were lost to follow-up . Abbreviations: TUS transabdominal ultrasound.