



Universiteit  
Leiden  
The Netherlands

## **Metabolomic analysis of *Strychnos nux-vomica*, *Strychnos icaia* and *Strychnos ignatii* extracts by $^1\text{H}$ nuclear magnetic resonance spectrometry and multivariate analysis techniques**

Frederich, M.; Choi, Y.H.; Angenot, L.; Harnischfeger, G.; Lefeber, A.W.M.; Erkelens, C.; Verpoorte, R.

### **Citation**

Frederich, M., Choi, Y. H., Angenot, L., Harnischfeger, G., Lefeber, A. W. M., Erkelens, C., & Verpoorte, R. (2004). Metabolomic analysis of *Strychnos nux-vomica*, *Strychnos icaia* and *Strychnos ignatii* extracts by  $^1\text{H}$  nuclear magnetic resonance spectrometry and multivariate analysis techniques. *Phytochemistry*, 65(13), 1993-2001.  
doi:10.1016/j.phytochem.2004.06.015

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/4107143>

**Note:** To cite this publication please use the final published version (if applicable).

# Metabolomic analysis of *Strychnos nux-vomica*, *Strychnos icaja* and *Strychnos ignatii* extracts by $^1\text{H}$ nuclear magnetic resonance spectrometry and multivariate analysis techniques

Michel Fr  d  rich <sup>a,b</sup>, Young Hae Choi <sup>a</sup>, Luc Angenot <sup>b</sup>, Goetz Harnischfeger <sup>c</sup>,  
Alfons W.M. Lefeber <sup>d</sup>, Robert Verpoorte <sup>a,\*</sup>

<sup>a</sup> Division of Pharmacognosy, Section Metabolomics, Institute of Biology, Leiden University, P.O. Box 9502, NL-2300 RA, Leiden, The Netherlands

<sup>b</sup> Laboratory of Pharmacognosy, Natural and Synthetic Drug Research Center, University of Li  ge, Avenue de l'H  pital, 1, B36, B-4000 Li  ge, Belgium

<sup>c</sup> Schafer and Br  mmer GmbH and Co KG, D-38251 Salzgitter, Germany

<sup>d</sup> Division of NMR, Institute of Chemistry, Gorlaeus Laboratories, P.O. Box 9502, NL-2300 RA, Leiden, The Netherlands

Received 26 January 2004; received in revised form 2 June 2004

Available online 20 July 2004

## Abstract

$^1\text{H}$  Nuclear magnetic resonance spectrometry and multivariate analysis techniques were applied for the metabolic profiling of three *Strychnos* species: *Strychnos nux-vomica* (seeds, stem bark, root bark), *Strychnos ignatii* (seeds), and *Strychnos icaja* (leaves, stem bark, root bark, collar bark). The principal component analysis (PCA) of the  $^1\text{H}$  NMR spectra showed a clear discrimination between all samples, using the three first components. The key compounds responsible for the discrimination were brucine, loganin, fatty acids, and *Strychnos icaja* alkaloids such as icajine and sungucine. The method was then applied to the classification of several "false angostura" samples. These samples were, as expected, identified as *S. nux-vomica* by PCA, but could not be clearly discriminated as root bark or stem bark samples after further statistical analysis.

   2004 Published by Elsevier Ltd.

**Keywords:** Loganiaceae; *Strychnos nux-vomica*; *Strychnos ignatii*; *Strychnos icaja*; Strychnine; Brucine; Metabolic profiling;  $^1\text{H}$  NMR; Multivariate analysis; PCA; Alkaloid

## 1. Introduction

*Strychnos* is a genus of small trees and climbing shrubs, members of the family Loganiaceae, distributed throughout Asia, Australia, Africa and South-America. In fact, among about 190 species, only seven would contain strychnine (*Strychnos nux-vomica* L. (Baser and Bisset, 1982; Bisset, 1976; Gadi-Biala et al., 1996; Se  covic et al., 1968), *Strychnos ignatii* P. Bergius (Bisset and Walker, 1974; Bratati and Bisset, 1990) and *Strychnos wallichiana* Steud ex DC (Bisset, 1976; Bisset and Choudhury, 1974) from Asia, *Strychnos lucida* R. Br (Bisset, 1976; Shaw and de la Lande, 1948) from

Australia, *Strychnos icaja* Baillon (Denoel, 1950; Kam-bu et al., 1980) from Africa, *Strychnos tabescana* and *Strychnos panamensis* (Krukoff et al., 1972; Marini-Bettolo et al., 1972) from South-America). In this paper, we have investigated the three most common species: *S. nux-vomica*, *S. ignatii* and *S. icaja* (Table 1). *S. nux-vomica* L. (Loganiaceae) is a tree from South-East Asia whose seeds have been used for the extraction of strychnine (**1**) and brucine (**2**) (Fig. 1), as well as in the preparation of galenicals. Strychnine (**1**) is a particularly toxic bulbar and medullar stimulant (LD<sub>50</sub> 0.2 mg/kg, parenteral) that has been intensively used both as a rat poison, as well as a tonic. The extracts of *S. nux-vomica* seeds are still used in allopathy (essentially in China and India) for a wide variety of disorders, including epilepsy, digestive disorders and debility (Ambasta, 1986; Dymock et al., 1891) and in homeopathy (in Western

\* Corresponding author. Tel.: +31-71-527-4528; fax: +31-71-527-4511.

E-mail address: [verpoort@chem.leidenuniv.nl](mailto:verpoort@chem.leidenuniv.nl) (R. Verpoorte).

Table 1  
*Strychnos* species and organs used for chemical fingerprinting by  $^1\text{H}$  NMR

Species	Organ	Origin	Batch code	Number of samples
<i>S. nux-vomica</i>	Seeds	Commercial	NV02	4
<i>S. nux-vomica</i>	Seeds	Commercial	NV04	4
<i>S. nux-vomica</i>	Stem bark	Chennai, India	NV01	3
<i>S. nux-vomica</i>	Stem bark	University Göttingen collections	NV05	3
<i>S. nux-vomica</i>	Bark	Falsification of angostura bark	NV08	3
		University Göttingen collections		
<i>S. nux-vomica</i>	Bark	Falsification of angostura bark	NV09	3
		University Göttingen collections		
<i>S. nux-vomica</i>	Bark	Falsification of angostura bark	NV10	3
		University Göttingen collections		
<i>S. nux-vomica</i>	Root bark	West Bengal, India	NV03	4
<i>S. ignatii</i>	Seeds	Commercial	IGN02	3
<i>S. ignatii</i>	Seeds	Commercial	IGN03	3
<i>S. icaja</i>	Leaves	Kikwit, Congo	ICA12	3
<i>S. icaja</i>	Leaves	Mukila, Congo	ICA13	3
<i>S. icaja</i>	Stem bark	Kikwit, Congo	ICA05	4
<i>S. icaja</i>	Collar bark	Kasongo-Lunda, Congo	ICA04	3
<i>S. icaja</i>	Root bark	Mukila, Congo	ICA03	3
<i>S. icaja</i>	Root bark	Kikwit, Congo	ICA14	3




Fig. 1. Chemical structure of strychnine and brucine.

countries). *S. ignatii* P. Bergius is a large shrub, up to 20-m high, found in Thailand, South-China, and the Philippines. The plant is used as a remedy for a number of ailments (Ambasta, 1986; Bisset, 1974; Dymock et al., 1891). *S. icaja* is a shrub that can become an important liana, up to 100-m long, and which is found all over central Africa. Its roots have a characteristic red outer bark. The plant is mainly known for its use as a hunting or ordeal (trial) poison. Nevertheless, the use of this *Strychnos* in traditional medicine, in particular, to treat chronic malaria, has also been reported (Iwu, 1993; Neuwinger, 1996). Recently, several anti-plasmodial bisindole alkaloids have been isolated from *S. icaja* roots, again drawing interest to this plant (Frédérich et al., 2001, 2000; Philippe et al., 2003).

The components of the metabolome can be viewed as the end products of gene expression and define the biochemical phenotype of a cell or tissue. Metabolic profiling could be used in a number of areas to provide biological information beyond the simple identification of plant constituents (Sumner et al., 2003). Among these areas is the chemical fingerprinting of species and plant organs for identification or taxonomic purposes. The method commonly used by industry for quality control of plant material is analysing the product for the presence of chemical markers known to be present in the plant. However, the presence of specific chemical makers does not always guarantee the identity of the plant material, especially if the product has been spiked with chemical markers. A better authentication of the plant material could be possible by a chemical fingerprinting of the botanical metabolites, implying a general overview of all plant constituents, and should be included into quality control methods (Bailey et al., 2002; Schaneberg et al., 2003). NMR associated to PCA could then also be used for chemotaxonomic analysis of plants, microbes and worms (Bundy et al., 2002).

In this work, high-resolution NMR spectroscopy was used to characterise the overall composition of three strychnine-containing species of *Strychnos*, aiming at identifying a wide range of metabolites. The principal aim was to explore the potential of  $^1\text{H}$  NMR spectroscopy method, coupled with multivariate analysis, to enable an efficient identification (metabolic fingerprinting) of the nature of *Strychnos* samples (species, organs). Secondly, the method was also applied to a series of samples of “false angostura bark”. The angostura bark (cortex *Angosturae*, *Galipea officinalis* St. Hilaire, Rutaceae) comes from a northern South American tropical

tree, and is used in traditional medicine to treat dysenteries, paralytic infections, and as a tonic and to make bitter liquors (Jacquemond-Collet et al., 1999; Tschirch, 1923). At first appearance, this bark resembles the poisonous bark of the *nux-vomica* tree. So it was not previously uncommon (early 19th century) to find portions of the latter mixed with angostura bark. This proved to be a most dangerous adulteration, causing several deaths between 1804 and 1806 in Germany, Switzerland and Hungary. As a result, *true* angostura fell into disfavour, (the *nux-vomica* being called *false* angostura) (Fl  ckiger and Hanbury, 1878; Grieve, 1971; Tschirch, 1923). We have analysed here four different samples of false angostura bark, to check if they were classified amongst *S. nux-vomica* samples.

## 2. Results and discussion

### 2.1. Extraction and analysis of plants

The extraction method used was very simple, requiring direct extraction of powdered, freeze-dried material by methanol-*d*<sub>4</sub> for NMR. The use of a deuterated solvent for extraction eliminated the need for evaporation and re-dissolution of extracts, which has the associated potential problem of loss of material and risk of chemical modifications. Trifluoroacetic acid was added just before NMR measurements to control the acidity and minimise the risks of signal shifting for acidic or basic compounds (alkaloids). Minimal shifts of signals between replicates are indeed essential to realise metabolic profiling. Three replicates were measured for each plant material studied.

### 2.2. Visual inspection of <sup>1</sup>H NMR spectra

Some differences were observed between the spectra of the various samples. In general, the <sup>1</sup>H NMR spectra obtained showed a dominance of signals in the carbohydrate (  2.5–  4.5) and, for seeds, in the aliphatic region (  0.5–  1.5) of the spectra. In addition to these signals, well-defined (but smaller) signals were present in the aromatic region of the spectra, and essentially attributed to alkaloids, caffeic acid, and loganin (seeds). Some differences could be visually observed between the different spectra. Strychnine was present in all extracts except *S. icaja* leaves, but the amount of the compound was extremely variable according to the species and batches: major alkaloids present in *S. nux-vomica* seeds or in *S. ignatii* seeds, but only present in traces in *S. icaja* leaves (unidentified in crude extract). Brucine was absent from *S. icaja* and very abundant in *S. nux-vomica* roots and in “false angostura” bark. Different substances were identified in the extracts. These identified substances are listed in Table 2. Signals were assigned using reference

Table 2  
Metabolites detected and identified from the <sup>1</sup>H NMR spectra of various crude extracts

Compounds	<i>S. nux-vomica</i> seeds	<i>S. nux-vomica</i> stem bark	<i>S. nux-vomica</i> “false angostura” root bark	<i>S. nux-vomica</i> root bark	<i>S. ignatii</i> seeds	<i>S. icaja</i> stem bark	<i>S. icaja</i> bark	<i>S. icaja</i> collar bark	<i>S. icaja</i> leaves
�-Colubrine									
�-Colubrine									
Brucine	+	+	+	+	+	+	+	+	
Caffeic acid ester	+	+	+	+	+	+	+	+	
Epoxy-novacine									
Fatty acids	+	+							
Fatty acids (unsaturated)	+								
Hydroxy-epoxy-novacine									
Icajine									
Isosungucine									
Loganin	+				+				
Novacine	+				+				
Strychnine	+	+			+	+	+	+	
Sucrose	+	+			+	+	+	+	
Sungucine									
Vomicine	+								

compounds and various 2D-NMR techniques (COSY, TOCSY, HSQC, and HMBC).

As chlorogenic acid is generally described as present in *S. nux-vomica*, we then compared our extracts with an authentic sample of chlorogenic acid by NMR, TLC and HPLC. The presence of a close compound showing the same fluorescence and close retention factor on TLC (probably an isomer of chlorogenic acid) was observed, but chlorogenic acid itself was not present. It was impossible to clearly identify this compound because of the lack of reference compounds and the crowded NMR spectra in the sugar part. Although there were clear visual differences between the spectra, for easier and non-biased interpretation of the results, and to reduce the dimensionality of the multivariate data obtained with the NMR results, we analysed the samples using principal component analysis (PCA) (see Table 3).

### 2.3. Principal component analysis of $^1\text{H}$ NMR spectra

PCA is an unsupervised method requiring no knowledge of the data set and which acts to reduce the dimensionality of multivariate data while preserving most of the variance within it (Eriksson et al., 2001). If the data are mean-centred with no scaling, then a covariance matrix is produced. But if the data are mean-centred, and the columns of the data matrix are scaled to unit variance, a correlation matrix is produced. An advantage of the covariance matrix is that the loadings retain the scale of the original data. For the correlation method, however, a weaker signal possessing a discriminatory power can be considered at the same level as stronger signals (Ward et al., 2003). In this study, both

methods were evaluated and the covariance method showed better separation results (Fig. 2). For the data set obtained from the analysis of *Strychnos* extract, a nine-component model explained 99% of the variance, with the first three components explaining 85.4%. As observed in Fig. 2(a), there is a clear discrimination possible between *S. nux-vomica* seeds and other samples, particularly *S. ignatii* seeds. The extracts of *S. ignatii* seeds, however, show quite similar metabolic patterns to those of *S. icaja* leaves. The *S. icaja* stem bark, collar bark and root bark are not easily distinguishable but the leaves present a clearly different metabolic content.

The seeds of *S. nux-vomica* are clearly separated from other samples by PC1 (Fig. 2(a)). The seeds of *S. nux-vomica* are located in the higher PC1 region. Lipids play an important role in this discrimination. The discriminating metabolites are distinguishably shown in loading plot of PC1 and PC2 (Fig. 3). It is obviously postulated that the seeds of *S. nux-vomica* contain higher amount of lipids and the major part of the variability of PC1 is explained by them and also by loganin ( $\delta$  7.38), also quite common in seeds (Fig. 3(a)). PC2 is essentially influenced by loganin at  $\delta$  7.36 and various *S. icaja* alkaloids such as icajine at  $\delta$  8.06 and sungucine at  $\delta$  8.34. For more separation, PC3 was additionally used (Fig. 4). In this case, the root extracts of *S. nux-vomica* are obviously separated from other samples by PC3 (Fig. 4(a)) but the differentiating metabolite was identified by loading plot of PC3 and found to be brucine almost alone (Fig. 4(b)). Two positive distinguishable  $^1\text{H}$  NMR signals in the loading plot of PC3 were identified as H-12 and H-9 of brucine, respectively.

The data from the false angostura samples were then added to these results. They were plotted in Fig. 5. As expected, these samples were plotted close to the *S. nux-vomica* samples, but more particularly, close to the *S. nux-vomica* root bark samples (Fig. 5(a)). For confirmation, ANOVA test was performed to PC1, PC2, and PC3 of *S. nux-vomica* samples (roots and stem barks) and false angostura. Unfortunately, the PC scores are not significantly different. The discriminating metabolites in the plot containing false angostura samples are distinguishably shown in Fig. 5(b) and (c) (loading plots of PC1 and PC2) and are identical to those previously described, although *S. icaja* alkaloids have here a negative influence on PC2.

### 2.4. Conclusions

We showed that it was possible to discriminate three different *Strychnos* species from various origins by multivariate analysis of  $^1\text{H}$  NMR spectra of crude extracts. Seeds of *S. ignatii* and *S. nux-vomica* are easily distinguishable. *S. icaja* and *S. nux-vomica* stem barks are much more similar, but could be distinguished using the brucine content (mostly influencing PC3). Finally,

Table 3  
Important chemical shifts influencing the PCA discrimination

PC	Chemical shift	Corresponding compound and signal
PC1	0.88	Fatty acids $\text{CH}_3$
PC1	0.90	Fatty acids $\text{CH}_3$
PC1	1.10	Loganin H10
PC1	1.32	Fatty acids $\text{CH}_2$
PC1	1.60	Fatty acids $^*\text{CH}_2\text{—CH}_2\text{—CO}$
PC1	2.02	Unsaturated fatty acids ( $^*\text{CH}_2\text{—CH=CH}$ )
PC1	2.26	Fatty acids $\text{CH}_2\text{—}^*\text{CH}_2\text{—CO}$
PC1, PC2, PC3	3.84	Brucine methoxy
PC1	4.18	Unidentified signal
PC1	5.34	Unsaturated fatty acids ( $\text{CH}_2\text{—}^*\text{CH=CH}$ )
PC1, PC2, PC3	7.02	Brucine H9
PC2	7.36	Sungucine + isosungucine + strychnine
PC1	7.38	Loganin H3
PC1, PC2, PC3	7.76	Brucine H12
PC2	8.06	Icajine H12
PC2	8.34	Sungucine H12'




Fig. 2. Score plot of principal component analysis of *Strychnos* extracts obtained by covariance (a) and correlation (b) method using PC1 and PC2. The ellipse represents the Hotelling T2 with 95% confidence in score plots. (1) Stem barks of *S. nux-vomica*, (2) seeds of *S. nux-vomica*, (3) roots of *S. nux-vomica*, (4) stem barks of *S. icaja*, (5) root barks of *S. icaja*, (6) collars of *S. icaja*, (7) leaves of *S. icaja*, (8) seeds of *S. ignatii*.

we analysed four collection samples from “false angostura”, which is a falsification of angostura bark (*Galipea officinalis*). In the literature, from the 19th and early 20th century, the false angostura was generally described as “bark from *S. nux-vomica* imported from India”, without more precision, the real origin of the samples remaining quite confused (Felter and Lloyd, 1898; Fl  ckiger and Hanbury, 1878; Grieve, 1971; Tschirch, 1923). According to our PCA results, the false angostura bark should be classified as an *S. nux-vomica* root bark sample. This classification is quite plausible, as stem and root barks of *S. nux-vomica* are still used in traditional medicine in India (Ambasta, 1986). Further statistical analysis (ANOVA) could not show significant differences between PCs of false angostura samples and

root bark or stem bark of *S. nux-vomica*. To confirm this point, as score plots of root and stem bark of *S. nux-vomica* were quite close, it would be useful to further analyse authenticated root bark and stem bark samples from *S. nux-vomica* (and samples from different origins). It is likely that the different false angostura samples, clearly plotted as *S. nux-vomica* in our PCA analysis, do not have a homogeneous origin (some could be stem bark and others root bark).

We then showed that the major compounds responsible for the discrimination were brucine, fatty acids, loganin and several *S. icaja* alkaloids (mainly icajine and sungucine). Strychnine, though present in various amounts in all extracts analysed, was not the key compound for the discrimination of samples.




Fig. 3. Loading plot of principal component analysis of *Strychnos* extracts obtained by covariance method. (a) PC1, (b) PC2.

### 3. Experimental

#### 3.1. Solvents and chemicals

Methanol- $d_4$  (99.80% D) and deuterated trifluoroacetic acid were purchased from Eurisotop (Gif-sur-Yvette, France). Trifluoroacetic acid, strychnine, brucine, caffeic acid, and chlorogenic acid were purchased from Aldrich (Steinheim, Germany). Methanol was from Merck Biosolve Ltd. (Valkenswaard, The Netherlands).

#### 3.2. NMR measurements

All spectra were recorded on a Bruker AV-400 NMR spectrometer operating at a proton NMR frequency of

400.13 MHz. For each sample, 256 scans were recorded with the following parameters: 0.126 Hz/point, pulse width (PW) = 30° (4.0  $\mu$ s) and relaxation delay (RD) = 1.0 s. FIDs were Fourier transformed with LB = 0.5 Hz and GB = 0. The spectra were referenced to solvent.

#### 3.3. Extraction of alkaloids

The powdered plant material (100.0 mg) was freeze-dried during 24 h and then extracted with 1.0 ml of methanol- $d_4$  under ultrasonication for 1 h at room temperature. The mixture was then centrifuged at 13,000 g for 10 min and 750  $\mu$ l of supernatant was mixed with trifluoroacetic acid (1%, v/v) for NMR measurement.




Fig. 4. Score plot of principal component analysis of *Strychnos* extracts obtained by covariance method using PC1 and PC3 (a) and loading plot of PC3 (b). The ellipse represents the Hotelling T2 with 95% confidence in score plots. (1) Stem barks of *S. nux-vomica*, (2) seeds of *S. nux-vomica*, (3) roots of *S. nux-vomica*, (4) stem barks of *S. icaja*, (5) root barks of *S. icaja*, (6) collars of *S. icaja*, (7) leaves of *S. icaja*, (8) seeds of *S. ignatii*.

### 3.4. Plant material

*S. nux-vomica* seeds and *S. ignatii* seeds were commercial samples obtained from Denolin (Braine l'Alleud, Belgium) and Longeval (Deux Acres, Belgium). Seeds of *S. nux-vomica* were authenticated according to the description in the Swiss and French Pharmacopoeia and those of *S. ignatii* following the microscopic description in Perrot and Gathercoal (Gathercoal and Wirth, 1947; Perrot, 1943). The samples of *S. icaja* were collected near Kasongo-Lunda, Mulika and Kikwit (Congo-Zaire). Voucher specimens of the plants (Duvigneaud H787, H864 and H900) have been deposited in the herbarium of the Pharmaceutical Institute in Liège,

and in the herbarium of the Belgian National Botanical Garden, in Meise. The samples of “false angostura bark” NV06 came from the laboratory collections and were identified by the late Professor Bisset (King's College, University of London.). The *S. nux-vomica* stems were collected in Chennai (India) by Phn. A. Kone (NV01, stems). A voucher specimen (NV01) of this plant has been deposited in the herbarium of the Pharmaceutical Institute in Liège. *S. nux-vomica* root barks (NV03) were collected at Midnapur, West Bengal, India, and identified by the late Professor N.G. Bisset (King's College, University of London). A voucher sample (NGB 23289) has been deposited in the Pharmaceutical Institute, University of Liège, Belgium. *S. nux-vomica*






Fig. 5. Score plot of principal component analysis of *Strychnos* and “false angostura” extracts obtained by covariance method using PC1 and PC2 (a), and loading plot of PC1 (b) and PC2 (c). The ellipse represents the Hotelling T2 with 95% confidence in score plots. (1) Stem barks of *S. nux-vomica*, (2) seeds of *S. nux-vomica*, (3) roots of *S. nux-vomica*, (4) stem barks of *S. icaja*, (5) root barks of *S. icaja*, (6) collars of *S. icaja*, (7) leaves of *S. icaja*, (8) seeds of *S. ignatii*, (9) stem barks of “false angostura” stems.

stem bark NV05 and the “false angostura” barks NV08, NV09, NV10 came from the collections of the University of G  ttingen (Herbarium Goet, Germany, Phar-

makognostische Sammlung 12/2001 Nos. 3294, 1112, 2670, and 3280, respectively). All species and their origin are reported in Table 1.

### 3.5. Data analysis

The  $^1\text{H}$  NMR spectra were automatically reduced to ASCII files using AMIX (v. 3.7, Bruker Biospin). Spectral intensities were scaled to 0.01% (v/v) of HMDS and reduced to integrated regions of equal width (0.02 ppm) corresponding to the region of  $\delta$  0.40– $\delta$  10.00. The region of  $\delta$  4.75– $\delta$  5.24 of HDO,  $\delta$  3.55– $\delta$  3.65 of residual ethanol,  $\delta$  3.25– $\delta$  3.50 of residual MeOH,  $\delta$  2.14– $\delta$  2.15 of residual acetone, and  $\delta$  1.14– $\delta$  1.19 of residual ethanol were removed for further analysis. PCA and ANOVA analysis were performed with the SIMCA-P software (Umetrics, Ume  , Sweden).

### 3.6. Assignment of peaks

To assign selected peaks, various two dimensional NMR methods such as  $^1\text{H}$ – $^1\text{H}$  COSY,  $^1\text{H}$ – $^{13}\text{C}$  COSY (Hetcor), and  $^1\text{H}$ – $^{13}\text{C}$  long range coupling (HMBC) were applied.

## Acknowledgements

The Post-doctoral Fellowship Program of the Korea Science Engineering Foundation (KOSEF) and the Belgian National Fund for Scientific Research (FNRS) are gratefully acknowledged for their support of Y.H. Choi and M. Fr  d  rich, respectively. The authors are indebted to the keeper of the herbarium Goet, Albrecht-von Haller-Institut f  r Pflanzenwissenschaften, Abteilung Systematische Botanik, G  ttingen for permission to take samples from the collections.

## References

- Ambasta, S.S.P., 1986. The Useful Plants of India. Publications and Information Directorate, New Delhi, pp. 604–606.
- Bailey, N.J.C., Sampson, J., Hylands, P.J., Nicholson, J.K., Holmes, E., 2002. Multi-component metabolic classification of commercial feverfew preparations via high-field  $^1\text{H}$  NMR spectroscopy and chemometrics. *Planta Medica* 68, 734–738.
- Baser, K.H.C., Bisset, N.G., 1982. Alkaloids of Sri Lankan *Strychnos nux-vomica*. *Phytochemistry* 21, 1423–1429.
- Bisset, N.G., 1974. The Asian species of *Strychnos*. Part III. The ethnobotany. *Lloydia* 37, 62–107.
- Bisset, N.G., 1976. The Asian species of *Strychnos*. Part IV. The alkaloids. *Lloydia* 39, 263–315.
- Bisset, N.G., Choudhury, A.K., 1974. Alkaloids from the leaves of *Strychnos wallichiana*. *Phytochemistry* 13, 259–263.
- Bisset, N.G., Walker, M.D., 1974. Alkaloids from the stem bark of *Strychnos ignatii*. *Phytochemistry* 13, 525–526.
- Bratati, D., Bisset, N.G., 1990. Alkaloids of *Strychnos ignatii*. *Planta Medica* 56, 133.
- Bundy, J.G., Spurgeon, D.J., Svendsen, C., Hankard, P.K., Osborn, D., Lindon, J.C., Nicholson, J.K., 2002. Earthworm species of the genus *Eisenia* can be phenotypically differentiated by metabolic profiling. *FEBS Letters* 521, 115–120.
- Denoel, A., 1950. Le *Strychnos icaja* Baillon du Congo Belge. *Journal de Pharmacie de Belgique* 5, 59–77.
- Dymock, W., Warden, C.J.H., Hooper, D., 1891. *Pharmacographia Indica. A History of the Principal Drugs of Vegetable Origin*. Trench, Tr  bner & Co, London, pp. 458–505.
- Eriksson, L., Johansson, E., Kettaneh-Wold, N., Wold, S., 2001. *Multi- and Megavariable Data Analysis*. Umetrics Academy, Ume  , Sweden.
- Felter, H.W., Lloyd, J.U., 1898. *King's American Dispensatory*. Ohio Valley Co., Cincinnati.
- Fl  ckiger, F.A., Hanbury, D., 1878. *Histoire des Drogues d'Origine V  g  tale*. Octave Doin, Paris.
- Fr  d  rich, M., De Pauw, M., Prosperi, C., Tits, M., Brandt, V., Penelle, J., Hayette, M., De Mol, P., Angenot, L., 2001. Strychnogucines A and B, two new antiparasitoid bisindole alkaloids from *Strychnos icaja*. *Journal of Natural Products* 64, 12–16.
- Fr  d  rich, M., Depauw-Gillet, M.C., Llabres, G., Tits, M., Hayette, M.P., Brandt, V., Penelle, J., De Mol, P., Angenot, L., 2000. New anti-malarial and cytotoxic sanguine derivatives from *Strychnos icaja* roots. *Planta Medica* 66, 262–269.
- Gadi-Biala, R., Tits, M., Wauters, J.N., Angenot, L., 1996. A new HPLC method for the assay of alkaloids in *Strychnos nux-vomica* and *Strychnos ignatii*. *Fitoterapia* 67, 163–165.
- Gathercoal, E.N., Wirth, E.H., 1947. *Pharmacognosy*. Lea and Fegiger, Philadelphia.
- Grieve, M., 1971. *A Modern Herbal*, vol. 1, Dover Publications, New York, pp. 51–53.
- Iwu, M., 1993. *Handbook of African Medicinal Plants*. CRC Press, Boca Raton, pp. 244–245.
- Jacquemond-Collet, I., Hannedouche, S., Fabre, N., Fouraste, I., Moulis, C., 1999. Two tetrahydroquinoline alkaloids from *Galipea officinalis*. *Phytochemistry* 51, 1167–1169.
- Kambu, K., Coune, C., Angenot, L., 1980. Nouveaux alcalo  ides des racines du *Strychnos icaja*. *Planta Medica* 37, 161–164.
- Krukoff, B.A., Marini-Bettolo, G.B., Bisset, N.G., 1972. American species of *Strychnos*. *Lloydia* 35, 193–271.
- Marini-Bettolo, G.B., Ciasca, M.A., Galeffi, C., Bisset, N.G., Krukoff, B.A., 1972. The occurrence of strychnine and brucine in an American species of *Strychnos*. *Phytochemistry* 11, 381–384.
- Neuwinger, H.D., 1996. *African Ethnobotany: Poisons and Drugs, Chemistry, Pharmacology, Toxicology*. Chapman & Hall, London.
- Perrot, E., 1943. *Mat  res Premi  res Usuelles du R  gne V  g  tal*. Masson, Paris.
- Philippe, G., De Mol, P., Zeches-Hanrot, M., Nuzillard, J.M., Tits, M.H., Angenot, L., Fr  d  rich, M., 2003. Indolomonoterpenic alkaloids from *Strychnos icaja* roots. *Phytochemistry* 62, 623–629.
- Schaneberg, B.T., Crockett, S., Bedir, E., Khan, I.A., 2003. The role of chemical fingerprinting: application to *Ephedra*. *Phytochemistry* 62, 911–918.
- Sefcovic, P., Dubravkova, L., Torto, F.G., 1968. Die Alkaloide der Bl  tter von *Strychnos nux-vomica* L. *Planta Medica* 16, 143–146.
- Shaw, F.H., de la Lande, J.S., 1948. The alkaloids of *Strychnos lucida*. *Australian Journal of Experimental Biological and Medicinal Sciences* 26, 199–203.
- Sumner, L.W., Mendes, P., Dixon, R.A., 2003. Plant metabolomics: large-scale phytochemistry in the functional genomics era. *Phytochemistry* 62, 817–836.
- Tschirch, A., 1923. *Handbuch der Pharmakognosie*. Verlag von Chr. Herm. Tauchnitz, Leipzig.
- Ward, J., Harris, C., Lewis, J., Beale, H., 2003. Assessment of  $^1\text{H}$  NMR spectroscopy and multivariate analysis as a technique for metabolite fingerprinting of *Arabidopsis thaliana*. *Phytochemistry* 62, 949–957.