



Universiteit
Leiden

The Netherlands

Advances in cardiac computed tomography angiography and characterization of cardiac disease

Kuneman, J.H.

Citation

Kuneman, J. H. (2024, December 12). *Advances in cardiac computed tomography angiography and characterization of cardiac disease*. Retrieved from <https://hdl.handle.net/1887/4172200>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4172200>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 8

Prognostic implications of change in left ventricular ejection fraction after transcatheter aortic valve implantation

Kuneman JH
Butcher SC
Singh GK
Wang X
Hirasawa K
van der Kley F
Leon MB
Knuuti J
Pibarot P
Ajmone Marsan A
Delgado V
Bax JJ

Am J Cardiol. 2022;177:90-99.



ABSTRACT

BACKGROUND: Reduced left ventricular (LV) systolic function is associated with worse prognosis in patients with severe aortic stenosis (AS) treated with transcatheter aortic valve implantation (TAVI). We aimed to examine the changes in LV ejection fraction (LVEF) after TAVI among patients with varying baseline LVEF. Moreover, variables associated with lack of LVEF improvement were identified. Finally, the association with long-term outcomes was investigated.

METHODS: Patients with severe AS who underwent transfemoral TAVI between 2007-2019 were selected. LVEF was assessed from transthoracic echocardiography at baseline (before TAVI) and at 6 and 12 months after TAVI. Patients were stratified according to baseline LVEF: 1) LVEF $\geq$ 50%, 2) LVEF 40%-49%, and 3) LVEF<40%. The clinical endpoint was $\geq$ 5% LVEF improvement. The primary outcome was all-cause mortality.

RESULTS: A total of 560 patients (age 80 ± 7 years, 53% male) were included. Patients with baseline LVEF<40% showed greater increase in LVEF compared to those with LVEF 40-49% and baseline LVEF $\geq$ 50% (from $33\pm 6\%$ to $43\pm 10\%$, $p<0.001$; from $45\pm 3\%$ to $52\pm 8\%$, $p<0.001$; and from $58\pm 5\%$ to $59\pm 7\%$, $p=0.012$, respectively, p for interaction <0.001). Coronary artery disease (CAD, OR: 1.80 (95%CI 1.06-3.06), $p=0.031$), myocardial infarction (OR: 2.07 (95%CI 1.19-3.61), $p=0.010$), and permanent pacemaker (OR: 1.93 (1.25-3.00), $p=0.003$) were independently associated with lack of $\geq$ 5% LVEF improvement. During a median follow-up of 3.8 [IQR 2.6-5.2] years, 176 patients died (31%). Patients with $\geq$ 5% LVEF improvement had similar outcomes compared to those with <5% LVEF improvement (log-rank $p=0.89$).

CONCLUSION: Patients with severe AS and baseline LVEF<40% had greatest improvement in LVEF at 1-year follow-up after TAVI. CAD, myocardial infarction, and permanent pacemaker were associated with lack of LVEF improvement. However, LVEF improvement at 12 months was not associated with long-term outcomes.

INTRODUCTION

Severe aortic stenosis (AS) may have a significant impact on left ventricular (LV) function. Increased afterload caused by AS may lead to LV remodeling resulting in LV hypertrophy and impaired LV function^{1,2}. Approximately one-third of patients with severe AS present with LV systolic dysfunction^{3,4}. Moreover, LV systolic dysfunction has been linked with worse prognosis in patients with severe AS who were treated medically or underwent surgical aortic valve replacement⁵⁻⁷. Transcatheter aortic valve implantation (TAVI) has become an alternative to surgical aortic valve replacement in patients with severe symptomatic AS⁸. Similarly, reduced baseline LV ejection fraction (LVEF) is associated with an increased risk of all-cause and cardiovascular mortality after TAVI⁹⁻¹¹. LVEF is an important parameter for the assessment of LV systolic function and is relevant in the management of patients with asymptomatic severe AS according to current guidelines^{12,13}. Previous studies showed that LV systolic function improved after TAVI in patients with reduced LVEF¹⁴⁻¹⁷. However, data on the change of LVEF after TAVI in patients with varying LVEF at baseline are scarce. In addition, the impact of LVEF improvement on outcomes after TAVI is not clear. Accordingly, the aims of this study were to investigate the potential improvement of LVEF after TAVI in patients with preserved baseline LVEF (LVEF $\geq 50\%$), midrange LVEF (LVEF 40-49%), and reduced LVEF (LVEF $< 40\%$). Moreover, the variables associated with lack of LVEF improvement were identified. Finally, the association between LVEF improvement with long-term outcomes after TAVI was evaluated.

METHODS

Study population

Patients with severe AS treated with TAVI at the Leiden University Medical Center (Leiden, the Netherlands) between 2007 and 2019 were included in this retrospective analysis. Patients who underwent transapical TAVI or a valve-in-valve procedure were excluded, or if echocardiographic data at baseline and/or follow-up was lacking. All patients underwent standard routine follow-up transthoracic echocardiography. Patients were stratified according to baseline (pre-TAVI) LVEF as appointed in the 2016 European Society of Cardiology heart failure guidelines¹⁸: patients with preserved LVEF (LVEF $\geq 50\%$), midrange LVEF (LVEF 40-49%), and reduced LVEF (LVEF $< 40\%$). Demographic and clinical characteristics at the time of TAVI were collected from the electronic patient records (EPD-vision, Leiden University Medical Center, Leiden, the Netherlands). Clinical characteristics included comorbidities, cardiac risk factors, previous coronary

revascularization, and medication use. The estimated glomerular filtration rate (eGFR) was calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formula and renal impairment was defined as an eGFR <60 mL/min/1.73 m². This retrospective analysis complies with the Declaration of Helsinki and was approved by the institutional review board which waived the need for written informed consent due to the retrospective study design.

Transcatheter aortic valve implantation

TAVI eligibility and feasibility as well as decision-making on the access route and valve type were decided by the local heart team. Multidetector-row computed tomography measurements of the aortic annulus were used to select the transcatheter heart valve size, as previously described¹⁹. The TAVI procedure was performed according to standard practice²⁰. Balloon- and self-expandable valves that were used include Edwards SAPIEN, SAPIEN XT, SAPIEN 3 (Edwards Lifesciences, Irvine, CA, USA), and Medtronic CoreValve Evolut R and Evolut Pro (Medtronic, MN, Minnesota, USA).

Echocardiography

Transthoracic 2-dimensional echocardiographic examinations were performed before TAVI (to evaluate AS severity, LV systolic function, and LV dimensions) and during routine follow-up: immediately after TAVI (prosthetic valve hemodynamics), and at 6 and 12 months follow-up (prosthetic valve hemodynamics, LV function, and LV dimensions). All echocardiographic examinations were acquired by experienced echocardiographers using commercially available ultrasound systems (Vivid-7, E9 or E95, General Electric Vingmed, Horten, Norway) equipped with 3.5 MHz or M5S transducers. The echocardiographic examinations were performed with patients at rest and data were obtained from the parasternal, apical, and subcostal views. All echocardiographic examinations were reported according to current recommendations^{21,22}. Echocardiographic analysis was performed offline using commercially available software (EchoPac version 113.0.3 and 203; GE Medical Systems, Horten, Norway). Peak and mean transvalvular gradients were calculated from continuous-wave Doppler recordings of the apical 3- or 5-chamber views according to the Bernoulli equation. Aortic valve area (AVA) was calculated using the continuity equation and indexed to body surface area (AVA_i). Severe AS was defined as an AVA <1.0 cm² or AVA_i <0.6 cm²/m² and a mean transvalvular pressure gradient ≥ 40 mmHg or peak aortic jet velocity ≥ 4 m/s at rest or during dobutamine stress echocardiography^{12,13}. LV volumes (end-diastolic and end-systolic) were measured using planimetry in the apical 2- and 4-chambers views and were indexed to body surface area. LVEF was calculated using Simpsons' biplane method. LV dimensions (end-diastolic

diameter, intraventricular septum thickness, and posterior wall thickness) were obtained in the parasternal long-axis views at end-diastole²². LV mass was calculated using the Devereux formula and indexed to body surface area²². The presence of post-procedural aortic regurgitation and paravalvular leakage was detected and severity was graded according to current recommendations: mild (grade 1), moderate (grade 2), moderate to severe (grade 3), and severe (grade 4)²³. Significant paravalvular leakage was defined by a grade ≥ 2 .

Clinical endpoints and follow-up

Changes in LVEF and LV volumes over time are presented as absolute numbers and LV volumes also as percentual reduction compared to baseline. Additional echocardiographic endpoints included the increase in LVEF $\geq 5\%$ and $\geq 10\%$ (as percent point) at follow-up. An increase of LVEF $\geq 5\%$ was the primary endpoint, as used previously²⁴. Variables linked with lack of $\geq 5\%$ LVEF improvement were identified. In addition, patients were followed up for the occurrence of all-cause mortality after TAVI and survival time was restricted to 5 years. Data on mortality were collected from the departmental electronic patient files which were linked with the Social Security Death Index and were acquired for all patients.

Statistical analysis

Continuous variables following a normal distribution are presented as mean $\pm$ standard deviation and were compared using the one-way ANOVA test. Non-normally distributed continuous variables are presented as median with 25%-75% interquartile range and were compared using the Kruskal Wallis test. Bonferroni's post-hoc analysis was performed to assess between-group differences in case of a significant difference in the overall 3 group comparison. Distribution of continuous variables was evaluated using histograms and Q-Q plots. Categorical variables are presented as absolute numbers and percentages and were compared using the χ^2 test or Fisher's exact test. General linear models with repeated measures analysis were used to evaluate changes in echocardiographic variables over time and to test differences between the groups of baseline LVEF. The Greenhouse-Geisser correction was used if the sphericity assumption was violated and Bonferroni's post-hoc analysis was performed in case of a significant difference in the overall 3 group comparison. Additional analyses were performed to correct for potential confounders (age, sex, hypertension, diabetes mellitus, coronary artery disease, previous myocardial infarction, concomitant moderate or severe mitral- or aortic regurgitation, moderate or severe paravalvular leakage, permanent pacemaker (before and after TAVI), atrial fibrillation, pre-TAVI AV mean gradient, and pre-TAVI LVEF)

on the change of the LV parameters over time, and were included as covariates in the general linear models²⁵. Uni- and multivariable binary logistic regression models were used to evaluate the variables that were associated with lack of $\geq 5\%$ LVEF improvement. All variables with a p-value < 0.10 in the univariable analysis were included in the multivariable model. Odds ratios (OR) and 95% confidence intervals (CI) were calculated and reported. Kaplan-Meier curves were generated to estimate the cumulative survival rates of all-cause mortality and the log-rank test was used to compare patients with $\geq 5\%$ LVEF improvement at follow-up. Multivariable Cox proportional hazards regression analysis was used to evaluate the association of improvement in LVEF (as continuous and dichotomous variables) at follow-up with all-cause mortality. Potential confounders including age, sex, cardiac risk factors, coronary artery disease, previous myocardial infarction, previous stroke/transient ischemic attack, peripheral vascular disease, chronic obstructive pulmonary disease, previous coronary revascularization, chronic kidney disease, pre-TAVI AV mean gradient, and pre-TAVI LVEF were incorporated in the multivariable Cox proportional hazard model. Hazard ratios (HR) and 95% CI were calculated and reported. A two-sided p-value < 0.05 was considered significant. Data analyses were performed with SPSS version 25.0 (IBM SPSS Statistics, IBM Corporation, Armonk, New York, USA).




FIGURE 1. Flowchart of the study population. AS = aortic stenosis, TAVI = transcatheter aortic valve implantation.

RESULTS

Patient and procedural characteristics

A total of 560 patients (age 80 ± 7 years, 53% male) with severe AS were included in the analysis (Figure 1). The distribution of the different groups of LV systolic function before TAVI and at 6 months and 12 months follow-up are shown in Figure 2. Before TAVI, 350 patients (62%) had preserved LVEF (LVEF $\geq 50\%$), 122 patients (22%) had midrange LVEF (LVEF 40–49%), and reduced LVEF (LVEF $<40\%$) was present in 88 patients (16%). Baseline (pre-TAVI) demographical and clinical characteristics of the overall population and according to baseline LVEF are presented in Table 1. Patients with reduced baseline LVEF had a more frequent history of myocardial infarction, NYHA class III–IV, concomitant moderate or severe mitral- or aortic regurgitation, and permanent pacemaker compared to patients with preserved and midrange baseline LVEF, whereas hypertension was less common. Accordingly, patients with reduced baseline LVEF were more frequently using diuretics, mineralocorticoid receptor antagonists, and oral anticoagulation and less calcium antagonists. In addition, patients with reduced baseline LVEF had a higher logistic EuroSCORE and worse kidney function compared to patients with midrange and preserved baseline LVEF. However, the frequency of reduced kidney function was comparable between the groups (50% vs. 52% vs. 45%, respectively, $p=0.42$).

TABLE 1. Baseline (pre-TAVI) demographical and clinical characteristics of the overall population and according to baseline left ventricular ejection fraction.

Variable	Overall population (n=560)	Left Ventricular Ejection Fraction $\geq 50\%$ (n=350)	Left Ventricular Ejection Fraction 40–49% (n=122)	Left Ventricular Ejection Fraction $<40\%$ (n=88)	p-value
Age (years)	80 ± 7	80 ± 7	80 ± 6	79 ± 9	0.46
Male sex	296 (53%)	178 (51%)	68 (56%)	50 (57%)	0.47
Body mass index (kg/m ²)	26.7 ± 4.6	26.7 ± 4.8	27.0 ± 4.2	26.4 ± 4.4	0.67
Logistic EuroSCORE	12.3 [8.6–20.1]	10.7 [7.9–17.1]	13.4 [9.5–21.6]*	21.9 [13.8–31.9]*†	<0.001
eGFR (ml/min)	61 ± 20	63 ± 19	$58 \pm 21^*$	$57 \pm 21^{*†}$	0.007
Hypertension	407 (73%)	266 (76%)	88 (72%)	53 (60%)	0.009
Hypercholesterolemia	338 (61%)	206 (59%)	80 (66%)	52 (59%)	0.43
Diabetes mellitus	159 (28%)	97 (28%)	35 (29%)	27 (31%)	0.86
Smoker	76 (14%)	47 (14%)	14 (12%)	15 (18%)	0.58
CAD	351 (63%)	208 (60%)	80 (66%)	63 (72%)	0.089
Previous myocardial infarction	112 (20%)	55 (16%)	30 (25%)	27 (31%)	0.003

Variable	Overall population (n=560)	Left Ventricular Ejection Fraction $\geq 50\%$ (n=350)	Left Ventricular Ejection Fraction 40-49% (n=122)	Left Ventricular Ejection Fraction $< 40\%$ (n=88)	p-value
Previous revascularization	293 (52%)	172 (49%)	67 (55%)	54 (61%)	0.099
PCI	173 (31%)	108 (31%)	36 (30%)	29 (33%)	0.22
CABG	120 (21%)	64 (18%)	31 (25%)	25 (28%)	
NYHA class					
I-II	231 (41%)	156 (45%)	50 (41%)	25 (28%)	0.020
III-IV	327 (59%)	192 (55%)	72 (59%)	63 (72%)	
Previous stroke / TIA	103 (19%)	62 (18%)	23 (19%)	18 (21%)	0.86
Peripheral vascular disease	100 (18%)	56 (16%)	24 (20%)	20 (23%)	0.28
Atrial fibrillation	147 (26%)	76 (22%)	42 (34%)	29 (33%)	0.007
Chronic obstructive pulmonary disease	101 (18%)	63 (18%)	17 (14%)	21 (24%)	0.19
Concomitant mitral / aortic regurgitation $\geq$ grade 2	148 (26%)	64 (18%)	45 (37%)	39 (44%)	<0.001
Permanent pacemaker	83 (15%)	37 (11%)	23 (19%)	23 (26%)	<0.001
Medication					
Beta-blocker	326 (58%)	207 (60%)	72 (59%)	47 (53%)	0.58
ACE-I / ARB II	290 (52%)	178 (51%)	61 (50%)	51 (58%)	0.46
Calcium antagonist	136 (24%)	81 (23%)	41 (34%)	14 (16%)	0.010
Diuretics	295 (53%)	159 (46%)	75 (62%)	61 (69%)	<0.001
MR antagonist	57 (10%)	27 (8%)	14 (12%)	16 (18%)	0.014
Statins	349 (63%)	222 (64%)	79 (65%)	48 (55%)	0.24
Antiplatelet	325 (58%)	204 (58%)	71 (58%)	50 (57%)	0.97
Anticoagulation	211 (38%)	111 (32%)	51 (43%)	49 (56%)	<0.001

Data are presented as mean $\pm$ SD, median [25-75% interquartile range] and n (%). ACE-I = angiotensin-converting enzyme, ARB II = angiotensin-II receptor blocker, CABG = coronary artery bypass grafting, CAD = coronary artery disease, CKD-EPI = chronic kidney disease epidemiology collaboration, eGFR = estimated glomerular filtration rate, MR = mineralocorticoid receptor, NYHA = New York Heart Association, PCI = percutaneous coronary intervention, TIA = transient ischemic attack.

* $p < 0.05$ vs group 'Left Ventricular Ejection Fraction $\geq 50\%$ ' with Bonferroni's post-hoc analysis.

† $p < 0.05$ vs group 'Left Ventricular Ejection Fraction 40-49%' with Bonferroni's post-hoc analysis

Baseline (pre-TAVI) echocardiographic data are presented in Table 2. LV volumes and LV mass index were larger in patients with reduced LVEF compared to those with midrange and preserved LVEF before TAVI. In contrast, AV peak and mean gradient as well as stroke volume index were lower in patients with reduced baseline LVEF. AVAi was comparable between patients with reduced, midrange, and preserved baseline LVEF.

All patients underwent TAVI via the transfemoral approach. The majority of patients received balloon-expandable valves: Edwards SAPIEN 3 341 (61%); SAPIEN XT 46 (8%); SAPIEN 36 (6%), and SAPIEN Ultra 3 (0,5%). Self-expandable valves were used in 133 patients (24%). Prosthesis size ranged from 23 mm to 31 mm, with 26 mm being most frequent in 221 patients (40%).

TABLE 2. Baseline (pre-TAVI) echocardiographic data of the overall population and according to baseline left ventricular ejection fraction.

Variable	Overall population (n=560)	Left Ventricular Ejection Fraction $\geq 50\%$ (n=350)	Left Ventricular Ejection Fraction 40-49% (n=122)	Left Ventricular Ejection Fraction $< 40\%$ (n=88)	p-value
LVEF (%)	51 $\pm$ 11	58 $\pm$ 5	45 $\pm$ 3*	33 $\pm$ 6*†	<0.001
AV peak gradient (mmHg)	66 $\pm$ 25	70 $\pm$ 25	63 $\pm$ 26*	55 $\pm$ 25*†	<0.001
AV mean gradient (mmHg)	42 $\pm$ 17	45 $\pm$ 16	41 $\pm$ 18	35 $\pm$ 15*†	<0.001
AVA (cm ²)	0.80 $\pm$ 0.28	0.82 $\pm$ 0.28	0.78 $\pm$ 0.27	0.75 $\pm$ 0.30	0.13
AVAi (cm ² /m ²)	0.43 $\pm$ 0.15	0.44 $\pm$ 0.15	0.42 $\pm$ 0.15	0.41 $\pm$ 0.16	0.19
LVEDVi (ml/m ²)	44 [36-54]	40 [34-48]	47 [36-54]*	60 [50-74]* †	<0.001
LVESVi (ml/m ²)	20 [15-29]	17 [14-20]	25 [20-30]*	41 [33-50]* †	<0.001
SVi (ml/m ²)	39 $\pm$ 12	41 $\pm$ 12	37 $\pm$ 12*	32 $\pm$ 10*†	<0.001
LV mass index (g/m ²)	119 $\pm$ 32	112 $\pm$ 27	126 $\pm$ 36*	138 $\pm$ 34*†	<0.001
LVEDD (mm)	47.7 $\pm$ 7.4	45.7 $\pm$ 6.5	49.0 $\pm$ 7.2*	53.9 $\pm$ 7.1*†	<0.001
IVST (mm)	13.1 $\pm$ 1.8	13.2 $\pm$ 1.7	13.2 $\pm$ 2.1	12.5 $\pm$ 1.7*†	0.004
PWT (mm)	12.2 $\pm$ 2.1	12.3 $\pm$ 2.1	12.5 $\pm$ 2.1	11.9 $\pm$ 2.0	0.12
RWT	0.52 $\pm$ 0.13	0.54 $\pm$ 0.13	0.51 $\pm$ 0.12	0.44 $\pm$ 0.10*†	<0.001

Data are presented as mean $\pm$ SD, median [25-75% interquartile range] and n (%). AV = aortic valve, AVA = aortic valve area, AVAi = indexed aortic valve area, IVST = intraventricular septum thickness, LV = left ventricular, LVEDVi = left ventricular end-diastolic volume index, LVEDD = left ventricular end-diastolic diameter, LVEF = left ventricular ejection fraction, LVESVi = left ventricular end-systolic volume index, PWT = posterior wall thickness, RWT = relative wall thickness, SVi = stroke volume index.

* p<0.05 vs group 'Left Ventricular Ejection Fraction $\geq 50\%$ ' with Bonferroni's post-hoc analysis.

† p<0.05 vs group 'Left Ventricular Ejection Fraction 40-49%' with Bonferroni's post-hoc analysis

Change in LVEF after TAVI

LVEF improved significantly during follow-up (from 51 $\pm$ 11% to 54 $\pm$ 10% at 6 months and to 55 $\pm$ 10% at 12 months, p<0.001). The changes in LVEF according to baseline LVEF are displayed in Figure 3. The largest increase in LVEF after TAVI was noted in patients with

reduced baseline LVEF versus those with midrange LVEF (from $33\pm 6\%$ to $42\pm 10\%$ at 6 months and to $43\pm 10\%$ at 12 months, $p<0.001$; and from $45\pm 3\%$ to $51\pm 8\%$ at 6 months and to $52\pm 8\%$ at 12 months, $p<0.001$, respectively), and a modest improvement in LVEF was noted in patients with preserved baseline LVEF (from $58\pm 5\%$ to $58\pm 7\%$ at 6 months and to $59\pm 7\%$ at 12 months, $p=0.012$; p for interaction <0.001). LV volumes reduced significantly after TAVI. The largest reduction in LV end-diastolic volume index (LVEDVi) was noted in patients with reduced LVEF as compared to patients with preserved baseline LVEF (from 60 [50 - 74] ml/m^2 to 51 [43 - 70] ml/m^2 at 6 months and to 55 [38 - 67] ml/m^2 at 12 months, $p<0.001$; and from 40 [34 - 48] ml/m^2 to 39 [32 - 47] ml/m^2 at 6 months and to 38 [30 - 45] ml/m^2 at 12 months, $p<0.001$, respectively), whereas LVEDVi was not significantly reduced among patients with midrange baseline LVEF (from 47 [36 - 54] ml/m^2 to 44 [38 - 55] ml/m^2 at 6 months and to 43 [34 - 50] ml/m^2 at 12 months, $p=0.055$; p for interaction $=0.001$). Similarly, patients with reduced LVEF had the largest reduction in LV end-systolic volume index after TAVI compared to those with midrange LVEF and preserved LVEF (reduced LVEF: from 41 [33 - 50] ml/m^2 to 32 [23 - 41] ml/m^2 at 6 months and to 30 [21 - 41] ml/m^2 at 12 months, $p<0.001$; midrange LVEF: from 25 [20 - 30] ml/m^2 to 22 [17 - 27] ml/m^2 at 6 months and to 20 [16 - 25] ml/m^2 at 12 months, $p<0.001$; and preserved LVEF: from 17 [14 - 20] ml/m^2 to 16 [13 - 20] ml/m^2 at 6 months and to 15 [11 - 19] ml/m^2 at 12 months, $p<0.001$; p for interaction <0.001). LV mass index regressed significantly after TAVI (from 119 ± 32 g/m^2 to 105 ± 26 g/m^2 at 6 months and to 99 ± 26 g/m^2 at 12 months, $p<0.001$) and was comparable between the varying groups of baseline LVEF over time (p for interaction $=0.48$). Prosthetic AV mean gradient was similar during follow-up (from 9 ± 4 mmHg immediately after TAVI to 10 ± 5 mmHg at 6 months and to 10 ± 5 mmHg at 12 months) without significant changes between groups over time (p for interaction $=0.48$). Additional analysis was performed to correct for confounders of the change in LVEF after TAVI and showed similar results (Supplemental Table 1). The absolute change in LVEF and LV volumes and mass at 6 and 12 months follow-up are shown in Supplemental Table 2.

LVEF improvement

An improvement in LVEF $\geq 5\%$ at follow-up was noted in 233 patients (42%) and was most frequently observed among patients with reduced LVEF (67%), followed by patients with midrange LVEF (58%), and those with preserved baseline LVEF (29%, $p<0.001$). Similarly, $\geq 10\%$ LVEF improvement was observed in 121 patients (22%) and most common in patients with reduced baseline LVEF (48% vs. 32% vs. 11%, respectively, $p<0.001$). A scatterplot of LVEF at baseline versus follow-up according to $\geq 5\%$ LVEF improvement is displayed in Figure 4.




FIGURE 2. Distribution of left ventricular ejection fraction before TAVI and at follow-up. Pie chart displaying the distribution of patients with preserved left ventricular ejection fraction (LVEF) $\geq 50\%$ (green), midrange LVEF 40-49% (blue), and reduced LVEF $< 40\%$ (red) at baseline and 6 and 12 months after transcatheter aortic valve implantation.




FIGURE 3. Evolution of left ventricular ejection fraction after transcatheter aortic valve implantation.

Changes in left ventricular ejection fraction (LVEF) of patients with preserved baseline LVEF $\geq 50\%$ (green line), midrange LVEF 40-49% (blue line), and patients with reduced LVEF $< 40\%$ (red line) from baseline to 6 months and 12 months follow-up after transcatheter aortic valve implantation.

* shows p-value for interaction between the varying groups of baseline LVEF over time. Error bars indicate 95% confidence intervals.




FIGURE 4. Scatterplot of left ventricular ejection fraction before transcatheter aortic valve implantation and at follow-up.

Blue dots represent patients with $\geq 5\%$ LVEF improvement; red dots represent patients with $< 5\%$ LVEF improvement.

Patients with $\geq 5\%$ LVEF improvement at follow-up had less hypertension compared to those with $< 5\%$ LVEF improvement (68% vs. 77%, $p=0.021$) as well as less coronary artery disease (55% vs. 68%, $p=0.001$), less myocardial infarction (15% vs. 24%, $p=0.007$), less previous percutaneous coronary interventions (25% vs. 35%, $p=0.016$), and less permanent pacemaker implantation (25% vs. 35%, $p=0.021$). In contrast, atrial fibrillation (32% vs. 22%, $p=0.012$) and concomitant moderate or severe mitral- or aortic regurgitation (31% vs. 23%, $p=0.026$) were more frequent in patients with $\geq 5\%$ LVEF improvement. Similar results were noted in the comparison of patients with $\geq 10\%$ LVEF improvement. In addition, the baseline clinical and echocardiographic characteristics of patients with $\text{LVEF} < 50\%$ at baseline and follow-up versus patients with preserved LVEF at baseline and follow-up among patients with $< 5\%$ LVEF improvement are summarized in Supplemental Table 3.

The baseline clinical and echocardiographic variables associated without $\geq 5\%$ LVEF improvement are displayed in Table 3. Baseline $\text{LVEF} < 50\%$ was independently associated with $\geq 5\%$ LVEF improvement at follow-up (OR for lack of $\geq 5\%$ LVEF improvement: 0.22 (95%CI 0.15-0.35), $p<0.001$), whereas coronary artery disease (OR: 1.80 (95%CI 1.06-3.06), $p=0.031$), myocardial infarction (OR: 2.07 (95%CI 1.19-3.61), $p=0.010$), and need for permanent pacemaker (OR: 1.93 (1.25-3.00), $p=0.003$) were independently associated with lack of $\geq 5\%$ LVEF improvement after TAVI.

Follow-up

During a median follow-up of 3.8 [IQR 2.6-5.2] years, 176 patients died (31%). Kaplan-Meier analysis demonstrated similar outcomes for patients with versus without $\geq 5\%$ LVEF improvement after TAVI (log-rank χ^2 : 0.19, $p=0.89$, Figure 5, Panel A). Moreover, long-term survival was comparable for patients with and without $\geq 5\%$ LVEF improvement after TAVI among patients with preserved baseline LVEF (log-rank χ^2 : 0.002, $p=0.97$, Figure 5, Panel B), baseline LVEF $<50\%$ ($n=210$; log-rank χ^2 : 0.27, $p=0.60$, Figure 5, Panel C), and among patients with reduced baseline LVEF (log-rank χ^2 : 0.18, $p=0.68$, Figure 5, Panel D). In addition, baseline LVEF (as a continuous variable, HR: 0.995 (95% CI: 0.980-1.009), $p=0.47$), a 1% improvement in LVEF after TAVI (HR: 0.996 (95% CI: 0.979-1.013), $p=0.64$), $\geq 5\%$ LVEF improvement (HR: 0.927 (95% CI: 0.670-1.283), $p=0.65$), and $\geq 10\%$ LVEF improvement (HR: 1.129 (95% CI: 0.776-1.643), $p=0.53$) were not associated with all-cause mortality after TAVI in multivariable cox regression analysis.




FIGURE 5. Kaplan-Meier curves demonstrating the event-free survival for all-cause mortality according to $\geq 5\%$ improvement in left ventricular ejection fraction (LVEF) of the overall population (Panel A) and among patients with baseline preserved LVEF (LVEF $>50\%$, Panel B), LVEF $<50\%$ (Panel C), and reduced LVEF (LVEF $<40\%$, Panel D).

TABLE 3. Uni- and multivariable logistic regression of clinical and echocardiographic parameters associated without $\geq 5\%$ LVEF improvement after transcatheter aortic valve implantation.

Variable	Univariable			Multivariable		
	OR	95% CI	p-value	OR	95% CI	p-value
Age	0.995	0.971-1.019	0.66			
Male sex	1.272	0.908-1.782	0.16			
Body mass index	1.021	0.983-1.060	0.28			
Logistic EuroSCORE	0.983	0.966-1.000	0.056	0.989	0.967-1.011	0.33
eGFR<60ml/min	1.210	0.864-1.696	0.27			
Hypertension	1.555	1.068-2.265	0.021	1.281	0.825-1.990	0.27
Hypercholesterolemia	1.198	0.850-1.689	0.30			
Diabetes mellitus	0.970	0.668-1.407	0.87			
CAD	1.776	1.254-2.515	0.001	1.796	1.055-3.057	0.031
Previous myocardial infarction	1.833	1.176-2.858	0.007	2.072	1.189-3.611	0.010
PCI	1.578	1.087-2.291	0.016	0.976	0.571-1.670	0.93
CABG	1.301	0.858-1.973	0.22			
Previous stroke / TIA	1.012	0.655-1.564	0.96			
Peripheral vascular disease	0.804	0.520-1.242	0.33			
Atrial fibrillation	0.618	0.423-0.902	0.013	0.761	0.486-1.190	0.23
Chronic obstructive pulmonary disease	0.827	0.536-1.276	0.39			
Concomitant mitral / aortic regurgitation $\geq$ grade 2	0.938	0.550-1.601	0.82			
Baseline LVEF<50%	0.257	0.179-0.368	<0.001	0.224	0.145-0.346	<0.001
Paravalvular leakage $\geq$ moderate (grade 2)	1.443	0.637-3.272	0.38			
Permanent pacemaker	1.548	1.066-2.249	0.022	1.931	1.245-2.995	0.003

CABG = coronary artery bypass grafting, CAD = coronary artery disease, CKD-EPI = chronic kidney disease epidemiology collaboration, eGFR = estimated glomerular filtration rate, LVEF = left ventricular ejection fraction, PCI = percutaneous coronary intervention, TIA = transient ischemic attack.

DISCUSSION

In patients with severe AS, LVEF improved significantly at 1-year follow-up after TAVI. Patients with reduced baseline LVEF showed the greatest improvement in LVEF. In addition, coronary artery disease, myocardial infarction, and need for a permanent

pacemaker were independently associated with lack of LVEF improvement after TAVI. However, improvement in LVEF at 1-year follow-up was not associated with long-term outcomes.

Changes in LVEF after TAVI

LV systolic dysfunction is common among patients with severe AS; this may be secondary to severe AS or co-existent cardiomyopathy. In severe AS, the LV myocardium may remodel and develop LV hypertrophy as a consequence of the chronic LV pressure overload associated with severe AS, in order to normalize wall stress and maintain cardiac output^{1,26}. However, with AS progression, this compensation mechanism may fail and subsequently lead to heart failure and mortality^{27,28}. Aortic valve replacement results in immediate relief of LV afterload and is associated with LV reverse remodeling and improvement in LV systolic function^{14,29}. The present study evaluated the development of LVEF after TAVI among patients with varying baseline LVEF. The greatest improvement in LVEF at 1-year follow-up was noted in patients with reduced LVEF (<40%), despite a higher prevalence of ischemic heart disease which potentially could limit LV reverse remodeling. Patients with reduced LV function may have the greatest potential for LV recovery after TAVI, as confirmed by the current results.

Ewe et al.¹⁵ evaluated the LV systolic function of 147 patients with severe AS who underwent TAVI with preserved baseline LVEF and impaired LVEF (<50%). In line with our findings, patients with impaired LVEF showed significant improvement in LVEF at follow-up. However, LVEF was comparable in patients with preserved baseline LVEF at follow-up, whereas in the present study, including more patients and longer echocardiographic follow-up, a modest but significant improvement in LVEF was noted in patients with preserved baseline LVEF at 1-year after TAVI. Similarly, a study including 505 patients with severe AS who underwent TAVI demonstrated that patients with severe LV dysfunction (<35%) had the greatest improvement in LVEF at 6-months follow-up compared to those with moderate LV dysfunction (35-50%)³⁰. Moreover, a meta-analysis including 26 articles reported significant improvement in LVEF among patients with severe AS and low baseline LVEF (<50%) at 30 days, 6 months, and at 1 year after TAVI¹¹.

Next, the variables related to lack of LVEF improvement were evaluated. Permanent pacemaker implantation, more specifically right ventricular pacing, is known to affect LV function and LV recovery after TAVI^{25,31}. The present analysis showed that the majority of patients with a permanent pacemaker did not improve in LVEF at follow-up. In addition, coronary artery disease and myocardial infarction were also associated with lack of LVEF improvement, whereas baseline LVEF <50% was linked with LVEF improvement similar to previous studies^{30,32}.

LVEF improvement versus long-term outcomes

LV systolic function yields important prognostic information in patients with severe AS who underwent surgical aortic valve replacement⁶. Similarly, it has been shown that baseline LVEF is an independent predictor of outcomes after TAVI⁹⁻¹¹. However, data on LVEF improvement versus long-term outcomes after TAVI are scarce. In a cohort of patients with severe AS and reduced LVEF (<45%) who underwent TAVI, Angelillis et al.³³ reported that a $\geq 10\%$ increase in LVEF at 1-month follow-up was associated with improved outcomes, although patients with previous coronary revascularization were excluded from the analysis. Similarly, Dauerman et al.³² evaluated early LV recovery after TAVI in patients with severe AS and reduced LVEF (<40%) with high operative risk. Patients with a $\geq 10\%$ increase in LVEF at 1-month follow-up had fewer adverse events, although not statistically different possibly due to small patient numbers. In the present study including more patients, an improvement of $\geq 5\%$ in LVEF at 6- or 12-months follow-up after TAVI was not associated with long-term outcomes, even in the patients with reduced baseline LVEF.

Limitations

Several limitations should be acknowledged. First, this is a single-center retrospective analysis with inherent limitations related to the study design. Second, patients without echocardiographic follow-up have been excluded from the analysis which may have caused selection bias. However, the baseline clinical characteristics of this population versus the study population are comparable (Supplemental Table 3). Third, the group of patients with reduced LVEF was limited. Finally, the current study used LVEF as a marker of LV systolic function, where LV strain may be a more sensitive marker to assess LV dysfunction.

CONCLUSION

LVEF improved significantly in patients with severe AS at 1-year follow-up after TAVI. Patients with reduced baseline LVEF revealed the greatest improvement in LVEF. Coronary artery disease, myocardial infarction, and the need for a permanent pacemaker were independently associated with lack of LVEF improvement. However, LVEF improvement at 12 months follow-up was not associated with long-term outcomes.

REFERENCES

1. Otto CM, Prendergast B. Aortic-valve stenosis--from patients at risk to severe valve obstruction. *N Engl J Med* 2014;371:744-756.
2. Grossman W, Jones D, McLaurin LP. Wall stress and patterns of hypertrophy in the human left ventricle. *J Clin Invest* 1975;56:56-64.
3. Baron SJ, Arnold SV, Herrmann HC et al. Impact of Ejection Fraction and Aortic Valve Gradient on Outcomes of Transcatheter Aortic Valve Replacement. *J Am Coll Cardiol* 2016;67:2349-2358.
4. Elmariah S, Palacios IF, McAndrew T et al. Outcomes of transcatheter and surgical aortic valve replacement in high-risk patients with aortic stenosis and left ventricular dysfunction: results from the Placement of Aortic Transcatheter Valves (PARTNER) trial (cohort A). *Circ Cardiovasc Interv* 2013;6:604-614.
5. Passeri JJ, Elmariah S, Xu K et al. Transcatheter aortic valve replacement and standard therapy in inoperable patients with aortic stenosis and low EF. *Heart* 2015;101:463-471.
6. Dahl JS, Eleid MF, Michelena HI et al. Effect of left ventricular ejection fraction on postoperative outcome in patients with severe aortic stenosis undergoing aortic valve replacement. *Circ Cardiovasc Imaging* 2015;8.
7. Halkos ME, Chen EP, Sarin EL et al. Aortic valve replacement for aortic stenosis in patients with left ventricular dysfunction. *Ann Thorac Surg* 2009;88:746-751.
8. Siontis GCM, Overtchouk P, Cahill TJ et al. Transcatheter aortic valve implantation vs. surgical aortic valve replacement for treatment of symptomatic severe aortic stenosis: an updated meta-analysis. *Eur Heart J* 2019;40:3143-3153.
9. Eleid MF, Goel K, Murad MH et al. Meta-Analysis of the Prognostic Impact of Stroke Volume, Gradient, and Ejection Fraction After Transcatheter Aortic Valve Implantation. *Am J Cardiol* 2015;116:989-994.
10. Furer A, Chen S, Redfors B et al. Effect of Baseline Left Ventricular Ejection Fraction on 2-Year Outcomes After Transcatheter Aortic Valve Replacement: Analysis of the PARTNER 2 Trials. *Circ Heart Fail* 2019;12:e005809.
11. Sannino A, Gargiulo G, Schiattarella GG et al. Increased mortality after transcatheter aortic valve implantation (TAVI) in patients with severe aortic stenosis and low ejection fraction: a meta-analysis of 6898 patients. *Int J Cardiol* 2014;176:32-39.
12. Vahanian A, Beyersdorf F, Praz F et al. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur J Cardiothorac Surg* 2021;60:727-800.
13. Otto CM, Nishimura RA, Bonow RO et al. 2020 ACC/AHA Guideline for the Management of Patients With Valvular Heart Disease: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2021;143:e72-e227.
14. Takagi H, Ando T, Umemoto T, Group A. A meta-analysis of effects of transcatheter versus surgical aortic valve replacement on left ventricular ejection fraction and mass. *Int J Cardiol* 2017;238:31-36.
15. Ewe SH, Ajmone Marsan N, Pepi M et al. Impact of left ventricular systolic function on clinical and echocardiographic outcomes following transcatheter aortic valve implantation for severe aortic stenosis. *Am Heart J* 2010;160:1113-1120.
16. Ribeiro HB, Lerakis S, Gilard M et al. Transcatheter Aortic Valve Replacement in Patients With Low-Flow, Low-Gradient Aortic Stenosis: The TOPAS-TAVI Registry. *J Am Coll Cardiol* 2018;71:1297-1308.

17. Kamperidis V, Joyce E, Debonnaire P et al. Left ventricular functional recovery and remodeling in low-flow low-gradient severe aortic stenosis after transcatheter aortic valve implantation. *J Am Soc Echocardiogr* 2014;27:817-825.
18. Ponikowski P, Voors AA, Anker SD et al. 2016 ESC Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure. *Rev Esp Cardiol (Engl Ed)* 2016;69:1167.
19. Delgado V, Ng AC, van de Veire NR et al. Transcatheter aortic valve implantation: role of multi-detector row computed tomography to evaluate prosthesis positioning and deployment in relation to valve function. *Eur Heart J* 2010;31:1114-1123.
20. Otto CM, Kumbhani DJ, Alexander KP et al. 2017 ACC Expert Consensus Decision Pathway for Transcatheter Aortic Valve Replacement in the Management of Adults With Aortic Stenosis: A Report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol* 2017;69:1313-1346.
21. Lancellotti P, Pibarot P, Chambers J et al. Recommendations for the imaging assessment of prosthetic heart valves: a report from the European Association of Cardiovascular Imaging endorsed by the Chinese Society of Echocardiography, the Inter-American Society of Echocardiography, and the Brazilian Department of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2016;17:589-590.
22. Lang RM, Badano LP, Mor-Avi V et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2015;28:1-39 e14.
23. Zoghbi WA, Asch FM, Bruce C et al. Guidelines for the Evaluation of Valvular Regurgitation After Percutaneous Valve Repair or Replacement: A Report from the American Society of Echocardiography Developed in Collaboration with the Society for Cardiovascular Angiography and Interventions, Japanese Society of Echocardiography, and Society for Cardiovascular Magnetic Resonance. *J Am Soc Echocardiogr* 2019;32:431-475.
24. Bax JJ, Poldermans D, Elhendy A et al. Improvement of left ventricular ejection fraction, heart failure symptoms and prognosis after revascularization in patients with chronic coronary artery disease and viable myocardium detected by dobutamine stress echocardiography. *J Am Coll Cardiol* 1999;34:163-169.
25. Ngo A, Hassager C, Thyregod HGH et al. Differences in left ventricular remodelling in patients with aortic stenosis treated with transcatheter aortic valve replacement with corevalve prostheses compared to surgery with porcine or bovine biological prostheses. *Eur Heart J Cardiovasc Imaging* 2018;19:39-46.
26. Dweck MR, Boon NA, Newby DE. Calcific aortic stenosis: a disease of the valve and the myocardium. *J Am Coll Cardiol* 2012;60:1854-1863.
27. Bing R, Cavalcante JL, Everett RJ, Clavel MA, Newby DE, Dweck MR. Imaging and Impact of Myocardial Fibrosis in Aortic Stenosis. *JACC Cardiovasc Imaging* 2019;12:283-296.
28. Hein S, Arnon E, Kostin S et al. Progression from compensated hypertrophy to failure in the pressure-overloaded human heart: structural deterioration and compensatory mechanisms. *Circulation* 2003;107:984-991.
29. Pai RG, Kapoor N, Bansal RC, Varadarajan P. Malignant natural history of asymptomatic severe aortic stenosis: benefit of aortic valve replacement. *Ann Thorac Surg* 2006;82:2116-2122.
30. Elhmidi Y, Bleiziffer S, Deutsch MA et al. Transcatheter aortic valve implantation in patients with LV dysfunction: impact on mortality and predictors of LV function recovery. *J Invasive Cardiol* 2014;26:132-138.
31. Urena M, Webb JG, Tamburino C et al. Permanent pacemaker implantation after transcatheter aortic valve implantation: impact on late clinical outcomes and left ventricular function. *Circulation* 2014;129:1233-1243.

32. Dauerman HL, Reardon MJ, Popma JJ et al. Early Recovery of Left Ventricular Systolic Function After CoreValve Transcatheter Aortic Valve Replacement. *Circ Cardiovasc Interv* 2016;9.
33. Angelillis M, Giannini C, De Carlo M et al. Prognostic Significance of Change in the Left Ventricular Ejection Fraction After Transcatheter Aortic Valve Implantation in Patients With Severe Aortic Stenosis and Left Ventricular Dysfunction. *Am J Cardiol* 2017;120:1639-1647.

SUPPLEMENTAL MATERIAL

SUPPLEMENTAL TABLE 1. Unadjusted and adjusted significance testing for changes in echocardiographic parameters over time between patients with preserved, midrange, and reduced baseline left ventricular ejection fraction.

Variable	p-value for interaction unadjusted	p-value for interaction adjusted*
LVEF	<0.001	0.010
LVEDVi	0.001	0.003
LVESVi	<0.001	0.15
LV mass index	0.48	0.88
Prosthetic mean gradient	0.48	0.99

LV = left ventricular, LVEDVi = left ventricular end-diastolic volume index, LVEF = left ventricular ejection fraction, LVESVi = left ventricular end-systolic volume index.

*Adjusted for the following variables: Age, sex, hypertension, diabetes mellitus, coronary artery disease, previous myocardial infarction, concomitant moderate or severe mitral – or aortic regurgitation, paravalvular leakage ≥moderate, permanent pacemaker (at baseline and after TAVI), atrial fibrillation, pre-TAVI aortic valve mean gradient, and pre-TAVI LVEF.

SUPPLEMENTAL TABLE 2. Change in left ventricular parameters at 6 and 12 months after transcatheter aortic valve implantation.

Variable		Overall population	Left Ventricular Ejection Fraction $\geq 50\%$	Left Ventricular Ejection Fraction 40-49%	Left Ventricular Ejection Fraction $< 40\%$	p-value
Absolute LVEF improvement (%)	6m (n=524)	3 $\pm$ 9	0 $\pm$ 8	6 $\pm$ 7*	9 $\pm$ 10*†	<0.001
	12m (n=487)	4 $\pm$ 9	1 $\pm$ 8	7 $\pm$ 8*	10 $\pm$ 10*†	<0.001
Absolute LVEDVi reduction (ml)	6m (n=524)	3 $\pm$ 12	2 $\pm$ 9	1 $\pm$ 13	8 $\pm$ 17*	<0.001
	12m (n=487)	4 $\pm$ 13	4 $\pm$ 11	4 $\pm$ 15	9 $\pm$ 18*	0.005
Absolute LVESVi reduction (ml)	6m (n=524)	3 $\pm$ 8	1 $\pm$ 5	3 $\pm$ 7*	10 $\pm$ 13*†	<0.001
	12m (n=487)	4 $\pm$ 9	2 $\pm$ 6	5 $\pm$ 8*	11 $\pm$ 15*†	<0.001
Absolute LV mass index regression (g/m ²)	6m (n=520)	14 $\pm$ 21	12 $\pm$ 20	15 $\pm$ 20	17 $\pm$ 28	0.21
	12m (n=485)	20 $\pm$ 24	18 $\pm$ 20	22 $\pm$ 30	22 $\pm$ 31	0.31
LVEDVi reduction as percentage of baseline (%)	6m (n=524)	2 $\pm$ 24	3 $\pm$ 22	-3 $\pm$ 27	11 $\pm$ 23*†	<0.001
	12m (n=487)	7 $\pm$ 25	6 $\pm$ 24	1 $\pm$ 29	12 $\pm$ 25†	0.018
LVESVi reduction as percentage of baseline (%)	6m (n=524)	7 $\pm$ 27	3 $\pm$ 27	9 $\pm$ 26	22 $\pm$ 25*†	<0.001
	12m (n=487)	11 $\pm$ 32	7 $\pm$ 34	14 $\pm$ 26	24 $\pm$ 25*	<0.001
LVMI reduction as percentage of baseline (%)	6m (n=435)	10 $\pm$ 16	10 $\pm$ 17	11 $\pm$ 15	11 $\pm$ 17	0.73
	12m (n=406)	15 $\pm$ 16	15 $\pm$ 16	16 $\pm$ 17	14 $\pm$ 18	0.75
$\geq 5\%$ LVEF reduction	6m (n=439)	75 (14%)	66 (20%)	7 (6%)	2 (2%)	<0.001
	12m (n=408)	70 (14%)	64 (21%)	6 (6%)	0	<0.001

Data are presented as mean $\pm$ SD in absolute change compared to baseline and as percentual reduction compared to baseline. LVEDVi = left ventricular end-diastolic volume index, LVEF = left ventricular ejection fraction, LVESVi = left ventricular end-systolic volume index, LVMI = left ventricular mass index.

* p<0.05 vs group 'Left Ventricular Ejection Fraction $\geq 50\%$ ' with Bonferroni's post-hoc analysis.

† p<0.05 vs group 'Left Ventricular Ejection Fraction 40-49%' with Bonferroni's post-hoc analysis

SUPPLEMENTAL TABLE 3. Baseline clinical and echocardiographic characteristics of patients with left ventricular ejection fraction <50% at baseline and follow-up vs. patients with preserved left ventricular ejection fraction at baseline and follow-up among patients with <5% improvement in left ventricular ejection fraction.

Variable	Left Ventricular Ejection Fraction <50% at baseline and follow-up (n=70)	Left Ventricular Ejection Fraction ≥50% at baseline and follow-up (n=213)	p-value
Age (years)	79 ± 7	80 ± 7	0.84
Male sex	46 (66%)	104 (49%)	0.014
Body mass index (kg/m ²)	27.0 ± 3.5	26.9 ± 4.8	0.81
Logistic EuroSCORE	18.4 [10.4-26.6]	10.1 [7.5-16.6]	<0.001
eGFR (ml/min)	55 ± 22	63 ± 19	0.001
Hypertension	51 (73%)	161 (76%)	0.56
Hypercholesterolemia	48 (69%)	129 (61%)	0.25
Diabetes mellitus	20 (29%)	56 (26%)	0.71
Smoker	7 (11%)	28 (14%)	0.50
CAD	56 (80%)	137 (65%)	0.016
Previous myocardial infarction	28 (40%)	36 (17%)	<0.001
Previous revascularization	50 (71%)	116 (55%)	0.012
PCI	27 (39%)	74 (35%)	0.020
CABG	23 (33%)	42 (20%)	
NYHA class			
I-II	27 (39%)	103 (49%)	0.15
III-IV	43 (61%)	109 (51%)	
Previous stroke / TIA	13 (19%)	42 (21%)	0.74
Peripheral vascular disease	14 (20%)	34 (16%)	0.44
Atrial fibrillation	18 (26%)	41 (19%)	0.25
Chronic obstructive pulmonary disease	12 (17%)	35 (17%)	0.88
Concomitant mitral / aortic regurgitation ≥grade 2	26 (37%)	33 (16%)	<0.001
Permanent pacemaker	21 (30%)	22 (10%)	<0.001
Medication			
Beta-blocker	47 (67%)	130 (61%)	0.38
ACE-I / ARB II	42 (60%)	111 (52%)	0.27
Calcium antagonist	21 (30%)	45 (21%)	0.13
Diuretics	49 (70%)	92 (43%)	<0.001
MR antagonist	10 (14%)	18 (6%)	0.16
Statins	51 (73%)	145 (68%)	0.48
Antiplatelet	43 (61%)	132 (62%)	0.94
Anticoagulation	35 (50%)	61 (29%)	0.002

Variable	Left Ventricular Ejection Fraction <50% at baseline and follow-up (n=70)	Left Ventricular Ejection Fraction ≥50% at baseline and follow-up (n=213)	p-value
Echocardiographic data			
LVEF (%)	41 ± 6	59 ± 5	<0.001
AV peak gradient (mmHg)	51 ± 24	69 ± 26	<0.001
AV mean gradient (mmHg)	32 ± 15	45 ± 17	<0.001
AVA (cm ²)	0.87 ± 0.29	0.82 ± 0.29	0.16
AVAi (cm ² /m ²)	0.46 ± 0.16	0.44 ± 0.15	0.19
LVEDVi (ml/m ²)	56 [47-70]	40 [35-48]	<0.001
LVESVi (ml/m ²)	32 [25-43]	16 [14-20]	<0.001
SVi (ml/m ²)	36 ± 11	42 ± 12	0.001
LV mass index (g/m ²)	129 ± 36	114 ± 28	<0.001
LVEDD (mm)	51.6 ± 7.0	45.6 ± 6.1	<0.001
IVST (mm)	12.6 ± 1.9	13.2 ± 1.7	0.015
PWT (mm)	12.2 ± 2.3	12.4 ± 2.0	0.35

Data are presented as mean ± SD, median [25-75% interquartile range] and n (%). ACE-I = angiotensin-converting enzyme, ARB II = angiotensin-II receptor blocker, AV = aortic valve, AVA = aortic valve area, AVAi = indexed aortic valve area, CABG = coronary artery bypass grafting, CAD = coronary artery disease, CKD-EPI = chronic kidney disease epidemiology collaboration, eGFR = estimated glomerular filtration rate, IVST = intraventricular septum thickness, LV = left ventricular, LVEDVi = left ventricular end-diastolic volume index, LVEDD = left ventricular end-diastolic diameter, LVEF = left ventricular ejection fraction, LVESVi = left ventricular end-systolic volume index, MR = mineralocorticoid receptor, NYHA = New York Heart Association, PCI = percutaneous coronary intervention, PWT = posterior wall thickness, RWT = relative wall thickness, SVi = stroke volume index, TIA = transient ischemic attack.

SUPPLEMENTAL TABLE 4. Baseline clinical and echocardiographic characteristics of the study population vs. the excluded population without echocardiographic follow-up.

Variable	Population with echocardiographic follow-up (n=560)	Population without echocardiographic follow-up (n=200)	p-value
Age (years)	80 ± 7	80 ± 7	0.40
Male sex	296 (53%)	102 (51%)	0.65
Body mass index (kg/m²)	26.7 ± 4.6	26.8 ± 5.2	0.92
Logistic EuroSCORE	12.3 [8.6-20.1]	12.9 [9.6-19.3]	0.47
eGFR (ml/min)	61 ± 20	57 ± 20	0.019
Hypertension	407 (73%)	140 (78%)	0.20
Hypercholesterolemia	338 (61%)	102 (57%)	0.37
Diabetes mellitus	159 (28%)	63 (35%)	0.092
Smoker	76 (14%)	28 (18%)	0.20
CAD	351 (63%)	112 (63%)	0.98
Previous myocardial infarction	112 (20%)	39 (22%)	0.58
Previous revascularization	293 (52%)	79 (52%)	0.88
PCI	173 (31%)	71 (40%)	0.006
CABG	120 (21%)	19 (11%)	
NYHA class			
I-II	232 (42%)	78 (44%)	0.63
III-IV	327 (59%)	101 (56%)	
Previous stroke / TIA	103 (19%)	25 (17%)	0.13
Peripheral vascular disease	100 (18%)	30 (17%)	0.72
Atrial fibrillation	147 (26%)	49 (33%)	0.13
Chronic obstructive pulmonary disease	101 (18%)	40 (23%)	0.15
Concomitant mitral / aortic regurgitation ≥grade 2	148 (26%)	49 (28%)	0.71
Permanent pacemaker	83 (15%)	13 (9%)	0.042
Medication			
Beta-blocker	326 (58%)	101 (56%)	0.54
ACE-I / ARB II	290 (52%)	93 (51%)	0.89
Calcium antagonist	136 (24%)	46 (25%)	0.78
Diuretics	295 (53%)	116 (64%)	0.008
MR antagonist	57 (10%)	26 (14%)	0.12
Statins	349 (63%)	112 (62%)	0.87
Antiplatelet	325 (58%)	83 (55%)	0.50
Anticoagulation	211 (38%)	82 (45%)	0.092
Echocardiographic data			
LVEF (%)	51 ± 11	54 ± 14	0.010

Variable	Population with echocardiographic follow-up (n=560)	Population without echocardiographic follow-up (n=200)	p-value
LVEF			
≥50%	350 (62%)	102 (66%)	0.15
40-49%	122 (22%)	23 (15%)	
<40%	88 (16%)	29 (19%)	
AV peak gradient (mmHg)	66 ± 25	66 ± 22	0.81
AV mean gradient (mmHg)	42 ± 17	42 ± 16	0.77
AVA (cm ²)	0.80 ± 0.28	0.84 ± 0.31	0.11
AVAi (cm ² /m ²)	0.43 ± 0.15	0.45 ± 0.17	0.071
LVEDVi (ml/m ²)	44 [36-54]	50 [38-66]	0.001
LVESVi (ml/m ²)	20 [15-29]	21 [14-33]	0.69
SVi (ml/m ²)	39 ± 12	40 ± 12	0.28
LV mass index (g/m ²)	119 ± 32	128 ± 40	0.023
LVEDD (mm)	47.7 ± 7.4	47.3 ± 7.3	0.53
IVST (mm)	13.1 ± 1.8	13.3 ± 3.7	0.57
PWT (mm)	12.2 ± 2.1	12.3 ± 2.3	0.84

Data are presented as mean ± SD, median [25-75% interquartile range] and n (%). ACE-I = angiotensin-converting enzyme, ARB II = angiotensin-II receptor blocker, AV = aortic valve, AVA = aortic valve area, AVAi = indexed aortic valve area, CABG = coronary artery bypass grafting, CAD = coronary artery disease, CKD-EPI = chronic kidney disease epidemiology collaboration, eGFR = estimated glomerular filtration rate, IVST = intraventricular septum thickness, LV = left ventricular, LVEDVi = left ventricular end-diastolic volume index, LVEDD = left ventricular end-diastolic diameter, LVEF = left ventricular ejection fraction, LVESVi = left ventricular end-systolic volume index, MR = mineralocorticoid receptor, NYHA = New York Heart Association, PCI = percutaneous coronary intervention, PWT = posterior wall thickness, RWT = relative wall thickness, SVi = stroke volume index, TIA = transient ischemic attack.