



Universiteit
Leiden
The Netherlands

Broken blue-tilted inflationary gravitational waves: a joint analysis of NANOGrav 15-year and BICEP/Keck 2018 data

Jiang, J-Q.; Cai, Y.; Ye, G.; Piao, Y-S.

Citation

Jiang, J. -Q., Cai, Y., Ye, G., & Piao, Y. -S. (2024). Broken blue-tilted inflationary gravitational waves: a joint analysis of NANOGrav 15-year and BICEP/Keck 2018 data. *Journal Of Cosmology And Astroparticle Physics*, 5. doi:10.1088/1475-7516/2024/05/004

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/4172549>

Note: To cite this publication please use the final published version (if applicable).

PAPER

Broken blue-tilted inflationary gravitational waves: a joint analysis of NANOGrav 15-year and BICEP/Keck 2018 data

To cite this article: Jun-Qian Jiang *et al* JCAP05(2024)004

View the [article online](#) for updates and enhancements.

You may also like

- [THE SPIN RATE OF PRE-COLLAPSE STELLAR CORES: WAVE-DRIVEN ANGULAR MOMENTUM TRANSPORT IN MASSIVE STARS](#)
Jim Fuller, Matteo Cantiello, Daniel Lecoanet et al.
- [The NANOGrav 15 yr Data Set: Search for Signals from New Physics](#)
Adeela Afzal, Gabriella Agazie, Akash Anumalapudi et al.
- [ANGULAR MOMENTUM TRANSPORT VIA INTERNAL GRAVITY WAVES IN EVOLVING STARS](#)
Jim Fuller, Daniel Lecoanet, Matteo Cantiello et al.

RECEIVED: January 18, 2024

REVISED: March 14, 2024

ACCEPTED: March 25, 2024

PUBLISHED: May 2, 2024

Broken blue-tilted inflationary gravitational waves: a joint analysis of NANOGrav 15-year and BICEP/Keck 2018 data

Jun-Qian Jiang^{a,b}, Yong Cai,^c Gen Ye^d and Yun-Song Piao^{a,b,e,f}

^a*School of Fundamental Physics and Mathematical Sciences,
Hangzhou Institute for Advanced Study, UCAS,
Hangzhou 310024, China*

^b*School of Physical Sciences, University of Chinese Academy of Sciences,
Beijing 100049, China*

^c*School of Physics and Microelectronics, Zhengzhou University,
Zhengzhou, Henan 450001, China*

^d*Leiden University, Instituut-Lorentz for Theoretical Physics,
2333CA, Leiden, Netherlands*

^e*International Center for Theoretical Physics Asia-Pacific,
Beijing/Hangzhou, China*

^f*Institute of Theoretical Physics, Chinese Academy of Sciences,
P.O. Box 2735, Beijing 100190, China*

E-mail: jiangjq2000@gmail.com, caiyong@zzu.edu.cn,
ye@lorentz.leidenuniv.nl, yspiao@ucas.ac.cn

ABSTRACT: Recently, the pulsar timing array (PTA) collaborations have reported the evidence for a stochastic gravitational wave background (SGWB) at nano-Hertz band. The spectrum of inflationary gravitational wave (IGW) is unknown, which might exhibit different power law at different frequency-bands, thus if the PTA signal is primordial, it will be significant to explore the underlying implications of current PTA and CMB data on IGW. In this Letter, we perform a joint Markov Chain Monte Carlo analysis for a broken power-law spectrum of IGW with the NANOGrav 15-year and BICEP/Keck 2018 data. It is found that though the bestfit spectral tilt of IGW at PTA band is $n_T^{\text{PTA}} = 2.42^{+0.32}_{-0.91}$, at CMB band the bestfit is $n_T^{\text{CMB}} = 0.55^{+0.37}_{-0.10}$ while a detectable amplitude of r with $n_T^{\text{CMB}} \simeq 0$ is still compatible. The implication of our results for inflation is also discussed.

KEYWORDS: Bayesian reasoning, gravitational waves / sources, gravitational waves and CMBR polarization, inflation

ARXIV EPRINT: [2307.15547](https://arxiv.org/abs/2307.15547)

Contents

1	Introduction	1
2	Dataset and method	2
3	Results	3
3.1	$n_{\text{T}}^{\text{CMB}}$ is free	3
3.2	$n_{\text{T}}^{\text{CMB}} = -r/8$	4
4	Discussion	5
4.1	Conclusion	5
4.2	Implications for inflation	7

1 Introduction

It is well-known that the detection of inflationary or primordial gravitational waves (IGW) will not only solidify our confidence on the inflation scenario [1–5], but also bring us significant insight into the physics of very early universe.

The ultra-low-frequency IGW with $f \sim 10^{-18} - 10^{-16}\text{Hz}$ will source the B-mode polarization in the cosmic microwave background (CMB) [6, 7], which has been still searched for by BICEP/Keck [8]. Recently, the PTA experiments [9–12], have found a stochastic GW background (SGWB) at $f \sim 10^{-10} - 10^{-8}\text{Hz}$ [13, 14]. Though such a SGWB is compatible with that brought by inspiralling supermassive black holes binaries [13–15], see also [16, 17] for the supermassive primordial black holes, it might be just IGW but with $n_{\text{T}} > 0$ [13, 14, 18–20].¹

However, the spectrum of IGW is actually unknown. The amplitude of IGW at CMB band is usually quantified as the tensor-to-scalar ratio $r = \frac{A_{\text{T}}}{A_{\text{s}}}$, we have

$$P_{\text{T}}(k) = r A_{\text{s}} \left(\frac{k}{k_{\text{pivot}}} \right)^{n_{\text{T}}}, \quad (1.1)$$

where k_{pivot} is the pivot scale. It is possible for inflation to yield a blue-tilted SGWB with $n_{\text{T}} > 0$ [20, 38–51], however, the primordial scalar perturbation is nearly scale-invariant at CMB band [52], which seems to be in favor of a slow-roll model of inflation with $n_{\text{T}} \simeq 0$ [53]. Inspired by [54, 55], see also [56–60], it might be more reliable to consider a broken power-law IGWB at $f \sim 10^{-18} - 10^{-8}\text{Hz}$,

$$P_{\text{T}}(k) = r A_{\text{s}} \left(\frac{k}{k_{\text{pivot}}} \right)^{n_{\text{T}}^{\text{CMB}}} \left(1 + \left(\frac{k}{k_{\text{break}}} \right) \right)^{-n_{\text{T}}^{\text{CMB}} + n_{\text{T}}^{\text{PTA}}}, \quad (1.2)$$

where $f_{\text{break}} \ll 10^{-8}\text{Hz}$ is the scale at which power-law is broken, and the conversion between k and f is

$$f = \frac{kc}{2\pi a_0} = 1.5 \times 10^{-15} \left(\frac{k}{\text{Mpc}^{-1}} \right) \text{Hz}. \quad (1.3)$$

¹In addition, see also recent other possibilities e.g. [21, 21, 22, 22–37].

In refs. [61–63] such a broken SGWB has been studied but with $f_{\text{break}} > 10^{-7}\text{Hz}$. According to eq. (1.2), we have $P_{\text{T}} = rA_s \left(\frac{k}{k_{\text{pivot}}}\right)^{n_{\text{T}}^{\text{CMB}}}$ when $k \ll k_{\text{break}}$, while

$$P_{\text{T}} = rA_s \left(\frac{k_{\text{break}}^{n_{\text{T}}^{\text{CMB}} - n_{\text{T}}^{\text{PTA}}}}{k_{\text{pivot}}^{n_{\text{T}}^{\text{CMB}}}}\right) k^{n_{\text{T}}^{\text{PTA}}}, \quad (1.4)$$

when $k \gg k_{\text{break}}$. Actually, a period of inflation might be complex so that at different bands of $f \sim 10^{-18} - 10^{-8}\text{Hz}$ we will have different power-law IGW, while eq. (1.2) is just the simplest of such possibilities.²

It has been found that for power-law IGW (1.1), recent NANOGrav data favors $n_{\text{T}} \simeq 2$. As a result, IGW at CMB band is negligible. Thus recent CMB data has not been included in the analysis of NANOGrav [13] (a hard prior $r \lesssim 0.03$ is actually sufficient [19]). However, this is not valid for broken power-law IGW (1.2), which allows a non-negligible r at CMB band. Thus it is significant to perform a joint analysis of both latest PTA and CMB data with full exact likelihood to explore the underlying impact of current data on IGW.

However, a joint Markov Chain Monte Carlo (MCMC) analysis of NANOGrav 15-year and BICEP/Keck 2018 data has been still open. In this Letter, we present the first such analysis, and find that the bestfit spectral tilt of IGW at PTA band is $n_{\text{T}}^{\text{PTA}} \simeq 2$, however, different from the results of NANOGrav [13], $n_{\text{T}}^{\text{CMB}} \simeq 2$ is not favored at CMB band, instead the bestfit is $n_{\text{T}}^{\text{CMB}} \simeq 0.5$ while a detectable amplitude of r with $n_{\text{T}}^{\text{CMB}} \simeq 0$ is still compatible. The implication of our results for inflation is also discussed.

2 Dataset and method

NANOGrav. We use the NANOGrav 15-year dataset [9] at PTA band, assuming that the signals observed in NANOGrav [9], EPTA [10], PPTA [11] and CPTA [12] are mutually consistent. The likelihoods are calculated with `ceffy1` [64].

BICEP/Keck (BK18). We use the BICEP/Keck 2018 official likelihood³ [8] at CMB band, taking dust, synchrotron and noise into account.

As in the analysis of NANOGrav [13], we fix the parameters of standard ΛCDM model to the bestfit values of the Planck 2018 baseline results:⁴ $\Omega_b h^2 = 0.02238$, $\Omega_c h^2 = 0.12011$, $100\theta_{MC} = 1.040909$, $\tau = 0.0543$, $\ln(10^{10}A_s) = 3.0448$, $n_s = 0.96605$. In addition to the nuisance parameters in the BICEP/Keck likelihood, our MCMC parameters set is $\{r_{0.05}, n_{\text{T}}^{\text{CMB}}, n_{\text{T}}^{\text{PTA}}, \log_{10} k_{\text{break}}\}$, where the subscript 0.05 for r indicates that it is calculated at the pivot scale $k = 0.05\text{Mpc}^{-1}$. The uniform priors are set, and the unit of k_{break} is Mpc^{-1} .

²Actually, beyond PTA band such an IGW spectrum will be conflicted with the BBN bound on relativistic components, however, we only consider (1.2) at $f \sim 10^{-18} - 10^{-8}\text{Hz}$, since at higher-frequency band (1.2) might have been modified [61–63], see also section 3.

³http://bicepkeck.org/bk18_2021_release.html.

⁴The corresponding bestfit values might shift in light of the early resolution of the Hubble tension [65], in particular the spectral index n_s of primordial scalar perturbation will shift towards $n_s = 1$ [66–68], however, such shifts will not essentially alter our results.

Parameter	Best fit	68% limits
$r_{0.05}$	0.0296	< 0.0549 (< 0.107)
$r_{0.01}$	0.0141	< 0.0199 (< 0.0338)
$n_{\text{T}}^{\text{CMB}}$	0.428	$0.55^{+0.37}_{-0.10}$
$n_{\text{T}}^{\text{PTA}}$	2.08	$2.42^{+0.32}_{-0.91}$
$\log_{10} k_{\text{break}}$	4.19	$5.0^{+2.0}_{-1.5}$

Table 1. The bestfit values and the 68% limit of posterior distributions of the parameters in the broken power-law model (1.2) of IGW with $n_{\text{T,CMB}}$ free. We also show the 95% limit for r in parentheses.

And we use CLASS [69] to calculate the evolutions of GWs and other components, such as photons and baryons, and cobaya [70] with the MCMC Metropolis sampler and oversampling the nuisance parameters and $\{n_{\text{T}}^{\text{PTA}}, k_{\text{break}}\}$ to speed up our calculation.

Here, we also need to calculate the energy spectrum $\Omega_{\text{GW}}(f)$ of IGW. As in the analysis of NANOGrav [13], around the PTA scales we have

$$\Omega_{\text{GW}}(f) = \frac{\Omega_{\text{r}}}{24} \left(\frac{g_*(f)}{g_*^0} \right) \left(\frac{g_{*,s}^0}{g_{*,s}(f)} \right)^{4/3} P_{\text{T}}(f), \quad (2.1)$$

where Ω_{r} is the ratio of radiation to critical energy density at present, and g_*^0 and $g_{*,s}^0$ are the effective number of relativistic degrees of freedom for energy and entropy, respectively, while $g_*(f)$ and $g_{*,s}(f)$ correspond to the quantities when the modes with comoving wavenumber $k = 2\pi a_0 f$ reentered the Hubble horizon, see e.g. ref. [71].

3 Results

In this section, we will focus on the broken power-law model (1.2) of IGW with $n_{\text{T}}^{\text{CMB}}$ free and $n_{\text{T}}^{\text{CMB}} = -r/8$, respectively.

3.1 $n_{\text{T}}^{\text{CMB}}$ is free

The results are presented in figure 1 and table 1. As expected, the bestfit value $\log_{10} k_{\text{break}} = 4.19$, which is just between the CMB and PTA bands. The posterior of the spectral index $n_{\text{T}}^{\text{PTA}}$ at PTA band is $n_{\text{T}}^{\text{PTA}} \simeq 2$, in agreement with the results of NANOGrav [13, 19],⁵ since it is constrained mainly by the PTA itself. According to eq. (1.4), we have $P_{\text{T}} \sim r k_{\text{T}}^{n_{\text{T}}^{\text{PTA}}} k_{\text{break}}^{n_{\text{T}}^{\text{CMB}} - n_{\text{T}}^{\text{PTA}}}$ for fixed k_{pivot} , which suggests that $n_{\text{T}}^{\text{CMB}}$ is correlated positively with $\log_{10} k_{\text{break}}$, as showed in figure 1.

The 95% upper limit of $r_{0.05}$ is $r_{0.05} < 0.107$, higher than that of Planck+BK18 (e.g. $r \lesssim 0.03$ in refs. [8, 72]), while the 95% upper limit of $r_{0.01}$ is $r_{0.01} < 0.0338$. However, this is a natural result, since for a blue-tilted IGW with the bestfit $n_{\text{T}}^{\text{CMB}} = 0.428$, the amplitude of IGW must be lower at larger scale (smaller k_{pivot}). Accordingly, the constraint will be notably different at different $k_{\text{pivot}} \lesssim 0.05 \text{ Mpc}^{-1}$.

⁵Here, our results corresponds to the $T_{\text{rh}} \gtrsim 1 \text{ GeV}$ part of ref. [13].




Figure 1. Marginalized posterior distributions of the parameters in the broken power-law model (1.2) of IGW with $n_{T,CMB}$ free.

However, it is significant to find that unlike that of NANOGrav [13] at CMB band $n_T^{CMB} \simeq 2$ is not favored, instead $n_T^{CMB} = 0.55^{+0.37}_{-0.10}$ with the bestfit $n_T^{CMB} = 0.428$, while $n_T^{CMB} \simeq 0$ is still in the 95% CL. range, which suggests that a slow-roll period of inflation at CMB band is not excluded. Thus it is interesting to see what if we fix $n_T^{CMB} = -r/8$.

3.2 $n_T^{CMB} = -r/8$

Next, we fix $n_{T,CMB} = -r/8$, i.e. standard slow-roll inflation is not broken at CMB scale. The results are presented in figure 2 and table 2. As expected, the broken scale is $k_{break} = 2.72^{+0.92}_{-0.40}$, which is well between the CMB and PTA bands, and the posterior of the spectral index n_T^{PTA} at PTA band is still $n_T^{PTA} \simeq 2$.

The 95% CL. range of r is $r_{0.05} = 0.019^{+0.020}_{-0.019}$, which is similar to that of $r_{0.01}$, since unlike in section 3.1, we have $n_T^{CMB} = -r/8 \simeq 0$ nearly scale-invariant. This upper limit, $r \lesssim 0.039$, is comparable with (but slightly higher than) that of Planck+BK18 [8, 72]. The




Figure 2. Marginalized posterior distributions of the parameters in the broken power-law model (1.2) with slow-roll $n_{\text{T,CMB}}$.

Parameter	Best fit	68% limits
$r_{0.05}$	0.0091	$0.0194^{+0.0069}_{-0.014}$ ($+0.020$ -0.019)
$r_{0.01}$	0.0086	$0.0184^{+0.0065}_{-0.013}$ ($+0.019$ -0.018)
$n_{\text{T}}^{\text{PTA}}$	2.198	$2.04^{+0.39}_{-0.34}$
$\log_{10} k_{\text{break}}$	3.01	$2.72^{+0.92}_{-0.40}$

Table 2. The bestfit values and the 68% limit of posterior distributions of the parameters of the broken power-law model (1.2) with slow-roll $n_{\text{T,CMB}}$. We also show the 95% limit for r in parentheses.

bestfit r is $r_{0.05} = 0.0091$, however, it is unexpected that $r = 0$ is at 2σ level. According to eq. (1.2), the bound on r seems to be indirectly affect the NANOGrav results ($n_{\text{T}}^{\text{PTA}}$, k_{break}).

4 Discussion

4.1 Conclusion

In conclusion, we find that though the bestfit spectral tilt of IGW at PTA band is $n_{\text{T}}^{\text{PTA}} \simeq 2$, unlike that of the NANOGrav [13] $n_{\text{T}}^{\text{CMB}} \simeq 2$ is not favored at CMB band, instead the bestfit is $n_{\text{T}}^{\text{CMB}} = 0.55^{+0.37}_{-0.10}$, while a detectable amplitude of r with $n_{\text{T}}^{\text{CMB}} \simeq 0$ is still compatible. In particular if we fix $n_{\text{T}}^{\text{CMB}} = -r/8$ in light of standard slow-roll inflation, the upper bound on r is $r \lesssim 0.039$, slightly higher than that of Planck+BK18 [8]




Figure 3. The bestfit energy spectrum $\Omega_{GW}(f)$ of IGWB with n_T^{CMB} free and $n_T^{\text{CMB}} = -r/8$, respectively. We also show the sensitivity curves for LiteBIRD, SKA (Square Kilometre Array), space-based laser interferometers: LISA and Taiji. Here, we use the transfer function showed in ref. [73], which can return to eq. (2.1) at the high frequency limit, see also refs. [61, 74–77]. As commented in section 4, when $f > 10^{-7}$ the highly-blue ($n_T^{\text{PTA}} \simeq 2$) tilt needs to be cut off, though we plot $\Omega_{GW}(f > 10^{-7})$ with a straight line, which but is actually model-dependent, e.g. [62, 63].

In figure 3, we show our bestfit energy spectrum $\Omega_{GW}(f)$ of IGWB with n_T^{CMB} free and $n_T^{\text{CMB}} = -r/8$, respectively. The sensitivity curves for LiteBIRD and SKA are based on the results of ref. [78]. Here, our bestfit $\Omega_{GW}(f)$ covers the band at $f \sim 10^{-18} - 10^{-8}$ Hz, which naturally offers a guide for building relevant models of inflation and exploring new physics at corresponding band, since such a bestfit $\Omega_{GW}(f)$ can simultaneously explain current observations at both CMB and PTA bands.

The GW energy density contributes to the effective number of relativistic degrees of freedom N_{eff} at the time of BBN, which can be constrained by observations (see e.g. [79, 80]):

$$N_{\text{eff}}^{\text{GW}} = \left(3.046 + \frac{8}{7} \left(\frac{11}{4} \right)^{4/3} \right) \frac{1}{12} \int d \ln k P_T \quad (4.1)$$

The IR cutoff for the integral is the comoving horizon at the time of BBN $f_{\text{IR}} \simeq 10^{-10}$ Hz. Our expression of the power spectrum is assumed to be valid until $f = 2.97 \times 10^{-8}$ Hz, which is the frequency of the 14th bin of the NANOGrav 15-yr result. We find a contribution of $\Delta N_{\text{eff}} = 0.012$ for a free n_T^{CMB} and $\Delta N_{\text{eff}} = 0.014$ for $n_T^{\text{CMB}} = -r/8$, which is consistent with current constraint $\Delta N_{\text{eff}} \lesssim 0.4$ (95% CL., see e.g. [52]).

However, beyond the PTA band the highly-blue ($n_T^{\text{PTA}} \simeq 2$) tilt must be cut off at a certain $k_{\text{cutoff}} = 2\pi f_{\text{cutoff}}$ to avoid the conflation with the BBN bound. For example, if the power spectrum maintains flat till some $f_{\text{UV}} \sim 10^{86}$ when $f > 10^{-7}$, it will lead to $\Delta N_{\text{eff}} = 1.01$ for a free n_T^{CMB} and $\Delta N_{\text{eff}} = 1.27$ for $n_T^{\text{CMB}} = -r/8$. We find a cutoff frequency $f_{\text{cutoff}} \simeq 10^{-3}$ can produce $\Delta N_{\text{eff}} \simeq 0.4$, which has not been excluded by the current constraint. Besides, if the power spectrum is kept flat till $f = 20\text{--}76.6$ Hz, the energy density of the GWs

⁶The choice of f_{UV} has many assumptions. Here we consider the value in refs. [79, 81].

(3.72×10^{-7} for a free n_T^{CMB} and 5.45×10^{-7} for $n_T^{\text{CMB}} = -r/8$) will violate the upper limit of the LIGO/Virgo constraint $\Omega_{\text{GW}} \lesssim 5.8 \times 10^{-9}$ in the band 20-76.6 Hz [82]. It also required the cutoff to not conflict with these results. In figure 3, what is P_T at $k > k_{\text{cutoff}}$ is not relevant with our MCMC results, which actually is model-dependent, e.g. [62, 63], see also discussion in section 4.2. Thus the spectrum of IGWB at that band is open, however, the space-based laser interferometers, LISA and Taiji, might detect the corresponding IGWB signal.

In next decade, with the accumulations of PTA and CMB data, SGWB at PTA band would be confirmed eventually, and also upcoming CMB observations, such as CMB-S4 [83], LiteBIRD [84] would bring us more information on IGWB at CMB band. In view of this, our work is suggesting that a joint MCMC analysis of PTA and CMB data might be indispensable, which would open a unforeseen door into our very early Universe.

4.2 Implications for inflation

It is interesting to see what our results will imply for inflation. As commented in ref. [19], a blue-tilted IGW suggests that inflation must violate null energy condition (NEC), i.e. $T_{\mu\nu}n^\mu n^\nu < 0$ (equivalently $\dot{H} > 0$ or $\epsilon = -\frac{\dot{H}}{H^2} < 0$), e.g. [38, 39], see also [43, 44], if the initial Bunch-Davis state of perturbation modes is not modified. Though it is not difficult to achieve $n_T^{\text{PTA}} \simeq 2$, which requires $\epsilon \ll -1$, e.g. [40, 56], such a highly blue tilt of IGWB implies that inflation must end at a low scale [13, 19], or else it will be conflicted with the BBN bound on relativistic components.

However, it might be possible that the violation of NEC happened before a (standard) slow-roll inflation, so that the blue-tilted P_T is cut off at certain k_{cutoff} beyond the PTA band (when $k > k_{\text{cutoff}}$ the spectrum is flat). In corresponding models, the energy spectrum of IGW will be shown as the black solid $\Omega_{\text{GW}}(f)$ curve in figure 3, and at CMB band for such a period of NEC violation the scale-invariant primordial scalar perturbation required by Planck observations can be harvested in light of [38, 43, 56–59, 85].

It is also possible that the violation of NEC might be just short-lived, which happened between two (slow-roll) periods of inflation with $H \simeq H_{\text{infl1}}$ and a higher scale $H \simeq H_{\text{infl2}} \gg H_{\text{infl1}}$, respectively. In corresponding model [54, 86, 87], we approximately have

$$P_T(k) \sim \frac{H_{\text{infl1}}^2 + \mathcal{A} \left(\frac{k}{k_{\text{cutoff}}} \right)^{n_T^{\text{PTA}}} H_{\text{infl2}}^2}{1 + \mathcal{A} \left(\frac{k}{k_{\text{cutoff}}} \right)^{n_T^{\text{PTA}}}}, \quad (4.2)$$

where $\mathcal{A} = \frac{\Gamma^2(\nu)}{\pi} \left(\frac{2\nu-1}{4} \right)^{1-2\nu} \sim \mathcal{O}(1)$ with $\nu = 1/2 + \frac{1}{1-\epsilon}$,

$$n_T^{\text{PTA}} = -\frac{2\epsilon}{1-\epsilon} \simeq 2, \quad \text{for } \epsilon \ll 1. \quad (4.3)$$

According to eq. (4.2), we have $P_T \simeq H_{\text{infl1}}^2$ for $k/k_{\text{cutoff}} \ll (H_{\text{infl1}}/H_{\text{infl2}})^{2/n_T^{\text{PTA}}} \ll 1$ corresponding the CMB band, while $P_T \sim k^{n_T^{\text{PTA}}} \simeq k^2$ for $(H_{\text{infl1}}/H_{\text{infl2}})^{2/n_T^{\text{PTA}}} \ll k/k_{\text{cutoff}} \ll 1$ corresponding the PTA band. Thus when $k/k_{\text{cutoff}} \ll 1$, we have

$$P_T(k) \sim H_{\text{infl1}}^2 \left[1 + \mathcal{A} \left(\frac{k}{k_{\text{cutoff}}} \right)^{n_T^{\text{PTA}}} \frac{H_{\text{infl2}}^2}{H_{\text{infl1}}^2} \right]. \quad (4.4)$$

It is just eq. (1.2) for $k_{\text{break}} \simeq k_{\text{cutoff}} (\frac{H_{\text{inf1}}}{H_{\text{inf2}}})^{2/n_{\text{T}}^{\text{PTA}}}$ and $n_{\text{T}}^{\text{CMB}} \simeq 0$. Thus the energy spectrum will be showed as the black dashed $\Omega_{\text{GW}}(f)$ curve in figure 3. Here, k_{cutoff} is a cutoff scale, see also section 4.1, beyond which $P_{\text{T}} \sim H_{\text{inf2}}^2$ is flat, or might be modified as in e.g. refs. [61–63]. Thus the IGWB at $k > k_{\text{cutoff}}$ ($f > 10^{-7}\text{Hz}$ in figure 3) is model-dependent, which is unknown and needs to be explored, which suggests that the requirement of BBN that inflation must end at a low scale [13, 19] is not workable.

It is also interesting to investigate other mechanisms, in which blue-tilted IGW at PTA band is sourced by other components during slow-roll inflation, such as the gauge fields [88–94], the non-Bunch-Davis initial states [95, 96], modified gravity, e.g. [97–100], and the collisions of bubbles nucleated [101, 102]. In light of our MCMC results on r , $n_{\text{T}}^{\text{CMB}}$ and $n_{\text{T}}^{\text{PTA}}$, and also bestfit $\Omega_{\text{GW}}(f)$ in figure 3, it is interesting to resurvey the relevant models.

Acknowledgments

YSP is supported by NSFC, No.12075246 and the Fundamental Research Funds for the Central Universities. Y. C. is supported in part by the National Natural Science Foundation of China (Grant No. 11905224), the China Postdoctoral Science Foundation (Grant No. 2021M692942) and Zhengzhou University (Grant No. 32340282). GY is supported by NWO and the Dutch Ministry of Education, Culture and Science (OCW) (grant VI.Vidi.192.069).

References

- [1] A.H. Guth, *The inflationary universe: a possible solution to the horizon and flatness problems*, *Phys. Rev. D* **23** (1981) 347 [INSPIRE].
- [2] A.D. Linde, *A new inflationary universe scenario: a possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems*, *Phys. Lett. B* **108** (1982) 389 [INSPIRE].
- [3] A. Albrecht and P.J. Steinhardt, *Cosmology for grand unified theories with radiatively induced symmetry breaking*, *Phys. Rev. Lett.* **48** (1982) 1220 [INSPIRE].
- [4] A.A. Starobinsky, *A new type of isotropic cosmological models without singularity*, *Phys. Lett. B* **91** (1980) 99 [INSPIRE].
- [5] A.D. Linde, *Chaotic inflation*, *Phys. Lett. B* **129** (1983) 177 [INSPIRE].
- [6] M. Kamionkowski, A. Kosowsky and A. Stebbins, *Statistics of cosmic microwave background polarization*, *Phys. Rev. D* **55** (1997) 7368 [astro-ph/9611125] [INSPIRE].
- [7] M. Kamionkowski, A. Kosowsky and A. Stebbins, *A probe of primordial gravity waves and vorticity*, *Phys. Rev. Lett.* **78** (1997) 2058 [astro-ph/9609132] [INSPIRE].
- [8] BICEP and KECK collaborations, *Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season*, *Phys. Rev. Lett.* **127** (2021) 151301 [arXiv:2110.00483] [INSPIRE].
- [9] NANOGrav collaboration, *The NANOGrav 15 yr data set: evidence for a gravitational-wave background*, *Astrophys. J. Lett.* **951** (2023) L8 [arXiv:2306.16213] [INSPIRE].
- [10] EPTA and INPTA: collaborations, *The second data release from the European Pulsar Timing Array — III. Search for gravitational wave signals*, *Astron. Astrophys.* **678** (2023) A50 [arXiv:2306.16214] [INSPIRE].

- [11] D.J. Reardon et al., *Search for an isotropic gravitational-wave background with the Parkes Pulsar Timing Array*, *Astrophys. J. Lett.* **951** (2023) L6 [[arXiv:2306.16215](#)] [[INSPIRE](#)].
- [12] H. Xu et al., *Searching for the nano-Hertz stochastic gravitational wave background with the Chinese pulsar timing array data release I*, *Res. Astron. Astrophys.* **23** (2023) 075024 [[arXiv:2306.16216](#)] [[INSPIRE](#)].
- [13] NANOGrav collaboration, *The NANOGrav 15 yr data set: search for signals from new physics*, *Astrophys. J. Lett.* **951** (2023) L11 [[arXiv:2306.16219](#)] [[INSPIRE](#)].
- [14] EPTA collaboration, *The second data release from the European Pulsar Timing Array: V. Implications for massive black holes, dark matter and the early universe*, [arXiv:2306.16227](#) [[INSPIRE](#)].
- [15] T. Broadhurst, C. Chen, T. Liu and K.-F. Zheng, *Binary supermassive black holes orbiting dark matter solitons: from the dual AGN in UGC4211 to nano-Hertz gravitational waves*, [arXiv:2306.17821](#) [[INSPIRE](#)].
- [16] H.-L. Huang et al., *Supermassive primordial black holes in multiverse: for nano-Hertz gravitational wave and high-redshift JWST galaxies*, [arXiv:2306.17577](#) [[INSPIRE](#)].
- [17] P.F. Depta, K. Schmidt-Hoberg, P. Schwaller and C. Tasillo, *Do pulsar timing arrays observe merging primordial black holes?*, [arXiv:2306.17836](#) [[INSPIRE](#)].
- [18] S. Vagnozzi, *Implications of the NANOGrav results for inflation*, *Mon. Not. Roy. Astron. Soc.* **502** (2021) L11 [[arXiv:2009.13432](#)] [[INSPIRE](#)].
- [19] S. Vagnozzi, *Inflationary interpretation of the stochastic gravitational wave background signal detected by pulsar timing array experiments*, *JHEAp* **39** (2023) 81 [[arXiv:2306.16912](#)] [[INSPIRE](#)].
- [20] I. Ben-Dayan, U. Kumar, U. Thattarampilly and A. Verma, *Probing the early universe cosmology with NANOGrav: possibilities and limitations*, *Phys. Rev. D* **108** (2023) 103507 [[arXiv:2307.15123](#)] [[INSPIRE](#)].
- [21] E. Madge et al., *Primordial gravitational waves in the nano-Hertz regime and PTA data — towards solving the GW inverse problem*, *JHEP* **10** (2023) 171 [[arXiv:2306.14856](#)] [[INSPIRE](#)].
- [22] G. Franciolini, J.A. Iovino, V. Vaskonen and H. Veermäe, *Recent gravitational wave observation by pulsar timing arrays and primordial black holes: the importance of non-gaussianities*, *Phys. Rev. Lett.* **131** (2023) 201401 [[arXiv:2306.17149](#)] [[INSPIRE](#)].
- [23] T. Ghosh et al., *Did we hear the sound of the universe boiling? Analysis using the full fluid velocity profiles and NANOGrav 15-year data*, [arXiv:2307.02259](#) [[INSPIRE](#)].
- [24] X. Niu and M.H. Rahat, *NANOGrav signal from axion inflation*, *Phys. Rev. D* **108** (2023) 115023 [[arXiv:2307.01192](#)] [[INSPIRE](#)].
- [25] R.A. Konoplya and A. Zhidenko, *Asymptotic tails of massive gravitons in light of pulsar timing array observations*, [arXiv:2307.01110](#) [[INSPIRE](#)].
- [26] S.-P. Li and K.-P. Xie, *Collider test of nano-Hertz gravitational waves from pulsar timing arrays*, *Phys. Rev. D* **108** (2023) 055018 [[arXiv:2307.01086](#)] [[INSPIRE](#)].
- [27] P. Di Bari and M.H. Rahat, *The split majoron model confronts the NANOGrav signal*, [arXiv:2307.03184](#) [[INSPIRE](#)].
- [28] S. Wang, Z.-C. Zhao and Q.-H. Zhu, *Constraints on scalar-induced gravitational waves up to third order from a joint analysis of BBN, CMB, and PTA data*, *Phys. Rev. Res.* **6** (2024) 013207 [[arXiv:2307.03095](#)] [[INSPIRE](#)].
- [29] X.K. Du, M.X. Huang, F. Wang and Y.K. Zhang, *Did the nHz gravitational waves signatures observed by NANOGrav indicate multiple sector SUSY breaking?*, [arXiv:2307.02938](#) [[INSPIRE](#)].

- [30] G. Ye and A. Silvestri, *Can the gravitational wave background feel wiggles in spacetime?*, *Astrophys. J. Lett.* **963** (2024) L15 [[arXiv:2307.05455](#)] [[INSPIRE](#)].
- [31] S. Balaji, G. Domènech and G. Franciolini, *Scalar-induced gravitational wave interpretation of PTA data: the role of scalar fluctuation propagation speed*, *JCAP* **10** (2023) 041 [[arXiv:2307.08552](#)] [[INSPIRE](#)].
- [32] Z. Zhang et al., *Nano-Hertz gravitational waves from collapsing domain walls associated with freeze-in dark matter in light of pulsar timing array observations*, *Phys. Rev. D* **108** (2023) 095037 [[arXiv:2307.11495](#)] [[INSPIRE](#)].
- [33] V.K. Oikonomou, *Flat energy spectrum of primordial gravitational waves versus peaks and the NANOGrav 2023 observation*, *Phys. Rev. D* **108** (2023) 043516 [[arXiv:2306.17351](#)] [[INSPIRE](#)].
- [34] A. Ashoorioon, K. Rezazadeh and A. Rostami, *NANOGrav signal from the end of inflation and the LIGO mass and heavier primordial black holes*, *Phys. Lett. B* **835** (2022) 137542 [[arXiv:2202.01131](#)] [[INSPIRE](#)].
- [35] K. Cheung, C.J. Ouseph and P.-Y. Tseng, *NANOGrav signal and PBH from the modified Higgs inflation*, [arXiv:2307.08046](#) [[INSPIRE](#)].
- [36] X.-F. Li, *Probing the high temperature symmetry breaking with gravitational waves from domain walls*, [arXiv:2307.03163](#) [[INSPIRE](#)].
- [37] S.A. Hosseini Mansoori, F. Felegray, A. Talebian and M. Sami, *PBHs and GWs from T^2 -inflation and NANOGrav 15-year data*, *JCAP* **08** (2023) 067 [[arXiv:2307.06757](#)] [[INSPIRE](#)].
- [38] Y.-S. Piao and Y.-Z. Zhang, *Phantom inflation and primordial perturbation spectrum*, *Phys. Rev. D* **70** (2004) 063513 [[astro-ph/0401231](#)] [[INSPIRE](#)].
- [39] Y.-S. Piao and E. Zhou, *Nearly scale invariant spectrum of adiabatic fluctuations may be from a very slowly expanding phase of the universe*, *Phys. Rev. D* **68** (2003) 083515 [[hep-th/0308080](#)] [[INSPIRE](#)].
- [40] Y.-S. Piao and Y.-Z. Zhang, *The primordial perturbation spectrum from various expanding and contracting phases*, *Phys. Rev. D* **70** (2004) 043516 [[astro-ph/0403671](#)] [[INSPIRE](#)].
- [41] M. Baldi, F. Finelli and S. Matarrese, *Inflation with violation of the null energy condition*, *Phys. Rev. D* **72** (2005) 083504 [[astro-ph/0505552](#)] [[INSPIRE](#)].
- [42] Z.-G. Liu, J. Zhang and Y.-S. Piao, *Phantom inflation with a steplike potential*, *Phys. Lett. B* **697** (2011) 407 [[arXiv:1012.0673](#)] [[INSPIRE](#)].
- [43] T. Kobayashi, M. Yamaguchi and J. Yokoyama, *G-inflation: inflation driven by the Galileon field*, *Phys. Rev. Lett.* **105** (2010) 231302 [[arXiv:1008.0603](#)] [[INSPIRE](#)].
- [44] T. Kobayashi, M. Yamaguchi and J. Yokoyama, *Generalized G-inflation: inflation with the most general second-order field equations*, *Prog. Theor. Phys.* **126** (2011) 511 [[arXiv:1105.5723](#)] [[INSPIRE](#)].
- [45] T. Fujita, S. Kuroyanagi, S. Mizuno and S. Mukohyama, *Blue-tilted primordial gravitational waves from massive gravity*, *Phys. Lett. B* **789** (2019) 215 [[arXiv:1808.02381](#)] [[INSPIRE](#)].
- [46] S. Endlich, A. Nicolis and J. Wang, *Solid inflation*, *JCAP* **10** (2013) 011 [[arXiv:1210.0569](#)] [[INSPIRE](#)].
- [47] D. Cannone, G. Tasinato and D. Wands, *Generalised tensor fluctuations and inflation*, *JCAP* **01** (2015) 029 [[arXiv:1409.6568](#)] [[INSPIRE](#)].
- [48] W. Giarè, F. Renzi and A. Melchiorri, *Higher-curvature corrections and tensor modes*, *Phys. Rev. D* **103** (2021) 043515 [[arXiv:2012.00527](#)] [[INSPIRE](#)].

- [49] Y. Cai, Y.-T. Wang and Y.-S. Piao, *Propagating speed of primordial gravitational waves and inflation*, *Phys. Rev. D* **94** (2016) 043002 [[arXiv:1602.05431](#)] [[INSPIRE](#)].
- [50] Y. Cai, Y.-T. Wang and Y.-S. Piao, *Is there an effect of a nontrivial c_T during inflation?*, *Phys. Rev. D* **93** (2016) 063005 [[arXiv:1510.08716](#)] [[INSPIRE](#)].
- [51] M. Baumgart, J.J. Heckman and L. Thomas, *CFTs blueshift tensor fluctuations universally*, *JCAP* **07** (2022) 034 [[arXiv:2109.08166](#)] [[INSPIRE](#)].
- [52] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [*Erratum ibid.* **652** (2021) C4] [[arXiv:1807.06209](#)] [[INSPIRE](#)].
- [53] PLANCK collaboration, *Planck 2018 results. X. Constraints on inflation*, *Astron. Astrophys.* **641** (2020) A10 [[arXiv:1807.06211](#)] [[INSPIRE](#)].
- [54] Y. Cai and Y.-S. Piao, *Intermittent null energy condition violations during inflation and primordial gravitational waves*, *Phys. Rev. D* **103** (2021) 083521 [[arXiv:2012.11304](#)] [[INSPIRE](#)].
- [55] H.W.H. Tahara and T. Kobayashi, *Nanohertz gravitational waves from a null-energy-condition violation in the early universe*, *Phys. Rev. D* **102** (2020) 123533 [[arXiv:2011.01605](#)] [[INSPIRE](#)].
- [56] Z.-G. Liu, J. Zhang and Y.-S. Piao, *A Galileon design of slow expansion*, *Phys. Rev. D* **84** (2011) 063508 [[arXiv:1105.5713](#)] [[INSPIRE](#)].
- [57] Z.-G. Liu and Y.-S. Piao, *A Galileon design of slow expansion: emergent universe*, *Phys. Lett. B* **718** (2013) 734 [[arXiv:1207.2568](#)] [[INSPIRE](#)].
- [58] S. Nishi and T. Kobayashi, *Generalized Galilean genesis*, *JCAP* **03** (2015) 057 [[arXiv:1501.02553](#)] [[INSPIRE](#)].
- [59] S. Nishi and T. Kobayashi, *Scale-invariant perturbations from null-energy-condition violation: a new variant of Galilean genesis*, *Phys. Rev. D* **95** (2017) 064001 [[arXiv:1611.01906](#)] [[INSPIRE](#)].
- [60] W. Giarè, M. Forconi, E. Di Valentino and A. Melchiorri, *Towards a reliable calculation of relic radiation from primordial gravitational waves*, *Mon. Not. Roy. Astron. Soc.* **520** (2023) 2 [[arXiv:2210.14159](#)] [[INSPIRE](#)].
- [61] S. Kuroyanagi, T. Takahashi and S. Yokoyama, *Blue-tilted tensor spectrum and thermal history of the universe*, *JCAP* **02** (2015) 003 [[arXiv:1407.4785](#)] [[INSPIRE](#)].
- [62] S. Kuroyanagi, T. Takahashi and S. Yokoyama, *Blue-tilted inflationary tensor spectrum and reheating in the light of NANOGrav results*, *JCAP* **01** (2021) 071 [[arXiv:2011.03323](#)] [[INSPIRE](#)].
- [63] M. Benetti, L.L. Graef and S. Vagnozzi, *Primordial gravitational waves from NANOGrav: a broken power-law approach*, *Phys. Rev. D* **105** (2022) 043520 [[arXiv:2111.04758](#)] [[INSPIRE](#)].
- [64] W.G. Lamb, S.R. Taylor and R. van Haasteren, *Rapid refitting techniques for Bayesian spectral characterization of the gravitational wave background using pulsar timing arrays*, *Phys. Rev. D* **108** (2023) 103019 [[arXiv:2303.15442](#)] [[INSPIRE](#)].
- [65] G. Ye, B. Hu and Y.-S. Piao, *Implication of the Hubble tension for the primordial Universe in light of recent cosmological data*, *Phys. Rev. D* **104** (2021) 063510 [[arXiv:2103.09729](#)] [[INSPIRE](#)].
- [66] G. Ye and Y.-S. Piao, *Is the Hubble tension a hint of AdS phase around recombination?*, *Phys. Rev. D* **101** (2020) 083507 [[arXiv:2001.02451](#)] [[INSPIRE](#)].
- [67] J.-Q. Jiang and Y.-S. Piao, *Toward early dark energy and $n_s = 1$ with Planck, ACT, and SPT observations*, *Phys. Rev. D* **105** (2022) 103514 [[arXiv:2202.13379](#)] [[INSPIRE](#)].

- [68] J.-Q. Jiang, G. Ye and Y.-S. Piao, *Return of Harrison-Zeldovich spectrum in light of recent cosmological tensions*, *Mon. Not. Roy. Astron. Soc.* **527** (2023) L54 [[arXiv:2210.06125](#)] [[INSPIRE](#)].
- [69] D. Blas, J. Lesgourgues and T. Tram, *The Cosmic Linear Anisotropy Solving System (CLASS) II: approximation schemes*, *JCAP* **07** (2011) 034 [[arXiv:1104.2933](#)] [[INSPIRE](#)].
- [70] J. Torrado and A. Lewis, *Cobaya: code for Bayesian analysis of hierarchical physical models*, *JCAP* **05** (2021) 057 [[arXiv:2005.05290](#)] [[INSPIRE](#)].
- [71] K. Saikawa and S. Shirai, *Precise WIMP dark matter abundance and standard model thermodynamics*, *JCAP* **08** (2020) 011 [[arXiv:2005.03544](#)] [[INSPIRE](#)].
- [72] M. Tristram et al., *Improved limits on the tensor-to-scalar ratio using BICEP and Planck data*, *Phys. Rev. D* **105** (2022) 083524 [[arXiv:2112.07961](#)] [[INSPIRE](#)].
- [73] T. Kite, J. Chluba, A. Ravenni and S.P. Patil, *Clarifying transfer function approximations for the large-scale gravitational wave background in Λ CDM*, *Mon. Not. Roy. Astron. Soc.* **509** (2021) 1366 [[arXiv:2107.13351](#)] [[INSPIRE](#)].
- [74] M.S. Turner, M.J. White and J.E. Lidsey, *Tensor perturbations in inflationary models as a probe of cosmology*, *Phys. Rev. D* **48** (1993) 4613 [[astro-ph/9306029](#)] [[INSPIRE](#)].
- [75] L.A. Boyle and P.J. Steinhardt, *Probing the early universe with inflationary gravitational waves*, *Phys. Rev. D* **77** (2008) 063504 [[astro-ph/0512014](#)] [[INSPIRE](#)].
- [76] W. Zhao and Y. Zhang, *Relic gravitational waves and their detection*, *Phys. Rev. D* **74** (2006) 043503 [[astro-ph/0604458](#)] [[INSPIRE](#)].
- [77] X.-J. Liu, W. Zhao, Y. Zhang and Z.-H. Zhu, *Detecting relic gravitational waves by pulsar timing arrays: effects of cosmic phase transitions and relativistic free-streaming gases*, *Phys. Rev. D* **93** (2016) 024031 [[arXiv:1509.03524](#)] [[INSPIRE](#)].
- [78] P. Campeti, E. Komatsu, D. Poletti and C. Baccigalupi, *Measuring the spectrum of primordial gravitational waves with CMB, PTA and laser interferometers*, *JCAP* **01** (2021) 012 [[arXiv:2007.04241](#)] [[INSPIRE](#)].
- [79] P.D. Meerburg, R. Hložek, B. Hadzhiyska and J. Meyers, *Multiwavelength constraints on the inflationary consistency relation*, *Phys. Rev. D* **91** (2015) 103505 [[arXiv:1502.00302](#)] [[INSPIRE](#)].
- [80] I. Ben-Dayan, B. Keating, D. Leon and I. Wolfson, *Constraints on scalar and tensor spectra from N_{eff}* , *JCAP* **06** (2019) 007 [[arXiv:1903.11843](#)] [[INSPIRE](#)].
- [81] G. Cabass et al., *Updated constraints and forecasts on primordial tensor modes*, *Phys. Rev. D* **93** (2016) 063508 [[arXiv:1511.05146](#)] [[INSPIRE](#)].
- [82] KAGRA et al. collaborations, *Upper limits on the isotropic gravitational-wave background from advanced LIGO and advanced Virgo's third observing run*, *Phys. Rev. D* **104** (2021) 022004 [[arXiv:2101.12130](#)] [[INSPIRE](#)].
- [83] CMB-S4 collaboration, *CMB-S4: forecasting constraints on primordial gravitational waves*, *Astrophys. J.* **926** (2022) 54 [[arXiv:2008.12619](#)] [[INSPIRE](#)].
- [84] M. Hazumi et al., *LiteBIRD: a satellite for the studies of B-mode polarization and inflation from cosmic background radiation detection*, *J. Low Temp. Phys.* **194** (2019) 443 [[INSPIRE](#)].
- [85] Y.-S. Piao, *Adiabatic spectra during slowly evolving*, *Phys. Lett. B* **701** (2011) 526 [[arXiv:1012.2734](#)] [[INSPIRE](#)].
- [86] Y. Cai and Y.-S. Piao, *Generating enhanced primordial GWs during inflation with intermittent violation of NEC and diminishment of GW propagating speed*, *JHEP* **06** (2022) 067 [[arXiv:2201.04552](#)] [[INSPIRE](#)].

- [87] Y. Cai, M. Zhu and Y.-S. Piao, *Primordial black holes from null energy condition violation during inflation*, [arXiv:2305.10933](#) [[INSPIRE](#)].
- [88] M.M. Anber and L. Sorbo, *Non-Gaussianities and chiral gravitational waves in natural steep inflation*, *Phys. Rev. D* **85** (2012) 123537 [[arXiv:1203.5849](#)] [[INSPIRE](#)].
- [89] J.L. Cook and L. Sorbo, *Particle production during inflation and gravitational waves detectable by ground-based interferometers*, *Phys. Rev. D* **85** (2012) 023534 [Erratum *ibid.* **86** (2012) 069901] [[arXiv:1109.0022](#)] [[INSPIRE](#)].
- [90] S. Mukohyama, R. Namba, M. Peloso and G. Shiu, *Blue tensor spectrum from particle production during inflation*, *JCAP* **08** (2014) 036 [[arXiv:1405.0346](#)] [[INSPIRE](#)].
- [91] R. Namba et al., *Scale-dependent gravitational waves from a rolling axion*, *JCAP* **01** (2016) 041 [[arXiv:1509.07521](#)] [[INSPIRE](#)].
- [92] E. Dimastrogiovanni, M. Fasiello and T. Fujita, *Primordial gravitational waves from axion-gauge fields dynamics*, *JCAP* **01** (2017) 019 [[arXiv:1608.04216](#)] [[INSPIRE](#)].
- [93] I. Obata, *Chiral primordial blue tensor spectra from the axion-gauge couplings*, *JCAP* **06** (2017) 050 [[arXiv:1612.08817](#)] [[INSPIRE](#)].
- [94] R.R. Caldwell and C. Devulder, *Axion gauge field inflation and gravitational leptogenesis: a lower bound on B-modes from the matter-antimatter asymmetry of the universe*, *Phys. Rev. D* **97** (2018) 023532 [[arXiv:1706.03765](#)] [[INSPIRE](#)].
- [95] A. Ashoorioon, K. Dimopoulos, M.M. Sheikh-Jabbari and G. Shiu, *Non-Bunch-Davis initial state reconciles chaotic models with BICEP and Planck*, *Phys. Lett. B* **737** (2014) 98 [[arXiv:1403.6099](#)] [[INSPIRE](#)].
- [96] S. Choudhury, *Single field inflation in the light of pulsar timing array data: quintessential interpretation of blue tilted tensor spectrum through non-Bunch-Davies initial condition*, *Eur. Phys. J. C* **84** (2024) 278 [[arXiv:2307.03249](#)] [[INSPIRE](#)].
- [97] V.K. Oikonomou, *Amplification of the primordial gravitational waves energy spectrum by a kinetic scalar in $F(R)$ gravity*, *Astropart. Phys.* **144** (2023) 102777 [[arXiv:2209.09781](#)] [[INSPIRE](#)].
- [98] V.K. Oikonomou, *Primordial gravitational waves predictions for GW170817-compatible Einstein-Gauss-Bonnet theory*, *Astropart. Phys.* **141** (2022) 102718 [[arXiv:2204.06304](#)] [[INSPIRE](#)].
- [99] S.D. Odintsov, V.K. Oikonomou and F.P. Fronimos, *Quantitative predictions for $f(R)$ gravity primordial gravitational waves*, *Phys. Dark Univ.* **35** (2022) 100950 [[arXiv:2108.11231](#)] [[INSPIRE](#)].
- [100] S.D. Odintsov, V.K. Oikonomou and R. Myrzakulov, *Spectrum of primordial gravitational waves in modified gravities: a short overview*, *Symmetry* **14** (2022) 729 [[arXiv:2204.00876](#)] [[INSPIRE](#)].
- [101] H.-H. Li, G. Ye and Y.-S. Piao, *Is the NANOGrav signal a hint of dS decay during inflation?*, *Phys. Lett. B* **816** (2021) 136211 [[arXiv:2009.14663](#)] [[INSPIRE](#)].
- [102] Y.-T. Wang, Y. Cai and Y.-S. Piao, *Phase-transition sound of inflation at gravitational waves detectors*, *Phys. Lett. B* **789** (2019) 191 [[arXiv:1801.03639](#)] [[INSPIRE](#)].