



Universiteit
Leiden
The Netherlands

European Society for Immunodeficiencies guidelines for the management of patients with congenital athymia

Kreins, A.Y.; Dhalla, F.; Flinn, A.M.; Howley, E.; Ekwall, O.; Villa, A.; ... ; European Soc Immunodeficiencies Clinical Working Party

Citation

Kreins, A. Y., Dhalla, F., Flinn, A. M., Howley, E., Ekwall, O., Villa, A., ... Davies, E. G. (2024). European Society for Immunodeficiencies guidelines for the management of patients with congenital athymia. *Journal Of Allergy And Clinical Immunology*, 154(6), 1391-1408. doi:10.1016/j.jaci.2024.07.031

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)
Downloaded from: <https://hdl.handle.net/1887/4195197>

Note: To cite this publication please use the final published version (if applicable).

European Society for Immunodeficiencies guidelines for the management of patients with congenital athymia



Alexandra Y. Kreins, MD, PhD,^{a,b} Fatima Dhalla, MD, PhD,^{c,d} Aisling M. Flinn, MD, PhD,^{e,f,g} Evey Howley, RN, MSc,^a Olov Ekwall, MD, PhD,^{h,i} Anna Villa, MD,^{j,k} Frank J. T. Staal, PhD,^{l,m} Graham Anderson, PhD,ⁿ Andrew R. Gennery, MD,^{e,f} Georg A. Holländer, MD,^{c,o,p} and E. Graham Davies, MD,^{a,b} on behalf of the European Society for Immunodeficiencies Clinical Working Party
Birmingham, London, Newcastle upon Tyne, and Oxford, United Kingdom; Crumlin, Ireland; Gothenburg, Sweden; Milan, Italy; Leiden, The Netherlands; and Basel, Switzerland

Congenital athymia is a life-limiting disorder due to rare inborn errors of immunity causing impaired thymus organogenesis or abnormal thymic stromal cell development and function.

Athymic infants have a T-lymphocyte-negative, B-lymphocyte-positive, natural killer cell-positive immunophenotype with profound T-lymphocyte deficiency and are susceptible to severe infections and autoimmunity. Patients variably display syndromic features. Expanding access to newborn screening for severe combined immunodeficiency and T lymphocytopenia and broad genetic testing, including next-generation sequencing technologies, increasingly facilitate their timely identification.

The recommended first-line treatment is allogeneic thymus transplantation, which is a specialized procedure available in Europe and the United States. Outcomes for athymic patients are best with early diagnosis and thymus transplantation before the development of infectious and inflammatory complications. These guidelines on behalf of the European Society for Immunodeficiencies provide a comprehensive review for clinicians who manage patients with inborn thymic stromal cell defects; they offer clinical practice recommendations focused on the diagnosis, investigation, risk stratification, and management of congenital athymia with the aim of improving patient outcomes. (J Allergy Clin Immunol 2024;154:1391-408.)

Key words: Congenital athymia, thymus transplantation, 22q11.2 deletion syndrome, DiGeorge syndrome, immunology guidelines

From ^athe Department of Immunology and Gene Therapy, Great Ormond Street Hospital for Children NHS Foundation Trust, London; ^bthe Infection Immunity and Inflammation Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London; ^cthe Department of Paediatrics and Institute of Developmental and Regenerative Medicine, University of Oxford, Oxford; ^dthe Department of Clinical Immunology, Oxford University Hospitals NHS Foundation Trust, Oxford; ^ethe Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne; ^fthe Paediatric Stem Cell Transplant Unit, Great North Children's Hospital, Newcastle upon Tyne; ^gthe Department of Paediatric Immunology, Children's Health Ireland at Crumlin, Crumlin; ^hthe Department of Pediatrics, Institute of Clinical Sciences, The Sahlgrenska Academy, University of Gothenburg, Gothenburg; ⁱthe Department of Rheumatology and Inflammation Research, Institute of Medicine, The Sahlgrenska Academy, University of Gothenburg, Gothenburg; ^jthe San Raffaele Telethon Institute for Gene Therapy, IRCCS San Raffaele Hospital, Milan; ^kthe Istituto di Ricerca Genetica e Biomedica, Consiglio Nazionale Delle Ricerche (IRGB-CNR), Milan; ^lthe Department of Pediatrics, Pediatric Stem Cell Transplantation Program, Willem-Alexander Children's Hospital, Leiden; ^mthe Department of Immunology, Leiden University Medical Center, Leiden; ⁿthe Institute of Immunology and Immunotherapy, Medical School, University of Birmingham, Birmingham; ^oPaediatric Immunology, Department of Biomedicine, University of Basel and University Children's Hospital Basel, Basel; and ^pthe Department of Biosystems Science and Engineering, ETH Zurich, Basel.

The first 3 authors contributed equally to this article, and all should be considered first author.

Received for publication February 8, 2024; revised July 6, 2024; accepted for publication July 15, 2024.

Available online September 18, 2024.

Corresponding author: Alexandra Y. Kreins, MD, PhD, UCL GOS Institute of Child Health, Zayed Centre for Research into Rare Disease in Children, 20c Guilford St, London WC1N 1DZ, United Kingdom. E-mail: a.kreins@ucl.ac.uk.

The CrossMark symbol notifies online readers when updates have been made to the article such as errata or minor corrections

0091-6749

© 2024 The Authors. Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.1016/j.jaci.2024.07.031>

To generate a diverse T-lymphocyte repertoire that is tolerant to self-antigens, bone marrow-derived T-lymphocyte progenitors complete their differentiation and selection in the thymus.^{1,2} The thymic stroma, composed of epithelial cells, various vascular cell types, mesenchymal cells, and fibroblasts, provides the intricate and highly specialized 3-dimensional microenvironment critical for this process.^{1,2} Thymic epithelial cells (TEC) originate from the endodermal lining of the third pharyngeal pouch, whereas the nonepithelial components of the thymic stroma descend from the surrounding mesoderm and neural crest-derived mesenchyme.²⁻⁶ The correct patterning of the third pharyngeal pouch and early thymus organogenesis are tightly controlled by the action of several transcription factors, including TBX (T-box transcription factor) 1, HOXA3 (homeobox protein A3), PAX (paired box) 1, and PAX9.⁷⁻¹² Following a first-fate commitment, differentiation, growth, and function of TEC are dependent on the expression of FOXP1 (forkhead box N1), a member of the forkhead family of transcription factors.^{7,13-18} TEC can be subdivided on the basis of anatomic, phenotypic, functional, and more recently transcriptomic characteristics into separate subpopulations and subtypes, defined respectively by their expression of cell-surface markers or by their transcriptomes.¹⁹⁻²⁶ TEC resident in the outer cortex of the thymus (designated cortical TEC) control the early stages of thymopoiesis, including the attraction of blood-borne lymphoid precursors to the thymus, their commitment to a T-lymphocyte fate, and their expansion and positive selection.^{13,23,27-30} The latter constitutes a process that enables

Abbreviations used

22q11.2DS: 22q11.2 deletion syndrome

CHARGE syndrome: Syndrome comprising coloboma, heart defects, atresia of nasal choanae, retardation of growth and development, genitourinary anomalies, and ear anomalies

CHD: Congenital heart disease

CHD7: Chromodomain helicase DNA binding protein 7

CMV: Cytomegalovirus

CSA: Cyclosporine A

DGS: DiGeorge syndrome

FOXI3: Forkhead box I3

FOXP1: Forkhead box N1

GOSH: Great Ormond Street Hospital

HCT: Hematopoietic cell transplantation

HLA: Human leukocyte antigen

HSC: Hematopoietic stem cell

HSPC: Hematopoietic stem and progenitor cells

IEI: Inborn error of immunity

IgRT: Immunoglobulin replacement therapy

NBS: Newborn screening

NK: Natural killer

OFCS2: Otofaciocervical syndrome type 2

PAX1/9: Paired box 1/9

SCID: Severe combined immunodeficiency

TBX1/2: T-box transcription factor 1/2

TCR: T-cell antigen receptor

TEC: Thymic epithelial cells

TREC: T-cell–receptor excision circle

WES: Whole exome sequencing

WGS: Whole genome sequencing

immature T lymphocytes (or thymocytes) that have successfully expressed a T-cell antigen receptor (TCR) with sufficient affinity for a peptide–major histocompatibility complex to receive signals that allow their further survival. Positively selected thymocytes are next subjected to another quality control of the cells' antigen specificity, a process known as central tolerance induction, which encompasses negative selection and regulatory T-lymphocyte development. Negative selection ensures that maturing T lymphocytes with a high affinity for self–peptide–major histocompatibility complex undergo programmed cell death (apoptosis) and are thus deleted from the repertoire because these cells are particularly prone to initiate autoimmunity. Alternatively, thymocytes with an intermediately high affinity for self-antigens may escape deletion and instead be diverted to a regulatory T-cell fate. To efficiently achieve this, TEC collectively express in a promiscuous fashion an almost comprehensive set of tissue-restricted antigens,^{24,31–34} some of which are under the control of transcriptional regulators, namely autoimmune regulator (AIRE) and family zinc finger 2 (FEZF2).^{24,27,28,31,33} While cortical TEC are singularly responsible for positive selection, negative selection can be mediated by TEC lineages as well as bone marrow–derived antigen presenting cells including dendritic cells and B lymphocytes.^{27,28,30,35–43} However, within the thymus, AIRE and FEZF2 are predominantly expressed in a subset of medullary TEC, rendering these cells especially competent to shape the TCR repertoire. Taken together, lymph–stromal interactions within the thymus instruct the commitment and maturation of T

lymphocytes and shape a diverse repertoire of TCR specificities that are tolerant to self yet reactive to foreign antigens.

Congenital thymic stromal cell disorders are inborn pathologies caused by abnormalities in thymic organogenesis and stromal cell development and/or function, resulting in deficient and/or dysregulated T-lymphocyte immunity.^{44–47} The severity of peripheral T lymphocytopenia correlates with the reduced size of the thymic stromal scaffold.⁴⁶ Thymic hypoplasia, depending on its severity, may therefore result in either mild T lymphocytopenia with little or no clinical consequence, or alternatively, it may be characterized by a clinically significant lack of peripheral T lymphocytes, reduced T-lymphocyte proliferative responses, an oligoclonal TCR repertoire, and secondary impaired humoral immunity.^{48–54} Complete thymic aplasia is rare and results in a T-lymphocyte–negative, B-lymphocyte–positive, natural killer cell–positive (T-B+NK+) immunophenotype with severe to complete absence of naive T lymphocytes, T-lymphocyte proliferative responses, and T-cell–receptor excision circles (TRECs), which are small circles of DNA created as a by-product of TCR gene rearrangement.^{44,50,52–58} A T-B+NK+ immunophenotype may of course also be seen in the context of severe combined immunodeficiency (SCID) due to various hematopoietic cell–intrinsic genetic defects.^{45,59}

Aberrant patterning of the third pharyngeal pouch during early embryogenesis may result in congenital athymia associated with defective development of other anatomic structures derived from the adjacent pharyngeal apparatus.^{44,60} Defects of thymic organogenesis may therefore variably be associated with additional anomalies affecting multiple organs, including craniofacial structures, the heart, great vessels, and the parathyroids.⁶¹ Such field defects occur in the context of DiGeorge syndrome (DGS), a clinical diagnosis defined by the triad of thymic hypoplasia/aplasia, hypoparathyroidism, and congenital heart disease (CHD), although the immunodeficiency and other features of DGS typically display substantial variability in their clinical penetrance.^{50,53,54,62} Multiple genetic and environmental etiologies can cause a DGS phenotype (Table I), with heterozygous chromosomal deletions at 22q11.2 (22q11.2 deletion syndrome, or 22q11.2DS) constituting the most frequent cause, with an estimated incidence of 1:4000 live births.⁶³ Athymia is uncommon, occurring in ≤1% of individuals with 22q11.2DS, in whom the condition has previously been referred to as complete DGS because of the severity of the ensuing immunodeficiency.^{50,56,64} The deletion is typically between 1.5 and 3.0 Mb in size, resulting in the loss of approximately 30 to 100 genes. The size of the deletion does not correlate with the clinical phenotype.⁶⁵ Among the deleted genes is *TBX1*, a T-box transcription factor that regulates almost 2000 genes⁶⁶ and plays a major role in the pharyngeal patterning defects seen in 22q11.2DS, including immunodeficiency.^{10,67,68} The terms “DGS” and “22q11.2DS” are often used interchangeably. However, other genetic and environmental causes of the DGS triad have been identified.⁴⁴ Thus, where known, reference should be made to the exact etiology. The second most frequent genetic cause of DGS comprises autosomal-dominant mutations in the *CHD7* (chromodomain helicase DNA binding protein 7) gene, which underlies CHARGE syndrome, or a syndrome with coloboma, heart defects, atresia of nasal choanae, retardation of growth and development, genitourinary anomalies, and ear anomalies.^{69,70} While some clinicians consider CHARGE syndrome a separate disease entity, in this guideline, we include CHARGE syndrome as one of the genetic

etiologies of the clinical DGS triad. The incidence of CHARGE syndrome is approximately 1:10,000–17,000 live births;⁷¹ however, the frequency of congenital athymia among these is not known. Other rare genetic causes of DGS include mutations in *TBX1*^{72–74} or *TBX2*;⁷⁵ 22q11.2 duplications;⁷⁶ haploinsufficiency of *FOXP3* (forkhead box I3) due to microdeletions at 2p11.2⁷⁷ or heterozygous loss-of-function single gene mutations;⁷⁸ and partial deletion of the short arm of chromosome 10.^{79–81} Although thymic hypoplasia and other features of DGS have been reported in 22q11.2 duplication, lymphopenia is not commonly present.^{82–85} In cases where DGS remains genetically undefined, it is important to consider whether *in utero* exposure in the first trimester to poorly controlled maternal diabetes,^{86,87} alcohol,⁸⁸ and overexposure^{89,90} or underexposure⁹¹ to retinoic acid may have occurred. Retinoic acid metabolism is impaired during pregestational diabetes⁹² and has been linked to *TBX1* signaling as well as to other transcription factors with a role in thymus organogenesis.^{90,93,94}

Congenital athymia has also been identified in other syndromic inborn errors of immunity (IEI) (Table 1),^{44,95} including otofaciocervical syndrome type 2 (OFCS2) caused by homozygous *PAX1* deficiency and nude “SCID” resulting from homozygous *FOXP1* deficiency.^{44,58,70,96–100} *PAX1* is an evolutionary conserved transcription factor, expressed in the pharyngeal pouches during embryogenesis, where it is important for patterning and the development of the thymic anlagen. Later, *PAX1* expression is maintained in a fraction of cortical TEC in a *FOXP1*-dependent manner.^{9,101} OFCS2 patients have ear anomalies, facial dysmorphism, and skeletal anomalies, and, depending on the degree of residual *PAX* transcriptional activity conferred by their mutation or mutations, they may present with congenital athymia or with a milder phenotype consisting of combined immunodeficiency with overlapping features with DGS.^{99,100,102–105} As the master transcriptional regulator,¹⁴ *FOXP1* is essential for TEC differentiation and maintenance, as well as the functional ability to support thymopoiesis by controlling the expression of key genes.¹³ The nude SCID phenotype, comprising congenital athymia, alopecia totalis, and nail dystrophy, was first reported in patients with homozygous complete loss-of-function mutations in *FOXP1*. These include early truncating mutations that likely result in nonsense mediated mRNA decay, and missense mutations that alter critical residues within the forkhead domain, abrogating its ability to bind DNA.^{44,96,97,106–108} Several studies have recently characterized monoallelic and compound heterozygous mutations, finding a range of functional consequences from complete to partial loss of function, including some mutants that exert a dominant negative effect on the wild-type allele.^{109–112} These studies have further revealed the dose dependency for *FOXP1* in TEC, such that the severity of the immunodeficiency appears to reflect the degree of residual *FOXP1*-induced transcriptional activity.¹⁰⁶

Left untreated, congenital athymia is incompatible with long-term survival. It is therefore crucial to identify and treat affected patients as early as possible. Congenital athymia can be treated by transplanting lymphodepleted thymus tissue, donated by infants undergoing cardiac surgery, into the quadriceps muscles of athymic recipients. This provides a functional stromal environment to generate recipient-derived T lymphocytes. The current tissue culture and implantation process has not changed much since its early medical applications.^{113,114} To date, thymus transplantation outcomes have been reported for more than 140 athymic patients treated at 2 centers, Duke University Medical Center

in the United States and Great Ormond Street Hospital (GOSH) in the United Kingdom.^{115–117} The overall survival of patients with athymia treated with thymus transplantation is approximately 75%, with mortality mainly due to preexisting infections or infections acquired before immune reconstitution is established.^{55,58,115,116,118} Although absolute T-lymphocyte counts usually remain below those seen in the normal population, they are sufficient to clear and prevent infections, enabling patients to participate in normal activities and thus improving their quality of life.^{115,116,118,119} Autoimmune manifestations, particularly affecting the thyroid gland and blood cells, are relatively common after thymus transplantation,^{115,116} suggesting an incomplete establishment of self-tolerance by the transplanted thymic stroma. Overall, thymus transplantation is a lifesaving procedure for congenital athymia, with superior outcomes compared to hematopoietic cell transplantation (HCT).¹²⁰ Poorer survival and severe graft-versus-host disease have been reported after HCT, particularly if no matched sibling donor is available.^{120,121} Therefore, it is widely agreed that thymus transplantation, as the most appropriate treatment for congenital athymia, should be the standard of care and that every effort should be made to ensure prompt access to treatment, where available.

Here we provide clinical guidelines to help identify, investigate, and manage patients with congenital athymia who should be considered for thymus transplantation. Our expert panel recommendations, summarized in Fig 1, focus on when to suspect a thymic stromal cell defect, how to diagnose and stratify patients according to risk in order to identify those with congenital athymia who might benefit from thymus transplantation, and their initial, supportive clinical management. While we also discuss the role of novel diagnostic approaches, the *status quo* of allogeneic thymus transplantation, its follow-up, and future directions, the emphasis of this guideline is to promote and facilitate early recognition of congenital athymia and prompt referral for specialist treatment to improve clinical outcomes. These guidelines are focused on the management of congenital athymia, but many of the recommendations, particularly those related to diagnosis and investigation, are applicable to congenital thymic hypoplasia. Specific guidance on the management of patients with thymic hypoplasia is available elsewhere.⁵⁶

WHEN TO SUSPECT CONGENITAL ATHYMIA

Infants with congenital athymia can come to medical attention via several different routes (Fig 2). Increasingly, this occurs in the context of newborn screening (NBS) programs for SCID and T lymphocytopenia.^{117,122} NBS for SCID started in some parts of the United States in 2008, and since 2018, all 50 states have universal screening programs in place.¹²³ Universal and pilot NBS programs for SCID are progressively being rolled out in a growing number of countries.^{122,124–128} These programs rely on the detection of TRECs in dried blood spot samples routinely taken from newborns shortly after birth. TRECs are stable, circular, episomal DNA excised from genomic sequences during TCR gene rearrangement and are a relative measure of thymic output because they mark newly generated T lymphocytes. Low or undetectable TRECs are a positive finding in SCID NBS, and these TREC-based screening programs also identify infants with thymic aplasia and hypoplasia.^{117,122,129,130}

TABLE I. Disorders known to cause congenital thymic aplasia or hypoplasia and associated clinical features

DGS etiology	Condition	Inheritance	Syndromic features
Genetic	22q11.2DS ⁴⁴	<i>De novo</i> (90-95%); AD (5-10%)	● Variable features of DGS
	CHARGE syndrome (CHD7 haploinsufficiency in majority) ⁵³	<i>De novo</i> (majority)	● Variable features of DGS ● Coloboma ● Choanal atresia ● Growth retardation ● Genitourinary abnormalities ● Ear anomalies ● Cranial nerve dysfunction
	TBX1 deficiency ⁶⁷	AD	● Variable features of DGS
	TBX2 deficiency ⁷⁵	AD	● Variable features of DGS
	22q11.2 duplication ⁷⁶	<i>AD/de novo</i>	● Variable features of DGS
	FOXI3 haploinsufficiency, including 2p11.2 microdeletions ⁷⁷ and heterozygous loss-of-function FOXI3 variants ⁷⁸	AD (majority)	● Variable features of DGS
	Partial monosomy 10p ⁷⁹⁻⁸¹	<i>De novo</i>	● Variable features of DGS ● Craniofacial malformation
	Diabetic embryopathy ^{86,87}	NA	● Variable features of DGS ● Renal agenesis ● Vertebral anomalies
	<i>In utero</i> overexposure to alcohol ⁸⁷	NA	● Variable features of DGS ● Variable features of foetal alcohol syndrome
	<i>In utero</i> overexposure to retinoic acid ^{89,90} <i>In utero</i> under exposure to retinoic acid ⁹¹ Other genetic syndromes associated with defective thymic development	NA NA	● Variable features of DGS ● Variable features of DGS
Environmental	Nude SCID (FOXN1 deficiency) ⁹⁶	AR	● Congenital alopecia totalis ● Nail dystrophy
	Hypomorphic FOXN1 deficiency, including heterozygous variants ¹¹⁰ and compound heterozygote mutations ¹¹¹	AD, AR	● Nail dystrophy ● Alopecia may be absent, hair thinning ● Eczema
	OFCS2 (PAX1 deficiency) ^{99,100}	AR	● Ear anomalies, preauricular pits, hearing impairment ● Branchial cysts/fistulas ● Facial dysmorphism ● Skeletal anomalies ● Intellectual disability
			● Can have overlapping features with DGS including hypoparathyroidism and congenital heart defects

Variable features of DGS indicates that DGS features include a wide variety of possible anomalies⁴⁴ but most commonly the following: congenital cardiac defects; parathyroid hypoplasia; facial dysmorphism; palatal, pharyngolaryngeal, and tracheobronchial defects; and developmental delay.

AD, Autosomal dominant; AR, autosomal recessive; NA, not applicable.

If not captured by NBS programs, infants with congenital athymia may present clinically with complications consequent to their severe T-lymphocyte deficiency,¹¹⁵⁻¹¹⁷ typically within the first few months of life, with failure to thrive as well as unusually persistent, severe, or opportunistic infections such as *Pneumocystis jirovecii* pneumonia, CMV pneumonitis, or disseminated bacillus Calmette-Guérin infection. Persistent respiratory tract infections—for example, those caused by respiratory syncytial virus, parainfluenza virus, or adenovirus—or persistent candidiasis should also prompt further immunologic investigation. Gastrointestinal infection and chronic diarrhea are common features; causes include viruses such as adenovirus, norovirus, or rotavirus (wild type or vaccine strain). Immune dysregulation may occur, which most commonly manifests with Omenn syndrome-like features, including erythroderma, diarrhea, hepatosplenomegaly, lymphadenopathy, elevated IgE, and eosinophilia.^{57,99,106,131} More rarely, patients present with symptoms of autoimmunity, particularly hematologic cytopenias.

In addition, clinical features related to an underlying syndrome are often present (Table I). Because athymia is most frequently encountered in the context of DGS, field defects affecting structures derived from the adjacent pharyngeal apparatus, including the parathyroids, aortic arch, cardiac outflow tract, thyroid, maxilla, mandible, and external/middle ear, are variably present (Table I). Parathyroid hypoplasia can manifest with hypocalcemia and neonatal seizures, and it may be the presenting feature of DGS. Cardiac conotruncal outflow tract defects lead to particular CHD, including tetralogy of Fallot, truncus arteriosus, interrupted aortic arch, double outlet right or left ventricle, transposition of the great arteries, and aortopulmonary septal defects.⁴⁴ In addition to the classical DGS triad, other clinical features may be present, including facial dysmorphism, palatal and/or pharyngolaryngeal defects, tracheobronchomalacia, gastrointestinal problems (ie, feeding difficulties, constipation, and gastroesophageal reflux disease), renal anomalies, skeletal problems (ie, scoliosis and talipes equinovarus), and developmental




FIG 1. Flowchart summarizing clinical guidelines for recognition, investigation, and management of patients with congenital athymia. *aCGH*, Array comparative genomic hybridization; *EBV*, Epstein-Barr virus; *FISH*, fluorescence *in situ* hybridization; *HHV6*, human herpesvirus 6; *NPA*, nasopharyngeal aspirate; *RTE*, recent thymic emigrant.

delay.^{49,132} In addition to DGS with underlying 22q11.2DS, other genetic and environmental etiologies of DGS have distinct clinical features that may be evident (Table I).⁴⁴ Patients with CHARGE syndrome may have coloboma, atresia choanae, retardation of growth and/or development, genitourinary and/or ear anomalies, and cranial nerve dysfunction.^{133,134} Infants born to diabetic mothers may have features of caudal dysplasia sequence, in particular renal agenesis and vertebral anomalies.^{86,135} Clinical

features of homozygous *FOXP1* deficiency include congenital alopecia totalis and nail dystrophy.^{96,97} *PAX1* deficiency underlies OFCS2 with preauricular pits and hearing impairment, facial dysmorphism, skeletal anomalies, and intellectual disability.^{99,100,102,103} While the aforementioned clinical features should alert clinicians to the possibility of immunodeficiency resulting from impaired thymic stromal cell development, the severity of the immunologic phenotype in an individual patient

WHEN TO SUSPECT CONGENITAL ATHYMIA

Very low/absent TRECs on newborn screening for SCID and T lymphocytopenia

Clinical features consequent to severe T lymphocyte deficiency (including opportunistic/severe/persistent infections and failure to thrive)

Omenn-like features

Autoimmunity (particularly cytopenias)

Lymphopenia in the context of other relevant syndromic features such as cardiac defects, hypoparathyroidism, facial dysmorphism, palatal defects, congenital *alopecia totalis* etc (see Table 1)

Family history of congenital athymia

Genetically undefined T-B+NK+ immunophenotype

Failure of naïve T lymphocyte reconstitution in patients treated with allogeneic HCT for genetically undefined T-B+NK+ immunophenotype

FIG 2. When to suspect congenital athymia.

cannot be predicted by the severity of the other features. Therefore, all children with suggestive clinical features should undergo immunologic and genetic investigations to exclude immunodeficiency—and, crucially, to identify those with life-threatening athymia who require corrective treatment.

Timely recognition of congenital athymia and early referral for treatment are the result of several factors. Awareness among clinicians has increased to recognize the syndromic features associated with athymia and is matched by expanding access to TREC-based NBS and next-generation sequencing to facilitate diagnosis of cases with a genetic etiology. Nonetheless, incomplete clinical penetrance, together with clinical and genetic heterogeneity, can still make timely recognition of athymia challenging. It may only be recognized after failure of naïve T-lymphocyte reconstitution despite adequate donor engraftment and reconstitution of other blood lineages in patients treated empirically with allogeneic HCT for genetically undefined suspected SCID, highlighting the importance of extensively investigating patients with a molecularly undefined T-B+NK+ immunophenotype to distinguish between hematopoietic cell-intrinsic SCID and congenital athymia before considering allogeneic HCT as a treatment strategy.^{99,136}

HOW TO INVESTIGATE INFANTS WITH SUSPECTED CONGENITAL ATHYMIA

Children with possible congenital athymia should undergo a series of investigations aimed at rapidly securing the diagnosis and etiology, as well as characterizing the severity of their underlying immune defect (Fig 3). The latter will guide immunologic management and, importantly, will identify those likely to derive benefit from allogeneic thymus transplantation.

Immunology

A finding of lymphocytopenia based on a full blood count with differential is often the initial finding in congenital athymia. Lymphocyte subsets enumerating T, B, and NK cells typically demonstrate a low CD3⁺ T-lymphocyte count with normal B- and NK-cell counts, corresponding to a T-B+NK+ immunophenotype. Athymic patients with Omenn syndrome-like clinical

features (previously also referred to as athymia with atypical features⁵⁶) have higher, or even normal, lymphocyte counts due to oligoclonal expansion of dysregulated T lymphocytes of memory phenotype (CD45RA⁻CD27⁺).^{57,131} More definitive immunologic assessment involves quantitative evaluation of thymic output. Thymic output can be assessed by enumeration of TRECs using real-time quantitative PCR on isolated peripheral blood mononuclear cells and provides a practical and accepted indicator of thymic output, with low or absent TRECs indicating lack of thymic naïve T-lymphocyte production.¹³⁷ This is distinct from the NBS assay, which measures TRECs in a dried blood spot sample.^{49,138} Although cutoff values vary in different NBS programs, a TREC value of <20 copies/μL will successfully identify SCID cases regardless of the underlying cause.^{139,140} However, because this assay is not routinely available in many clinical laboratories, naïve T lymphocytes (CD45RA⁺CD27⁺) or recent thymic emigrants (CD45RA⁺CD31⁺) can alternatively be measured by flow cytometry. Their frequencies strongly correlate with TREC levels, suggesting either can be used as a marker for thymic output.⁴³ In congenital athymia, naïve T lymphocytes are profoundly reduced (<50 cells/μL or <5% of the total T lymphocytes).^{56,141} Flow cytometric immunophenotyping is a fast and widely available technique but can be associated with significant variability among different centers in sample processing, immunostaining, instrument setup, and data analysis. Standardized protocols have been developed to generate reproducible and reliable results and have been shown to be highly useful in the diagnosis of conditions including SCID, alongside immunologic functional and genetic testing.¹⁴²

If the above investigations are consistent with athymia, qualitative T-lymphocyte tests are of limited value and are not routinely necessary. In patients who have measurable peripheral T lymphocytes, qualitative assessment can be performed by measuring diversity of the TCR repertoire and by assessing T-lymphocyte function. Because the diversity of the TCR repertoire is almost completely reflective of the naïve T-lymphocyte compartment, patients with reduced thymic output have a restricted TCR repertoire.^{138,143} In cases with an Omenn syndrome-like phenotype, oligoclonality is seen.^{116,131} Assessment of TCR diversity can be performed using flow cytometry to quantify TCR Vβ usage, or more reliably using spectratyping,

INVESTIGATION OF PATIENTS WITH SUSPECTED CONGENITAL ATHYMIA

Test	Congenital athymia
Basic immunological assessment	
Full blood count	Lymphocytopenia (variable)
Lymphocyte subsets	Low CD3 ⁺ T lymphocytes, normal B and NK cells. Normal /high CD3 ⁺ T lymphocytes if Omenn-like features or maternal engraftment present
TRECs	Very low / absent
Naïve T lymphocytes / RTEs	Profoundly reduced (<50 cells/μL or <5% of the total T lymphocytes)
Immunoglobulins	IgM and IgA very low, IgG may be normal depending on patient age and presence of maternal IgG IgE may be elevated if Omenn-like features present
Qualitative T lymphocytes tests (if measurable T lymphocytes present)	
TCR repertoire diversity	Restricted TCR repertoire, oligoclonality if Omenn-like features
T lymphocyte proliferation	Absent/very low responses to mitogens Patients with Omenn-like features can have partial or normal proliferative mitogen responses but impaired responses to specific antigens
Genetic	
Cytogenetic studies for chromosomal abnormalities (aCGH or FISH)	Del22q11.2, Dup22q11.2, Del10p13-14, Del2p11.2
WGS or WES (typically trio or proband only)	
IEI gene panel testing	Including <i>CHD7</i> , <i>TBX1</i> , <i>TBX2</i> , <i>FOXI3</i> , <i>FOXN1</i> , <i>PAX1</i>
Single gene sequencing	For highly suggestive phenotypes or positive family history
Research Diagnostic assays	
Reporter assays to assess pathogenicity of novel genetic variants in transcription factors	
<i>Ex vivo</i> T lymphocyte differentiation assay to differentiate thymic stromal cell defects from primary hematopoietic defects	

FIG 3. How to investigate infants with suspected congenital athymia. *aCGH*, Array comparative genomic hybridization; *Del*, deletion; *Dup*, duplication; *FISH*, fluorescence *in situ* hybridization; *RTE*, recent thymic emigrant.

a molecular technique that measures the length distribution of complementarity determining region 3 (aka CDR3).¹⁴⁴ T-lymphocyte function can be evaluated by measuring *in vitro* proliferative capacity to mitogenic stimulants such as phytohemagglutinin or anti-CD3/28. While patients with congenital athymia normally have absent or very low responses to mitogens, athymic patients with oligoclonal T-lymphocyte expansion may demonstrate normal proliferative mitogen responses but lack response to specific antigens. However, these assays are usually only performed in specialized laboratories and can be unreliable in lymphopenic patients, reflecting the reduced number of proliferating T lymphocytes rather than intrinsic T-lymphocyte dysfunction.⁵⁶ IgM and IgA are usually quite low, whereas, depending on the age of the patient, IgG may be normal as a result of maternal transplacental

IgG transfer. Further investigation of humoral immunity by assessing specific antibody production is time-consuming and unnecessary in the diagnosis of athymia but is recommended in milder cases.

Genetics

Attempts should be made to genetically define suspected congenital thymic aplasia/hypoplasia; selection of the most appropriate genetic investigations may be guided by the patient's clinical phenotype (Table I). Chromosomal abnormalities underlie the majority of DGS cases but are routinely missed by whole exome sequencing (WES). Cytogenetic studies should therefore be performed, particularly in patients with a DGS phenotype. In this context, array comparative genomic hybridization is

preferable to fluorescence *in situ* hybridization or karyotyping because it allows for high-resolution genome-wide screening for genetic copy number variation and has been shown to increase diagnostic yield in the context of suspected DGS.^{145,146} Where a monogenic disorder is suspected, the choice of molecular genetic tests is likely to be influenced by what is locally available. Broad next-generation sequencing approaches are preferred, including gene panel testing, where it is imperative to ensure that candidate genes (Table 1) are included in the panel being used, and WES or whole genome sequencing (WGS). With the latter 2 modalities, a patient's genetic variants are usually first filtered against a list of genes known to cause ICI before more agnostic approaches are applied that have the potential to identify novel genetic causes.^{147,148} For WES and WGS, although only sequencing of the proband can be performed, the inclusion of unaffected (typically parents) and, where available, affected, family members strengthens diagnostic power. WGS offers several advantages over WES because of its ability to detect noncoding and structural variants, as well as superior coverage of coding regions.^{149,150} Single gene sequencing might suffice where the clinical phenotype is highly suggestive of a specific genetic diagnosis or if there is a positive family history, such as in nude SCID (*FOXN1*) or OFCS2 (*PAX1*), particularly if it is the most accessible and most quickly available diagnostic modality. When a genetic cause is identified, the family should be referred for genetic counseling and, if indicated, further genetic and clinical assessments. With regard specifically to 22q11.2DS, although most cases occur *de novo*, 5% to 10% of patients inherit the microdeletion from a parent.^{50,151} Therefore, the risk of recurrence should be considered, with genetic testing extended to relatives and future progeny as appropriate.

Increasing access to next-generation sequencing and NBS for SCID has the potential to uncover genetic variation of uncertain clinical significance as well as to broaden the spectrum of disease associated with known ICI genes. The latter is exemplified by the recent description of hypomorphic *FOXN1* variants and consequent attenuated clinical phenotypes.^{106,109-112} In contrast to nude "SCID"—causing null mutations, these hypomorphic variants may lead to delayed presentation, atypical, and/or milder clinical and immunologic phenotypes, which may improve with age and are unlikely to necessitate thymus transplantation.^{106,110,111} Despite increasing access to comprehensive genetic testing, approximately 10% of all SCID patients remain genetically undefined.^{152,153} Similarly, 13% of athymic patients who were recently referred to GOSH for thymus transplantation did not have a genetic diagnosis.^{117,130} Patients with a T-B+NK+ immunophenotype in whom the underlying cause could either be a primary hematopoietic defect or congenital athymia require additional investigations, including an evolving array of research assays to assist their clinical management.

Other diagnostic investigations, including research assays

In genetically undefined patients, it is not possible to distinguish between hematopoietic cell intrinsic SCID and congenital athymia solely on the bases of immunophenotyping and proliferation assays. Imaging for thymus tissue does not differentiate between the 2 conditions, as an absent or greatly reduced thymic shadow can be seen in both and may be misleading when thymus tissue is ectopically positioned.¹⁵⁴

Research assays are increasingly used to assist in the diagnosis of thymic stromal cell defects and in therapeutic decision-making. For example, to functionally assess the pathogenicity of novel genetic variants in transcription factors, reporter assays can be used to test the ability of the mutant transcription factor to bind to its target promoter DNA sequence and activate expression of a reporter protein. In the context of thymic stromal cell defects, such assays have been utilized to assess the pathogenicity of novel genetic variants in *FOXN1*, *PAX1*, *TBX1*, and *FOXI3*.^{78,99,100,106,110,112,155} For genetically undefined T lymphocytopenia, the use of *ex vivo* T-lymphocyte differentiation research assays has been proposed to distinguish patients with primary hematopoietic defects from those with thymic stromal cell defects and thus direct the most appropriate form of corrective treatment—HCT versus thymus transplantation, respectively.^{55,156,157} Patient CD34⁺ hematopoietic stem and progenitor cells (HSPCs) are cocultured with stromal cell lines engineered to express the human Notch ligands DLL-1 or DLL-4 (delta-like ligand 1/4) in the presence of growth factors to promote T-lymphocyte lineage commitment and differentiation, either in 2-dimensional monolayer cultures¹⁵⁸ or in 3-dimensional artificial thymic organoids or reaggregate thymus organ cultures.¹⁵⁹ Feeder cell-free systems are also available for *ex vivo* T lymphopoiesis.¹⁶⁰ On the one hand, in principle, successful production *ex vivo* of CD4⁺ and CD8⁺ double-positive and TCR⁺ CD3⁺ stages from patient-derived HSPCs after 6 to 8 weeks of coculture argues against a hematopoietic defect and is instead suggestive of a possible thymic stromal cell defect. On the other hand, HSPCs from patients with primary hematopoietic defects are expected to be intrinsically impaired in their ability to differentiate to these stages.^{156,157,160} There are, however, exceptions to this, and HSPCs from patients with a number of hematopoietic stem cell (HSC)-intrinsic defects, such as hypomorphic variants in *RAG* genes or *IL2RG*, have been shown to differentiate *ex vivo* beyond their expected developmental block into double-positive and CD3⁺TCR αβ⁺ stages; and those with adenosine deaminase deficiency defy expectations because *in vitro* T-lymphocyte development is normal.^{156,157} The sensitivity and specificity of these assays are not well established, and there is a lack of standardization across different research laboratories. Despite these limitations, *ex vivo* T-lymphocyte differentiation assays are helpful when facing the therapeutic dilemma of HCT versus thymus transplantation in patients with molecularly undefined, selective T-lymphocyte deficiency.¹⁶¹ To avoid treatment delays, research assays to assist clinical decision-making are best arranged in coordination with a thymus transplantation team with experience in their clinical interpretation.

HOW TO IDENTIFY PATIENTS WHO NEED CORRECTIVE TREATMENT

Patients with congenital thymic stromal cell defects can have variable degrees of thymic hypoplasia and consequently thymic output. This means that the overall immunologic consequences can range from normal T-lymphocyte immunity to a T-B+NK+ immunophenotype. It is therefore important to stratify patients according to the extent of their immunodeficiency to distinguish those with complete athymia requiring thymus transplantation from those with thymic hypoplasia and residual thymic function, requiring just supportive care. This stratification is mainly based on immunophenotyping with quantification of thymic output and on clinical presentation (Fig 4).

HOW TO IDENTIFY PATIENTS THAT NEED CORRECTIVE TREATMENT	
Congenital athymia	→ Corrective treatment
Low CD3 ⁺ T lymphocyte counts (<50 cells/μl) Low naïve T lymphocytes (<50 cells/μl; <5% of total CD3 ⁺) Negligible TRECs	
Congenital athymia & autologous immune dysregulation (Omenn-like features)	→ Corrective treatment
CD3 ⁺ T lymphocyte counts can be normal or increased (predominant memory/activated phenotype) Low naïve T lymphocytes (<5% of total CD3 ⁺) Negligible TRECs Oligoclonal TCR repertoire Usually impaired T lymphocyte proliferation	
Thymic hypoplasia	→ Supportive care
Milder T lymphopenia (<1500 cells/μl during the first 3 years of life) Naïve T lymphocytes and TRECs reduced but not absent	
Borderline cases	→ Careful follow-up & consideration
Do not meet the immunological criteria for athymia, but display clinically relevant immunodeficiency / immune dysregulation	

FIG 4. How to identify patients who need corrective treatment.

Immunology in patients with complete athymia

Patients with complete athymia typically have low total lymphocyte counts, although they can be normal as a result of increased B- and/or NK-cell numbers or as a result of oligoclonal T-lymphocyte expansions. More rarely, elevated T-lymphocyte counts can also be found as a result of maternal T-lymphocyte engraftment. Here, the engrafted T lymphocytes are usually predominantly CD4⁺ or CD8⁺ and have an activated or memory phenotype. All athymic patients, regardless of T-lymphocyte counts, have negligible TRECs and less than 5% of T lymphocytes displaying a naïve phenotype, reflecting their absent thymic output.^{56,141} Their immunodeficiency is life-limiting, and they need to be referred for corrective treatment without delay.

Immunology in patients with thymic hypoplasia

After exclusion of congenital athymia, certain patients are diagnosed with significant thymic hypoplasia, which is associated with milder T lymphocytopenia, defined as a CD3⁺ T-lymphocyte count of <1500 cells/μL during the first 3 years of life.⁴⁹ Typical findings include reduced CD3⁺, CD4⁺, and CD8⁺ T lymphocytes (respectively, <400 and <200 cells/μL), naïve T lymphocytes, and TRECs reflecting reduced but not absent thymic output.^{49,56} T-lymphocyte numbers improve with age, reaching levels similar to healthy adult controls as a result of homeostatic proliferation with accumulation of memory cells and resultant skewing of the TCR repertoire.^{49,51} Naïve T lymphocytes and TRECs, however, remain reduced at all ages compared to age-matched controls, indicating persistently reduced thymic function.⁵¹ T-lymphocyte function in patients with thymic hypoplasia is largely intact with normal mitogen responses. In patients with marked T lymphocytopenia, responses to specific antigens may be reduced, although this is likely solely due to low numbers rather than an intrinsic functional defect.⁵¹ Regulatory T cells

are reduced in number and frequency with defective suppressive capacity, which may contribute to the increased incidence of autoimmunity.^{64,162} Abnormalities in humoral immunity may include low immunoglobulins, most commonly IgM or IgA deficiency, poor specific antibody responses, and occasionally hypogammaglobulinemia requiring immunoglobulin replacement therapy (IgRT).^{163,164} The panel does not recommend administration of live vaccines in patients with CD4⁺ T lymphocytes <400/μL and CD8⁺ T lymphocytes <200/μL, or in patients with nonprotective IgG titers after tetanus immunization.⁵⁶ These patients may require supportive treatment, but they are not eligible for corrective treatment with thymus transplantation.

Borderline cases

Some cases do not meet the immunologic criteria for athymia but nevertheless display clinically relevant immunodeficiency with a history of significant infections and/or immune dysregulation. These patients require careful clinical and immunologic follow-up because their phenotype may evolve over time. For patients falling into this category case-by-case consideration regarding the most appropriate therapeutic strategy is best achieved in conjunction with a thymus transplantation center. This is particularly relevant when considering treatment options for patients with novel or ultrarare defects.^{105,117,119} Moreover, if patients develop Omenn syndrome-like features, this should be considered as a marker of athymia, indicating the need for corrective treatment.

HOW TO MANAGE PATIENTS WITH CONGENITAL ATHYMIA

These guidelines are focused on the management of infants with congenital athymia (Fig 5). Guidance for the management of patients with thymic hypoplasia is available elsewhere.^{56,165-167}

HOW TO MANAGE PATIENTS WITH CONGENITAL ATHYMIA

Corrective treatment

Refer to specialist centre for allogeneic thymus transplantation

T cell replete allogeneic HCT from a fully matched donor in certain circumstances (see main text)

Supportive care as per local SCID protocol

Isolation

Anti-microbial prophylaxis

Immunoglobulin replacement therapy

No live vaccines

Irradiated CMV negative blood products

Withhold breastfeeding from CMV seropositive mothers

Immunosuppression if Omenn-like phenotype

Family support

Monitoring

Microbial surveillance including viral PCRs on blood, stool and NPA as per local SCID protocol Investigations and management of other co-morbidities as appropriate

Monitoring for complications and immune reconstitution after corrective treatment

FIG 5. How to manage patients with congenital athymia. *NPA*, Nasopharyngeal aspirate.

Ideally, thymus transplantation should be considered as the first-line treatment in patients with congenital athymia, although geographical and financial constraints might limit access. Akin to outcomes after HCT for hematopoietic cell–intrinsic SCID,¹⁶⁸ outcomes after thymus transplantation are better when patients are treated early,¹¹⁷ before they acquire infections.^{115–117} Treatment at a younger age additionally seems to be associated with better initial immune reconstitution as a result of higher thymic output earlier after thymus transplantation.¹¹⁷ Therefore, there should be no delay in referring patients to a thymus transplantation center while instituting comprehensive supportive measures locally.

Supportive measures and monitoring for complications

Patients are best managed by a local specialist pediatric immunology unit in conjunction with a thymus transplantation center. While awaiting corrective treatment, all efforts should be made to ensure that the patient remains free from infection and in the best possible overall clinical state because these factors influence outcome. Preventative care should be promptly instigated once the diagnosis of athymia is suspected, and SCID management protocols can be co-opted to guide comprehensive supportive care as well as to monitor for and treat complications.^{56,169} Although there is variability among centers in such protocols, all athymic patients should be subject to reverse isolation measures, avoidance of ill contacts, and restricted non-staff caregivers.¹⁶⁹ Antimicrobial prophylaxis therapy should be initiated in line with local SCID protocols, including *P jirovecii* pneumonia prophylaxis (ie, trimethoprim–sulfamethoxazole), an azole antifungal (preferably fluconazole or itraconazole; itraconazole requires regular therapeutic drug monitoring), IgRT, and seasonal anti–respiratory syncytial virus prophylaxis using passive immunization with monoclonal antibody. Antimycobacterial prophylaxis with azithromycin should be considered in at-

risk patients, particularly if there is likely to be a lengthy wait for thymus transplantation.¹⁷⁰ Live immunizations, such as bacillus Calmette–Guérin, rotavirus, and oral polio vaccines, are contraindicated, and patients should only receive irradiated, CMV-negative blood products. Breast-feeding should be withheld until maternal CMV status is known and discontinued if the mother is found to be CMV seropositive. Monitoring for viral infections is recommended, including regular PCRs on blood (for CMV, Epstein–Barr virus, adenovirus, and human herpesvirus 6), stool, and nasopharyngeal aspirates as per local SCID protocols. Additional microbiologic investigations may be required, guided by clinical presentation.

If a patient develops Omenn syndrome–like features, it is important to confirm this complication by documenting the occurrence of oligo-clonal T-lymphocyte expansions by immunophenotyping and/or spectratyping, and, if applicable, by showing spongiosis with T-lymphocyte infiltration on analysis of skin biopsy samples.¹³¹ Patients developing an Omenn syndrome–like phenotype should be treated with cyclosporine A (CSA), with careful therapeutic monitoring of drug levels, typically aiming for trough levels of 150 to 200 µg/L. They also require careful skin management with emollients and, for troublesome skin symptoms, additional topical corticosteroids. Expert nutritional support is also essential. In patients with more severe clinical features, it may be necessary to temporarily treat with systemic steroids before immunosuppression while awaiting therapeutic CSA levels to be established. Systemic steroids should be reduced and stopped when the inflammation is under control and CSA levels optimal. In severe cases, immunosuppression with antithymocyte globulin may be considered. Alemtuzumab should be avoided before thymus transplantation because of its potential for depleting dendritic cells.¹⁷¹ CSA treatment should be continued until after thymus transplantation.

Because athymia is often part of a wider congenital syndrome, patients may have comorbidities that require acute medical

attention, and some of these will require stabilization before treatment with thymus transplantation can be contemplated.^{117,119,172} These include patients with CHD, who may need cardiac surgery before thymus transplantation; airway stabilization with positive airway pressure or tracheostomy due to underlying anatomic anomalies; and correction of hypocalcemia due to hypoparathyroidism. In patients with very wasted quadriceps muscles resulting from failure to thrive, implantation of thymus tissue may be difficult, and a period of nutritional support may be required to achieve muscle gain before the procedure. In some patients with life-limiting comorbidities, in particular cardiac and/or neurologic, palliative care is considered in discussion with the parents and multidisciplinary teams.^{117,119} For all patients and their families, provision of adequate psychosocial support is essential.¹¹⁹

Overall, athymic patients often require complex clinical care, which benefits from multispecialty coordination and early involvement of the thymus transplantation team.^{119,172}

Thymus transplantation

Thymus transplantation programs have been established in the United States at Duke University Medical Center since 1993¹¹⁵ and in the United Kingdom at GOSH since 2009.¹¹⁶ In the United States, cultured thymus tissue implantation, was approved as a regenerative medicinal product (Rethymic by Enzyvant, now Summito Pharma) by the US Food and Drug Administration in 2021 for the treatment of congenital athymia at Duke Medical Center. Conversely, in the United Kingdom, thymus transplantation is not considered a medicinal product but is offered as a transplantation procedure as part of a nationally commissioned transplantation service regulated by the Human Tissue Authority. Both transplantation services are available to patients outside of the United States or the United Kingdom, although the service offered by GOSH is significantly less expensive than the commercialized treatment in the United States. At GOSH, treatment of patients from the European Union is currently still funded through reciprocal health care agreements with the United Kingdom, and those outside of the European Union can access thymus transplantation through the International and Private Patients service.¹⁷³

Although the treatment access pathways are different, tissue preparation and implantation are broadly similar across both centers. Thymus tissue is donated by immunocompetent infants undergoing median sternotomy for cardiac surgery if tissue needs to be removed to improve access to the surgical field.^{55,141} The tissue is processed into slices, which are then cultured for 13 to 19 days to deplete donor thymocytes while preserving the thymic stroma.¹¹³ Once thymus tissue is in culture, rapid transfer of the recipient to the thymus transplantation center needs to be arranged. Donors and recipients do not need to be tissue type matched for human leukocyte antigens (HLAs), given the ability of donor thymus to induce tolerance,^{174,175} but screening for anti-HLA antibodies in the recipient is recommended before proceeding with implantation of donor tissue. After microbiologic and histopathologic assessment to confirm safety and suitability,¹⁷⁶ cultured thymus tissue is implanted bilaterally into the quadriceps muscles of the athymic recipient.

Immunosuppression before transplantation is only required in patients with Omenn syndrome-like features and oligoclonal T-lymphocyte expansions. This is achieved with antithymocyte

globulin serotherapy (Genzyme, 2 mg/kg once daily, 3 doses) in the days just before allograft implantation, in addition to previously established CSA treatment (trough levels of 150–200 µg/L). CSA is subsequently continued after transplantation until initial thymic output is evidenced by a frequency of naive CD4⁺ T lymphocytes of >10% within total peripheral CD4⁺ T lymphocytes, after which it is slowly decreased over 8 weeks.

Patients are usually transferred back to the referring center 2 to 4 weeks after the thymus transplantation procedure. Immune reconstitution after thymus transplantation is slow,^{115,116} typically taking 5 or 6 months before naive T lymphocytes can be found in the peripheral blood—and longer in particular in patients with certain risk factors, such as systemic viral infections or ongoing cardiorespiratory instability. Therefore, patients initially remain severely immunocompromised after transplantation and require continued isolation, monitoring, and unchanged antimicrobial prophylaxis until satisfactory immune reconstitution is achieved. Avoidance of procedures requiring (prolonged) general anaesthesia and invasive ventilation in the weeks after thymus transplantation is recommended, if possible, because these may compromise revascularization of the allograft.¹⁷⁷ In the posttransplantation period, care is also needed in avoiding potential adverse consequences of treatments for comorbidities, such as receipt of corticosteroid therapy for airway issues and receipt of testosterone, which has negative trophic effects on the thymus,^{178,179} for treatment of micropenis when present in CHARGE syndrome.

On revascularization of the implanted thymus tissue, the allograft will be repopulated with recipient-derived T-lymphocyte precursors, which then undergo stepwise maturation into functional T lymphocytes before egressing into the peripheral circulation. While this takes several months, thymopoiesis can typically be detected within the thymic grafts by 2 to 3 months after the procedure.¹⁸⁰ The panel recommends histopathologic assessment of thymic graft biopsy samples approximately 3 months after transplantation. This is best done by specialist surgeons, requiring a short readmission to the thymus transplantation center. Knowing the status of thymopoiesis in the thymic graft allows more informed decisions when managing potential complications such as autoimmune or inflammatory disease before initial recovery of peripheral T-lymphocyte immunity.

MANAGEMENT OF EARLY COMPLICATIONS AFTER THYMUS TRANSPLANTATION (BEFORE IMMUNE RECONSTITUTION)

The majority of deaths occur in the first year after thymus transplantation.^{115,116} Infection is the most frequent cause of mortality, including preexisting infections and posttransplantation infections acquired before immune reconstitution has been achieved. Systemic viral infections are particularly challenging to manage in these patients because recovery of T-lymphocyte immunity after thymus transplantation is slow.

In the first months after thymus transplantation, patients should be carefully monitored for inflammatory complications. Before immune reconstitution, patients may still develop Omenn syndrome-like features as a result of their underlying condition, necessitating immunosuppression with CSA (targeting trough levels of 150–200 µg/L). At the time of immune reconstitution, preexisting infections or previously administered bacillus Calmette-Guérin can provoke an immune reconstitution inflammatory

response.¹¹⁶ Life-threatening immune reconstitution inflammatory response requires treatment with high-dose steroids, which inhibits thymopoiesis and thus delays immune reconstitution. Early inflammatory complications are often transient, and if mild, steroid-sparing strategies should be explored to protect early thymopoiesis in the thymic allografts. A significant proportion of patients develop transient autoimmunity early after thymus transplantation, including autoimmune cytopenias and nephropathy.^{115,116} Steroid-sparing strategies such as immunomodulation with high-dose intravenous immunoglobulin and/or B-lymphocyte depletion with rituximab are preferable. When faced with these various inflammatory and autoimmune complications, prior documentation of the status of thymopoiesis on thymic graft biopsy samples is helpful to aid therapeutic decision making, allowing the risks of these complications to be balanced against those posed by steroid-related toxicity within the developing allograft.¹¹⁹ In a small number of patients, early donor T-lymphocyte engraftment has been found,¹¹⁵ but without any adverse effects such as graft-versus-host disease. The level of this engraftment diminishes over time.

LONG-TERM FOLLOW-UP AFTER THYMUS TRANSPLANTATION AND LATE COMPLICATIONS

After the first emergence of naive T lymphocytes to the periphery, T-lymphocyte counts progressively increase over time, peaking 1 to 2 years after thymus transplantation.^{115,116} In most patients, the absolute numbers of total T lymphocytes, naive T lymphocytes, and TRECs (quantified on peripheral blood T lymphocytes) remain below the 10th percentile for age. Nevertheless, thymic output is sustained with at least 10% of T lymphocytes continuing to have a naive phenotype, the TCR repertoire becomes diverse, and T-lymphocyte proliferative responses to mitogens and antigens normalize. All this, taken together, suggests that despite suboptimal counts, thymic output is sufficient for satisfactory T-lymphocyte immunity, allowing clearance of existing and new infections. Antimicrobial prophylaxis and IgRT can usually be discontinued.

Continuation of IgRT for 1 or 2 years after thymus transplantation until there is evidence of increasing and sustained recovery of T-lymphocyte immunity is recommended. The choice of home therapy with subcutaneous immunoglobulin infusions should be provided to families, if available, via the child's primary hospital with appropriate training and support for parents.¹⁸¹ Immunizations, following national immunization schedules, should be commenced 3 months after discontinuation of IgRT if IgG levels are maintained within normal ranges and the patient remains clinically well. Once immunizations begin, it is important to document protective antibody titers against inactivated vaccines before proceeding with administration of live attenuated vaccines. The latter have been administered to most transplanted patients without any adverse events despite low CD8⁺ T-lymphocyte counts in some patients. Long-term clinical and immunologic outcomes at >2 years after thymus transplantation have not been published for most patients, and multicenter data collection is limited by the absence of congenital athymia as an entity in IIE registries.¹¹⁹ A standardized immunophenotyping protocol for monitoring is recommended, including assessment of naive T lymphocytes and/or recent thymic emigrants at regular intervals after thymus transplantation. The long-term monitoring of thymic output also benefits from

TREC analysis on sorted T lymphocytes and thymus donor engraftment studies.

Thymus transplantation is aimed as a one-off therapy, and only one patient has been reported to require a second thymus transplantation procedure after the first graft failed in the context of sepsis shortly after implantation.¹¹⁶ To date, few late deaths have been reported after thymus transplantation^{115,117} and no late complications have occurred, such as significant infections due to hypothetical thymic graft exhaustion or malignancies, although overall follow-up is still relatively short.

Ongoing autoimmunity is relatively common after thymus transplantation,^{55,115,116} suggesting that the allografts may generate suboptimal recipient-specific central tolerance. While chronic cytopenias have been reported in a tiny number of patients, autoimmunity after thymus transplantation mainly seems to be restricted against the thyroid, with autoimmune thyroiditis observed at significantly higher rates than in patients with thymic hypoplasia.⁴⁹ It is therefore important to regularly monitor thyroid function in patients and, if impaired, to test for antithyroid autoantibodies.

The panel recommends multidisciplinary long-term follow-up. Syndromic comorbidities do not resolve after thymus transplantation, but their management becomes easier after satisfactory immune reconstitution.¹⁷² A degree of predisposition to recurrent infections may persist in patients with coexisting anatomic or functional abnormalities, particularly of the airways. However, infection risk overall decreases, and patients (and their families) can proceed with socialization, thus improving their overall quality of life with better access to health, education, and social services.¹¹⁹ Access to ongoing familial support can be provided through immunology clinical nurse specialists and dedicated clinical nurse specialists at the thymus transplantation center.¹¹⁹ This support encompasses emotional and psychological support, education and training, and signposting. There are no established patient advocacy groups for congenital athymia, but clinical teams may be able to offer peer support from previously treated families.

ROLE FOR HCT IN CONGENITAL ATHYMIA

Although thymus transplantation is recognized as the most appropriate treatment for congenital athymia, T-cell-replete HCT from a fully HLA-matched donor can be attempted under certain circumstances.^{74,120,182} These include situations where geographic and/or financial constraints prevent access to thymus transplantation, and in the context of preexisting systemic viral infection (ie, disseminated CMV disease and adenofibroma). In such situations, HCT can confer some degree of T-lymphocyte immunity through the transfer of postthymic mature T lymphocytes. Overall, HCT in congenital athymia yields poor outcomes, with a survival rate below 50% and a high risk of graft-versus-host disease.^{99,120} One study reported improved outcomes if a matched sibling donor is available, with an initial survival rate of 60% (8/13), and 25% (2/8) for those who received HCT from HLA-matched unrelated donors.^{120,121} The quality of immune reconstitution after HCT is also inferior to that achieved with thymus transplantation, as it relies on homeostatic expansion of transferred mature T lymphocytes rather than generation of new T lymphocytes. Naive T-lymphocyte counts are therefore lower and the TCR repertoire remains restricted after HCT.¹¹⁷ If HCT is from a matched family donor, it is recommended that no

conditioning be provided, and this should allow immune reconstitution to a degree capable of controlling viral infections significantly more rapidly compared to after thymus transplantation.^{74,182} It may be possible to subsequently proceed with a thymus transplantation procedure to improve immunity, provided there have been no severe complications after HCT, as has been reported in 5 patients.^{105,115} Thymus transplantation after HCT requires careful selection of donor thymus tissue with partial tissue type matching with the HSC donor at any alleles that were mismatched with the recipient.^{105,115} The development of more effective antiviral medications and advances in adoptive virus-specific T-lymphocyte therapies may also provide temporary solutions for patients before thymus transplantation.^{183,184}

TRANSLATIONAL RESEARCH AND FUTURE DIRECTIONS FOR NOVEL TREATMENT STRATEGIES

We have already emphasized the growing use of research assays, in particular transcription factor reporter assays and *ex vivo* T-lymphocyte differentiation assays in the clinical diagnosis and therapeutic management of patients with a genetically undefined T-B+NK+ immunophenotype. These *in vitro* assays also play a role in disease modeling of novel defects associated with impaired T lymphopoiesis, including novel thymic stromal cell defects, by delineating their likely HSC-intrinsic or HSC-extrinsic nature.^{78,100,156,157,160,185,186} Additionally, induced pluripotent stem cells can be differentiated into thymic epithelial progenitor cells,¹⁸⁷⁻¹⁸⁹ making it possible to specifically investigate the impact of novel variants in known disease-causing genes or in candidate genes on TEC development and function, when starting from patient-derived and/or gene-edited induced pluripotent stem cell lines.^{99,185,190} Together, these approaches facilitate the characterization of novel defects, contributing to strengthening the diagnostic pathway on validation of more disease-causing genes.

Research is also focused on optimizing the current treatment approach using cultured postnatal thymus tissue, and on developing novel treatment strategies.⁵⁵ How HLA-mismatched thymic allografts are able to support thymopoiesis remains poorly understood, yet the incomplete recovery of T-lymphocyte immunity and relatively common autoimmune manifestations after thymus transplantation suggest that T-lymphocyte development and induction of central tolerance are suboptimal. To date, no beneficial effect has been reported for fortuitous partial HLA matching between thymus donor and athymic recipient.^{116,191} This needs to be reassessed in the growing cohorts of transplanted patients, but in theory at least, partial tissue type matching may contribute to improving outcomes. Generation of a thymus tissue bank would be required for partial tissue type matching to be feasible, and encouragingly, preclinical data suggest that cryopreserved thymus tissue can support T-lymphocyte development after culture and transplantation into athymic mice.^{192,193}

Considerable progress has been made recently toward producing engineered thymic stroma suitable for clinical applications.⁵⁵ Natural decellularized extracellular matrix from human thymus can be obtained and seeded with human thymus stromal progenitor cells that can be expanded *in vitro*.^{20,26} Larger amounts of human thymus stroma could be produced with this approach, and because stromal progenitors can be banked, partial tissue type matching would also be feasible.

CONCLUSIONS

Congenital athymia is a life-limiting disorder, requiring corrective treatment, ideally by transplanting lymphodepleted donor thymus tissue. We have provided expert guidance on the diagnosis, investigation, and management of patients with congenital athymia, with the aim of improving their outcomes. Our panel recommendations are summarized in Fig 1.

Outcomes are best with early diagnosis and institution of both supportive and definitive management, before infectious and inflammatory complications have developed. We have therefore emphasized the scenarios in which congenital athymia can present. Namely, patients might present clinically with syndromic features, infections, or inflammatory complications; increasingly, patients are identified in the context of TREC-based NBS programs for SCID; and finally, failed T-lymphocyte reconstitution after allogeneic HCT for genetically undefined suspected SCID should alert to the possibility of an absent thymic niche. These scenarios should prompt the completion of the set of investigations recommended herein.

Increasing access to NBS and comprehensive genetic testing are promoting early diagnosis and treatment, resulting in better outcomes after thymus transplantation, but have also revealed a particular challenge with respect to the diagnosis of athymia versus hematopoietic cell-intrinsic defects in molecularly undefined congenital T-lymphocyte deficiency, where knowing the cellular etiology influences treatment choice between thymus transplantation and HCT. With higher morbidity and mortality rates after HCT in athymic patients, clinical translation of research assays to assist in diagnosis and therapeutic decision-making is essential.

Thymus transplantation, the recommended first-line treatment for congenital athymia, is a highly specialized treatment, currently only available in 2 centers worldwide. Patients treated with thymus transplantation have good outcomes overall, with durable T-lymphocyte immunity and improved quality of life. Mortality is mostly due to infections occurring before treatment or before successful immune reconstitution, which is typically slow after thymus transplantation. Autoimmune complications are relatively common, in particular transient autoimmune cytopenias and persistent autoimmune thyroiditis. The panel recommends centralized recording of long-term clinical and immunologic outcomes after thymus transplantation to adequately prioritize strategies aimed at improving outcomes for athymic patients yet is complicated by the geographical spread of the patients and the lack of national and international registries for congenital athymia. Cost and availability of lifesaving novel gene and cell therapies for rare conditions are a concern even in high-income countries, and while in Europe accessible and timely treatment with thymus transplantation is available, continued efforts are necessary to overcome geographic and economic challenges and to promote equitable and timely access to treatment. In the future, the further expansion of NBS programs, the creation of dedicated registries, the development of new approaches for thymus replacement therapy, and the support of initiatives promoting sustained accessibility to novel therapies may address some of these challenges.

DISCLOSURE STATEMENT

A.Y.K. is supported by the Wellcome Trust (222096/Z/20/Z). F.D. is supported by an NIHR Academic Clinical Lectureship

and an Academy for Medical Sciences Starter Grant. E.H. and E.G.D. are supported by a grant from the Great Ormond Street Hospital Children's Charity. O.E.'s laboratory is supported by grants from the Swedish Research Council (2018-02752 and 2022-00781) and the Swedish state under an ALF agreement between the Swedish government and the county councils (ALFGBG-965795). A.V. is supported by a core grant from the Telethon Foundation. F.J.T.'s laboratory is supported in part by EU H2020 grant RECOMB (755170–^b) and has received funding from the European Union Horizon 2020 research and innovation program as well as from reNEW, the Novo Nordisk Foundation for Stem Cell Research (NNF21CC0073729). G.A. is supported by an MRC Programme Grant to GA (MR/T029765/a). and G.H. is supported by the Wellcome Trust (211944/Z/18/Z).

Disclosure of potential conflict of interest: The authors declare that they have no relevant conflicts of interest.

The European Society for Immunodeficiencies Clinical Working Party collaborators are as follows: Siobhan O. Burns, Institute of Immunity and Transplantation, University College London, and Department of Immunology, The Royal Free London NHS Foundation Trust, London, United Kingdom; Maria Carrabba, Internal Medicine Department, RITA-ERN Center, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico di Milano, Milan, Italy; Ann Gardulf, Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden, and Faculty of Social and Health Sciences, Inland Norway University of Applied Sciences, Elverum, Norway; Filomeen Haerynck, Primary Immune Deficiency Research Laboratory, Department of Internal Diseases and Pediatrics, Ghent University, and Department of Pediatric Pulmonology, Infectious Diseases and Immune Deficiency, Centre for Primary Immune Deficiency Ghent, Jeffrey Modell Diagnosis and Research Centre, Ghent University Hospital, Ghent, Belgium; Fabian Hauck, Division of Pediatric Immunology and Rheumatology, Department of Pediatrics, Dr von Hauner Children's Hospital, University Hospital, Ludwig-Maximilians-Universität München, Munich, Germany; Peter Jandus, Division of Clinical Immunology and Allergy, Department of Medicine, University Hospital and Medical Faculty, University of Geneva, Geneva, Switzerland; Adam Klocperk, Department of Immunology, 2nd Faculty of Medicine Charles University, University Hospital in Motol, Prague, Czechia; Isabelle Meyts, Department of Pediatrics, University Hospital Leuven, and Laboratory for Inborn Errors of Immunity, Department of Microbiology, Immunology and Transplantation, KU Leuven, and FWO Vlaanderen and JMF Diagnostic and Research Center Leuven, Leuven, Belgium; Bénédicte Neven, Pediatric Hematology-Immunology and Rheumatology Department, Assistance Publique Hôpitaux de Paris, and Necker University Hospital, and Laboratory of Immunogenetics of Pediatric Autoimmune Diseases, Imagine Institute, INSERM UMR-S-1163, and Université Paris Cité, Paris, France; Malgorzata Pac, Department of Immunology, The Children's Memorial Health Institute, Warsaw, Poland; Martine Pergent, The International Patient Organisation for Primary Immunodeficiencies, Brussels, Belgium; Anna Sediva, Department of Immunology, 2nd Faculty of Medicine Charles University, University Hospital in Motol, Prague, Czechia; Pere Soler-Palacín, Pediatric Infectious Diseases and Immunodeficiencies Unit, University Hospital Vall d'Hebron, and Institut de Recerca Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Catalonia, Spain; Margarita Velcheva, Kenes Group, Amsterdam, Netherlands; Klaus Warnatz, Department of Rheumatology and Clinical Immunology and Center for Chronic Immunodeficiency, Medical Center—University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany.

REFERENCES

- Nitta T, Suzuki H. Thymic stromal cell subsets for T cell development. *Cell Mol Life Sci* 2016;73:1021-37.
- Nitta T, Takayanagi H. Non-epithelial thymic stromal cells: unsung heroes in thymus organogenesis and T cell development. *Front Immunol* 2020;11:620894.
- Manley NR, Richie ER, Blackburn CC, Condie BG, Sage J. Structure and function of the thymic microenvironment. *Front Biosci (Landmark Ed)* 2011;16:2461-77.
- Gordon J, Wilson VA, Blair NF, Sheridan J, Farley A, Wilson L, et al. Functional evidence for a single endodermal origin for the thymic epithelium. *Nat Immunol* 2004;5:546-53.
- Bockman DE, Kirby ML. Dependence of thymus development on derivatives of the neural crest. *Science* 1984;223(4635):498-500.
- Itoi M, Tsukamoto N, Yoshida H, Amagai T. Mesenchymal cells are required for functional development of thymic epithelial cells. *Int Immunol* 2007;19:953-64.
- Farley AM, Morris LX, Vroegindewij E, Depreter ML, Vaidya H, Stenhouse FH, et al. Dynamics of thymus organogenesis and colonization in early human development. *Development* 2013;140:2015-26.
- Manley NR, Capecchi MR. The role of Hoxa-3 in mouse thymus and thyroid development. *Development* 1995;121:1989-2003.
- Wallin J, Eibel H, Neubuser A, Wiltling J, Koseki H, Balling R. *Pax1* is expressed during development of the thymus epithelium and is required for normal T-cell maturation. *Development* 1996;122:23-30.
- Jerome LA, Papaioannou VE. DiGeorge syndrome phenotype in mice mutant for the T-box gene, *Tbx1*. *Nat Genet* 2001;27:286-91.
- Hetzer-Egger C, Schorpp M, Haas-Assenbaum A, Balling R, Peters H, Boehm T. Thymopoiesis requires *Pax9* function in thymic epithelial cells. *Eur J Immunol* 2002;32:1175-81.
- Xu H, Cerrato F, Baldini A. Timed mutation and cell-fate mapping reveal reiterated roles of *Tbx1* during embryogenesis, and a crucial function during segmentation of the pharyngeal system via regulation of endoderm expansion. *Development* 2005;132:4387-95.
- Zuklys S, Handel A, Zhanybekova S, Govani F, Keller M, Maio S, et al. *Foxn1* regulates key target genes essential for T cell development in postnatal thymic epithelial cells. *Nat Immunol* 2016;17:1206-15.
- Romano R, Palamaro L, Fusco A, Giardino G, Gallo V, Del Vecchio L, et al. *FOXN1*: a master regulator gene of thymic epithelial development program. *Front Immunol* 2013;4:187.
- Blackburn CC, Augustine CL, Li R, Harvey RP, Malin MA, Boyd RL, et al. The *nu* gene acts cell-autonomously and is required for differentiation of thymic epithelial progenitors. *Proc Natl Acad Sci U S A* 1996;93:5742-6.
- Nehls M, Pfeifer D, Schorpp M, Hedrich H, Boehm T. New member of the winged-helix protein family disrupted in mouse and rat nude mutations. *Nature* 1994;372(6501):103-7.
- Chen L, Xiao S, Manley NR. *Foxn1* is required to maintain the postnatal thymic microenvironment in a dosage-sensitive manner. *Blood* 2009;113:567-74.
- Wortis HH, Nehlsen S, Owen JJ. Abnormal development of the thymus in "nude" mice. *J Exp Med* 1971;134(3 pt 1):681-92.
- Park JE, Botting RA, Dominguez Conde C, Popescu DM, Lavaert M, Kunz DJ, et al. A cell atlas of human thymic development defines T cell repertoire formation. *Science* 2020;367(6480).
- Campinoti S, Gjinovci A, Ragazzini R, Zanieri L, Ariza-McNaughton L, Catucci M, et al. Reconstitution of a functional human thymus by postnatal stromal progenitor cells and natural whole-organ scaffolds. *Nat Commun* 2020;11:6372.
- Bautista JL, Cramer NT, Miller CN, Chavez J, Berrios DI, Byrnes LE, et al. Single-cell transcriptional profiling of human thymic stroma uncovers novel cellular heterogeneity in the thymic medulla. *Nat Commun* 2021;12:1096.
- Baran-Gale J, Morgan MD, Maio S, Dhalla F, Calvo-Asensio I, Deadman ME, et al. Ageing compromises mouse thymus function and remodels epithelial cell differentiation. *Elife* 2020;9:e56221.
- Klein F, Veiga-Villauriz C, Borsch A, Maio S, Palmer S, Dhalla F, et al. Combined multidimensional single-cell protein and RNA profiling dissects the cellular and functional heterogeneity of thymic epithelial cells. *Nat Commun* 2023;14:4071.
- Dhalla F, Baran-Gale J, Maio S, Chappell L, Hollander GA, Ponting CP. Biologically indeterminate yet ordered promiscuous gene expression in single medullary thymic epithelial cells. *EMBO J* 2020;39:e101828.
- Givony T, Leshkowitz D, Del Castillo D, Nevo S, Kadouri N, Dassa B, et al. Thymic mimetic cells function beyond self-tolerance. *Nature* 2023;622(7981):164-72.
- Ragazzini R, Boeig S, Zanieri L, Green M, D'Agostino G, Bartolovic K, et al. Defining the identity and the niches of epithelial stem cells with highly pleiotropic multilineage potency in the human thymus. *Dev Cell* 2023;58:2428-46.e9.
- Anderson G, Takahama Y. Thymic epithelial cells: working class heroes for T cell development and repertoire selection. *Trends Immunol* 2012;33:256-63.
- Takaba H, Takayanagi H. The mechanisms of T cell selection in the thymus. *Trends Immunol* 2017;38:805-16.
- Watson SA, Javanmardi Y, Zanieri L, Shahreza S, Ragazzini R, Bonfanti P, et al. Integrated role of human thymic stromal cells in hematopoietic stem cell extravasation. *Bioeng Transl Med* 2023;8:e10454.
- Klein L, Robey EA, Hsieh CS. Central CD4⁺ T cell tolerance: deletion versus regulatory T cell differentiation. *Nat Rev Immunol* 2019;19:7-18.

31. Derbinski J, Schulte A, Kyewski B, Klein L. Promiscuous gene expression in medullary thymic epithelial cells mirrors the peripheral self. *Nat Immunol* 2001;2:1032-9.
32. Sansom SN, Shikama-Dorn N, Zhanybekova S, Nusspaumer G, Macaulay IC, Deadman ME, et al. Population and single-cell genomics reveal the Aire dependency, relief from Polycomb silencing, and distribution of self-antigen expression in thymic epithelia. *Genome Res* 2014;24:1918-31.
33. Takaba H, Morishita Y, Tomofuji Y, Danks L, Nitta T, Komatsu N, et al. Fezf2 orchestrates a thymic program of self-antigen expression for immune tolerance. *Cell* 2015;163:975-87.
34. Yano M, Kuroda N, Han H, Meguro-Horike M, Nishikawa Y, Kiyonari H, et al. Aire controls the differentiation program of thymic epithelial cells in the medulla for the establishment of self-tolerance. *J Exp Med* 2008;205:2827-38.
35. Klein L, Hinterberger M, von Rohrscheidt J, Aichinger M. Autonomous versus dendritic cell-dependent contributions of medullary thymic epithelial cells to central tolerance. *Trends Immunol* 2011;32:188-93.
36. Hadeiba H, Lahl K, Edalati A, Oderup C, Habtezion A, Pachynski R, et al. Plasmacytoid dendritic cells transport peripheral antigens to the thymus to promote central tolerance. *Immunity* 2012;36:438-50.
37. Yamano T, Nedjic J, Hinterberger M, Steinert M, Koser S, Pinto S, et al. Thymic B cells are licensed to present self antigens for central T cell tolerance induction. *Immunity* 2015;42:1048-61.
38. Yamano T, Steinert M, Klein L. Thymic B cells and central T cell tolerance. *Front Immunol* 2015;6:376.
39. Inaba M, Inaba K, Hosono M, Kumamoto T, Ishida T, Muramatsu S, et al. Distinct mechanisms of neonatal tolerance induced by dendritic cells and thymic B cells. *J Exp Med* 1991;173:549-59.
40. Frommer F, Heinen TJ, Wunderlich FT, Yorgev N, Buch T, Roers A, et al. Tolerance without clonal expansion: self-antigen-expressing B cells program self-reactive T cells for future deletion. *J Immunol* 2008;181:5748-59.
41. Frommer F, Waisman A. B cells participate in thymic negative selection of murine auto-reactive CD4⁺ T cells. *PLoS One* 2010;5:e15372.
42. Kleindienst P, Chretien I, Winkler T, Brocker T. Functional comparison of thymic B cells and dendritic cells *in vivo*. *Blood* 2000;95:2610-6.
43. Adams SP, Kricke S, Ralph E, Gilmour N, Gilmour KC. A comparison of TRECs and flow cytometry for naive T cell quantification. *Clin Exp Immunol* 2018;191:198-202.
44. Kreins AY, Maio S, Dhalla F. Inborn errors of thymic stromal cell development and function. *Semin Immunopathol* 2021;43:85-100.
45. Tangye SG, Al-Herz W, Bousfiha A, Cunningham-Rundles C, Franco JL, Holland SM, et al. Human inborn errors of immunity: 2022 update on the classification from the International Union of Immunological Societies Expert Committee. *J Clin Immunol* 2022;42:1473-507.
46. Dou Y, Schindewolf E, Crowley TB, McGinn DM, Moldenhauer JS, Coleman B, et al. The association of fetal thymus size with subsequent T cell counts in 22q11.2 deletion syndrome. *J Clin Immunol* 2020;40:783-5.
47. Giardino G, Borzacchiello C, De Luca M, Romano R, Prencipe R, Cirillo E, et al. T-cell immunodeficiencies with congenital alterations of thymic development: genes implicated and differential immunological and clinical features. *Front Immunol* 2020;11:1837.
48. Wong MT, Lambeck AJ, van der Burg M, la Bastide-van Gemert S, Hogendorf LA, van Ravenswaaij-Arts CM, et al. Immune dysfunction in children with CHARGE syndrome: a cross-sectional study. *PLoS One* 2015;10:e0142350.
49. Giardino G, Radwan N, Koletsi P, Morrogh DM, Adams S, Ip W, et al. Clinical and immunological features in a cohort of patients with partial DiGeorge syndrome followed at a single center. *Blood* 2019;133:2586-96.
50. Ryan AK, Goodship JA, Wilson DI, Philip N, Levy A, Seidel H, et al. Spectrum of clinical features associated with interstitial chromosome 22q11 deletions: a European collaborative study. *J Med Genet* 1997;34:798-804.
51. Piliero LM, Sanford AN, McDonald-McGinn DM, Zackai EH, Sullivan KE. T-cell homeostasis in humans with thymic hypoplasia due to chromosome 22q11.2 deletion syndrome. *Blood* 2004;103:1020-5.
52. Davies EG. Immunodeficiency in DiGeorge syndrome and options for treating cases with complete athymia. *Front Immunol* 2013;4:322.
53. Jyonouchi S, McDonald-McGinn DM, Bale S, Zackai EH, Sullivan KE. CHARGE (coloboma, heart defect, atresia choanae, retarded growth and development, genital hypoplasia, ear anomalies/deafness) syndrome and chromosome 22q11.2 deletion syndrome: a comparison of immunologic and nonimmunologic phenotypic features. *Pediatrics* 2009;123:e871-7.
54. Sullivan KE. The clinical, immunological, and molecular spectrum of chromosome 22q11.2 deletion syndrome and DiGeorge syndrome. *Curr Opin Allergy Clin Immunol* 2004;4:505-12.
55. Kreins AY, Bonfanti P, Davies EG. Current and future therapeutic approaches for thymic stromal cell defects. *Front Immunol* 2021;12:655354.
56. Mustillo PJ, Sullivan KE, Chinn IK, Notarangelo LD, Haddad E, Davies EG, et al. Clinical practice guidelines for the immunological management of chromosome 22q11.2 deletion syndrome and other defects in thymic development. *J Clin Immunol* 2023;43:247-70.
57. Gennery AR, Slatter MA, Rice J, Hoefsloot LH, Barge D, McLean-Tooke A, et al. Mutations in *CHD7* in patients with CHARGE syndrome cause T-B + natural killer cell + severe combined immune deficiency and may cause Omenn-like syndrome. *Clin Exp Immunol* 2008;153:75-80.
58. Markert ML, Devlin BH, Alexieff MJ, Li J, McCarthy EA, Gupton SE, et al. Review of 54 patients with complete DiGeorge anomaly enrolled in protocols for thymus transplantation: outcome of 44 consecutive transplants. *Blood* 2007;109:4539-47.
59. Dvorak CC, Haddad E, Heimall J, Dunn E, Buckley RH, Kohn DB, et al. The diagnosis of severe combined immunodeficiency (SCID): the Primary Immune Deficiency Treatment Consortium (PIDTC) 2022 definitions. *J Allergy Clin Immunol* 2023;151:539-46.
60. Bhalla P, Wysocki CA, van Oers NSC. Molecular insights into the causes of human thymic hypoplasia with animal models. *Front Immunol* 2020;11:830.
61. Wurdak H, Ittner LM, Sommer L. DiGeorge syndrome and pharyngeal apparatus development. *Bioessays* 2006;28:1078-86.
62. Cancrini C, Puliafito P, Digilio MC, Soresina A, Martino S, Rondelli R, et al. Clinical features and follow-up in patients with 22q11.2 deletion syndrome. *J Pediatr* 2014;164:1475-80.e2.
63. Tezenas Du Montcel S, Mendizabai H, Ayme S, Levy A, Philip N. Prevalence of 22q11 microdeletion. *J Med Genet* 1996;33:719.
64. McLean-Tooke A, Barge D, Spickett GP, Gennery AR. Immunologic defects in 22q11.2 deletion syndrome. *J Allergy Clin Immunol* 2008;122:362-7.e1-4.
65. Rozas MF, Benavides F, Leon L, Repetto GM. Association between phenotype and deletion size in 22q11.2 microdeletion syndrome: systematic review and meta-analysis. *Orphanet J Rare Dis* 2019;14:195.
66. Rauch A, Zink S, Zweier C, Thiel CT, Koch A, Rauch R, et al. Systematic assessment of atypical deletions reveals genotype-phenotype correlation in 22q11.2. *J Med Genet* 2005;42:871-6.
67. Yagi H, Furutani Y, Hamada H, Sasaki T, Asakawa S, Minoshima S, et al. Role of *TBX1* in human del22q11.2 syndrome. *Lancet* 2003;362(9393):1366-73.
68. Fulcoli FG, Huynh T, Scambler PJ, Baldini A. *Tbx1* regulates the BMP-Smad1 pathway in a transcription independent manner. *PLoS One* 2009;4:e6049.
69. Writzl K, Cale CM, Pierce CM, Wilson LC, Hennekam RC. Immunological abnormalities in CHARGE syndrome. *Eur J Med Genet* 2007;50:338-45.
70. Wong MT, Scholvinck EH, Lambeck AJ, van Ravenswaaij-Arts CM. CHARGE syndrome: a review of the immunological aspects. *Eur J Hum Genet* 2015;23:1451-9.
71. Janssen N, Bergman JE, Swertz MA, Tranebjaerg L, Lodahl M, Schoots J, et al. Mutation update on the *CHD7* gene involved in CHARGE syndrome. *Hum Mutat* 2012;33:1149-60.
72. Hasegawa K, Tanaka H, Higuchi Y, Hayashi Y, Kobayashi K, Tsukahara H. Novel heterozygous mutation in *TBX1* in an infant with hypocalcemic seizures. *Clin Pediatr Endocrinol* 2018;27:159-64.
73. Ogata T, Niihori T, Tanaka N, Kawai M, Nagashima T, Funayama R, et al. *TBX1* mutation identified by exome sequencing in a Japanese family with 22q11.2 deletion syndrome-like craniofacial features and hypocalcemia. *PLoS One* 2014;9:e91598.
74. Chitty-Lopez M, Duff C, Vaughn G, Trotter J, Monforte H, Lindsay D, et al. Case report: unmanipulated matched sibling donor hematopoietic cell transplantation in *TBX1* congenital athymia: a lifesaving therapeutic approach when facing a systemic viral infection. *Front Immunol* 2021;12:721917.
75. Liu N, Schoch K, Luo X, Pena LDM, Bhavana VH, Kukulich MK, et al. Functional variants in *TBX2* are associated with a syndromic cardiovascular and skeletal developmental disorder. *Hum Mol Genet* 2018;27:2454-65.
76. Ensenaer RE, Adeyinka A, Flynn HC, Michels VV, Lindor NM, Dawson DB, et al. Microduplication 22q11.2, an emerging syndrome: clinical, cytogenetic, and molecular analysis of thirteen patients. *Am J Hum Genet* 2003;73:1027-40.
77. Bernstock JD, Totten AH, Elkahoul AG, Johnson KR, Hurst AC, Goldman F, et al. Recurrent microdeletions at chromosome 2p11.2 are associated with thymic hypoplasia and features resembling DiGeorge syndrome. *J Allergy Clin Immunol* 2020;145:358-67.e2.
78. Ghosh R, Bosticardo M, Singh S, Similuk M, Delmonte OM, Pala F, et al. *FOXP3* haploinsufficiency contributes to low T-cell receptor excision circles and T-cell lymphopenia. *J Allergy Clin Immunol* 2022;150:1556-62.
79. Daw SC, Taylor C, Kraman M, Call K, Mao J, Schuffenhauer S, et al. A common region of 10p deleted in DiGeorge and velocardiofacial syndromes. *Nat Genet* 1996;13:458-60.
80. Berger R, Larroche JC, Toubas PL. Deletion of the short arm of chromosome no. 10. *Acta Paediatr Scand* 1977;66:659-62.

81. Van Esch H, Groenen P, Fryns JP, Van de Ven W, Devriendt K. The phenotypic spectrum of the 10p deletion syndrome versus the classical DiGeorge syndrome. *Genet Couns* 1999;10:59-65.
82. Sun D, Lee J, Heimall J, Jyonouchi S. Immunodeficiency in 22q11.2 duplication syndrome. *J Allergy Clin Immunol Pract* 2021;9:996-8.e3.
83. Chen C, Zhang C, Deng Y, Du S, Wang H, Li D. Thymic hypoplasia induced by copy number variations contributed to explaining sudden infant death based on forensic autopsies. *Forensic Sci Int* 2022;336:111323.
84. Mary L, Lavillaureix A, Perrot A, Loget P, Launay E, Leborgne AS, et al. Prenatal phenotype of 22q11 micro-duplications: a systematic review and report on 12 new cases. *Eur J Med Genet* 2022;65:104422.
85. Bhattarai D, McGinn DE, Crowley TB, Giunta V, Gaiser K, Zackai EH, et al. Immunologic, molecular, and clinical profile of patients with chromosome 22q11.2 duplications. *J Clin Immunol* 2023;43:794-807.
86. Wilson TA, Blethen SL, Vallone A, Alenick DS, Nolan P, Katz A, et al. DiGeorge anomaly with renal agenesis in infants of mothers with diabetes. *Am J Med Genet* 1993;47:1078-82.
87. Gosseye S, Golaire MC, Verellen G, Van Lierde M, Claus D. Association of bilateral renal agenesis and DiGeorge syndrome in an infant of a diabetic mother. *Helv Paediatr Acta* 1982;37:471-4.
88. Ammann AJ, Wara DW, Cowan MJ, Barrett DJ, Stiehm ER. The DiGeorge syndrome and the fetal alcohol syndrome. *Am J Dis Child* 1982;136:906-8.
89. Coberly S, Lammer E, Alashari M. Retinoic acid embryopathy: case report and review of literature. *Pediatr Pathol Lab Med* 1996;16:823-36.
90. Roberts C, Ivins SM, James CT, Scambler PJ. Retinoic acid down-regulates *Tbx1* expression *in vivo* and *in vitro*. *Dev Dyn* 2005;232:928-38.
91. Vermot J, Niederreither K, Garnier JM, Chambon P, Dolle P. Decreased embryonic retinoic acid synthesis results in a DiGeorge syndrome phenotype in newborn mice. *Proc Natl Acad Sci U S A* 2003;100:1763-8.
92. Lee LM, Leung MB, Kwok RC, Leung YC, Wang CC, McCaffery PJ, et al. Perturbation of retinoid homeostasis increases malformation risk in embryos exposed to pregestational diabetes. *Diabetes* 2017;66:1041-51.
93. Yao H, Hill SF, Skidmore JM, Sperry ED, Swiderski DL, Sanchez GJ, et al. *CHD7* represses the retinoic acid synthesis enzyme *ALDH1A3* during inner ear development. *JCI Insight* 2018;3(4).
94. Mulder GB, Manley N, Maggio-Price L. Retinoic acid-induced thymic abnormalities in the mouse are associated with altered pharyngeal morphology, thymocyte maturation defects, and altered expression of *Hoxa3* and *Pax1*. *Teratology* 1998;58:263-75.
95. Bosticardo M, Notarangelo LD. Human thymus in health and disease: recent advances in diagnosis and biology. *Semin Immunol* 2023;66:101732.
96. Rota IA, Dhalla F. *FOXP1* deficient nude severe combined immunodeficiency. *Orphanet J Rare Dis* 2017;12:6.
97. Frank J, Pignata C, Panteleyev AA, Prowse DM, Baden H, Weiner L, et al. Exposing the human nude phenotype. *Nature* 1999;398(6727):473-4.
98. Pignata C, Fiore M, Guzzetta V, Castaldo A, Sebastio G, Porta F, et al. Congenital alopecia and nail dystrophy associated with severe functional T-cell immunodeficiency in two sibs. *Am J Med Genet* 1996;65:167-70.
99. Yamazaki Y, Urrutia R, Franco LM, Giliani S, Zhang K, Alazami AM, et al. *PAX1* is essential for development and function of the human thymus. *Sci Immunol* 2020;5(44).
100. Yakici N, Kreins AY, Catak MC, Babayeva R, Erman B, Kenney H, et al. Expanding the clinical and immunological phenotypes of *PAX1*-deficient SCID and CID patients. *Clin Immunol* 2023;255:109757.
101. Thompson B, Davidson EA, Liu W, Nebert DW, Bruford EA, Zhao H, et al. Overview of *PAX* gene family: analysis of human tissue-specific variant expression and involvement in human disease. *Hum Genet* 2021;140:381-400.
102. Pohl E, Aykut A, Beleggia F, Karaca E, Durmaz B, Keupp K, et al. A hypofunctional *PAX1* mutation causes autosomal recessively inherited otofaciocervical syndrome. *Hum Genet* 2013;132:1311-20.
103. Patil SJ, Das Bhowmik A, Bhat V, Satidevi Vineeth V, Vasudevamurthy R, Dalal A. Autosomal recessive otofaciocervical syndrome type 2 with novel homozygous small insertion in *PAX1* gene. *Am J Med Genet A* 2018;176:1200-6.
104. Paganini I, Sestini R, Capone GL, Putignano AL, Contini E, Giotti I, et al. A novel *PAX1* null homozygous mutation in autosomal recessive otofaciocervical syndrome associated with severe combined immunodeficiency. *Clin Genet* 2017;92:664-8.
105. Kreins AY, Worth A, Ghosh S, Mohammed RW, Davies EG. First use of thymus transplantation in *PAX1* deficiency. *J Clin Immunol* 2023;43:1127-30.
106. Giardino G, Sharapova SO, Ciznar P, Dhalla F, Maragliano L, Radha Rama Devi A, et al. Expanding the nude SCID/CID phenotype associated with *FOXP1* homozygous, compound heterozygous, or heterozygous mutations. *J Clin Immunol* 2021;41:756-68.
107. Markert ML, Marques JG, Neven B, Devlin BH, McCarthy EA, Chinn IK, et al. First use of thymus transplantation therapy for *FOXP1* deficiency (nude/SCID): a report of 2 cases. *Blood* 2011;117:688-96.
108. Chou J, Massaad MJ, Wakim RH, Bainter W, Dbabo G, Geha RS. A novel mutation in *FOXP1* resulting in SCID: a case report and literature review. *Clin Immunol* 2014;155:30-2.
109. Rota IA, Handel AE, Maio S, Klein F, Dhalla F, Deadman ME, et al. *FOXP1* forms higher-order nuclear condensates displaced by mutations causing immunodeficiency. *Sci Adv* 2021;7:eabj9247.
110. Bosticardo M, Yamazaki Y, Cowan J, Giardino G, Corsino C, Scalia G, et al. Heterozygous *FOXP1* variants cause low TRECs and severe T cell lymphopenia, revealing a crucial role of *FOXP1* in supporting early thymopoiesis. *Am J Hum Genet* 2019;105:549-61.
111. Du Q, Huynh LK, Coskun F, Molina E, King MA, Raj P, et al. *FOXP1* compound heterozygous mutations cause selective thymic hypoplasia in humans. *J Clin Invest* 2019;129:4724-38.
112. Moses A, Bhalla P, Thompson A, Lai L, Coskun FS, Seroogy CM, et al. Comprehensive phenotypic analysis of diverse *FOXP1* variants. *J Allergy Clin Immunol* 2023.
113. Markert ML, Kostyu DD, Ward FE, McLaughlin TM, Watson TJ, Buckley RH, et al. Successful formation of a chimeric human thymus allograft following transplantation of cultured postnatal human thymus. *J Immunol* 1997;158:998-1005.
114. Hong R, Santosham M, Schulte-Wissermann H, Horowitz S, Hsu SH, Winkelstein JA. Reconstitution of B and T lymphocyte function in severe combined immunodeficiency disease after transplantation with thymic epithelium. *Lancet* 1976;2(7998):1270-2.
115. Markert ML, Gupta SE, McCarthy EA. Experience with cultured thymus tissue in 105 children. *J Allergy Clin Immunol* 2022;149:747-57.
116. Davies EG, Cheung M, Gilmour K, Maimaris J, Curry J, Furmanski A, et al. Thymus transplantation for complete DiGeorge syndrome: European experience. *J Allergy Clin Immunol* 2017;140:1660-70.e16.
117. Howley E, Golwala Z, Buckland M, Barzaghi F, Ghosh S, Hackett S, et al. Impact of newborn screening for SCID on the management of congenital athymia. *J Allergy Clin Immunol* 2024;153:330-4.
118. Markert ML, Devlin BH, McCarthy EA. Thymus transplantation. *Clin Immunol* 2010;135:236-46.
119. Howley E, Davies EG, Kreins AY. Congenital athymia: unmet needs and practical guidance. *Ther Clin Risk Manag* 2023;19:239-54.
120. Janda A, Sedlacek P, Honig M, Friedrich W, Champagne M, Matsumoto T, et al. Multicenter survey on the outcome of transplantation of hematopoietic cells in patients with the complete form of DiGeorge anomaly. *Blood* 2010;116:2229-36.
121. Land MH, Garcia-Lloret MI, Borzy MS, Rao PN, Aziz N, McGhee SA, et al. Long-term results of bone marrow transplantation in complete DiGeorge syndrome. *J Allergy Clin Immunol* 2007;120:908-15.
122. Speckmann C, Nennstiel U, Honig M, Albert MH, Ghosh S, Schuetz C, et al. Prospective newborn screening for SCID in Germany: a first analysis by the Pediatric Immunology Working Group (API). *J Clin Immunol* 2023;43:965-78.
123. Puck JM. Newborn screening for severe combined immunodeficiency and T-cell lymphopenia. *Immunol Rev* 2019;287:241-52.
124. Wakamatsu M, Kojima D, Muramatsu H, Okuno Y, Kataoka S, Nakamura F, et al. TREC/KREC newborn screening followed by next-generation sequencing for severe combined immunodeficiency in Japan. *J Clin Immunol* 2022;42:1696-707.
125. Barreiros LA, Sousa JL, Geier C, Leiss-Piller A, Kanegae MPP, Franca TT, et al. SCID and other inborn errors of immunity with low TRECs—the Brazilian experience. *J Clin Immunol* 2022;42:1171-92.
126. Lev A, Sharir I, Simon AJ, Levy S, Lee YN, Frizinsky S, et al. Lessons learned from five years of newborn screening for severe combined immunodeficiency in Israel. *J Allergy Clin Immunol Pract* 2022;10:2722-31.e9.
127. Argudo-Ramirez A, Martin-Nalda A, Gonzalez de Aledo-Castillo JM, Lopez-Gallera R, Marin-Soria JL, Pajares-Garcia S, et al. Newborn screening for SCID. Experience in Spain (Catalonia). *Int J Neonatal Screen* 2021;7(3).
128. Heather N, de Hora M, Brothers S, Grainger P, Knoll D, Webster D. Introducing newborn screening for severe combined immunodeficiency—the New Zealand experience. *Int J Neonatal Screen* 2022;8(2).
129. Martin-Nalda A, Cueto-Gonzalez AM, Argudo-Ramirez A, Marin-Soria JL, Martinez-Gallo M, Colobran R, et al. Identification of 22q11.2 deletion syndrome via newborn screening for severe combined immunodeficiency. Two years' experience in Catalonia (Spain). *Mol Genet Genomic Med* 2019;7:e1016.
130. Howley E, Soomann M, Kreins AY. Parental engagement in identifying information needs after newborn screening for families of infants with suspected athymia. *J Clin Immunol* 2024;44:79.

131. Markert ML, Alexieff MJ, Li J, Sarzotti M, Ozaki DA, Devlin BH, et al. Complete DiGeorge syndrome: development of rash, lymphadenopathy, and oligoclonal T cells in 5 cases. *J Allergy Clin Immunol* 2004;113:734-41.
132. Homans JF, Tromp IN, Colo D, Schlosser TPC, Krut MC, Deeney VFX, et al. Orthopaedic manifestations within the 22q11.2 deletion syndrome: a systematic review. *Am J Med Genet A* 2018;176:2104-20.
133. Pagon RA, Graham JM Jr, Zonana J, Yong SL. Coloboma, congenital heart disease, and choanal atresia with multiple anomalies: CHARGE association. *J Pediatr* 1981;99:223-7.
134. van Ravenswaaij-Arts C, Martin DM. New insights and advances in CHARGE syndrome: diagnosis, etiologies, treatments, and research discoveries. *Am J Med Genet C Semin Med Genet* 2017;175:397-406.
135. Dentici ML, Placidi S, Francalanci P, Capolino R, Rinelli G, Marino B, et al. Association of DiGeorge anomaly and caudal dysplasia sequence in a neonate born to a diabetic mother. *Cardiol Young* 2013;23:14-7.
136. Heimall J, Keller M, Saltzman R, Bunin N, McDonald-McGinn D, Zakai E, et al. Diagnosis of 22q11.2 deletion syndrome and artemis deficiency in two children with T⁺B⁻NK⁺ immunodeficiency. *J Clin Immunol* 2012;32:1141-4.
137. Mensen A, Ochs C, Stroux A, Wittenbecher F, Szyska M, Imberti L, et al. Utilization of TREC and KREC quantification for the monitoring of early T- and B-cell neogenesis in adult patients after allogeneic hematopoietic stem cell transplantation. *J Transl Med* 2013;11:188.
138. Mengrelis K, Kucera F, Shahid N, Watt E, Ross S, Lau CI, et al. T cell phenotype in paediatric heart transplant recipients. *Pediatr Transplant* 2021;25:e13930.
139. Adams SP, Rashid S, Premachandra T, Harvey K, Ifederu A, Wilson MC, et al. Screening of neonatal UK dried blood spots using a duplex TREC screening assay. *J Clin Immunol* 2014;34:323-30.
140. van der Spek J, Groenwold RH, van der Burg M, van Montfrans JM. TREC based newborn screening for severe combined immunodeficiency disease: a systematic review. *J Clin Immunol* 2015;35:416-30.
141. Kreins AY, Graham Davies E. Replacing defective thymus function. *Curr Opin Allergy Clin Immunol* 2020;20:541-8.
142. van der Burg M, Kalina T, Perez-Andres M, Vlkova M, Lopez-Granados E, Blanco E, et al. The EuroFlow PID orientation tube for flow cytometric diagnostic screening of primary immunodeficiencies of the lymphoid system. *Front Immunol* 2019;10:246.
143. Douek DC, Vescio RA, Betts MR, Brenchley JM, Hill BJ, Zhang L, et al. Assessment of thymic output in adults after haematopoietic stem-cell transplantation and prediction of T-cell reconstitution. *Lancet* 2000;355(9218):1875-81.
144. Amrolia PJ, Muccioli-Casadei G, Huls H, Adams S, Duret A, Gee A, et al. Adoptive immunotherapy with allodepleted donor T-cells improves immune reconstitution after haploidentical stem cell transplantation. *Blood* 2006;108:1797-808.
145. Bahamat AA, Assidi M, Lary SA, Almughamsi MM, Peer Zada AA, Chaudhary A, et al. Use of array comparative genomic hybridization for the diagnosis of DiGeorge syndrome in Saudi Arabian population. *Cytogenet Genome Res* 2018;154:20-9.
146. Bahamat AA, Lary SA, Peer Zada AA, Al-Qahtani MH. The evaluation of DiGeorge syndrome gene deletion using molecular cytogenetic techniques. *BMC Genomics* 2014;15(suppl 2).
147. Thaventhiran JED, Lango Allen H, Burren OS, Rae W, Greene D, Staples E, et al. Whole-genome sequencing of a sporadic primary immunodeficiency cohort. *Nature* 2020;583(7814):90-5.
148. Pagnamenta AT, Camps C, Giacomuzzi E, Taylor JM, Hashim M, Calpena E, et al. Structural and non-coding variants increase the diagnostic yield of clinical whole genome sequencing for rare diseases. *Genome Med* 2023;15:94.
149. Meyts I, Bosch B, Bolze A, Boisson B, Itan Y, Belkadi A, et al. Exome and genome sequencing for inborn errors of immunity. *J Allergy Clin Immunol* 2016;138:957-69.
150. Souche E, Beltran S, Brosens E, Belmont JW, Fossum M, Riess O, et al. Recommendations for whole genome sequencing in diagnostics for rare diseases. *Eur J Hum Genet* 2022;30:1017-21.
151. McDonald-McGinn DM, Zackai EH. Genetic counseling for the 22q11.2 deletion. *Dev Disabil Res Rev* 2008;14:69-74.
152. Cirillo E, Cancrini C, Azzari C, Martino S, Martire B, Pession A, et al. Clinical, immunological, and molecular features of typical and atypical severe combined immunodeficiency: report of the Italian Primary Immunodeficiency Network. *Front Immunol* 2019;10:1908.
153. Dvorak CC, Haddad E, Buckley RH, Cowan MJ, Logan B, Griffith LM, et al. The genetic landscape of severe combined immunodeficiency in the United States and Canada in the current era (2010-2018). *J Allergy Clin Immunol* 2019;143:405-7.
154. Fukushima T, Suzuki S, Ohira T, Shimura H, Midorikawa S, Ohtsuru A, et al. Prevalence of ectopic intrathyroidal thymus in Japan: the Fukushima health management survey. *Thyroid* 2015;25:534-7.
155. Zweier C, Sticht H, Aydin-Yaylagul I, Campbell CE, Rauch A. Human *TBX1* missense mutations cause gain of function resulting in the same phenotype as 22q11.2 deletions. *Am J Hum Genet* 2007;80:510-7.
156. Bosticardo M, Pala F, Calzoni E, Delmonte OM, Dobbs K, Gardner CL, et al. Artificial thymic organoids represent a reliable tool to study T-cell differentiation in patients with severe T-cell lymphopenia. *Blood Adv* 2020;4:2611-6.
157. Bifsha P, Leiding JW, Pai SY, Colamartino ABL, Hartog N, Church JA, et al. Diagnostic assay to assist clinical decisions for unclassified severe combined immune deficiency. *Blood Adv* 2020;4:2606-10.
158. Six EM, Benjelloun F, Garrigue A, Bonhomme D, Morillon E, Rouiller J, et al. Cytokines and culture medium have a major impact on human *in vitro* T-cell differentiation. *Blood Cells Mol Dis* 2011;47:72-8.
159. Seet CS, He C, Bethune MT, Li S, Chick B, Gschwend EH, et al. Generation of mature T cells from human hematopoietic stem and progenitor cells in artificial thymic organoids. *Nat Methods* 2017;14:521-30.
160. Iancu O, Allen D, Knop O, Zehavi Y, Breier D, Arviv A, et al. Multiplex HDR for disease and correction modeling of SCID by CRISPR genome editing in human HSPCs. *Mol Ther Nucleic Acids* 2023;31:105-21.
161. Golwala Z, Devi Moirangthem R, Evans G, Lizot S, de Koning C, Garrigue A, et al. *Ex vivo* T-lymphopoiesis assays assisting corrective treatment choice for genetically undefined T-lymphocytopenia. Preprint, <https://doi.org/10.21203/rs.3.rs-4668549/v1>
162. Marcovecchio GE, Bortolomai I, Ferrua F, Fontana E, Imberti L, Conforti E, et al. Thymic epithelium abnormalities in DiGeorge and Down syndrome patients contribute to dysregulation in T cell development. *Front Immunol* 2019;10:447.
163. Gennery AR, Barge D, O'Sullivan JJ, Flood TJ, Abinun M, Cant AJ. Antibody deficiency and autoimmunity in 22q11.2 deletion syndrome. *Arch Dis Child* 2002;86:422-5.
164. Patel K, Akhter J, Kobrynski L, Benjamin Gathmann MA, Davis O, Sullivan KE, et al. Immunoglobulin deficiencies: the B-lymphocyte side of DiGeorge syndrome. *J Pediatr* 2012;161:950-3.
165. Morsheimer M, Brown Whitehorn TF, Heimall J, Sullivan KE. The immune deficiency of chromosome 22q11.2 deletion syndrome. *Am J Med Genet A* 2017;173:2366-72.
166. Bassett AS, McDonald-McGinn DM, Devriendt K, Digilio MC, Goldenberg P, Habel A, et al. Practical guidelines for managing patients with 22q11.2 deletion syndrome. *J Pediatr* 2011;159:332-9.e1.
167. Fung WL, Butcher NJ, Costain G, Andrade DM, Boot E, Chow EW, et al. Practical guidelines for managing adults with 22q11.2 deletion syndrome. *Genet Med* 2015;17:599-609.
168. Thakar MS, Logan BR, Puck JM, Dunn EA, Buckley RH, Cowan MJ, et al. Measuring the effect of newborn screening on survival after haematopoietic cell transplantation for severe combined immunodeficiency: a 36-year longitudinal study from the Primary Immune Deficiency Treatment Consortium. *Lancet* 2023;402(10396):129-40.
169. Dorsey MJ, Wright NAM, Chaimowitz NS, Davila Saldana BJ, Miller H, Keller MD, et al. Infections in infants with SCID: isolation, infection screening, and prophylaxis in PIDTC centers. *J Clin Immunol* 2021;41:38-50.
170. Hicks ED, Agada NO, Yates TR, Kelly MS, Tam JS, Ferdman RM, et al. Case report: nontuberculous mycobacterial infections in children with complete DiGeorge anomaly. *Front Immunol* 2023;14:1078976.
171. Buggins AG, Mufti GJ, Salisbury J, Codd J, Westwood N, Arno M, et al. Peripheral blood but not tissue dendritic cells express CD52 and are depleted by treatment with alemtuzumab. *Blood* 2002;100:1715-20.
172. Gupton SE, McCarthy EA, Markert ML. Care of children with DiGeorge before and after cultured thymus tissue implantation. *J Clin Immunol* 2021;41:896-905.
173. Great Ormond Street Hospital (GOSH) International and Private Care. Thymus transplants. Available at: <https://www.gosh.com/kw/thymus-transplants>.
174. Fitch ZW, Kang L, Li J, Knechtle SJ, Turek JW, Kirk AD, et al. Introducing thymus for promoting transplantation tolerance. *J Allergy Clin Immunol* 2022;150:549-56.
175. Kreins AY, Junghanns F, Mifsud W, Somana K, Sebire N, Rampling D, et al. Correction of both immunodeficiency and hypoparathyroidism by thymus transplantation in complete DiGeorge syndrome. *Am J Transplant* 2020;20:1447-50.
176. Hale LP, Neff J, Cheatham L, Cardona D, Markert ML, Kurtzberg J. Histopathologic assessment of cultured human thymus. *PLoS One* 2020;15:e0230668.
177. Markert ML, Sarzotti M, Ozaki DA, Sempowski GD, Rhein ME, Hale LP, et al. Thymus transplantation in complete DiGeorge syndrome: immunologic and safety evaluations in 12 patients. *Blood* 2003;102:1121-30.
178. Olsen NJ, Kovacs WJ. Evidence that androgens modulate human thymic T cell output. *J Invest Med* 2011;59:32-5.
179. Taves MD, Ashwell JD. Effects of sex steroids on thymic epithelium and thymocyte development. *Front Immunol* 2022;13:975858.

180. Markert ML, Li J, Devlin BH, Hoehner JC, Rice HE, Skinner MA, et al. Use of allograft biopsies to assess thymopoiesis after thymus transplantation. *J Immunol* 2008;180:6354-64.
181. Paris K, Wall LA. The treatment of primary immune deficiencies: lessons learned and future opportunities. *Clin Rev Allergy Immunol* 2023;65:19-30.
182. Ip W, Zhan H, Gilmour KC, Davies EG, Qasim W. 22q11.2 deletion syndrome with life-threatening adenovirus infection. *J Pediatr* 2013;163:908-10.
183. Kreins AY, Roux E, Pang J, Cheng I, Charles O, Roy S, et al. Favipiravir induces HuNoV viral mutagenesis and infectivity loss with clinical improvement in immunocompromised patients. *Clin Immunol* 2024;259:109901.
184. Bekegnran CP, Driouich JS, Breuer J, Barthelemy K, Giocanti M, de Lamballerie X, et al. Simultaneous quantitation of favipiravir and its hydroxide metabolite in human plasma and hamster matrices using a UPLC-MS/MS method. *Biomed Chromatogr* 2023;37:e5689.
185. Volpi S, Yamazaki Y, Brauer PM, van Rooijen E, Hayashida A, Slavotinek A, et al. *EXTL3* mutations cause skeletal dysplasia, immune deficiency, and developmental delay. *J Exp Med* 2017;214:623-37.
186. Gardner CL, Pavel-Dinu M, Dobbs K, Bosticardo M, Reardon PK, Lack J, et al. Gene editing rescues *in vitro* T cell development of RAG2-deficient induced pluripotent stem cells in an artificial thymic organoid system. *J Clin Immunol* 2021;41:852-62.
187. Chhatta AR, Cordes M, Hanegraaf MAJ, Vloemans S, Cupedo T, Cornelissen JJ, et al. *De novo* generation of a functional human thymus from induced pluripotent stem cells. *J Allergy Clin Immunol* 2019;144:1416-9.e7.
188. Ramos SA, Morton JJ, Yadav P, Reed B, Alizadeh SI, Shilleh AH, et al. Generation of functional human thymic cells from induced pluripotent stem cells. *J Allergy Clin Immunol* 2022;149:767-81.e6.
189. Ramos SA, Armitage LH, Morton JJ, Alzofon N, Handler D, Kelly G, et al. Generation of functional thymic organoids from human pluripotent stem cells. *Stem Cell Reports* 2023;18:829-40.
190. Karvonen E, Krohn KJE, Ranki A, Hau A. Generation and characterization of iPS cells derived from APECED patients for gene correction. *Front Endocrinol (Lausanne)* 2022;13:794327.
191. Markert ML, Devlin BH, Chinn IK, McCarthy EA, Li YJ. Factors affecting success of thymus transplantation for complete DiGeorge anomaly. *Am J Transplant* 2008;8:1729-36.
192. Ross S, Cheung M, Lau CI, Sebire N, Burch M, Kilbride P, et al. Transplanted human thymus slices induce and support T-cell development in mice after cryopreservation. *Eur J Immunol* 2018;48:716-9.
193. Chawda MM, Ross S, Lau CI, Yanez DC, Rowell J, Kilbride P, et al. Cryopreservation of mouse thymus depletes thymocytes but supports immune reconstitution on transplantation. *Eur J Immunol* 2023;53:e2350546.