



Universiteit
Leiden
The Netherlands

Unraveling the drivers of antimicrobial pharmacokinetic variability in individuals with obesity and hospitalized patients with multimorbidity

Rhee, K.P. van

Citation

Rhee, K. P. van. (2025, February 19). *Unraveling the drivers of antimicrobial pharmacokinetic variability in individuals with obesity and hospitalized patients with multimorbidity*. Retrieved from <https://hdl.handle.net/1887/4195405>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4195405>

Note: To cite this publication please use the final published version (if applicable).

Part I

CHAPTER 1



Introduction



Infectious diseases are common among hospitalized patients as 1 in 3 patients receives treatment with at least one antimicrobial agent¹. Efficacious treatment of infectious diseases hinges on prompt administration of the right dose of the right drug. In the clinical setting pharmacological therapy is initiated immediately when a (severe) infectious disease is suspected. Selecting the right empirical antimicrobial drug is done by choosing an agent that covers the most likely pathogens given the local epidemiology. After the most appropriate drug is selected the right dose needs to be determined. But how does one determine the right dose for each patient? To determine the optimal dose, knowledge on both pharmacokinetics and pharmacodynamics are needed.

The studies described in this thesis aim at elucidating whether and how various patient characteristics influence concentrations of antimicrobial agents in plasma and how variation in these concentrations translate into effect. In this regard, focus lies on elucidating the influence of (morbid) obesity, (morbid) obesity & critical illness and mucosal barrier injury on drug concentration in plasma. The drugs of interest are the antifungal agent fluconazole and the antibacterial agent ciprofloxacin. For both drugs plasma concentrations are known to correlate with clinical- and microbiological cure and both antimicrobials are widely used in the hospital and community. At present it remains unknown how the pharmacokinetics of these two drugs are influenced by morbid obesity and multimorbidity. As a result, clear guidance for dose adjustment in these special patient populations is unavailable.

Microbes and antimicrobial agents – Fluconazole and Ciprofloxacin

In hospitalized patients infections of the respiratory-, urinary-, and gastro-intestinal tract are frequently encountered and may be caused by a wide range of pathogens.

These causes of infectious diseases remained unknown until Louis Pasteur disproved the spontaneous generation theory of spoilage and fermentation in 1861 when he revealed that invisible microorganisms were the cause². In 1881 this work was followed up by a number of papers describing the germ theory of disease which is the currently accepted scientific theory for infectious diseases³. In the meantime, the Norwegian physician Gerhard Armauer Hansen was the first to discover a bacterial cause for an infectious disease, leprosy, when he identified *Mycobacterium leprae* in 1873. The German microbiologist Robert Koch subsequently discovered *Bacillus anthracis* and *Mycobacterium tuberculosis* in 1882 and *Vibrio cholerae* in 1884, which were major causes of infectious diseases at the time⁴. Although the cause of infectious diseases was slowly unraveled, efficacious treatment was impossible because antimicrobial agents were not yet discovered.

The first antimicrobial agent was the arsenic based antibacterial agent arsphenamide which was discovered by Paul Ehrlich and Sahachiro Hata in 1909⁴. Various classes of antibacterial agents have been discovered ever since. The fluoroquinolones are a class of antibacterial agents derived from nalidixic acid which was discovered by George Leshner during an attempt to synthesize chloroquine in 1962. The second-generation fluoroquinolone ciprofloxacin was licensed in 1987⁵. It inhibits bacterial DNA synthesis by blocking DNA-gyrase and topo-isomerase IV which both play a vital role in folding, unfolding, stretching and relaxing of bacterial DNA. Blocking these enzymes ultimately kills the bacterium. Ciprofloxacin is a broad spectrum antibiotic and is mostly used for the treatment of aerobic gram-negative bacteria, especially *Enterobacteriales*⁶.

The first antifungal agent is nystatin which was licensed in 1954. Since then, similar to antibacterial agents, various classes of antifungals reached the market. Fluconazole is a member of the triazoles and was first licensed in 1990⁷. It induces structural deformation of the fungus by blocking ergosterol synthesis through CYP51 inhibition. Ergosterol is a sterol vital for maintaining cellular integrity of various fungi; its physiological function is similar to cholesterol in mammalian cells. Fluconazole has a fungistatic effect with a broad antimycotic spectrum and is mainly employed for the treatment of *Candida* and *Cryptococcal* infections⁸.

Pharmacokinetics, -dynamics and -metrics

The rationale for dosing antimicrobial agents is to aim for exposure that is high enough to treat the causative pathogen while preventing overexposure and thus minimize toxicity⁹.

The field of pharmacokinetics studies how drugs move through the body over time. Four processes can be distinguished: absorption, distribution, metabolism and elimination. Absorption of drugs after oral administration takes place in the gastro-intestinal tract and may be influenced by concomitant food or drug intake or mucosal barrier injury as was demonstrated for posaconazole¹⁰. Distribution of drugs depends on chemical properties like lipophilicity, ionization, protein binding and blood to plasma ratio¹¹. Drug distribution may be influenced by the presence of excess adipose tissue which negatively influences the penetration of ciprofloxacin, cefazoline and metronidazole into skin and soft tissue while meningitis increases the penetration of beta-lactam antibiotics into the central nervous system¹²⁻¹⁵. Similarly, patients admitted to the intensive care can be hemodynamically unstable. This requires aggressive fluid resuscitation in order to maintain the blood pressure within acceptable limits which may also influence drug distribution¹⁶. Metabolism describes the process of chemical reactions altering the molecular structure of a drug which mainly takes place in the liver. Drug metabolism can be inhibited or enhanced as a result of altered enzymatic activity for instance due

to pathophysiological alterations or drug-drug interactions¹⁷. Elimination describes how the drug and/or its metabolites are removed from the body by for instance renal clearance or intestinal excretion. The rate at which drugs are cleared from plasma may be influenced by patient characteristics like body weight, renal function, hepatic function or (critical) illness^{11, 16–18}.

Pharmacodynamics describes the balance between (drug) concentration and effect, for instance the killing of bacteria when an antibiotic is present or the increase in heart rate when adrenaline is released into the bloodstream. The likelihood of a patient being cured using a certain dosing regimen depends on both the exposure to the antimicrobial agent, the susceptibility of the causative pathogen towards the employed antimicrobial agent and the capacity of the host immune system. The susceptibility is expressed as the minimal inhibitory concentration (MIC) i.e. the minimal drug concentration at which in vitro pathogen growth is inhibited within a defined period of time⁹.

The MIC is commonly correlated to an exposure target that is predictive for clinical or microbiological cure. Three different exposure/response-relations, often referred to as pharmacokinetic/ pharmacodynamic (PK/PD) targets, are distinguished for the prediction of clinical outcome 1) The percentage of time within a 24 hour period the concentration is above the MIC ($T_{>MIC}$) or time dependent antimicrobials 2) The ratio of the peak plasma concentration and the MIC (C_{max}/MIC) or concentration-dependent antimicrobials 3) The ratio of the area under the concentration-time curve (AUC) and the MIC (AUC/MIC) or concentration and time-dependent antimicrobials⁹.

Which exposure/response relation is best associated with clinical outcome may be studied in animal, in-vitro or patient studies. Based on this relation a PK/PD target is formulated that takes both exposure and pathogen susceptibility into account. This PK/PD target can ultimately be used to design dosing regimens for patient populations that have altered absorption, clearance or volume of distribution. For fluconazole an unbound (*f*) fluconazole $fAUC_{24h}/MIC > 100$ is associated with clinical outcome for the treatment of invasive candidiasis¹⁹. For ciprofloxacin an $AUC/MIC > 125$ is associated with probabilities of microbiological- and clinical cure of 86% and 81%, respectively²⁰. Even though it remains unknown if these PK/PD targets are readily applicable to individuals with obesity or hospitalized patients with multimorbidity we propose exposure matching as a strategy. Doses are adapted in a special patient population of interest, aiming at an exposure that is similar to a reference population. This strategy has been adopted for tobramycin, gentamycin and vancomycin in individuals with obesity and even critically ill individuals with obesity in case of gentamicin and vancomycin^{21–25}.

Although many antimicrobial agents generally have favorable toxicity profiles, the above described PK/PD target rationale provides a lower-limit of exposure in order to optimize efficacy. Nevertheless, potential toxicity should also be taken into account when designing dosing strategies. Exposure thresholds (expressed as AUC or trough-concentration) may correlate with the development of toxicity and can be considered upper-limits of safety and should also be taken into account.

Pharmacometrics is the interdisciplinary field at the intersection of pharmacology, mathematics, statistics, and computer science, focused on quantifying and optimizing drug therapy through modelling approaches. It is an established tool for analyzing population pharmacokinetic data as it utilizes mathematical models to describe the pharmacokinetic processes within the human body. A population pharmacokinetic analysis is required for new drug applications to quantify pharmacokinetic variability within a population^{26–28}.

A pharmacokinetic model consists of algebraic equations for parameters such as bioavailability, volume of distribution and clearance while accounting for different sources of random variability (interindividual and residual variability). The main aim is to build a model that both describes and predicts the observations made in the study population, which could be focused on pharmacokinetic endpoints describing drug concentration in plasma or tissue over time or pharmacodynamic endpoints like killing of bacteria or fungi. During the modelling process, patient characteristics of interest (e.g. body weight, renal function, hepatic functioning, age or gender) are tested for associations with model parameters. A stepwise approach with regard to testing patient characteristics allows for discrimination between, and characterization of, the influence of for instance renal function on clearance and body weight on clearance.

If the pharmacokinetic model is improved using certain patient characteristics, these can subsequently be used to guide dosing for individual patients. To this end, simulations with the final pharmacokinetic model are made for (large numbers of) virtual patients with a distribution in the patient characteristics that matches the patient characteristics of the studied population. Using this approach, various dosing regimens can be evaluated and those that result in an acceptable exposure, i.e. that provide sufficient likelihood of therapeutic success while assuring the likelihood of toxicity stays within acceptable limits, can be employed in patient care.

In short, pharmacometric approaches can be used to determine if dosing regimens need to be adapted to specific patient characteristics by integrating data and information from clinical trials, preclinical experiments, and real-world patients. Pharmacometricians develop models to predict exposure in diverse patient populations. These models


enable the design of optimal dosing strategies, especially in special patient populations like children, elderly, individuals with obesity, renal- or hepatic impairment or other (multi) morbid pathophysiological conditions.

Obesity and multimorbidity – implications for optimal dosing regimens

Obesity is characterized by a disproportionate body weight for height with an excessive accumulation of adipose tissue that is usually accompanied by mild systemic inflammation²⁹. The global obesity prevalence reached the questionable milestone of 1 billion people in 2022 (~13% of the global population). It's prevalence tripled since 1975 for women while it quadrupled over the same period for men^{28, 29}. Although the prevalence increases globally, large regional differences occur as is presented in figure 1.

Individuals with obesity are more likely to develop comorbidities like hypertension, diabetes, coronary artery disease, heart failure, atrial fibrillation, various types of cancer, stroke and infectious diseases²⁹. Overweight or obesity may be quantified using the body mass index (BMI) which is calculated by dividing body weight in kilograms by the square of the height in meters. A BMI 25-30kg/m² is considered overweight (or Class 1 obesity) while a BMI 30-35 kg/m² is considered moderate-risk obesity (Class 2). The high-risk group are the individuals with a BMI >40kg/m², or >35kg/m² in the presence of obesity-related comorbidities like diabetes mellitus type II, hypertension, cardiorespiratory conditions (Class 3)³¹. Although this classification only goes up to class 3, obesity may even be viewed on a continuous scale as BMI's up to, and even beyond, 70kg/m² may be encountered.

The increased fat mass in individuals with obesity may have implications for selecting the optimal dosing regimen. Increased fat mass may influence pharmacokinetics, especially when these patients also have other comorbidities like reduced renal or hepatic functioning or critical illness which may independently influence metabolism and elimination^{16, 24, 25}. Nevertheless, multimorbidity may also be encountered in normal weight individuals, for instance in patients with hematological malignancies with mucosal barrier injury which could influence the absorption of orally administered drugs¹⁰.

Figure 1

The global prevalence of obesity (BMI >30kg/m²) combined for men and women shows large regional differences. The figure is based on data from 2022 from the NCD Risk Factor Collaboration³⁰.

To determine the optimal dose for a specific patient (population) it is vital to capture the influence of the specific health condition(s) on the pharmacokinetic process of interest^{27, 28}. To guide dosing in individuals with obesity, patients undergoing bariatric surgery are often studied to determine the influence of body weight on pharmacokinetics. An important strength of studying this population is that it allows dense sampling as these patients are already managed with an intravenous line and study procedures can be planned upfront. Moreover individuals with class 3 obesity are frequently encountered in the bariatric surgery population while patients with extreme body weights are relatively scarce among the clinical population. Collecting PK-data in individuals with these extreme body weights allows for precise characterization of the influence of body weight on pharmacokinetic parameters. Bariatric patients are typically otherwise healthy, have good renal function and are often relatively young. Carrying out pharmacokinetics studies in this population prevents any influence of for instance renal replacement therapy which may be encountered among hospitalized patients with obesity and infectious diseases. A theoretical drawback to using bariatric surgery patients to test the influence of obesity on pharmacokinetics is the fact that these patients require surgery. However, as blood loss is minimal (usually <100ml) during these short laparoscopic interventions (± 45 minutes), and blood pressure is tightly

monitored during the procedure, the influence of the surgery itself on pharmacokinetics is considered minimal.

Generalization of results from bariatric surgery patients to hospitalized patients with obesity and multimorbidity may be complicated by differences in the presence or distribution of covariates¹⁶. To ensure applicability of results found in bariatric surgery patients with class 3 obesity to the clinical population of patients with obesity, the model may be enriched with data from obese individuals with multimorbidity that are representative for the clinical population. The results from bariatric surgery patients can be used as a starting point when testing the influence of other patient characteristics on pharmacokinetic parameters based on the data from the clinical population. This approach facilitates assessment of the influence of other potentially relevant covariates like renal function, and admission to the ward or ICU^{24, 25}.

Aims and scope of this thesis

In this thesis we aim to characterize the influence of obesity on fluconazole and ciprofloxacin pharmacokinetics. Guidance on whether, and how, doses should be adjusted in this population is currently lacking. Ultimately, the results for these two antimicrobial agents are viewed in a broader context comparing our results with previous studies on the influence of body weight on the pharmacokinetics of other antimicrobial agents. Ciprofloxacin pharmacokinetics was subsequently studied in two other subpopulations with multimorbidity, i.e. patients with mucosal barrier injury secondary to the treatment of a hematological malignancy and patients with obesity admitted to the intensive care unit. As for both mucosal barrier injury and ICU admission, the influence on pharmacokinetic parameters is unknown it remains unclear if applying the standard dosing regimens result in sufficient exposure. Within the studies described in this thesis guidance for both fluconazole and ciprofloxacin were designed based on the identified drivers of pharmacokinetic variability.

Results from studies in individuals with (morbid) obesity undergoing bariatric surgery and in non-obese healthy volunteers are described in the first part of this thesis. The influence of body weight on the pharmacokinetics of fluconazole and ciprofloxacin was characterized in two rich sampling studies after both oral and intravenous administration. The results for fluconazole are presented in **Chapter 2**. The results for ciprofloxacin are presented in **Chapter 3**. Individuals with obesity are not currently recognized as a special patient population during drug development. Our view on the clinical implications resulting from the considerable heterogeneity in the influence of weight on clearance between drugs and the importance of taking the labeled dosing strategy (fixed dose versus mg/kg body weight) into account are described in **Chapter 4**.

Studies in hospitalized patients with multimorbidity are described in the second part of this thesis. A cohort of patients with (morbid) obesity admitted to the intensive care unit treated with ciprofloxacin was recruited to determine if additional patient characteristics like renal dysfunction may be needed to predict, and optimize, exposure to ciprofloxacin in the real-world setting of hospitalized individuals with obesity; results are presented in **Chapter 5**.

Besides empirical treatment, ciprofloxacin as oral formulation is often used for gram-negative infection prophylaxis during neutropenia and gastrointestinal mucositis in patients with hematological malignancies. We explored if bioavailability or clearance in this population is affected by mucosal barrier injury which may develop as a complication after preconditioning. Besides studying exposure, we explored whether biomarkers can be used to predict potential differences. Results of this study are presented in **Chapter 6**.

Finally, we summarize and discuss (clinical) implications of our findings, reflect on lessons learned and provide recommendations for future research in **Chapter 7**. Using simulations, we provide a framework for how the recommended future research may be shaped.

References

- 1 Versporten A, Zarb P, Caniaux I, Gros MF, Drapier N, Miller M, Jarlier V, Nathwani D, Goossens H; Global-PPS network. Antimicrobial consumption and resistance in adult hospital inpatients in 53 countries: results of an internet-based global point prevalence survey. *Lancet Glob Health*. 2018 Jun;6(6):e619-e629. Erratum in: *Lancet Glob Health*. 2018 Sep;6(9):e968.
- 2 PORTER JR. Louis PASTEUR; achievements and disappointments, 1861. *Bacteriol Rev*. 1961 Dec;25(4):389-403.
- 3 Pasteur. On the germ theory. *Science*. 1881 Sep 3;2(63):420-2.
- 4 Aminov RI. A brief history of the antibiotic era: lessons learned and challenges for the future. *Front Microbiol*. 2010 Dec 8;1:134.
- 5 Emmerson AM, Jones AM. The quinolones: decades of development and use. *J Antimicrob Chemother*. 2003 May;51 Suppl 1:13-20.
- 6 Ciprofloxacin access data. Food and Drug Administration, 30/7/2016. Accessed 4/3/2024 https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/019537s086lbl.pdf
7. Vanreppelen G, Wuyts J, Van Dijck P, Vandecruys P. Sources of Antifungal Drugs. *J Fungi (Basel)*. 2023 Jan 28;9(2):171.
- 8 Fluconazole access data. Food and drug administration, 28/02/2019. Accessed 4/3/2024 https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/019949s065,020090s047lbl.pdf
- 9 Johan W. Mouton, Michael N. Dudley, Otto Cars, Hartmut Derendorf, George L. Drusano, Standardization of pharmacokinetic/pharmacodynamic (PK/PD) terminology for anti-infective drugs: an update, *Journal of Antimicrobial Chemotherapy*, Volume 55, Issue 5, May 2005, Pages 601–607,
- 10 Jansen AME, Muilwijk EW, van der Velden WJFM, Maertens JA, Aerts R, Colbers A, Burger D, Verweij PE, Ter Heine R, Blijlevens NMA, Brüggemann RJM. Posaconazole bioavailability of the solid oral tablet is reduced during severe intestinal mucositis. *Clin Microbiol Infect*. 2022 Jul;28(7):1003-1009.
- 11 Smit C, De Hoogd S, Brüggemann RJM, Knibbe CAJ. Obesity and drug pharmacology: a review of the influence of obesity on pharmacokinetic and pharmacodynamic parameters. *Expert Opin Drug Metab Toxicol*. 2018 Mar;14(3):275-285.
- 12 van Rhee KP, Smit C, Wasmann RE, et al. Ciprofloxacin pharmacokinetics after oral and intravenous administration in (morbidly) obese and non-obese individuals; a prospective clinical study. *Clin Pharmacokinet*. 2022;61:1167–75.
- 13 Brill MJE, Houwink API, Schmidt S, et al. Reduced subcutaneous tissue distribution of ceftazidime in morbidly obese versus non-obese patients determined using clinical microdialysis. *J Antimicrob Chemother*. 2014;69:715–23.
- 14 Dorn C, Petroff D, Stoelzel M, et al. Perioperative administration of ceftazidime and metronidazole in obese and non-obese patients: a pharmacokinetic study in plasma and interstitial fluid. *J Antimicrob Chemother*. 2021;76:2114–20.
- 15 Jager NGL, van Hest RM, Lipman J, Roberts JA, Cotta MO. Antibiotic exposure at the site of infection: principles and assessment of tissue penetration. *Expert Rev Clin Pharmacol*. 2019 Jul;12(7):623-634.
- 16 Morales Castro D, Dresser L, Granton J, Fan E. Pharmacokinetic Alterations Associated with Critical Illness. *Clin Pharmacokinet*. 2023 Feb;62(2):209-220. doi: 10.1007/s40262-023-01213-x. Epub 2023 Feb 2.

- 17 Ogu CC, Maxa JL. Drug interactions due to cytochrome P450. *Proc (Bayl Univ Med Cent)*. 2000 Oct;13(4):421-3.
- 18 Masereeuw R, Russel FG. Mechanisms and clinical implications of renal drug excretion. *Drug Metab Rev*. 2001 Aug-Nov;33(3-4):299-351.
- 19 Rodríguez-Tudela JL, Almirante B, Rodríguez-Pardo D, Laguna F, Donnelly JP, Mouton JW, Pahissa A, Cuenca-Estrella M. Correlation of the MIC and dose/MIC ratio of fluconazole to the therapeutic response of patients with mucosal candidiasis and candidemia. *Antimicrob Agents Chemother*. 2007 Oct;51(10):3599-604.
- 20 Forrest A, Nix DE, Ballow CH, Goss TF, Birmingham MC, Schentag JJ. Pharmacodynamics of intravenous ciprofloxacin in seriously ill patients. *Antimicrob Agents Chemother*. 1993 May;37(5):1073-81.
- 21 Smit C, Wasmann RE, Wiezer MJ, van Dongen HPA, Mouton JW, Brüggemann RJM, Knibbe CAJ. Tobramycin Clearance Is Best Described by Renal Function Estimates in Obese and Non-obese Individuals: Results of a Prospective Rich Sampling Pharmacokinetic Study. *Pharm Res*. 2019 May 30;36(8):112. Erratum in: *Pharm Res*. 2020 Jan 9;37(2):32.
- 22 Smit C, Wasmann RE, Gouloze SC, Hazebroek EJ, Van Dongen EPA, Burgers DMT, Mouton JW, Brüggemann RJM, Knibbe CAJ. A Prospective Clinical Study Characterizing the Influence of Morbid Obesity on the Pharmacokinetics of Gentamicin: Towards Individualized Dosing in Obese Patients. *Clin Pharmacokinet*. 2019 Oct;58(10):1333-1343.
- 23 Smit C, Wasmann RE, Gouloze SC, Wiezer MJ, van Dongen EPA, Mouton JW, Brüggemann RJM, Knibbe CAJ. Population pharmacokinetics of vancomycin in obesity: Finding the optimal dose for (morbidly) obese individuals. *Br J Clin Pharmacol*. 2020 Feb;86(2):303-317.
- 24 Smit C, van Schip AM, van Dongen EPA, Brüggemann RJM, Becker ML, Knibbe CAJ. Dose recommendations for gentamicin in the real-world obese population with varying body weight and renal (dys)function. *J Antimicrob Chemother*. 2020 Nov 1;75(11):3286-3292.
- 25 Zhang T, Krekels EHJ, Smit C, van Dongen EPA, Brüggemann RJM, Knibbe CAJ. How to Dose Vancomycin in Overweight and Obese Patients with Varying Renal (Dys)function in the Novel Era of AUC 400-600 mg·h/L-Targeted Dosing. *Clin Pharmacokinet*. 2024 Jan;63(1):79-91.
- 26 Population pharmacokinetics, Guidance for Industry. Food and Drug administration, 28/2/2022. Accessed 6/3/2024 <https://www.fda.gov/media/128793/download>
- 27 Pharmacokinetics in patients with impaired renal function – study design, data analysis, and impact on dosing, Guidance for industry. Food an Drug Administration, 3/2024. Accessed 22/3/2024. <https://www.fda.gov/media/78573/download>
- 28 Reflection paper on investigation of pharmacokinetics in the obese population. European medicines agency. 4/12/2023. Accessed 6/3/2024 Microsoft Word - 04.12.2023 Reflection paper on pk investigations in obese population (europa.eu)
- 29 González-Muniesa P, Martínez-González MA, Hu FB, Després JP, Matsuzawa Y, Loos RJF, Moreno LA, Bray GA, Martínez JA. Obesity. *Nat Rev Dis Primers*. 2017 Jun 15;3:17034.
- 30 NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222 million children, adolescents, and adults. *Lancet*. 2024 Feb 29;S0140-6736(23)02750-2.
- 31 Overweight & Obesity. Centers for Disease Control and Prevention (CDC), 21/9/2023. Accessed 6/3/2024 <https://www.cdc.gov/obesity/index.html>