

Targeted interventions in mechanically ventilated patients Wal, L.I. van der

Citation

Wal, L. I. van der. (2025, April 4). *Targeted interventions in mechanically ventilated patients*. Retrieved from https://hdl.handle.net/1887/4210561

Version: Publisher's Version

Licence agreement concerning inclusion of doctoral

License: thesis in the Institutional Repository of the University

of Leiden

Downloaded from: https://hdl.handle.net/1887/4210561

Note: To cite this publication please use the final published version (if applicable).



Perspectives of ICU Patients on Deferred Consent in the Context of Post-ICU Quality of Life - a substudy of a randomized clinical trial

L. Imeen van der Wal, Chloe C.A. Grim, Michael R. del Prado, David J. van Westerloo, Marcus J. Schultz, Hendrik J.F. Helmerhorst Martine C. de Vries, Evert de Jonge, for the ICONIC investigators.

Critical Care Medicine, 2024 May 1;52(5):694-703.

Published online 2024 Jan 5.

Supplemental digital content of this article is available online

ABSTRACT

Objectives

Deferred consent enables research to be conducted in the intensive care unit (ICU) when patients are unable to provide consent themselves, and there is insufficient time to obtain consent from surrogates before commencing (trial) treatment. The aim of this study was to evaluate how former ICU patients reflect on their participation in a study with deferred consent and examine whether their opinions are influenced by the quality of life (QoL) following hospital discharge.

Design

Survey study by questionnaire

Setting

Eight ICUs in the Netherlands

Patients

Former ICU patients that participated in the ICONIC trial, a multicenter randomized clinical trial that evaluated oxygenation targets in mechanically ventilated ICU patients.

Interventions

Participants enrolled in the ICONIC trial in one of the eight participating centers in the Netherlands received a questionnaire six months after randomization. The questionnaire included 12 close-ended questions on their opinion about the deferred consent procedure. QoL was measured using the EQ-5D-5L questionnaire. By calculating the EQ-5Dindex, patients were divided in 4 QoL quartiles, where Q1 reflects the lowest and Q4 the highest.

Measurements and main results

Of 362 participants who were contacted, 197 responded (54%). More than half of the respondents (59%) were unaware of their they participation in the ICONIC study. In total 61% was content with the deferred consent procedure, 1% not content, 25% neutral, 9% did not know and 9% answered "other". Those with a higher QoL were more likely to be content (P=0.02). In all QoL groups the legal representative was the most often preferred individual to provide consent.

Conclusion

Former ICU patients who participated in the ICONIC study often did not remember their participation but were predominantly positive regarding the use of deferred consent. Those with a higher QoL were most likely to be content.

INTRODUCTION

Informed consent is an ethical cornerstone of medical research (2). In the Intensive Care Unit (ICU), however, patients are often unable to provide informed consent due to their critical condition (3). An alternative would be to ask a proxy or other legal representative for consent, although clinical practice often shows that a representative is either not available or overwhelmed by the situation and therefore not able to make a well-considered decision in the narrow time window of inclusion (3, 4). For such cases, deferred consent procedures have been developed in which patients can participate in medical research before obtaining informed consent under the condition that consent is sought from the subject or their legal representative as soon as circumstances allow it (5). Ethical concerns, however, have been raised due to the fact that patients cannot express their preferences in real-time, possibly impacting their autonomy.

In recent years, the use of deferred consent procedures in clinical studies has increased considerably. Some studies have demonstrated the feasibility and acceptability of deferred consent in the ICU setting (6). In a small study from Finland, 9 of 11 patients who had survived after participating in a study on therapeutic hypothermia after cardiac arrest agreed to research in emergency settings without consent of the patient or proxy (7). Nearly all ICU patients that participated in the NICE-Sugar study would have consented to study participation if asked for consent before enrollment (8).

An important factor that may influence patients' opinion on deferred consent procedures is their overall quality of life (QoL). While QoL can be seriously impaired after ICU stay (9), the influence of QoL on patients' opinion of deferred consent has not been evaluated. A prior study evaluating patients perspectives on Exception from Informed Consent (EFIC) in the "Progesterone for the treatment of Traumatic Brain Injury" (ProTECT III) trial found that the acceptance of the use of EFIC was generally high, however, patients with unfavorable outcomes were less accepting of their EFIC inclusion compared to those with favorable outcomes (10, 11). The ICONIC study (1, 12), a multicenter randomized controlled trial in ICU patients, comparing two oxygenation targets, allowed inclusion without prior consent and provided a population to evaluate this question. The aim of this substudy was to evaluate patients' perspectives on deferred consent and explore the influence of QoL. We hypothesized that patients with an impaired QoL after ICU are less likely to accept participating in studies without prior consent.

MATERIALS AND METHODS

Participants and setting

This is a substudy of the ICONIC trial (12), an international, multicenter, randomized, parallel-group trial, in which 664 patients were enrolled between November 2018 and November 2021. In addition, 125 patients were initially enrolled with deferred consent but subsequently excluded because consent was declined by the patient or his/her representative (appendix 1). The original trial was conducted in 8 ICUs in the Netherlands and 1 ICU in Italy. Ethical approval was granted on October 25 2018 for all centres by the Medical Ethical Committee of Leiden, The Hague and Delft (approval number: NL65236.058.18, study title: "ICONIC: Arterial oxygenation targets in mechanically ventilated patients in the intensive care unit, a randomized controlled trial"). A detailed description of the ICONIC study regulations can be found in the published protocol (1). In short, adult patients with an expected mechanical ventilation time of 24 hours or more were screened and randomized within two hours after intubation to either the low-oxygenation group (PaO2 55-80 mmHg or SpO2 91-94%) or the high-oxygenation group (PaO2 110-150 mmHg or SpO2 96-100%). Due to the emergency setting of this trial, the majority of the patients were included by deferred consent. The aim was to obtain delayed informed consent as soon as possible from either the patient or the representative. If this had not been achieved within 5 days after the study's commencement, patients were excluded from the study. In this trial no differences in mortality or other relevant clinical endpoints were observed between both groups.

For this substudy, participants were eligible if they were proficient in Dutch and if they were enrolled in the ICONIC study in one of the Dutch ICUs by deferred consent. Patients were excluded if informed consent was obtained before randomisation. The Medical Ethical Committee of Leiden The Hague and Delft reviewed and approved the study. Written informed consent was obtained from all patients or from their legal representatives. Patients were contacted for this study between May 2019 and November 2022. This study was conducted in accordance with the Declaration of Helsinki.

Questionnaire

To assess patient perspectives and experiences on the deferred consent procedure, a questionnaire used in a previous trial was modified and translated (appendix 2) (8). The questionnaire included 12 closed-end questions, and three options to provide a textual response to the choice "other". Participants were asked whether they were aware of their participation, if they provided consent themselves or if consent was provided by their legal representative, and if they would have participated if we could have asked them before the start of the study. Responses were "Yes", "No" or "I don't know". For questions regarding the most suitable substitute decision maker, if they

were content with the decision made on their behalf, whether this was similar to the decision they would have made, and whether participating in the study would help future intensive care patients, a 5-point Likert scale ranging from "Strongly agree" to "Strongly disagree" was used. Participants were given eight response options to indicate their preferred decision-maker. Additionally, in order to evaluate Quality of Life the EQ-5D-5L questionnaire was used (13) (appendix 3).

Procedures

At six months after enrollment in the ICONIC study, a research nurse checked in the electronic patient record if the patient was still alive and reviewed whether the patient consented to participate in the current study. Upon confirmation, patients received a questionnaire on deferred consent and the EQ-5D-5L (after 6 months) either digitally or by post, based on their preference. Reminder telephone calls were made to patients who did not respond within two weeks, and the questionnaire was resent if necessary. If patients still did not respond, a final reminder was sent out, either by e-mail or telephone, 3 weeks after the initial reminder. Patients who failed to respond within nine months of enrollment were excluded from the study. The same procedure was followed after twelve months to collect the second EQ-5D-5L questionnaire. All responses were automatically or manually registered in an electronic case report form (eCRF designed with Castor EDC) (14).

Statistical analysis

Data were extracted from Castor EDC (14) and analysed using R language and environment for statistical computing (R Foundation for Statistical Computing, Vienna, Austria, version 4.0.3). We conducted a comparative analysis of the responses obtained through the questionnaires, and aimed to evaluate acceptance of the deferred consent procedure by patients, whether patients could remember who gave consent and the process involving the substitute decision maker. Responses were presented for the different quality of life groups. Our primary focus was to evaluate whether respondents found deferred consent acceptable and whether this was influenced by quality of life. Continuous variables were presented as means and standard deviations (SDs), or as medians and interquartile ranges (IQRs) depending on the data distribution. Differences between groups were assessed using a Mann-Whitney U test. Categorical variables were presented as frequencies and percentages, and differences were evaluated using a chi-squared test or Fisher's exact test. Statistical significance was considered to be at a P-value of <0.05. The free text comments in the option "Other" were categorized by one investigator, and checked by a co-author. Differences were resolved by consensus.

In order to summarise the different health states of the EQ-5D-5L questionnaire the EQ-5D index value was calculated, including the five dimensions of health included in

the EQ-5D-5L questionnaire: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression (15). Each individual dimension can be scored from 1 (no problems) to 5 (extreme problems). In order to calculate the EQ-5D index predefined weights are assigned to each answer of the individual EQ-5D dimensions. The EQ-5D index value ranges from 0 (worst health) to 1 (full health). To categorize QoL, patients were divided in QoL quartiles based on the calculated EQ-5Dindex, where Q1 reflects the lowest and Q4 the highest. The EQ-5D-5L questionnaire also includes the EQ-VAS score which is a visual analogue scale allowing patients to provide a global assessment of their health status, ranging from 0 (worst imaginable health) to 100 (best imaginable health). To examine the responses to the deferred consent questionnaire in relation to QoL, we performed an ordinal or regular logistic regression, considering EQ5D-index, age, and sex as the independent factors. In order to create an ordinal scale, the answer options "I don't know", "other" and "not applicable", were omitted from the analysis. For one question in which the different answers could not be represented as ordinal items, a chi square test was performed.




Figure 1. The screening and enrollment process for patients who were enrolled in the ICONIC study.

RESULTS

Participants and Quality of Life

Between November 19, 2018 and November 21, 2021, a total of 664 patients were enrolled in the ICONIC study, of which 362 (55%) were eligible to participate in this substudy because deferred consent was obtained. Questionnaires were completed by 197 respondents, resulting in a response rate of approximately 54% (Figure 1). The median time from enrollment until completion of the questionnaire was 29 weeks (IQR, 26-33). The median EQVAS score on subjective health status of all respondents was 80 (IQR, 60-90). The median EQ-5D index score of all respondents was 0.85 (IQR, 0.70-1.00). Baseline characteristics of respondents, non-responders and the total ICONIC population are presented in appendix 4. Table 1 presents the baseline characteristics of respondents categorized by QoL group. Patients who reported a lower QoL had a longer hospital stay (P=0.04). The remaining baseline characteristics were similar between groups.

Deferred consent procedure

Details of the answers to the questions in the four QOL groups are listed in Table E2 in appendix 5. Most patients were either content (61%) or neutral (25%), when asked how they felt about the ICONIC study starting without having been able to give consent (Figure 2). Patients with a higher EQ-5D index were more likely to be content (P=0.02). Only one person (in the highest QoL group) reported not to be content. In addition to the multiple choice answers, two respondents stated they felt forced to consent because the study had already started (appendix 6). When respondents were asked if they knew they had participated in the ICONIC study, the majority of the respondents answered "No" (59%), regardless of their QoL. If consent could have been asked before start of the study, the majority of the respondents would have given consent to participate (89%). These results were similar across QoL groups. Almost all respondents either agreed (55%) or strongly agreed (35%) with the statement "Participation in the ICONIC study will help intensive care patients in the future". These results were independent of QoL.

Table 1. Characteristics of respondents per Quality of life group. To assess QoL life patients were divided in 4 QoL quartiles (Q1, Q2, Q3, Q4) based on the calculated EQ-5Dindex. Q1 reflects the lowest QoL, Q4 the highest. Differences between QoL quartiles were not significant with the exception of hospital length of stay (p=0.04).

Variable	Q1 (N=49)	Q2 (N=49)	Q3 (N=49)	Q4 (N=49)
Age (median (IQR))	63 (50, 68)	62 (53, 73)	63 (51, 72)	67 (56, 72)
Sex = female (%)	19 (39)	14 (29)	15 (31)	16 (33)
Time from randomization to informed consent (days) (median (IQR))	3 (1, 5)	2 (1, 4)	2 (1.75, 5.25)	1 (1, 4)
Apache IV score on admission (median (IQR))	75 (57, 92)	77 (56, 94)	73 (59, 89)	77 (61, 91)
SOFA admission score (median (IQR))	8 (6, 10)	8 (6, 10)	9 (7, 11)	8 (7, 9)
Type of admission (%)				
Medical	34 (70)	39 (80)	36 (74)	34 (69)
Emergency surgery	10 (20)	8 (16)	10 (20)	11 (22)
Elective surgery	5 (10)	2 (4)	3 (6)	4 (8)
Admission diagnosis (%)				
Sepsis	6 (12)	9 (18)	7 (14)	2 (4)
Pneumonia	9 (18)	7 (14)	9 (18)	5 (10)
Cardiac arrest	8 (16)	16 (33)	15 (31)	25 (51)
Abdominal	8 (16)	1 (2)	3 (6)	2 (4)
Neurologic	6 (12)	4 (8)	3 (6)	1 (2)
Trauma	3 (6)	3 (6)	2 (4)	1 (2)
Other	9 (18)	9 (18)	10 (20)	13 (27)
ICU length of stay (days) (median (IQR))	6.6 (4, 16)	5.2 (3, 12)	4.6 (3, 0)	4.5 (3, 8)
Hospital length of stay (days) (median (IQR))	22 (12, 45)	17 (8, 29)	16 (10, 23)	15 (8, 21)
Randomization group = High oxygenation target (110-150 mmHg) (%)	20 (41)	27 (55)	25 (51)	25 (51)
Highest level of education completed (%)				
None	1 (2)	0 (0)	0 (0)	1 (2)
Primary school	4 (8)	5 (10)	3 (6)	4 (8)
Pre-vocational secondary education	13 (27)	12 (25)	9 (18)	9 (18)
Secondary vocational education	17 (35)	17 (35)	22 (45)	18 (37)
Senior general secondary education/ pre-university education	4 (8)	5 (10)	4 (8)	4 (8)
Higher professional education	5 (10)	8 (16)	5 (10)	11 (22)
University	5 (10.2)	2 (4.1)	6 (12)	2 (4)
EQ-5D-index at 6 months (median (IQR))	0.47 (0.29, 0.56)	0.79 (0.74, 0.81)	0.88 (0.85, 0.89)	1 (1, 1)




Figure 2. Results of the level of satisfaction regarding the deferred consent procedure. In the figure, the responses to the question: "How do you feel about the study starting without being able to give consent" are presented an stratified by Quality of Life (QoL) quartiles. Q1 reflects the lowest QoL, Q4 the highest.

Recollection of consent


For 197 respondents, consent was given by a representative only in 136 cases (69%), by the patient only in 55 cases (28%) and by both a representative and the patient in 6 cases (3%). More information on recollection of consent is shown in figure 3. In total, 61 patients had provided written consent themselves. However, when these respondents were asked if they provided consent themselves, 21 (34%) answered "Yes", 21 (34%) answered "No", and 19 (31%) answered "I don't know". The 136 respondents who did not provide their own consent, 9 (7%) erroneously believed they did, while 97 (71%) remembered correctly, and 30 (22%) could not remember. For 142 patients, consent was provided by a legal representative. Among those 142, 104 (73%) could remember correctly (Figure 3). In the 55 cases where a representative did not provide consent, 27 patients (48%) believed they did, 18 (32%) could not remember and 11 (20%) remembered correctly.

Substitute decision maker


Details of responses to questions about substitute decision makers are listed in

Table E2 in appendix 5. In response to the question who participants would prefer to make a decision on their behalf, the majority of respondents preferred the same legal representative who received information about their medical situation during the ICU admission (84%), irrespective of QoL. Most respondents strongly agreed (49%) or agreed (39%) that the doctors asked the right person to provide consent. Patients with a higher QoL were more likely to agree (p=0.005). When asked whether the person who provided consent on their behalf made the same decision as they would have made, most respondents strongly agreed (41%) or agreed (47%). A higher QoL was associated with being more likely to agree (p=0.005). The majority of the patients either strongly agreed (33%) or agreed (55%) with the decision made on their behalf, and patients with a higher QoL were more likely to agree (p<0.001). Only one respondent (in QoL group Q2) disagreed.

Did you provide consent yourself for your participation in the ICONIC study?



Did your legal representative provide consent for your participation in the ICONIC study?



C. Representative gave consent (N=142)

D. Representative did not give consent (N=55)

Figure 3. Patients' memory of who gave consent. In the upper panel 61 patients who had given consent (A) and 136 patients who had not given consent (B) answered the question if they had provided consent themselves to participate in the ICONIC study. The lower panel shows answers to the question if a representative had provided consent for them for 142 patients for whom a representative had given consent (C) and for 55 patients for whom no consent was provided by a representative (D).

DISCUSSION

This study assessed the perspectives of ICU patients on the deferred consent procedure as used in the ICONIC study, and the influence of QoL on these perspectives. Despite many patients being unaware of their participation in the ICONIC study, even though deferred consent had been obtained in the process, most patients were positive regarding the use of deferred consent. Patients with a higher QoL were most likely to be content. In all QoL groups legal representatives were the most preferred individuals to provide consent, and overall, our findings suggest general acceptance of the deferred consent procedure among ICU patients, with a trend of higher acceptance in patients with a higher QoL.

Over the years, literature has shown high levels of patient acceptance of the deferred consent procedure, with acceptance rates ranging from 82-95.6% (7, 8, 16-18). The level of acceptance can be influenced by several factors (6). In the ESCAPE trial, a trial investigating endovascular thrombectomy for acute stroke patients, 78% of the patients disagreed with the use of deferred consent likely due to the high risk nature of the intervention (19). In the ProTECT trial, a trial in which EFIC was used, patients and surrogates of patients with unfavorable clinical outcomes were less accepting compared to patients with favorable outcomes (10, 11). Factors that increased the level of acceptance regarding the use of deferred consent were: perceived benefit of the research, the time-critical nature of the event, and the impact of the condition and emergency situation on the ability to provide consent (6). Other factors that were presumed to affect the level of acceptability of deferred consent were age, ethnicity, previous ICU or research experience, and gender (6). This is the first study to show that QoL affects the level of acceptability.

We hypothesized that patients with an impaired QoL after ICU were less likely to accept having participated in a study without their explicit consent. This was confirmed by the results of our study showing that patient with a higher QoL were more likely to be content with the deferred consent procedure compared to patients with a lower QoL. However, we found that it is difficult to evaluate the effect of QoL on patients' attitudes regarding the use of deferred consent when the vast majority of the patients were content with the procedure. Furthermore, in our study the median EQ-value and median EQ-VAS score were higher after 6 months compared to previous studies evaluating functional status and QoL after ICU stay (20, 21). Therefore, we cannot rule out that results will differ in patients with a severely impaired QoL. To add, some responses, such as 'I don't know', 'Not applicable', and 'other', were excluded from the analysis in order to create an ordinal scale. While a multinomial regression including these answers showed similar results (data not shown), it is something that needs to

be considered in the interpretation of our findings.

Despite patients being mostly positive regarding the use of the deferred consent procedure, our study showed that patients were generally poor at remembering their participation, which is in line with the results of a study evaluating the deferred consent procedure in obstetric emergency research (22). It is important to note that if patients do not recall giving consent, interpreting their attitudes towards enrollment, as assessed through questionnaire responses, becomes challenging. One could argue that it is not surprising that patients do not remember participating in the study because the majority of the patients did not give consent themselves. However, also in patients who did provide consent themselves, only a third of them could remember correctly. Even more remarkable, participants were given a detailed description of the ICONIC study as part of the introduction of the questionnaire, and still they struggled to recall their participation. These findings highlight the importance of effective poststudy communication methods to improve patients' awareness of study participation. Consent is not a one-off event, but needs to be a continuous process. Therefore, in the future, we need to focus on strategies for communicating with participants after enrollment to ensure they understand what they have been part of.

Our study found that a small percentage (4.1%) of patients post hoc disagreed with study participation. These results are consistent with earlier studies indicating similar low numbers (4, 8). Even though this proportion is very low, it is important to consider when performing studies with deferred consent. In line with another study (23) a few but considerable number of respondents reported feeling pressured when asked to provide consent because the study had already started. Careful and open communication about the procedure and about the research components they can still decide about is important when they are able to give consent themselves.

The following study strengths and limitations should be considered. First, this trial is the first to integrate a QoL assessment into the evaluation of patients' opinion of deferred consent procedures. Second, our trial had a relatively high inclusion rate compared to previous studies in this area. Additionally, earlier studies with larger sample sizes were mainly based on hypothetical scenarios with deferred consent, and did not include patients who had actual experience with the procedure. Therefore, a strength of our study is that we included critically ill patients with real-life experience with the deferred consent procedure in the ICONIC trial.

The response rate of 54% may limit the generalizability of our findings to all eligible patients who survived after participating in the ICONIC trial. Although respondents and non-responders were comparable in most baseline characteristics, it is possible

that non-responders may hold different opinions on the deferred consent procedure. Additionally, the opinions of patients who died within 6 months after inclusion or those who declined to consent were not obtained, therefore a group of patients that might have objections to deferred consent could not be included in the analysis. Also, the time from enrollment until responding to the questionnaire for the present study was 6 months and may be considered relatively long. Opinions on having participated in a study with deferred consent may change over time. We choose for studying opinions at 6 months because we anticipated that administering the QoL questionnaire immediately upon or shortly after hospital discharge might result in a less accurate reflection of the actual QoL. Finally, it is important to emphasize that this analysis only included patients from the Netherlands, and it should be noted that the ICONIC trial is classified as a low risk study. Therefore, the results from this trial may be confined to this specific cultural population and the context of a low risk study.

CONCLUSIONS

The present study provides more insight on perspectives of Dutch ICU survivors on their participation in research with deferred consent. It appears that the majority of ICU patients who took part in the ICONIC trial were positive regarding the use of the deferred consent procedure, with patients with a higher QoL status 6 months post-ICU discharge being most likely to be content with the deferred consent procedure. These findings confirm that deferred consent is a suitable option for obtaining consent from ICU patients.

REFERENCES

- Grim CCA, Van Der Wal LI, Helmerhorst HJF, et al. ICONIC study—conservative versus conventional oxygenation targets in intensive care patients: study protocol for a randomized clinical trial. Trials. 2022;23(1).
- World Medical Association Declaration of Helsinki. JAMA. 2013;310(20):2191.
- Burns KE, Zubrinich C, Marshall J, Cook D. The 'Consent to Research' paradigm in critical care: challenges and potential solutions. Intensive Care Med. 2009;35(10):1655-8.
- Kompanje EJO, Maas AIR, Hilhorst MT, Slieker FJA, Teasdale GM. Ethical considerations on consent procedures for emergency research in severe and moderate traumatic brain injury. Acta Neurochirurgica. 2005;147(6):633-40.
- CCMO. CCMO Memorandum Flowcharts deferred consent for medical research in emergency situations. (Available from: https://english.ccmo.nl/publications/publications/2020/04/07/ccmo-memorandum-flowcharts-deferred-consent-for-medical-research-in-emergency-situations.) Accessed June 1, 2023.
- Fitzpatrick A, Wood F, Shepherd V. Trials using deferred consent in the emergency setting: a systematic review and narrative synthesis of stakeholders' attitudes. Trials. 2022;23(1).
- Kamarainen A, Silfvast T, Saarinen S, Virta J, Virkkunen I. Conduct of emergency research in patients unable to give consent--experiences and perceptions of patients, their consent providing next of kin, and treating physicians following a prehospital resuscitation trial. Resuscitation. 2012;83(1):81-5.
- Potter JE, McKinley S, Delaney A. Research participants' opinions of delayed consent for a randomised controlled trial of glucose control in intensive care. Intensive Care Medicine. 2013;39(3):472-80.
- Cuthbertson BH, Roughton S, Jenkinson D, Maclennan G, Vale L. Quality of life in the five years after intensive care: a cohort study. Critical Care. 2010;14(1):R6.
- Dickert NW, Scicluna VM, Baren JM, et al. Patients' Perspectives of Enrollment in Research Without Consent. Critical Care Medicine. 2015;43(3):603-12.
- Whitesides LW, Baren JM, Biros MH, et al. Impact of individual clinical outcomes on trial participants' perspectives on enrollment in emergency research without consent. Clinical Trials. 2017;14(2):180-6.
- 12. van der Wal LI, Grim CCA, Del Prado MR, et

- al. Conservative versus Liberal Oxygenation Targets in Intensive Care Unit Patients (ICONIC): A Randomized Clinical Trial. Am J Respir Crit Care Med. 2023.
- Feng Y-S, Kohlmann T, Janssen MF, Buchholz

 Psychometric properties of the EQ-5D-5L: a systematic review of the literature. Quality of Life Research. 2021;30(3):647-73.
- Castor EDC. Castor Electronic Data Capture 2019 (27 Aug. 2019). Available from: https:// castoredc.com.
- EuroQol Research Foundation. EQ-5D-5L User Guide, 2019. (Available from: https://euroqol. org/publications/user-guides). Accessed June 1, 2023.
- Campwala I, Guyette FX, Brown JB, et al. Patient and surrogate attitudes via an interviewer-administered survey on exception from informed consent enrollment in the Prehospital Air Medical Plasma (PAMPer) trial. BMC Emergency Medicine. 2020;20(1).
- Scicluna VM, Biros M, Harney DK, et al. Patient and Surrogate Postenrollment Perspectives on Research Using the Exception From Informed Consent: An Integrated Survey. Annals of Emergency Medicine. 2020;76(3):343-9.
- Van Den Bos N, Van Den Berg SA, Caupain CM, et al. Patient and proxies' attitudes towards deferred consent in randomised trials of acute treatment for stroke: A qualitative survey. European Stroke Journal. 2021;6(4):395-402.
- Shamy MCF, Dewar B, Chevrier S, et al. Deferral of Consent in Acute Stroke Trials. Stroke. 2019;50(4):1017-20.
- Granja C, Teixeira-Pinto A, Costa-Pereira A. Quality of life after intensive care – evaluation with EQ-5D questionnaire. Intensive Care Medicine. 2002;28(7):898-907.
- Kang J, Yun S, Hong J. Health-related quality of life measured with the EQ-5D-5L in critical care survivors: A cross-sectional study. Intensive Crit Care Nurs. 2022;72:103252.
- Sweeney L, Lanz D, Daru J, et al. Deferred consent in emergency obstetric research: findings from qualitative interviews with women and recruiters in the ACROBAT pilot trial for severe postpartum haemorrhage. BMJ Open. 2022;12(5):e054787.
- Tonnerre EJ, Smith JL, Spencer WS, Date PA, Taylor DM. Patient perceptions of participation in emergency medicine research projects. Emergency Medicine Australasia. 2020;32(4):570-2.