



Universiteit
Leiden
The Netherlands

A hell of a phase curve: mapping the surface and atmosphere of a lava planet

Dang, L.; Zieba, S.; Nguyen, G.; Hammond, M.; Lewis, N.; Buchem, C.P.A. van; ... ; Kreidberg, L.

Citation

Dang, L., Zieba, S., Nguyen, G., Hammond, M., Lewis, N., Buchem, C. P. A. van, ... Kreidberg, L. (2024). A hell of a phase curve: mapping the surface and atmosphere of a lava planet. *Bulletin Of The American Astronomical Society*, 103.02. Retrieved from <https://hdl.handle.net/1887/4213005>

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/4213005>

Note: To cite this publication please use the final published version (if applicable).

Bulletin of the AAS

A Hell of a Phase Curve: Mapping the Surface and Atmosphere of a Lava Planet

**Lisa Dang¹ Sebastian Zieba² Giang Nguyen³ Mark Hammond⁴
Neil Lewis⁵ Christiaan Van Buchem⁶ Mantas Zilinskas⁷ Roxana Lupu⁸
Nicolas Cowan³ Ray Pierrehumbert⁹ Yamila Miguel¹⁰
Laura Kreidberg²**

¹Universite de Montreal, ²Max Planck Institute for Astronomy, ³McGill University,
⁴University of Oxford, ⁵University of Exeter, ⁶Leiden University, ⁷Leiden Observatory,
⁸Eureka Scientific, ⁹Oxford Univ/Univ Chicago, ¹⁰Leiden Observatory / SRON

Published on: Apr 03, 2024

URL: <https://baas.aas.org/pub/2024n4i103p02>

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Lava planets are ultra-short-period planets with bulk densities consistent with terrestrial composition, but dayside temperatures hot enough to melt—and vaporize—rock. Of these planets, the most favorable candidate for atmospheric and surface characterization is the super-Earth, K2-141b with a density of 8 g/cm^3 and an orbital period of 6.7 hrs around a $J = 9$ star. Near-Infrared Spitzer and optical K2 photometric phase curves of K2-141b hints at a tenuous rock vapor atmosphere. The combination of the strong signal expected from K2-141b and the large contrast at long wavelengths makes mid-IR phase curve observations of K2-141b ideal for imposing the most stringent constraint on the dayside and nightside temperature of a lava planet. We present the full-orbit spectroscopic MIRI/LRS (5-12 μm) phase curve of K2-141b to map the planet's thermal emission as a function of longitude. In addition to constraints on the phase offset, Bond albedo, and heat recirculation efficiency from the phase curve, the emission and transmission spectrum provide further evidence into the presence, or lack-of a global atmosphere on a lava planet. Our observations shed light on an extreme outcome of planet formation and provide insights into atmosphere-interior interactions on a rocky planet under extreme irradiation.