



**Universiteit
Leiden**
The Netherlands

Evaluating lung cancer care in the Netherlands: staging, treatment and surgical quality assurance

Hoeijmakers, F.

Citation

Hoeijmakers, F. (2025, May 8). *Evaluating lung cancer care in the Netherlands: staging, treatment and surgical quality assurance*. Retrieved from <https://hdl.handle.net/1887/4245487>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4245487>

Note: To cite this publication please use the final published version (if applicable).



PART I



**NATIONAL QUALITY
REGISTRIES**





CHAPTER 2

Impact of health care organization on surgical lung cancer care

D.J. Heineman, F. Hoeijmakers, N. Beck,
C. Dickhoff, J.M. Daniels, W.H. Schreurs, E. Jakobsen

Lung Cancer. 2019;135:181-7

ABSTRACT

Objectives

Organization and governance of national healthcare might play an important role in decision-making and outcomes in patients with lung cancer. Both Denmark and the Netherlands have a high level of healthcare but a different financial coverage, governance and level of centralization. By using both national databases we analyzed the consequences of these differences on patterns of care and outcomes with a focus on morbidity, mortality and clinical staging.

Materials and methods

General numbers on both healthcare systems were requested. All patients who had surgery for lung cancer from 2013-2016 were included. Mortality, morbidity and clinical staging were analyzed for patients with NSCLC without metastases, only one operation and no neo-adjuvant therapy.

Results

In 2016 annual budget as share of gross national product was 10.4% for both countries. In Denmark 4 hospitals performed lung surgery in 2016, compared to 43 hospitals in the Netherlands. We included 4,030 Danish and 8,286 Dutch patients. In the subgroup 30-day mortality was 1.5% in Denmark compared to 1.9% in the Netherlands. The percentage of patients with a complicated course was 24.4% and 34.8% respectively ($p < 0.05$). Accuracy between cTNM and pTNM was 53.0% in Denmark and 52.9% in the Netherlands.

Conclusion

Surgery for lung cancer is at a high level in both countries, reflected by low mortality-rates. Centralization has been implemented successfully in Denmark, which might explain the lower rate of patients with a complicated post-operative course, although different definitions preclude firm conclusions. In both countries correct clinical staging of lung cancer remains a challenge.

INTRODUCTION

Lung cancer is the most common cause of cancer related death worldwide. Accurate diagnosis and staging are crucial to direct the individual patient towards the optimal treatment. Ideally, all patients with lung cancer should be diagnosed, staged and treated in a uniform fashion. However, it is known that socio-economic status, gender, race and multidisciplinary tumor board discussion, are all factors that have the potential to influence the probability to receive a certain treatment.¹⁻⁴ In addition, to what extent the organization of national healthcare and governance might have an impact on daily practice and outcomes for patients with lung cancer, is currently unknown.

Although both Denmark and the Netherlands have high quality national healthcare systems according to the Euro Health Consumer Index (EHCI)⁵, healthcare organization differs significantly on several aspects: (1) Financial coverage: in Denmark healthcare is financed from taxes for all inhabitants. In contrast, in the Netherlands, healthcare insurance is obligatory and inhabitants pay insurance costs to insurance companies. (2) Governance: in both countries, politicians govern the healthcare system. In the Netherlands, however, the hospitals (and doctors) are independent caregivers, where insurance companies can contract healthcare for their patients. (3) Centralization: one of the results of the difference in governance is that lung cancer care (including surgery) is rigorously centralized in Denmark (4 hospitals performing lung surgery in 2016), a process that has just started in the Netherlands (from 79 in 2005 to 42 hospitals performing lung surgery in 2018).

Both Denmark and the Netherlands have a national audit in which data regarding lung cancer is systematically collected.^{6,7} To improve lung cancer care in Denmark, the Danish Lung Cancer Group (DLCG) was founded in 1991 and the Danish Lung Cancer Registry (DLCR) opened for registration in 2000.⁶ In the Netherlands, a database on lung cancer surgery was initiated in 2011 (Dutch Lung Surgery Audit (DLSA)). To cover and audit all aspects of lung cancer, the radiotherapists and the pulmonologists joined the audit in 2016 and 2017 respectively, and from then on the audit is called Dutch Lung Cancer Audit (DLCA, and DLCA-S for the surgical part of the database).^{7,8}

In this study, the aim was to analyze the differences in healthcare organization and governance and the possible impact on daily practice and outcome, with a focus on morbidity, mortality and clinical staging.

MATERIALS AND METHODS

General data and patients

To compare both countries in general, variables such as lung cancer incidence, number of inhabitants, lung cancer resection rates and life expectancy were requested from the Danish and Dutch Cancer Registries, and from the Organisation for Economic Co-operation and Development (OECDstat).

All patients who had surgery for lung cancer from January 1st 2013, until December 31st 2016 were identified (Group I) and after comparing data definitions, predefined data were compared.

Outcomes

At first, general characteristics of all patients who had surgery for lung cancer from January 1st 2013, until December 31st 2016 were analyzed (Group I). After selection, a subgroup (Group II) was defined to enable proper comparison of patients from both countries, without possible bias from previous (neo-adjuvant) therapy or metastases. This group consisted of patients with primary surgery for NSCLC, without neo-adjuvant therapy and without metastases at time of diagnosis (see *Figure 1*: Flowchart). For patients in Group II, outcomes were analyzed such as morbidity, mortality and risk adjusted mortality rates (RAMR), and the agreement between clinical stage (cTNM) and pathological stage (pTNM), as a measure of clinical staging accuracy in patients with NSCLC.




FIGURE 1. Flowchart of study population and analyses.

Statistical analysis

Descriptive statistics were used for all analysis. Chi-square tests were used to analyze dichotomous variables. To calculate the RAMR for both countries, the expected mortality per country was calculated using multivariable logistic regression analysis. Factors included in the regression model were sex, age, Eastern Cooperative Oncology Group (ECOG) performance status, histopathology, type of resection, type of entry in the thorax and year of surgery. Histopathology was removed from the definitive correction model because it was not significant in multivariable analysis. Statistical analysis was performed in SPSS version 25.

RESULTS

In *Table 1*, the basic characteristics of lung cancer care in Denmark and the Netherlands in 2016 are presented. The number of inhabitants was approximately 3 times higher in the Netherlands and life expectancy at birth was comparable. Incidences of airway tract

cancer were slightly higher in Denmark. In 2016, there were 4 hospitals performing lung surgery in Denmark (average number of resections per hospital per year: 252), compared to 43 hospitals in the Netherlands (average number of resections per hospital per year: 48). The annual budget for health care as a share of gross national product was exactly the same for both countries in 2016: 10.4%. Resection rates for NSCLC were comparable in both countries.

TABLE 1. Basic characteristics of lung cancer care in Denmark and the Netherlands in 2016.

	Denmark	Netherlands
Number of inhabitants ^{9, 10}	5,707,251	16,979,120
Life expectancy female at birth	82.8 years	83.2 years
Life expectancy male at birth	79.0 years	80.0 years
Annual budget for health care as share of gross national product ¹¹	10.4%	10.4%
Number of practicing physicians in (head count, 2015)	20,902	59,073
Number of hospitals performing lung surgery ^{12,13}	4	43
Average volume per hospital per year* (median (min-max))	252 (144-324)	48 (25-109)
Incidence airway tract cancer (lung + trachea + mesothelioma + thymus)/100.000 ^{14,15}	90	79
Incidence lung cancer overall/100.000 ^{14,15}	82	75
Incidence NSCLC/100.000 ^{12,15}	68	55
Resection rate NSCLC ^{12,13}	23.2%	22.8%

* Calculated over 2013-2016

NSCLC – Non-Small Cell Lung Cancer

Outcomes Group I

Between January 1st 2013 and December 31st 2016, a total of 4,030 patients with resected lung cancer were registered in the Danish DLCR and 8,286 patients in the Dutch DLCA-S (*Supplement 1*). The mean age at time of surgery differed slightly between Denmark (68 years) and the Netherlands (65.6 years), however age-distribution was different ($p < 0.05$). In Denmark 54.2% of patients were younger than 70 years compared with 62.5% in the Netherlands. Most surgically treated patients had clinical stage IA (40.2% in Denmark and 35.8% in the Netherlands). Primary surgery in patients with clinical N2 disease was uncommon: 3.6% in Denmark and 4.9% in the Netherlands. Neo-adjuvant treatment was used in 4.6% of patients in Denmark and 6.8% of patients in the Netherlands ($p < 0.05$). Surgery was predominantly performed by thoracoscopic approach in both countries:

64.7% in Denmark and 56.7% in the Netherlands ($p < 0.05$). In the Netherlands, however, in 10.9% of patients a thoracoscopy was converted to an open procedure (conversion rate in Denmark is unknown; converted procedures are recorded as thoracotomy). When analyzing type of resection, lobectomy was performed in 71.5% in Denmark and 74.8% in the Netherlands. In Denmark, 17.0% of patients had a wedge resection compared to 8.5% in the Netherlands, and 10.9% of the Danish patients had two or more surgical procedures compared to 2.4% in the Netherlands ($p < 0.05$).

Outcomes Group II

Table 2 presents the characteristics of Group II. In Denmark, 2,489 patients were registered who met these criteria, in the Netherlands 5,449 patients. In this selected group, clinical stage IA NSCLC remained the most frequent indication for surgery (42.5% versus 39.8% for Denmark and the Netherlands, respectively). Primary surgery for stage IIIA and IIIB comprised 7.8% of all surgical patients in Denmark, compared with 9.1% in the Netherlands. In Denmark 63.8% of patients were operated by thoracoscopic approach, 59.1% of patients in the Netherlands was operated this way ($p < 0.05$). However, in 11.9% of patients operations were started thoracoscopically in the Netherlands but were then converted to an open procedure (converted procedures recorded as thoracotomy in Denmark). Lobectomy remained the preferred procedure in this subgroup: 81.2% of the Danish and 81.4% of the Dutch patients underwent lobectomy. Wedge resection was the performed procedure in 8.1% of patients in Denmark and in 3.2% of patients in the Netherlands.

In *Table 3* the recorded morbidity and mortality of Group II is presented: in Denmark the percentage of patients with a complicated course after surgery was 24.4%, the 30-day mortality was 1.5% and the risk adjusted mortality 1.7% (95%-CI: 1.3-2.4%). For the Netherlands, these numbers were 34.8%, 1.9% and 1.8% (95%-CI: 1.4-2.2%), respectively ((risk adjusted) mortality: not significant, complicated course: $p < 0.05$).

Table 4 presents the clinical and pathological TNM for Group II. Accuracy of staging is highest in early stage lung cancer: in Denmark, this was 59.2% for stage IA and 56.5% for stage IB, in the Netherlands 65.9% and 49.8%, for stage IA and IB, respectively. Overall accuracy of the clinical staging process was 53.0% in Denmark and 52.9% in the Netherlands.

TABLE 2. Operations on patients with NSCLC, no metastasis, one operation and no neo-adjuvant therapy 2013-2016 (Group II).

	Denmark (n)	Denmark (%)	Netherlands (n)	Netherlands (%)	p-value (χ^2)
Total	2,489	100.0%	5,449	100.0%	
Year of surgery					< 0.05
2013	486	19.5%	1,179	21.6%	
2014	593	23.8%	1,118	20.5%	
2015	668	26.8%	1,551	28.5%	
2016	742	29.8%	1,601	29.4%	
Gender					< 0.05
Female	1,207	48.5%	2,414	44.3%	
Male	1,282	51.5%	3,035	55.7%	
Age in categories					< 0.05
<=59	380	15.3%	1,088	20.0%	
60-69 year	886	35.6%	2,109	38.7%	
70-79 year	985	39.6%	1,922	35.3%	
>=80 year	238	9.6%	330	6.1%	
ECOG score					< 0.05
ecog 0	1,505	60.5%	2,813	51.6%	
ecog 1	621	24.9%	1,331	24.4%	
ecog 2+	122	4.8%	176	3.2%	
unknown/missing	241	9.7%	1,129	20.7%	
Clinical TNM stage*					< 0.05
Stage IA	1,057	42.5%	2,168	39.8%	
Stage IB	630	25.3%	1,205	22.1%	
Stage IIA	295	11.9%	877	16.1%	
Stage IIB	312	12.5%	699	12.8%	
Stage IIIA	180	7.2%	471	8.6%	
Stage IIIB	15	0.6%	29	0.5%	
Clinical T stage					0.91
T1a-b (+ To-is)	1,431	45.4%	2,435	44.7%	
T2a-b	932	37.4%	2,054	37.7%	

Clinical N stage									
T3	359	14.4%	813	14.9%					
T4	67	2.7%	147	2.7%					< 0.05
No	2,185	87.8%	4,487	82.3%					
N1	238	9.6%	782	14.4%					
N2	56	2.2%	164	3.0%					
N3	10	0.4%	16	0.3%					
Histopathology									
Adenocarcinoma	1,459	58.6%	3,204	58.8%					< 0.05
Squamous cell carcinoma	720	28.9%	1,979	36.3%					
Different NSCLC	310	12.5%	266	4.9%					< 0.05**
Type of entry thorax									
Thoracotomy	901	36.2%	1,462	26.8%					
VATS/RATS	1,588	63.8%	3,223	59.1%					
Conversion**	-	-	649	11.9%					
Unknown/missing	0	0.0%	115	2.2%					
Type of resection									
Wedge resection	201	8.1%	174	3.2%					< 0.05
Segmentectomy	62	2.5%	101	1.9%					
Lobectomy	2,024	81.2%	4,437	81.4%					
Bilobectomy	101	4.1%	322	5.9%					
Pneumonectomy	101	4.1%	415	7.6%					

*According to TNM-7

** In the Danish data the conversion rate is unknown, conversion of a VATS/RATS was registered as thoracotomy. For calculating the p-value, conversions in the Dutch data are counted in the group thoracotomy.

ECOG – Eastern Cooperative Oncology Group Performance Score

NSCLC – Non-Small Cell Lung Cancer

VATS – Video-Assisted Thoracoscopic Surgery

RATS – Robotic-Assisted Thoracoscopic Surgery

TABLE 3. Mortality and morbidity for Group II.

	Denmark (n)	Denmark (%)	Netherlands (n)	Netherlands (%)	p-value (χ ²)
Total	2,489	100.0%	5,449	100.0%	
Mortality (30 days)	38	1.5%	102	1.9%	0.28
Complicated course (30 days)*	608	24.4%	1,894	34.8%	< 0.05
Bleeding with reoperation	29	1.2%	65	1.2%	0.92
Prolonged air leakage**	229	9.2%	456	8.4%	0.22
Pneumonia	132	5.3%	603	11.1%	< 0.05
Wound infection	6	0.2%	39	0.7%	< 0.05
Empyema	22	0.9%	90	1.7%	< 0.05
Bronchopleural fistula	2	0.1%	18	0.3%	< 0.05
Atelectasis	24	1.0%	143	2.6%	< 0.05
Myocardial ischemia/infarction	6	0.2%	23	0.4%	0.21
Atrial fibrillation	91	3.7%	266	4.9%	< 0.05
Pulmonary embolism	5	0.2%	42	0.8%	< 0.05

* Patients can have multiple complications

** Prolonged air leakage Denmark = more than 7 days, NL = more than 5 days

TABLE 4. Comparison between clinical stage and pathological stage for Group II.

Denmark	pStage										Total	% correct
	Stage IA	Stage IB	Stage IIA	Stage IIB	Stage IIIA	Stage IIIB	Stage IIIA	Stage IIIA	Stage IIIA	Stage IIIB		
cStage	Stage IA	626	271	72	43	45	0	1,057				59.2%
	Stage IB	81	356	87	51	54	1	630				56.5%
	Stage IIA	19	54	109	36	76	1	295				36.9%
	Stage IIB	18	41	48	137	65	3	312				43.9%
	Stage IIIA	18	17	16	35	90	4	180				50.0%
	Stage IIIB	3	2	5	2	2	1	15				6.7%
Total		765	741	337	304	332	10	2,489				53.0%
Netherlands	pStage										Total	% correct
cStage	Stage IA	1,429	419	151	48	118	3	2,168				65.9%
	Stage IB	160	600	223	91	128	3	1,205				49.8%
	Stage IIA	92	158	326	142	153	6	877				37.2%
	Stage IIB	55	64	116	292	170	2	699				41.8%
	Stage IIIA	36	38	63	91	231	12	471				49.0%
	Stage IIIB	4	3	6	7	6	3	29				10.3%
Total		1,776	1,282	885	671	806	29	5,449				52.9%
cStage – clinical stage												
pStage – pathological stage												

DISCUSSION

Clinical auditing was designed to assess quality of medical care and to benchmark treatment outcome. It can be used to analyze variation in patterns of care and outcomes. Benchmarking can be used as a feedback tool for hospitals or individual caregivers to provide insight in ways to improve quality of care.¹⁶⁻¹⁹ This analysis is the first using two national registries (DLCR and DLCA-S) to compare patterns of care and outcomes for lung cancer patients treated with surgery. Both countries are located in Western Europe and have access to high quality health care, but have a different national healthcare organization in terms of financial coverage, governance and level of centralization. Where lung cancer surgery is highly centralized in Denmark, centralization in the Netherlands is an ongoing process. Despite the differences in financial coverage and governance, it is remarkable that the annual budget for health care as share of gross national product is exactly the same in both countries. Several findings from this analysis deserve a closer look.

Centralization

In Denmark, lung surgical operations were performed in 4 hospitals during the entire inclusion period. In the Netherlands, there were 43 hospitals performing lung surgical operations in 2016 (last year of inclusion period). This leads to a considerable variety in number of procedures per hospital per year: an average of 252 procedures per hospital per year in Denmark, compared to 48 in the Netherlands during the analyzed period. Several studies report improved outcomes in high volume centers compared to low volume centers, however, the cut-off value for volume currently is an ongoing subject of debate.²⁰⁻²⁴ It is hard to conclude from our data whether differences in outcomes, as discussed in the following paragraphs, are in more or lesser extent attributable to the different level of centralization.

Characteristics Group I

A relative lower number of patients operated for NSCLC were included in the Dutch registration when compared with Denmark. There are several possible explanations for this finding. First, registration of pulmonary procedures was voluntary at the beginning of the analyzed period in the Netherlands. However, it became obligatory for all institutes performing lung surgery in 2015, resulting in an increased number of registered resections in the DLCA-S (from 1,870 in the year 2013 to 2,349 in 2016). Secondly, the slightly lower incidence of lung cancer in the Netherlands might be part of the explanation why there were not a 3-fold of patients registered in the DLCA-S compared to the DLCR. Another possible explanation is that patients in the Netherlands presented in higher clinical stages, not amenable for surgery. Finally, radiotherapy might have been the preferred choice of treatment in (elderly) patients instead of surgery. These possible explanations warrant further research by analyzing radiotherapy and pulmonology data.

When looking at age, in Denmark 54.2% of patients were younger than 70 years, compared to 62.5% of patients in the Netherlands. The distribution between age categories differed significantly between the two countries, probably explained by the higher centralization rate in Denmark, resulting in more resections for patients with higher ages.²⁵ In addition, the use of radiotherapy instead of surgery in higher aged patients might also contribute to this finding.

Resection type

In Denmark, 17% of patients were treated primarily with a wedge resection, predominantly for diagnostic purposes or metastasectomies, which appeared to be primary lung cancer on final pathological examination. In some of these patients, the wedge resection was followed by a completion lobectomy, explaining the high number of patients that had multiple surgeries in the Danish database. However, even after exclusion of these patients, still 8.1% of patients was treated by wedge resection for NSCLC (Group II) in Denmark. It would be interesting to know patient characteristics and long-term survival of these patients, as a wedge resection is considered an inferior treatment when compared to anatomical parenchymal resections.^{26,27}

Characteristics Group II

When analyzing Group II in more detail, clinical stage IA NSCLC was the most frequent indication for surgery and video-assisted thoracic surgery (VATS) was the preferred surgical approach in both countries. Although minimally invasive surgery does not improve survival compared to open procedures, its benefits are shorter length of stay, less postoperative pain and a better quality of life in the first year after surgery.²⁸⁻³²

Morbidity and mortality Group II

In Denmark, 24.4% of the patients had a complicated post-operative course compared to 34.8% in the Netherlands ($p < 0.05$). Regarding the type of complications, pneumonia, wound infection, empyema, fistula, atelectases, atrial fibrillation and pulmonary embolism were recorded more frequently in the Netherlands than in Denmark ($p < 0.05$). Although data definitions were compared, it is difficult to interpret these data because details regarding definitions of complications are lacking. For example, it is unclear whether the definitions and diagnosis of pneumonia or wound infection are the same in Denmark and the Netherlands. One clear difference in definition is prolonged air leakage: in Denmark this is defined as persistent air leakage longer than 7 days post-operatively, where in the Netherlands this is 5 days. A cardiopulmonary morbidity rate of 18.7% was found by Brunelli et al analyzing the European Society of Thoracic Surgeons (ESTS) database, and although this group published the data definitions of the Society of Thoracic Surgeons (STS) and ESTS database extensively in an earlier publication, it is

still very difficult to compare their reported morbidity rates with these found from our analysis due to the lack of well-defined data definitions.^{33,34}

Although there was an absolute difference in 30-day mortality of 0.4% in favor of Denmark compared with the Netherlands ($p=0.28$), risk adjusted mortality rates narrowed this difference (1.7 versus 1.8% respectively, not significant). Both mortality rates compare favorable with those reported in the literature. In a large retrospective ESTS analysis of 47,960 patients who underwent an anatomic lung resection from 2007-2015, the 30-day mortality was 2.7%.³⁴ When interpreting these data, however, one should realize that the nature of the data from both our dataset as well as the one used by the ESTS (e.g. retrospective analysis, voluntary participation (ESTS), not all hospitals (beginning of the DLCA-S)/countries (ESTS) participating), is subject to bias and true numbers may differ.

Accuracy of clinical staging Group II

The calculated overall staging accuracy of surgically treated NSCLC was 53.0% in Denmark and 52.9% in the Netherlands, with staging of early stages being more accurate. Although these numbers can be considered very low, other authors have previously reported numbers of this range.³⁵⁻³⁷ In Denmark, an accuracy of 91.3% has been reported previously.⁶ The algorithm that was used to calculate this impressive percentage assumed accordance was reached when there was no change of treatment for the patient. A change of treatment was made in those patients in whom a resectable tumor turned out to be T₄ pathologically, when unforeseen N₂ or N₃ nodes were discovered or when unexpected metastases were proven in the pathological TNM. The problem with this algorithm is that it assumes that there is no difference in treatment between a T_{1a} or T₃, or a N₀ or N₁. This used to be true when only surgical treatment could be used as treatment in resectable NSCLC. Currently there are many emerging therapies, from stereotactic ablative body radiotherapy (SABR) in operable stage I disease, chemoradiotherapy as induction therapy in locally advanced NSCLC, to new developments in using immunotherapy as neo-adjuvant therapy. It is therefore of utmost importance to obtain a correct clinical stage before any treatment starts. We do believe that difficulty in clinical staging will be an important limitation in current ongoing studies that evaluate neo-adjuvant therapy. From a Dutch cohort, it was already shown that in surgically treated clinical stage I disease, more than 22% of patients was upstaged to a pathological stage II or higher.³⁸

Limitations

The most important limitation is that comparing two national databases means comparing two sets with different definitions of endpoints. Databases are developed and designed for benchmarking purposes, and to compare hospitals or caregivers within a country. When comparing two countries, it is very important that data definitions

correspond, which is unclear in some of the parameters in this study because of lack of definition. Another important limitation is that it is not clear what impact governance and centralization have on lung cancer care: survival data are lacking, so the effect of centralization could not be extracted from this dataset. This merits further research, particularly since recent studies from the UK and Germany report on shorter length of stay and lower risk of re-admission and death in high volume centers.^{21,22,24} It is interesting to evaluate the different patterns of care in both countries, but differences in datasets hamper drawing firm conclusions on who perform best, and what recommendations would improve health care. Finally, the way data is gathered in the databases can cause unreliable results: in Denmark data registration is linked to the financial administration, assuming hospitals have a correct financial system and thus registered data are correct. In the Netherlands physicians are responsible for data registration and do it themselves or have supporting staff that register the data. Data-verification in the Netherlands is done by an external organization.^{7,39} Both ways of data collection have their own flaws and might cause bias.

Future Perspectives

Due to different data definitions used in both audits, it is very difficult to draw conclusions from the presented data. To compare healthcare and outcome between countries, to identify positive and negative outliers, and learn from each other how to improve outcomes in surgical lung cancer care, we recommend applying comparable data definitions for outcomes and complications or even synchronize registries throughout Europe (and if possible the world).

CONCLUSION

In this study we compared surgical lung cancer care in two western European countries with a high level of healthcare by using their respective national databases. Surgery for lung cancer is of good quality in both Denmark and the Netherlands, which is demonstrated by low mortality numbers. Centralization has been implemented successfully in Denmark, which might explain the lower rates of patients with a complicated course, although different definitions of endpoints in the databases preclude firm conclusions. In both Denmark and the Netherlands correct clinical staging of lung cancer remains a challenge. Implementation of uniform definitions of clinical endpoints on an international level is a prerequisite for comparing datasets. Relating organization and governance of national healthcare systems to clinically relevant endpoints may very well deliver the necessary tools to improve surgical lung cancer care on a global scale.

SUPPLEMENT**SUPPLEMENT 1.** Operations on patients with known age and sex and resection of lung cancer of any type 2013-2016 (Group I).

	Denmark (n)	Denmark (%)	Netherlands (n)	Netherlands (%)	p-value (χ^2)
Total	4,030	100.0%	8,286	100.0%	
Year of surgery					< 0.05
2013	768	19.1%	1,870	22.6%	
2014	1,140	28.3%	1,767	21.3%	
2015	1,030	25.6%	2,300	27.8%	
2016	1,092	27.1%	2,349	28.3%	
Gender					0.07
Female	1,887	46.8%	3,735	45.1%	
Male	2,143	53.2%	4,551	54.9%	
Age					< 0.05
=< 59 years	716	17.8%	1,986	24.0%	
60-69 years	1,467	36.4%	3,187	38.5%	
70-79 years	1,532	38.0%	2,698	32.6%	
=>80 years	315	7.8%	415	5.0%	
ECOG score					< 0.05
Ecog 0	2,329	57.8%	4,188	50.5%	
Ecog 1	972	24.1%	2,001	24.1%	
Ecog 2+	209	5.2%	282	3.4%	
Unknown/missing	520	12.9%	1,815	21.9%	
Clinical TNM stage*					< 0.05

Stage o/Occult	16	0.4%	157	1.9%
Stage IA	1,621	40.2%	2,969	35.8%
Stage IB	859	21.3%	1,504	18.2%
Stage IIA	402	10.0%	1,074	13.0%
Stage IIB	432	10.7%	982	11.9%
Stage IIIA	324	8.0%	833	10.1%
Stage IIIB	37	0.9%	86	1.0%
Stage IV	181	4.5%	86	1.0%
Unknown/missing	158	3.9%	595	7.2%
				< 0.05
Clinical T stage				
T1a-b (+ To-is)	1,827	45.3%	3,535	42.7%
T2a-b	1,352	33.5%	2,729	32.9%
T3	566	14.0%	1,297	15.7%
T4	165	4.1%	323	3.9%
Unknown/Tx/Missing	120	3.0%	402	4.9%
				< 0.05
Clinical N stage				
N0	3,335	82.8%	6,445	77.8%
N1	360	8.9%	1,063	12.8%
N2	144	3.6%	405	4.9%
N3	33	0.8%	50	0.6%
Unknown/missing	158	3.9%	323	3.9%

SUPPLEMENT 1. Continuous

Histopathology	Small cell carcinoma	55	1.4%	85	1.0%	< 0.05***	
	Adenocarcinoma	2,209	54.8%	4,190	50.6%		
	Squamous cell carcinoma	985	24.4%	2,405	29.0%		
	Different NSCLC	739	18.3%	813	9.8%		
	Benign	0	0.0%	473	5.7%		
	Missing/other	42	1.0%	320	3.9%		
	Neoadjuvant treatment						
	No	3,845	95.4%	7,726	93.2%		< 0.05
	Yes	185	4.6%	560	6.8%		
	Type of entry thorax						
Thoracotomy	1,422	35.3%	2,506	30.2%	< 0.05***		
VATS / RATS	2,606	64.7%	4,698	56.7%			
Conversion rate***	-	-	905	10.9%			
Unknown/missing	2	0.0%	177	2.2%			
Type of resection							
Explorative thoracotomy	64	1.6%	123	1.5%	< 0.05		
Wedge resection	684	17.0%	701	8.5%			
Segmentectomy	106	2.6%	187	2.3%			
Lobectomy	2,881	71.5%	6,195	74.8%			
Bilobectomy	150	3.7%	469	5.7%			

Pneumonectomy	139	3.4%	602	7.3%
Unknown/missing/ different	6	0.1%	9	0.1%
More than 1 surgery?				< 0.05
Yes	439	10.9%	202	2.4%
No	3,591	89.1%	8,084	97.6%

*According to TNM-7

** For calculating the p-value the groups 'benign' and 'missing/other' were merged.

*** In the Danish data the conversion rate is unknown, conversion of a VATS/RATS was registered as thoracotomy. For calculating the p-value, conversions in the Dutch data are counted in the thoracotomy group

ECOG – Eastern Cooperative Oncology Group Performance Score

NSCLC – Non-Small Cell Lung Cancer

VATS – Video-Assisted Thoracoscopic Surgery

RATS – Robotic-Assisted Thoracoscopic Surgery

REFERENCES

1. Bryant AS, Cerfolio RJ: Impact of race on outcomes of patients with non-small cell lung cancer. *J Thorac Oncol.* 2008;3:711-715.
2. Wouters MW, Siesling S, Jansen-Landheer NL, et al: Variation in treatment and outcome in patients with non-small cell lung cancer by region, hospital type and volume in the Netherlands. *Eur J Surg Oncol.* 2010;36:583-92.
3. Johnson AM, Hines RB, Johnson JA 3rd, et al: Treatment and survival disparities in lung cancer: the effect of social environment and place of residence. *Lung Cancer.* 2014;83:401-407.
4. Bilfinger TV, Albano D, Perwaiz M, et al: Survival outcomes among lung cancer patients treated using a multidisciplinary team approach. *Clin Lung Cancer.* 2018;19:346-351.
5. Euro Health Consumer Index report 2017, <https://healthpowerhouse.com/media/EHCI-2017/EHCI-2017-report.pdf>, accessed 11th December 2018.
6. Jakobsen E, Green A, Oesterlind K, et al: Nationwide quality improvement in lung cancer care: the role of the Danish Lung Cancer Group and Registry. *J Thorac Oncol.* 2013;8:1238-1247.
7. Ten Berge M, Beck N, Heineman DJ, et al: Dutch Lung Surgery Audit: a national audit comprising lung and thoracic surgery patients. *Ann Thorac Surg.* 2018;106:390-397.
8. Beck N, Hoeijmakers F, Wiegman EM, et al: Lessons learned from the Dutch Institute for Clinical Auditing: the Dutch model for quality assurance in lung cancer treatment. *J Thorac Dis.* 2018;10:53472-3485.
9. <https://www.dst.dk/da/Statistik/emner/befolkning-og-valg/befolkning-og-befolkningsfremskrivning/folketal>, accessed 5th December 2018.
10. <https://opendata.cbs.nl/statline/#/CBS/nl/dataset/37296ned/table?ts=1544007181060>, accessed 5th December 2018.
11. Organisation for economic co-operation and development, <https://stats.oecd.org/>, accessed 11th December 2018.
12. <http://www.lungecancer.dk/documents/B5F266A7-A049-4947-802C-D384A668720C.pdf>, accessed 5th December 2018.
13. <http://dica.nl/jaarrapportage-2016/dlca>, accessed 5th December 2018.
14. <https://sundhedsdatastyrelsen.dk/da/tal-og-analyser/analyser-og-rapporter/sygdomme/cancerregisteret>, accessed 5th December 2018.
15. <https://www.cijfersoverkanker.nl>, accessed 5th December 2018.
16. Pählman L, Bohe M, Cermak B, et al: The Swedish rectal cancer registry. *Br J Surg.* 2007;94:1285-1292.
17. Dreyer NA, Garner S: Registries for robust evidence. *JAMA.* 2009;302:790-791.
18. Ozhathil DK, Li Y, Smith JK, et al: Colectomy performance improvement with NSQIP 2005-2008. *J Surg Res.* 2011;171:e9-13.
19. Van Leersum NJ, Snijders HS, Henneman D, et al: The Dutch surgical colorectal audit. *Eur J Surg Oncol.* 2013;39:1063-70.
20. Kozower BD, Stukenborg GJ: Lung cancer resection volume: is procedure volume really an indicator of quality? *Semin Thorac Cardiovasc Surg.* 2012;24:93-98.
21. Lüchtenborg M, Riaz SP, Coupland VH, et al: High procedure volume is strongly associated with improved survival after lung cancer surgery. *J Clin Oncol.* 2013;31:3141-3146.

22. Møller H, Riaz SP, Holmberg L, et al: High lung cancer surgical procedure volume is associated with shorter length of stay and lower risks of re-admission and death: National cohort analysis in England. *Eur J Cancer*. 2016;64:32-43.
23. Smith CB, Wolf A, Mhango G, et al: Impact of surgeon volume on outcomes of older stage I lung cancer patients treated via video-assisted thoracic surgery. *Semin Thorac Cardiovasc Surg*. 2017;29:223-230.
24. Hoffman H, Passlick B, Ukena D, et al: Surgical therapy for lung cancer: why it should be performed in high volume centres. *Zentralbl Chir*. 2019;144:62-70.
25. Li WW, Visser O, Ubbink DT, et al: The influence of provider characteristics on resection rates and survival in patients with localized non-small cell lung cancer. *Lung cancer*. 2018;60:441-451.
26. Zhang Y, Sun Y, Wang R, et al: Meta-analysis of lobectomy, segmentectomy, and wedge resection for stage I non-small cell lung cancer. *J Surg Oncol*. 2015;111:334-340.
27. Cao C, Chandrakumar D, Gupta S, et al: Could less be more?-A systematic review and meta-analysis of sublobar resections versus lobectomy for non-small cell lung cancer according to patient selection. *Lung Cancer*. 2015;89:121-132.
28. Sugi K, Kaneda Y, Esato K: Video-assisted thoracoscopic lobectomy achieves a satisfactory long-term prognosis in patients with clinical stage IA lung cancer. *World J Surg*. 2000;24:27-30.
29. Cao C, Zhu ZH, Yan TD, et al: Video-assisted thoracic surgery versus open thoracotomy for non-small-cell lung cancer: a propensity score analysis based on a multi-institutional registry. *Eur J Cardiothorac Surg*. 2013;44:849-854.
30. Cai YX, Fu XN, Xu QZ, et al: Thoracoscopic lobectomy versus open lobectomy in stage I non-small cell lung cancer: a meta-analysis. *PLoS One*. 2013;8:e82366.
31. Cao C, Manganas C, Ang SC, et al: Video-assisted thoracic surgery versus open thoracotomy for non-small cell lung cancer: a meta-analysis of propensity score-matched patients. *Interact Cardiovasc Thorac Surg*. 2013;16:244-249.
32. Bendixen M, Jørgensen OD, Kronborg C, et al: Postoperative pain and quality of life after lobectomy via video-assisted thoracoscopic surgery or anterolateral thoracotomy for early stage lung cancer: a randomised controlled trial. *Lancet Oncol*. 2016;17:836-844.
33. Fernandez FG, Falcoz PE, Kozower BD, et al: The Society of Thoracic Surgeons and the European Society of Thoracic Surgeons general thoracic surgery databases: joint standardization of variable definitions and terminology. *Ann Thorac Surg*. 2015;99:368-376.
34. Brunelli A, Salati M, Rocco G, et al: European risk models for morbidity (EuroLung₁) and mortality (EuroLung₂) to predict outcome following anatomic lung resections: an analysis from the European Society of Thoracic Surgeons database. *Eur J Cardiothorac Surg*. 2017;51:490-497.
35. López-Encuentra A, García-Luján R, Rivas JJ, Rodríguez-Rodríguez J, Torres-Lanza J, Varela-Simo G. Comparison between clinical and pathological staging in 2,994 cases of lung cancer. *Ann Thorac Surg*. 2005;79:974-9.
36. Macia I, Moya J, Escobar I, Ramos R, Masuet C, Gamez C, Llatjos R, Martinez-Ballarín I. Quality study of a lung cancer committee: study of agreement between preoperative and pathological staging. *Eur J Cardiothorac Surg*. 2010;37:540-5.
37. Heineman DJ, ten Berge MG, Daniels JM, et al: The quality of staging non-small cell lung cancer in the Netherlands: data from the Dutch Lung Surgery Audit. *Ann Thorac Surg*. 2016;102:1622-1629.
38. Heineman DJ, ten Berge MG, Daniels JM, et al: Clinical staging of stage I non-small cell lung cancer in the Netherlands – Need for improvement in an era with expanding nonsurgical treatment options: data from the Dutch Lung Surgery Audit. *Ann Thorac Surg*. 2016;102:1615-1621.
39. Hoeijmakers F, Beck N, Wouters MWJM, et al: National quality registries: how to improve the quality of data? *J Thorac Dis*. 2018;10:S3490-3499