



Universiteit
Leiden
The Netherlands

Eradication of community-onset Methicillin-resistant *Staphylococcus aureus* carriage: a narrative review

Westgeest, A.C.; Hanssen, J.L.J.; Boer, M.G.J. de; Schippers, E.F.; Lambregts, M.M.C.

Citation


Westgeest, A. C., Hanssen, J. L. J., Boer, M. G. J. de, Schippers, E. F., & Lambregts, M. M. C. (2025). Eradication of community-onset Methicillin-resistant *Staphylococcus aureus* carriage: a narrative review. *Clinical Microbiology And Infection*, 31(2), 173-181.
doi:10.1016/j.cmi.2024.01.003

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/4247960>

Note: To cite this publication please use the final published version (if applicable).



Review

Eradication of community-onset Methicillin-resistant *Staphylococcus aureus* carriage: a narrative reviewAnnette C. Westgeest^{1,*}, Jaap L.J. Hanssen¹, Mark G.J. de Boer^{1,2}, Emile F. Schippers^{1,3}, Merel M.C. Lambregts¹¹ Department of Infectious Diseases, Leiden University Medical Center, Leiden, the Netherlands² Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands³ Department of Internal Medicine, Haga Teaching Hospital, The Hague, the Netherlands

ARTICLE INFO

Article history:

Received 31 October 2023

Received in revised form

14 December 2023

Accepted 6 January 2024

Available online 11 January 2024

Editor: E.J. Kuipers

Keywords:

Colonization

Eradication

MRSA

ABSTRACT

Background: Methicillin-resistant *Staphylococcus aureus* (MRSA) colonization increases infection risk in both patients and healthy individuals. Decolonization therapy has been proven to reduce *S. aureus* infections, but data on the effectiveness of individual decolonization strategies in community-onset MRSA carriage are scarce.

Objectives: The aim of this narrative review was to summarize the evidence on strategies for the elimination of MRSA colonization in community-onset MRSA carriers.

Sources: PubMed database was searched for studies on MRSA eradication, from inception to July 2023. **Content:** Topical therapy is proven to be effective in nasal-only carriage and in temporary load reduction. Mupirocin nasal ointment in combination with chlorhexidine body wash is highly effective in nasal-only MRSA carriers in the community as well. In patients with extra-nasal colonization, addition of orally administered antibiotics likely increases success rates compared with topical therapy alone. Studies on systemic treatment of extra-nasal MRSA decolonization are subject to a high heterogeneity of antimicrobial agents, treatment duration, and control groups. The majority of evidence supports the use of a combination of topical therapy with rifampin and another antimicrobial agent. Decolonization treatment with probiotics is a promising novel non-antibiotic strategy. However, achieving long-term decolonization is more likely in countries with low MRSA prevalence, given the risk of recolonization in a context of high MRSA prevalence.

Implications: The decision to pursue community-onset MRSA eradication treatment in the individual patient should be based on the combination of the treatment objective (short-term bacterial load reduction in health care settings vs. long-term eradication in community settings), and the likelihood of successful decolonization. The latter is influenced by both individual risk factors for treatment failure, and the risk of recolonization. The addition of a combination of systemic antibiotics is rational for extra-nasal long-term decolonization. To determine the most effective systemic antimicrobial agents in MRSA decolonization, more research is needed. **Annette C. Westgeest, Clin Microbiol Infect 2025;31:173**

© 2024 The Author(s). Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Methicillin-resistant *Staphylococcus aureus* (MRSA) is the leading cause of mortality attributable to antimicrobial resistance [1]. The pathogen is notorious for its nosocomial transmission and

hospital outbreaks. On top of that, community-onset MRSA (CO-MRSA) has emerged over the past decades and has become endemic in large parts of the world [2]. Although often carried asymptomatically in the anterior nares, skin lesions, and elsewhere, *S. aureus* is an important cause of severe infections such as bacteraemia. Isolates cultured from blood and the nares are identical in the large majority of patients with *S. aureus* bacteraemia, suggesting an endogenous infection route [3]. Colonization with MRSA increases infection risk even more than colonization with methicillin-susceptible *S. aureus* (MSSA), in both patients and healthy individuals [4–7]. In a North-American cohort of almost

* Corresponding author. Annette C. Westgeest, Department of Infectious Diseases, Leiden University Medical Center, C5-P, P.O. Box 9600, 2300 RC, Leiden, the Netherlands.

E-mail address: a.c.westgeest@lumc.nl (A.C. Westgeest).

30 000 patients who underwent MRSA screening at hospital admission, MRSA carriers had a 20-fold increased odds of developing MRSA bacteraemia compared with non-carriers [8]. In healthy athletes and soldiers, CO-MRSA colonization was associated with a notable increased risk for developing skin and soft tissue infections [4,9]. Decolonization therapy has been proven to reduce *S. aureus* infections in hospitalized patients, most pronounced in surgical patients [10–13]. Although evidence is limited, a 1-year survival benefit of *S. aureus* decolonization before clean surgical procedures is reported [14], as well as cost-effectiveness of active surveillance and decolonization at hospital admission [15].

However, data on the effectiveness of individual decolonization strategies in CO-MRSA carriage are scarce. This review discusses the evidence concerning strategies for elimination of MRSA colonization, with particular emphasis on CO-MRSA.

Methods

We searched PubMed from inception to 31 July 2023, using a combination of keywords to capture MRSA, colonization, and decolonization (search strategy in supplement). In addition, we hand-searched key references and international guidelines to identify citations not captured in the PubMed search. Screening was performed by one reviewer, and in case of uncertainty, a second reviewer was consulted. We screened 1335 titles and abstracts, and 129 articles were selected for a comprehensive full-text review. Studies published in languages other than English were excluded in the full-text review phase. Finally, 66 studies were included in this review. All studies were compiled in EndNote.

Results

Determining eligibility for eradication treatment

An important but complex question remains, which MRSA carriers should undergo eradication treatment. Worldwide differences in policies and attitudes towards MRSA carriage in the community exist between non-endemic and endemic areas. In countries with high MRSA prevalence, e.g. the United States, eradication treatment is not routinely recommended [16]. Some countries with low MRSA prevalence, e.g. the Netherlands and Denmark, successfully implemented a nationwide ‘search and destroy’ policy in the 1980s, targeting MRSA colonization [17,18]. This policy consists of screening and pre-emptive isolation of patients with an increased risk of MRSA carriage when hospitalized and subsequent decolonization treatment when persistent carriage is found. Two years after eradication treatment, 87% of CO-MRSA carriers in a non-endemic setting remained MRSA negative [19].

A major limitation in the generalizability of a ‘search and destroy’ approach to regions with high MRSA prevalence in the community is the high risk of recolonization. Currently, in countries with endemic MRSA, short-term *S. aureus* load reduction is often pursued to reduce infection risk in intensive care unit and surgical patients, either universally or targeted at MRSA carriers (or both MRSA and MSSA carriers) after screening [20]. This temporary suppression of MRSA is efficient in presurgical circumstances [21], but to prevent CO-MRSA transmission, complete eradication is desirable.

At an individual level, risk factors for failure of decolonization therapy can be a reason to refrain from pursuing this goal. Known risk factors for failure are indwelling catheters or medical devices, skin lesions, colonization of household contacts, chronic pulmonary disease, and an immunocompromised status [22,23].

As a result, two main factors should guide the decision for eradication therapy in an individual patient. First, the treatment

goal, which can be either long-term eradication to prevent community transmission and infections, or short-term load reduction to prevent nosocomial infections and transmission. Second, the likelihood of long-term success of decolonization treatment, influenced by both the presence of individual risk factors for failure and the prevalence of MRSA in the environment, driving the risk of recolonization (Fig. 1).

Lastly, when considering eradication treatment, potential adverse effects should be weighed in. This includes well-known effects such as (hepato-)toxicity and risk of *Clostridioides difficile* infection, but also newer insights such as potential disruption of the human microbiome [24].

Strategies for eradication therapy

MRSA eradication therapy usually exists of either topical—i.e. nasal ointment and skin wash—therapy alone or a combination of topical and systemic anti-staphylococcal agents. Topical therapy is proven to be effective in nasal-only carriage and in temporary (presurgical) load reduction [25,26]. In contrast, in patients with other body sites positive for MRSA, eradication with mupirocin and chlorhexidine skin wash is reported to be insufficient [27–29]. In a randomized controlled trial (RCT) of hospitalized patients colonized with MRSA on multiple body sites in a hospital with endemic MRSA, mupirocin was only marginally effective [26]. In particular, throat carriage is associated with failure of topical eradication treatment [30]. In a small study on Swedish outpatients with MRSA throat carriage, topical therapy led to successful eradication in only 13%, as compared with 61% when topical therapy was combined with systematic antibiotics [31]. Positive household contacts were simultaneously treated. A similar outcome was reported in outpatient MRSA carriers in Canada initially; however, after 1 year, success rates with and without systemic antibiotics were found to be equal [32]. Canada is a high-endemic area, and because no screening of household contacts or genotyping was performed, it remains undetermined whether this outcome resulted from recolonization with a different strain, or long-term failure of eradication treatment.

Discriminating between nasal-only and extra-nasal MRSA colonization to guide optimal eradication therapy seems appropriate considering the abovementioned studies and from a pathophysiologic perspective. This distinction is also made in the Dutch MRSA eradication guideline, where mupirocin-sensitive, nasal-only MRSA carriage with intact skin is considered ‘uncomplicated’ and is recommended to be treated with topical agents only. MRSA carriers with extra-nasal colonization or other risk factors for (topical) treatment failure, e.g. active skin lesions and foreign body material, are considered ‘complicated’ and are treated with additional systemic antimicrobial agents [33]. This specific approach led to sustained decolonization in 85% of carriers after 1 year of follow-up [23].

MRSA carriage of household members was the most frequently encountered risk factor for CO-MRSA infections in Denmark between 1999 and 2006 [34], and was associated with failure of eradication treatment [22]. This emphasizes the need for screening and simultaneous eradication of all positive household members, especially in case of treatment failure.

In general, infection prevention and control measures are crucial in preventing further spread of MRSA [35], but are not included in this review.

Efficacy of topical decolonization therapy

The most commonly used topical treatment for *S. aureus* decolonization is mupirocin nasal ointment, which achieves its




Fig. 1. Factors of influence on the decision for eradication treatment in an individual MRSA carrier. The decision to start eradication therapy in an MRSA carrier should be guided by the treatment goal and the likelihood of long-term success of decolonization treatment, influenced by both the presence of individual risk factors for failure and the prevalence of MRSA in the environment, driving the risk of recolonization. MRSA, methicillin-resistant *Staphylococcus aureus*.

antimicrobial effect by inhibiting bacterial protein synthesis. It is often combined with daily antiseptic body wash. Mupirocin nasal ointment was proven to be effective in MSSA decolonization in the 1980s and 1990s [36–45]. In a systematic review that included studies analysing both MSSA and MRSA colonization, mupirocin resulted in negative MRSA cultures in 94% of patients after 1 week [25]. This percentage decreased to 65% after (mid- to long-term) follow-up.

All RCTs on topical MRSA eradication treatment are summarized in Table 1 [26,27,29,46–53].

Very high MRSA decolonization success rates have been reported with mupirocin treatment in a prospective study in hospitalized patients (98%), and an RCT involving long-term care facility residents (93%) [53,54]. Furthermore, in a retrospective analysis of MRSA-colonized patients who were readmitted during the study period, mupirocin was associated with being MRSA negative at readmission, compared with no treatment [55,56].

Focusing specifically on MRSA eradication in the community, little evidence is available on the effectiveness of mupirocin [57]. In an RCT involving 134 healthy MRSA-colonized American soldiers, mupirocin led to 88% nasal eradication compared with 65% with placebo after 8 weeks of follow-up [51]. Similarly, in 87 German hospital workers with nasal MRSA colonization, who were withdrawn from work until MRSA free, treatment with mupirocin nasal ointment and antiseptic mouth rinse and body wash resulted in successful eradication in 84% at 3 months of follow-up [58]. Prolonged mupirocin decolonization treatment (twice monthly for 5 days during 6 months) after discharge in patients that had been

hospitalized in the United States with MRSA infections led to a higher decolonization rate compared with placebo (OR of colonization = 0.44) [46].

Conflicting results on the effectiveness of mupirocin in CO-MRSA have been reported in regions with high MRSA prevalence, which may be indicative of an increased risk of recolonization rather than treatment failure. In an RCT comparing topical with systemic treatment in patients treated at a dedicated MRSA outpatient clinic, initial decolonization was achieved in 13 of 25 patients who received topical treatment, but this decreased to three after 12 months [32]. The vast majority of patients in this study were colonized at multiple body sites. Seven days of mupirocin nasal ointment combined with antiseptic body wash compared with placebo did not improve decolonization rate in 49 outpatients living with HIV in a RCT [50]. In addition, in a study involving 223 households with ambulatory MRSA skin and soft tissue infections, persistent MRSA colonization was similar in households with and without topical decolonization after 6 months of follow-up [49].

A concern with the use of mupirocin is the emergence of mupirocin resistance [59]. The prevalence of mupirocin resistance varies widely and is reported to be associated with its increased use [60]. Remarkably, a *post-hoc* analysis of the REDUCE-MRSA trial showed an overall low prevalence of mupirocin-resistant isolates and no increase after mupirocin decolonization treatment [61].

Given the risk of the emergence of mupirocin resistance, alternative topical therapies have been evaluated. Medical-grade honey was only marginally inferior to mupirocin in decolonizing nasal MRSA colonization in a small RCT [47]. Topical therapy with tea tree

Table 1

Randomized trials on topical MRSA decolonization treatment

Author, year	Country	No. of patients	Population	Treatments	Treatment duration ^a	Follow-up ^a	Culture site(s) ^b	Decolonized at end of follow-up	Other outcome
Miller et al., 2023 [46]	United States	2121	Inpatients, post-discharge	1. Education 2. + mupirocin + chx	Twice monthly 5 d for 6 mo	270	N, T, A, G, W	1. 57% 2. 73% (p < 0.01)	
Poovelikunnel et al., 2018 [47]	Ireland	100	14% outpatients 86% inpatients	1. Medical-grade honey + Tricolsan 2. Mupirocin + Tricolsan	5	Short term	N, G, W	1. 43% 2. 57% (p 0.20)	Received two treatment courses: 1. 78% 2. 20%
Landelle et al., 2016 [48]	Switzerland	146	Inpatients	1. Polyhexanide 2. Placebo	10	28	N, G	1. 34% 2. 29% (p 0.56)	
Cluzet et al., 2016 [49]	United States	149	Households with SSTI	1. Education 2. + mupirocin + chx 3. + mupirocin + chx + reminders	7	180	N, A, G	1. ±80% 2. ±80% 3. ±80%	Time to clearance: 1. 19 d 2 + 3. 23 d
Weintrob et al., 2015 [50]	United States	49	Outpatients with HIV	1. Mupirocin + chx 2. Placebo	7	180	N, A, G, T, P	1. 67% 2. 67%	
Fritz et al., 2011 [29]	United States	300	Patients with SSTI + MSSA/MRSA colonization	1. Education 2. + mupirocin 3. + mupirocin + chx 4. + mupirocin + bleach baths	5	120	N, A, G	1. 48% 2. 56% (p 0.40) 3. 54% (p 0.51) 4. 71% (p 0.02)	Nasal decolonization: 1. 50% 2. 77% (p < 0.01) 3. 76% (p < 0.01) 4. 85% (p < 0.01)
Ellis et al., 2007 [51]	United States	134	Healthy soldiers	1. Mupirocin 2. Placebo	5	56	N	1. 88% 2. 65%	
Wendt et al., 2007 [27]	Germany	114	In- and outpatients, nursing home residents	1. Mupirocin + chx 2. Mupirocin + placebo	5	30	N, T, G, P	1. 8% (p 0.47) 2. 13%	Groin decolonization: 1. 93% (p < 0.01) 2. 82%
Dryden et al., 2004 [52]	UK	224	Inpatients	1. Mupirocin + chx 2. Tea tree oil	5	14	N, T, G, S, W	1. 49% 2. 42% (p 0.03)	Nasal decolonization: 1. 86% 2. 58% (p < 0.01)
Mody et al., 2003 [53]	United States	127	Long-term care facility residents with MRSA/MSSA colonization	1. Mupirocin 2. Placebo	14	30	N, W	1. 88% (p < 0.01) 2. 13%	
Harbarth et al., 1999 [26]	Switzerland	98	Inpatients	1. Mupirocin + chx 2. Placebo + chx	5	26	N, G, U, W	1. 25% (p 0.40) 2. 18%	Nasal decolonization: 1. 44% 2. 23%

Chx, chlorhexidine; MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin-susceptible *Staphylococcus aureus*; SSTI, skin and soft tissue infection.^a In days.^b N = nasal, A = axilla, G = groin, T = throat, P = perineum/rectum, W = wounds/skin lesions, S = sputum, U = urine.

Table 2
Randomized trials on systemic MRSA decolonization treatment

Author, year	Country	No. of patients	Population	Treatments	Treatment duration ^a	Follow-up ^a	Culture site(s) ^b	Decolonized at end of follow-up
Eum et al., 2021 [32]	Canada	98	Outpatients and inpatients	1. Mupirocin + chx 2. Mupirocin + chx + rifampin + doxycycline	7	365	N, P, W	1. 32% 2. 50% (p 0.04)
Lindgren et al., 2018 [31]	Sweden	52	Outpatients with throat colonization	1. Mupirocin + chx + rifampin + clindamycin/SXT 2. Mupirocin + chx	7	180	N, T, P, W	1. 61% 2. 13% (p < 0.01)
Simor et al., 2007 [63]	Canada	146	Inpatients	1. Mupirocin + chx + rifampin + doxycycline 2. No treatment	7	90	N, P, W, D	1. 74% 2. 32% (p < 0.01)
Chang et al., 2000 [64]	Taiwan	16	ICU patients	1. Fusidic acid 2. No treatment	7	28	N, T, W, S	1. 33% 2. 50% (p 0.95)
Parras et al., 1995 [65]	Spain	84	13% HCW and 87% inpatients	1. Mupirocin + chx 2. SXT + topical fusidic acid + chx	5	28	N	1. 96% 2. 95% (p > 0.05)
Muder et al., 1994 [66]	United States	35	Long-term care facility residents	1. Rifampin 2. Minocycline 3. Rifampin + minocycline 4. No treatment	5	90	N, U, W	1. 67% 2. 38% 3. 50% 4. 14%
Walsh et al., 1993 [67]	United States	94	HCWs and inpatients	1. Novobiocin + rifampin 2. SXT + rifampin	7	14	N, G, W, S	1. 67% 2. 53% (p 0.18)
Peterson et al., 1990 [68]	United States	21	Inpatients	1. Ciprofloxacin + rifampin 2. SXT + rifampin	14	180	N, G, W	1. 27% 2. 40% (p > 0.1)

Chx, chlorhexidine; HCW, health care worker; ICU, intensive care unit; MRSA, methicillin-resistant *Staphylococcus aureus*; SXT, trimethoprim/sulfamethoxazole.

^a In days.

^b N = nasal, A = axilla, G = groin, T = throat, P = perineum/rectum, W = wounds/skin lesions, S = sputum, U = urine, D = medical device or catheter exit site.

preparations was significantly less effective compared with mupirocin-based topical therapy for the clearance of intranasal MRSA colonization [52]. Polyhexanide was not effective in MRSA decolonization compared with placebo in an RCT [48], and inferior to mupirocin and chlorhexidine in a retrospective analysis [62].

Efficacy of decolonization therapy with addition of systemic antibiotics

Using systemic antibiotics in addition to the topical treatment for MRSA decolonization is common practice in case of extra-nasal colonization in some countries, reserved for cases of topical treatment failure in others, and seldom or never employed in a third category of countries. Most studies on systemic treatment for MRSA decolonization have been performed in health care settings, with a high heterogeneity of treatment agents and control groups. All RCTs on systemic MRSA eradication treatment are summarized in Table 2 [31,32,63–68].

The combination treatment consisting of antiseptic body wash, intranasal mupirocin, rifampin, and trimethoprim/sulfamethoxazole or doxycycline was highly effective in MRSA decolonization of hospitalized patients [63,69,70]. In a small RCT in long-term care facilities in the United States, rifampin monotherapy was superior to no treatment, as well as to minocycline monotherapy. Combination therapy with rifampin and minocycline was not superior to rifampin alone. The majority of patients had decubitus and indwelling catheters, and after 3 months only half of the treated patients remained MRSA negative [66]. Moreover, the risk of emerging resistance when using rifampin monotherapy makes this an inappropriate approach. Another randomized trial on oral fusidic acid monotherapy or no treatment showed no difference in MRSA decolonization rate in 16 intensive care unit patients. However, the study was terminated because of emergence of fusidic acid-resistant strains [64]. Two cohort studies on trimethoprim/sulfamethoxazole plus rifampin in hospitalized patients resulted in 64–66% successful MRSA decolonization [71,72]. Oral vancomycin, combined with topical therapy, was effective in eradicating MRSA-colonized staff and residents of a nursing home during an outbreak, although 80% experienced side effects [73].

Compared with topical therapy with mupirocin only, the combination of oral trimethoprim/sulfamethoxazole plus topical fusidic acid (without mupirocin) performed marginally worse in MRSA eradication in hospitalized patients and personnel after 14 days [65]. Rifampin plus novobiocin resulted in a non-significant higher decolonization rate after 14 days compared with rifampin plus trimethoprim-sulfamethoxazole (respectively 67% vs. 53%) in an RCT on MRSA-colonized patients and personnel in the United States. Decolonization in both groups was significantly more often achieved in colonization sites other than wounds [67]. However, novobiocin has since been withdrawn from the market. Rifampin with ciprofloxacin was more effective compared with rifampin with trimethoprim/sulfamethoxazole in an RCT on MRSA-colonized patients (50% vs. 37% eradicated after 6 months of follow-up). Only 21 patients were enrolled when the study was terminated because of emergence of ciprofloxacin resistance in the hospital, unrelated to the study [68].

Few studies have been published specifically on systemic MRSA decolonization in the community, mainly from countries with low MRSA prevalence. The previously mentioned Swedish study randomly assigned 52 outpatients with MRSA throat carriage between chlorhexidine, nasal mupirocin, rifampin, and either clindamycin or trimethoprim/sulfamethoxazole (group 1) and chlorhexidine and nasal mupirocin only (group 2). At 6 months of follow-up, 61% of systemically treated vs. 13% of topical treated patients were successfully decolonized (p < 0.01) [31]. In a cohort of




Fig. 2. Implications of prevalence of MRSA carriage for the approach of the community and individual. In a low CO-MRSA prevalent setting, sustained decolonization of CO-MRSA is feasible and can prevent further spread in the community. This supports the ‘search and destroy’ policy, in which carriers are identified, household contacts screened, and decolonization is attempted. In this setting, this policy is effective in maintaining a low MRSA prevalence. In contrast, in high-endemic regions, there is high risk of recolonization. Consequently, routine eradication treatment of CO-MRSA aiming at achieving a non-carrier state for a prolonged period of time is less likely to be successful. In this setting, a standard ‘search and destroy’ policy is not likely to reduce the high MRSA prevalence, and an individualized approach is more rational. CO-MRSA, community-onset methicillin-resistant *Staphylococcus aureus*.

Dutch outpatients with extra-nasal MRSA colonization, decolonization treatment combination of chlorhexidine body wash, mupirocin ointment intranasally, and a combination of two systemic antibiotics (mostly rifampin with trimethoprim or doxycycline) was successful in 85% of patients and the vast majority was still negative after 1 year of follow-up [23]. Two Danish cohort studies did not show a benefit of adding clindamycin to decolonization treatment of MRSA throat carriage [74,75].

In the previously discussed Canadian study, a country with high MRSA prevalence, 98 outpatients with MRSA colonization at any site were randomized between a 7-day course of topical treatment alone or supplemented with oral rifampin and doxycycline [32]. The initial success rate was higher in the systemic treatment arm, but this difference had disappeared after 12 months of follow-up. As said, no genotyping was performed to elucidate whether this was because of long-term treatment failure or recolonization with a different strain.

Future perspectives

Concerns about emerging resistance and the impact on the microbiome resulting from current treatment strategies drive the search for alternative, non-antibiotic, decolonization therapies. A recently published phase-two trial showed promising results of oral probiotics for nasal and intestinal *S. aureus* decolonization, with a 95% reduction of *S. aureus* colonization without notable changes in the microbiota [76]. Ongoing research is focused on engineering a skin probiotic to selectively combat MRSA colonization [77]. In addition, novel non-antibiotic drugs are being evaluated for their potential in *S. aureus* eradication, including the porphyrin drug XF-73, the LL-37-derived peptide P10 and SAAP-148 [78–80], and bacteriophage therapy [77].

Despite multiple attempts, vaccines to prevent *S. aureus* infections have so far not been proven clinically effective [81]. However, the high burden of disease provides grounds to continue the search.

Discussion and conclusion

MRSA decolonization has been proven to reduce infections in both patients and healthy individuals. However, determining eligible treatment candidates and applying experiences and results from countries with low MRSA prevalence to countries with high MRSA prevalence continue to be challenging. In general, eradication studies in high prevalence areas are hampered by the indistinguishability of failing eradication treatment vs. recolonization. The likelihood of successful long-term decolonization is lower in a high endemicity setting compared with a low endemicity setting, because of the heightened risk of recolonization (Fig. 2). Thus, both treatment goal (short-term bacterial load reduction in health care settings vs. long-term eradication in community settings), and likelihood of successful prolonged eradication should guide the eligibility for CO-MRSA decolonization treatment in the individual patient.

Although highly effective in decolonization of nasal MRSA carriage, the combination of mupirocin and antiseptic body wash appears to be insufficient in patients with extra-nasal MRSA colonization. The addition of systemic antibiotics is a rational approach in this patient category, but studies on systemic treatment of extra-nasal MRSA decolonization are subject to a high heterogeneity of treatment agents and comparator groups. Most evidence support a combination of topical therapy with rifampin and another antimicrobial agent for extra-nasal MRSA eradication. Future research would gain clinical applicability from reporting the carrier status of household contacts, long-term follow-up cultures, and reporting genotyping in case of failure. Eradication treatment with probiotics holds promise as a novel non-antibiotic strategy.

Author contributions

ACW initially screened all titles and abstracts. JIJH was consulted in case of uncertainty. ACW and JIJH screened all full-text articles and wrote the first draft of the manuscript. EFS, MGJdB, and MMCL contributed to the interpretation of the results and the revision of the manuscript.

Transparency declaration

MMCL is principle investigator in Leiden of the E.MBRACE study, a clinical vaccine study for the prevention of invasive *E. coli* disease by Janssen pharmaceuticals. The remaining authors declare that they have no conflicts of interest.

This project was partially funded by the Antimicrobial Resistance Network 'Holland West'.

Acknowledgements

We would like to thank Manon Zuurmond for her excellent help in creating the figures and Jan Schoones for his invaluable help in establishing the search strategy.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cmi.2024.01.003>.

References

- [1] Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* 2022;399:629–55. [https://doi.org/10.1016/s0140-6736\(21\)02724-0](https://doi.org/10.1016/s0140-6736(21)02724-0).
- [2] Baud O, Giron S, Aumeran C, Mouly D, Bardon G, Besson M, et al. First outbreak of community-acquired MRSA USA300 in France: failure to suppress prolonged MRSA carriage despite decontamination procedures. *Eur J Clin Microbiol Infect Dis* 2014;33:1757–62. <https://doi.org/10.1007/s10096-014-2127-6>.
- [3] von Eiff C, Becker K, Machka K, Stammer H, Peters G. Nasal carriage as a source of *Staphylococcus aureus* bacteremia. Study Group. *N Engl J Med* 2001;344:11–6. <https://doi.org/10.1056/nejm200101043440102>.
- [4] Ellis MW, Hospenthal DR, Dooley DP, Gray PJ, Murray CK. Natural history of community-acquired methicillin-resistant *Staphylococcus aureus* colonization and infection in soldiers. *Clin Infect Dis* 2004;39:971–9. <https://doi.org/10.1086/423965>.
- [5] Quezada Joaquin NM, Diekema DJ, Perencevich EN, Bailey G, Winokur PL, Schweizer ML. Long-term risk for readmission, methicillin-resistant *Staphylococcus aureus* (MRSA) infection, and death among MRSA-colonized veterans. *Antimicrob Agents Chemother* 2013;57:1169–72. <https://doi.org/10.1128/aac.01968-12>.
- [6] Safdar N, Bradley EA. The risk of infection after nasal colonization with *Staphylococcus aureus*. *Am J Med* 2008;121:310–5. <https://doi.org/10.1016/j.amjmed.2007.07.034>.
- [7] Davis KA, Stewart JJ, Crouch HK, Florez CE, Hospenthal DR. Methicillin-resistant *Staphylococcus aureus* (MRSA) nares colonization at hospital admission and its effect on subsequent MRSA infection. *Clin Infect Dis* 2004;39:776–82. <https://doi.org/10.1086/422997>.
- [8] Marzec NS, Bessesen MT. Risk and outcomes of methicillin-resistant *Staphylococcus aureus* (MRSA) bacteremia among patients admitted with and without MRSA nares colonization. *Am J Infect Control* 2016;44:405–8. <https://doi.org/10.1016/j.ajic.2015.11.006>.
- [9] Karanika S, Kinamon T, Grigoras C, Mylonakis E. Colonization with methicillin-resistant *Staphylococcus aureus* and risk for infection among asymptomatic athletes: a systematic review and metaanalysis. *Clin Infect Dis* 2016;63:195–204. <https://doi.org/10.1093/cid/ciw240>.
- [10] Perl TM, Cullen JJ, Wenzel RP, Zimmerman MB, Pfaller MA, Sheppard D, et al. Intranasal mupirocin to prevent postoperative *Staphylococcus aureus* infections. *N Engl J Med* 2002;346:1871–7. <https://doi.org/10.1056/NEJMoa003069>.
- [11] van Rijen M, Bonten M, Wenzel R, Kluytmans J. Mupirocin ointment for preventing *Staphylococcus aureus* infections in nasal carriers. *Cochrane Database Syst Rev* 2008;2008:CD006216. <https://doi.org/10.1002/14651858.CD006216.pub2>.
- [12] Bode LG, Kluytmans JA, Wertheim HF, Bogaers D, Vandenbroucke-Grauls CM, Roosendaal R, et al. Preventing surgical-site infections in nasal carriers of *Staphylococcus aureus*. *N Engl J Med* 2010;362:9–17. <https://doi.org/10.1056/NEJMoa0808939>.
- [13] Huang SS, Singh R, McKinnell JA, Park S, Gombosov A, Eells SJ, et al. Decolonization to reduce postdischarge infection risk among MRSA carriers. *N Engl J Med* 2019;380:638–50. <https://doi.org/10.1056/NEJMoa1716771>.
- [14] Bode LG, van Rijen MM, Wertheim HF, Vandenbroucke-Grauls CM, Troelstra A, Voss A, et al. Long-term mortality after rapid screening and decolonization of *Staphylococcus aureus* carriers: observational follow-up study of a randomized, placebo-controlled trial. *Ann Surg* 2016;263:511–5. <https://doi.org/10.1097/sla.0000000000001060>.
- [15] Nelson RE, Samore MH, Smith KJ, Harbarth S, Rubin MA. Cost-effectiveness of adding decolonization to a surveillance strategy of screening and isolation for methicillin-resistant *Staphylococcus aureus* carriers. *Clin Microbiol Infect* 2010;16:1740–6. <https://doi.org/10.1111/j.1469-0691.2010.03324.x>.
- [16] Liu C, Bayer A, Cosgrove SE, Daum RS, Fridkin SK, Gorwitz RJ, et al. Clinical practice guidelines by the Infectious Diseases Society of America for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children: executive summary. *Clin Infect Dis* 2011;52:285–92. <https://doi.org/10.1093/cid/cir034>.
- [17] Vos MC, Behrendt MD, Melles DC, Mollema FP, de Groot W, Parlevliet G, et al. 5 years of experience implementing a methicillin-resistant *Staphylococcus aureus* search and destroy policy at the largest university medical center in The Netherlands. *Infect Control Hosp Epidemiol* 2009;30:977–84. <https://doi.org/10.1086/605921>.
- [18] Böcher S, Skov RL, Knudsen MA, Guardabassi L, Mølbak K, Schouenborg P, et al. The search and destroy strategy prevents spread and long-term carriage of methicillin-resistant *Staphylococcus aureus*: results from the follow-up screening of a large ST22 (E-MRSA 15) outbreak in Denmark. *Clin Microbiol Infect* 2010;16:1427–34. <https://doi.org/10.1111/j.1469-0691.2009.03137.x>.
- [19] Longtin Y, Sudre P, François P, Schrenzel J, Aramburu C, Pastore R, et al. Community-associated methicillin-resistant *Staphylococcus aureus*: risk factors for infection, and long-term follow-up. *Clin Microbiol Infect* 2009;15:552–9. <https://doi.org/10.1111/j.1469-0691.2009.02715.x>.
- [20] Huang SS, Septimus E, Kleinman K, Moody J, Hickok J, Avery TR, et al. Targeted versus universal decolonization to prevent ICU infection. *N Engl J Med* 2013;368:2255–65. <https://doi.org/10.1056/NEJMoa1207290>.
- [21] Coia JE, Wilson JA, Bak A, Marsden GL, Shimonovich M, Loveday HP, et al. Joint healthcare infection society (HIS) and infection prevention society (IPS) guidelines for the prevention and control of methicillin-resistant *Staphylococcus aureus* (MRSA) in healthcare facilities. *J Hosp Infect* 2021;118:S1–39. <https://doi.org/10.1016/j.jhin.2021.09.022>.
- [22] Ammerlaan HS, Kluytmans JA, Berkhout H, Buiting A, de Brauwier EI, van den Broek PJ, et al. Eradication of carriage with methicillin-resistant *Staphylococcus aureus*: determinants of treatment failure. *J Antimicrob Chemother* 2011;66:2418–24. <https://doi.org/10.1093/jac/ckr250>.
- [23] Westgeest AC, Schippers EF, Delfos NM, Ellerbroek LJ, Koster T, Hira V, et al. Complicated carriage with methicillin-resistant *Staphylococcus aureus*: evaluation of the effectiveness of decolonization regimens advised in the Dutch National Guideline. *Antimicrob Agents Chemother* 2021;65:e0025721. <https://doi.org/10.1128/aac.00257-21>.
- [24] Fritz SA, Wylie TN, Gula H, Hogan PG, Boyle MG, Muenkes CE, et al. Longitudinal dynamics of skin bacterial communities in the context of *Staphylococcus aureus* decolonization. *Microbiol Spectr* 2022;10:e0267221. <https://doi.org/10.1128/spectrum.02672-21>.
- [25] Ammerlaan HS, Kluytmans JA, Wertheim HF, Nouwen JL, Bonten MJ. Eradication of methicillin-resistant *Staphylococcus aureus* carriage: a systematic review. *Clin Infect Dis* 2009;48:922–30. <https://doi.org/10.1086/597291>.
- [26] Harbarth S, Dharan S, Liassine N, Herrault P, Auckenthaler R, Pittet D. Randomized, placebo-controlled, double-blind trial to evaluate the efficacy of mupirocin for eradicating carriage of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1999;43:1412–6. <https://doi.org/10.1128/aac.43.6.1412>.
- [27] Wendt C, Schinke S, Württemberger M, Oberdorfer K, Bock-Hensley O, von Baum H. Value of whole-body washing with chlorhexidine for the eradication of methicillin-resistant *Staphylococcus aureus*: a randomized, placebo-controlled, double-blind clinical trial. *Infect Control Hosp Epidemiol* 2007;28:1036–43. <https://doi.org/10.1086/519929>.
- [28] Wertheim HF, Verveer J, Boelens HA, van Belkum A, Verbruggen HA, Vos MC. Effect of mupirocin treatment on nasal, pharyngeal, and perineal carriage of *Staphylococcus aureus* in healthy adults. *Antimicrob Agents Chemother* 2005;49:1465–7. <https://doi.org/10.1128/aac.49.4.1465-1467.2005>.
- [29] Fritz SA, Camins BC, Eisenstein KA, Fritz JM, Epplin EK, Burnham CA, et al. Effectiveness of measures to eradicate *Staphylococcus aureus* carriage in patients with community-associated skin and soft-tissue infections: a randomized trial. *Infect Control Hosp Epidemiol* 2011;32:872–80. <https://doi.org/10.1086/661285>.
- [30] Gilpin DF, Small S, Bakshi S, Kearney MP, Cardwell C, Tunney MM. Efficacy of a standard methicillin-resistant *Staphylococcus aureus* decolonisation protocol in routine clinical practice. *J Hosp Infect* 2010;75:93–8. <https://doi.org/10.1016/j.jhin.2009.11.025>.
- [31] Lindgren AK, Nilsson AC, Åkesson P, Gustafsson E, Melander E. Eradication of methicillin-resistant *Staphylococcus aureus* (MRSA) throat carriage: a randomised trial comparing topical treatment with rifampicin-based systemic therapy. *Int J Antimicrob Agents* 2018;51:642–5. <https://doi.org/10.1016/j.ijantimicag.2017.08.021>.
- [32] Eum LY, Materniak S, Duffley P, El-Bailey S, Golding GR, Webster D. Randomized controlled trial of chlorhexidine gluconate, intranasal mupirocin, rifampin, and doxycycline versus chlorhexidine gluconate and intranasal mupirocin alone for the eradication of methicillin-resistant *Staphylococcus aureus* (MRSA) colonization. *J Assoc Med Microbiol Infect Dis Can* 2021;6:296–306. <https://doi.org/10.3138/jammi-2020-0049>.
- [33] Wertheim HF, Ammerlaan HS, Bonten MJ, van den Broek PJ, Troelstra A, Vandenbroucke-Grauls CM, et al. [Optimisation of the antibiotic policy in The Netherlands. XII. The SWAB guideline for antimicrobial eradication of MRSA in carriers]. *Ned Tijdschr Geneesk* 2008;152:2667–71.
- [34] Larsen AR, Stegger M, Böcher S, Sorum M, Monnet DL, Skov RL. Emergence and characterization of community-associated methicillin-resistant

- Staphylococcus aureus* infections in Denmark, 1999 to 2006. J Clin Microbiol 2009;47:73–8. <https://doi.org/10.1128/jcm.01557-08>.
- [35] Humphreys H, Grundmann H, Skov R, Lucet JC, Cauda R. Prevention and control of methicillin-resistant *Staphylococcus aureus*. Clin Microbiol Infect 2009;15:120–4. <https://doi.org/10.1111/j.1469-0691.2009.02699.x>.
 - [36] Bulanda M, Gruszka M, Heczko B. Effect of mupirocin on nasal carriage of *Staphylococcus aureus*. J Hosp Infect 1989;14:117–24. [https://doi.org/10.1016/0195-6701\(89\)90114-x](https://doi.org/10.1016/0195-6701(89)90114-x).
 - [37] Casewell MW, Hill RL. Elimination of nasal carriage of *Staphylococcus aureus* with mupirocin ('pseudomonic acid')—a controlled trial. J Antimicrob Chemother 1986;17:365–72. <https://doi.org/10.1093/jac/17.3.365>.
 - [38] Doebbeling BN, Breneman DL, Neu HC, Aly R, Yangco BG, Holley HP, et al. Elimination of *Staphylococcus aureus* nasal carriage in health care workers: analysis of six clinical trials with calcium mupirocin ointment. The Mupirocin Collaborative Study Group. Clin Infect Dis 1993;17:466–74. <https://doi.org/10.1093/clinids/17.3.466>.
 - [39] Soto NE, Vaghjimal A, Stahl-Avicoli A, Protic JR, Lutwick LI, Chapnick EK. Bacitracin versus mupirocin for *Staphylococcus aureus* nasal colonization. Infect Control Hosp Epidemiol 1999;20:351–3. <https://doi.org/10.1086/501633>.
 - [40] Fernandez C, Gaspar C, Torrellas A, Vindel A, Saez-Nieto JA, Cruzet F, et al. A double-blind, randomized, placebo-controlled clinical trial to evaluate the safety and efficacy of mupirocin calcium ointment for eliminating nasal carriage of *Staphylococcus aureus* among hospital personnel. J Antimicrob Chemother 1995;35:399–408. <https://doi.org/10.1093/jac/35.3.399>.
 - [41] Leigh DA, Joy G. Treatment of familial staphylococcal infection—comparison of mupirocin nasal ointment and chlorhexidine/neomycin (Naseptin) cream in eradication of nasal carriage. J Antimicrob Chemother 1993;31:909–17. <https://doi.org/10.1093/jac/31.6.909>.
 - [42] Martin JN, Perdreau-Remington F, Kartalija M, Pasi OG, Webb M, Gerberding JL, et al. A randomized clinical trial of mupirocin in the eradication of *Staphylococcus aureus* nasal carriage in human immunodeficiency virus disease. J Infect Dis 1999;180:896–9. <https://doi.org/10.1086/314949>.
 - [43] Doebbeling BN, Reagan DR, Pfaller MA, Houston AK, Hollis RJ, Wenzel RP. Long-term efficacy of intranasal mupirocin ointment. A prospective cohort study of *Staphylococcus aureus* carriage. Arch Intern Med 1994;154:1505–8.
 - [44] Scully BE, Briones F, Gu JW, Neu HC. Mupirocin treatment of nasal staphylococcal colonization. Arch Intern Med 1992;152:353–6.
 - [45] Reagan DR, Doebbeling BN, Pfaller MA, Sheetz CT, Houston AK, Hollis RJ, et al. Elimination of coincident *Staphylococcus aureus* nasal and hand carriage with intranasal application of mupirocin calcium ointment. Ann Intern Med 1991;114:101–6. <https://doi.org/10.7326/0003-4819-114-2-101>.
 - [46] Miller LG, Singh R, Eells SJ, Gillen D, McKinnell JA, Park S, et al. Chlorhexidine and mupirocin for clearance of methicillin-resistant *Staphylococcus aureus* colonization after hospital discharge: a secondary analysis of the changing lives by eradicating antibiotic resistance trial. Clin Infect Dis 2023;76:e1208–16. <https://doi.org/10.1093/cid/ciac402>.
 - [47] Poovelikunnel TT, Gethin G, Solanki D, McFadden E, Codd M, Humphreys H. Randomized controlled trial of honey versus mupirocin to decolonize patients with nasal colonization of methicillin-resistant *Staphylococcus aureus*. J Hosp Infect 2018;98:141–8. <https://doi.org/10.1016/j.jhin.2017.10.016>.
 - [48] Landelle C, von Dach E, Hausteint T, Agostinho A, Renzi G, Renzoni A, et al. Randomized, placebo-controlled, double-blind clinical trial to evaluate the efficacy of polyhexanide for topical decolonization of MRSA carriers. J Antimicrob Chemother 2016;71:531–8. <https://doi.org/10.1093/jac/ckv331>.
 - [49] Cluzet VC, Gerber JS, Metlay JP, Nachamkin I, Zaoutis TE, Davis MF, et al. The effect of total household decolonization on clearance of colonization with methicillin-resistant *Staphylococcus aureus*. Infect Control Hosp Epidemiol 2016;37:1226–33. <https://doi.org/10.1017/ice.2016.138>.
 - [50] Weintrob A, Bebu I, Agan B, Diem A, Johnson E, Lalani T, et al. Randomized, double-blind, placebo-controlled study on decolonization procedures for methicillin-resistant *Staphylococcus aureus* (MRSA) among HIV-infected adults. PLOS ONE 2015;10:e0128071. <https://doi.org/10.1371/journal.pone.0128071>.
 - [51] Ellis MW, Griffith ME, Dooley DP, McLean JC, Jorgensen JH, Patterson JE, et al. Targeted intranasal mupirocin to prevent colonization and infection by community-associated methicillin-resistant *Staphylococcus aureus* strains in soldiers: a cluster randomized controlled trial. Antimicrob Agents Chemother 2007;51:3591–8. <https://doi.org/10.1128/aac.01086-06>.
 - [52] Dryden MS, Dailly S, Crouch M. A randomized, controlled trial of tea tree topical preparations versus a standard topical regimen for the clearance of MRSA colonization. J Hosp Infect 2004;56:283–6. <https://doi.org/10.1016/j.jhin.2004.01.008>.
 - [53] Mody L, Kauffman CA, McNeil SA, Gaclecki AT, Bradley SF. Mupirocin-based decolonization of *Staphylococcus aureus* carriers in residents of 2 long-term care facilities: a randomized, double-blind, placebo-controlled trial. Clin Infect Dis 2007;37:1467–74. <https://doi.org/10.1086/379325>.
 - [54] Dupeyron C, Campillo B, Bordes M, Faubert E, Richardet JP, Mangeney N. A clinical trial of mupirocin in the eradication of methicillin-resistant *Staphylococcus aureus* nasal carriage in a digestive disease unit. J Hosp Infect 2002;52:281–7. <https://doi.org/10.1053/jhin.2002.1287>.
 - [55] Robicsek A, Beaumont JL, Thomson RB, Govindarajan G, Peterson LR. Topical therapy for methicillin-resistant *Staphylococcus aureus* colonization: impact on infection risk. Infect Control Hosp Epidemiol 2009;30:623–32. <https://doi.org/10.1086/597550>.
 - [56] Nelson RE, Jones M, Rubin MA. Decolonization with mupirocin and subsequent risk of methicillin-resistant *Staphylococcus aureus* carriage in Veterans Affairs hospitals. Infect Dis Ther 2012;1:1. <https://doi.org/10.1007/s40121-012-0001-3>.
 - [57] Bradley SF. MRSA colonisation (eradicating colonisation in people without active invasive infection). BMJ Clin Evid 2015;2015: 0923.
 - [58] Kniehl E, Becker A, Forster DH. Bed, bath and beyond: pitfalls in prompt eradication of methicillin-resistant *Staphylococcus aureus* carrier status in healthcare workers. J Hosp Infect 2005;59:180–7. <https://doi.org/10.1016/j.jhin.2004.06.016>.
 - [59] Poovelikunnel T, Gethin G, Humphreys H. Mupirocin resistance: clinical implications and potential alternatives for the eradication of MRSA. J Antimicrob Chemother 2015;70:2681–92. <https://doi.org/10.1093/jac/ckv169>.
 - [60] Hetem DJ, Bonten MJ. Clinical relevance of mupirocin resistance in *Staphylococcus aureus*. J Hosp Infect 2013;85:249–56. <https://doi.org/10.1016/j.jhin.2013.09.006>.
 - [61] Hayden MK, Lolans K, Haffenreffer K, Avery TR, Kleinman K, Li H, et al. Chlorhexidine and mupirocin susceptibility of methicillin-resistant *Staphylococcus aureus* isolates in the REDUCE-MRSA trial. J Clin Microbiol 2016;54:2735–42. <https://doi.org/10.1128/jcm.01444-16>.
 - [62] Hamson C, Bignardi GE. MRSA decolonization with Prontoderm compared with chlorhexidine and mupirocin. J Hosp Infect 2010;75:142–3. <https://doi.org/10.1016/j.jhin.2010.01.009>.
 - [63] Simor AE, Phillips E, McGeer A, Konvalinka A, Loeb M, Devlin HR, et al. Randomized controlled trial of chlorhexidine gluconate for washing, intranasal mupirocin, and rifampin and doxycycline versus no treatment for the eradication of methicillin-resistant *Staphylococcus aureus* colonization. Clin Infect Dis 2007;44:178–85. <https://doi.org/10.1086/510392>.
 - [64] Chang SC, Hsieh SM, Chen ML, Sheng WH, Chen YC. Oral fusidic acid fails to eradicate methicillin-resistant *Staphylococcus aureus* colonization and results in emergence of fusidic acid-resistant strains. Diagn Microbiol Infect Dis 2000;36:131–6. [https://doi.org/10.1016/s0732-8893\(99\)00116-9](https://doi.org/10.1016/s0732-8893(99)00116-9).
 - [65] Parras F, Guerrero MC, Bouza E, Blázquez MJ, Moreno S, Menarguez MC, et al. Comparative study of mupirocin and oral co-trimoxazole plus topical fusidic acid in eradication of nasal carriage of methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 1995;39:175–9. <https://doi.org/10.1128/aac.39.1.175>.
 - [66] Muder RR, Boldin M, Brennen C, Hsieh M, Vickers RM, Mitchum K, et al. A controlled trial of rifampicin, minocycline, and rifampicin plus minocycline for eradication of methicillin-resistant *Staphylococcus aureus* in long-term care patients. J Antimicrob Chemother 1994;34:189–90. <https://doi.org/10.1093/jac/34.1.189>.
 - [67] Walsh TJ, Standiford HC, Reboli AC, John JF, Mulligan ME, Ribner BS, et al. Randomized double-blinded trial of rifampin with either novobiocin or trimethoprim-sulfamethoxazole against methicillin-resistant *Staphylococcus aureus* colonization: prevention of antimicrobial resistance and effect of host factors on outcome. Antimicrob Agents Chemother 1993;37:1334–42. <https://doi.org/10.1128/aac.37.6.1334>.
 - [68] Peterson LR, Quick JN, Jensen B, Homann S, Johnson S, Tenquist J, et al. Emergence of ciprofloxacin resistance in nosocomial methicillin-resistant *Staphylococcus aureus* isolates. Resistance during ciprofloxacin plus rifampin therapy for methicillin-resistant *S aureus* colonization. Arch Intern Med 1990;150:2151–5.
 - [69] Fung SK, Louie M, Simor AE. Combined topical and oral antimicrobial therapy for the eradication of methicillin-resistant *Staphylococcus aureus* (MRSA) colonization in hospitalized patients. Can J Infect Dis 2002;13:287–92. <https://doi.org/10.1155/2002/567090>.
 - [70] Roccaforte JS, Bittner MJ, Stumpf CA, Preheim LC. Attempts to eradicate methicillin-resistant *Staphylococcus aureus* colonization with the use of trimethoprim-sulfamethoxazole, rifampin, and bacitracin. Am J Infect Control 1988;16:141–6. [https://doi.org/10.1016/0196-6553\(88\)90024-7](https://doi.org/10.1016/0196-6553(88)90024-7).
 - [71] Asensio A, Guerrero A, Quereda C, Lizán M, Martínez-Ferrer M. Colonization and infection with methicillin-resistant *Staphylococcus aureus*: associated factors and eradication. Infect Control Hosp Epidemiol 1996;17:20–8. <https://doi.org/10.1086/647184>.
 - [72] Ellison III RT, Fn Judson, Peterson LC, Cohn DL, Ehret JM. Oral rifampin and trimethoprim/sulfamethoxazole therapy in asymptomatic carriers of methicillin-resistant *Staphylococcus aureus* infections. West J Med 1984;140:735–40.
 - [73] Maraha B, van Halteren J, Verzijl JM, Wintermans RG, Buiting AG. Decolonization of methicillin-resistant *Staphylococcus aureus* using oral vancomycin and topical mupirocin. Clin Microbiol Infect 2002;8:671–5. <https://doi.org/10.1046/j.1469-0691.2002.00404.x>.
 - [74] Petersen IS, Christensen JM, Zeuthen AB, Madsen PB. Danish experience of methicillin-resistant *Staphylococcus aureus* eradication with emphasis on nose-throat colonization and supplementary systemic antibiotic treatment. J Hosp Infect 2019;103:461–4. <https://doi.org/10.1016/j.jhin.2019.09.005>.
 - [75] Bagge K, Benfield T, Westh H, Bartels MD. Eradicating MRSA carriage: the impact of throat carriage and Panton-Valentine leukocidin genes on success rates. Eur J Clin Microbiol Infect Dis 2019;38:683–8. <https://doi.org/10.1007/s10096-019-03474-6>.
 - [76] Piewngam P, Khongthong S, Roekngam N, Theapparatt Y, Sunpawaravong S, Faroongsarn D, et al. Probiotic for pathogen-specific *Staphylococcus aureus* decolonisation in Thailand: a phase 2, double-blind, randomised, placebo-controlled trial. Lancet Microbe 2023;4:e75–83. [https://doi.org/10.1016/s2666-5247\(22\)00322-6](https://doi.org/10.1016/s2666-5247(22)00322-6).

- [77] Kaspar U, de Haro Sautto JA, Molinaro S, Peters G, Iidelevich EA, Becker K. The novel phage-derived antimicrobial agent HY-133 is active against livestock-associated methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2018;62:e00385-18. <https://doi.org/10.1128/aac.00385-18>.
- [78] Yendewa GA, Griffiss JM, Jacobs MR, Fulton SA, O'Riordan MA, Gray WA, et al. A two-part phase 1 study to establish and compare the safety and local tolerability of two nasal formulations of XF-73 for decolonisation of *Staphylococcus aureus*: a previously investigated 0.5mg/g viscosified gel formulation versus a modified formulation. *J Glob Antimicrob Resist* 2020;21:171–80. <https://doi.org/10.1016/j.jgar.2019.09.017>.
- [79] Nibbering PH, Göblyös A, Adriaans AE, Cordfunke RA, Ravensbergen B, Rietveld MH, et al. Eradication of methicillin-resistant *Staphylococcus aureus* from human skin by the novel LL-37-derived peptide P10 in four pharmaceutical ointments. *Int J Antimicrob Agents* 2019;54:610–8. <https://doi.org/10.1016/j.ijantimicag.2019.07.014>.
- [80] de Breij A, Riool M, Cordfunke RA, Malanovic N, de Boer L, Koning RI, et al. The antimicrobial peptide SAAP-148 combats drug-resistant bacteria and biofilms. *Sci Transl Med* 2018;10:eaat5731. <https://doi.org/10.1126/scitranslmed.aan4044>.
- [81] Clegg J, Soldaini E, McLoughlin RM, Rittenhouse S, Bagnoli F, Phogat S. *Staphylococcus aureus* vaccine research and development: the past, present and future, including novel therapeutic strategies. *Front Immunol* 2021;12:705360. <https://doi.org/10.3389/fimmu.2021.705360>.