

Destroy, create, transform and sublimate: laboratory dissociation studies on polycyclic aromatic hydrocarbons and analogues

Kamer, J.

Citation

Kamer, J. (2025, October 22). *Destroy, create, transform and sublimate: laboratory dissociation studies on polycyclic aromatic hydrocarbons and analogues*. Retrieved from https://hdl.handle.net/1887/4273690

Version: Publisher's Version

License: License agreement concerning inclusion of doctoral thesis

in the Institutional Repository of the University of Leiden

Downloaded from: https://hdl.handle.net/1887/4273690

Note: To cite this publication please use the final published version (if applicable).

IR SPECTRA OF CATIONIC 1,5,9-TRIAZACORONENE AND TWO OF ITS CATIONIC DERIVATIVES

Infrared emission features are observed towards diverse astronomical objects in the interstellar medium (ISM). Generally, the consensus is that these IR features originate from polycyclic aromatic hydrocarbons (PAHs) and are hence named aromatic infrared bands (AIBs). More recently, it has been suggested that nitrogen substituted PAHs (PANHs) contribute to the AIBs as well and it has even been shown that nitrogen inclusion in PAHs can improve the match with the AIBs, specifically around the 6.2 μ m feature. In order to determine which specific molecules or functional groups are at the origin of the AIBs, IR spectra of various PA(N)Hs are experimentally and computationally studied. In this work we expand on the spectroscopic investigation of PANHs by presenting the gas-phase mid-IR spectra of 1,5,9triazacoronene $^{\bullet+}$ (TAC $^{\bullet+}$, m/z 303), a threefold nitrogenated congener of coronene $^{\bullet+}$, its protonated derivative \mathbf{TACH}^+ (m/z 304) and the product that forms when water adds to dehydrogenated $TAC^{\bullet+}$ $[TAC-H+H_2O]^+$ (m/z 320). We analyze the mid-IR spectra by comparing them with vibrational modes calculated at the B3LYP/6-311++G(d,p) level a theory and we present a possible geometry for $[TAC-H+H_2O]^+$. The $TAC^{\bullet+}$ mid-IR spectrum is compared to that of coronene • + and is demonstrated to be remarkably similar. We put TAC^{•+} and TACH⁺ into astronomical context by comparing their recorded mid-IR spectra to observed ISM spectra of four spectral classes. From this we conclude that TAC^{•+} and TACH⁺ could both contribute to the AIBs, with TACH⁺ being a more likely contributor than $\mathbf{TAC}^{\bullet+}$.

Kamer, J., Schleier, D., Jiao, A., Schneider, G. F., Martens, J., Berden, G., Oomens, J., Bouwman, J., 2024, Physical Chemistry Chemical Physics, 26, 27912-27921.

60 4.1. Introduction

4.1 Introduction

Prominent infrared (IR) emission features at 3.3, 6.2, 7.7, 8.6, 11.2 and 12.7 μ m are observed towards a large variety of astronomical objects in the interstellar medium (ISM), such as protoplanetary disks, HII regions and reflection nebulae (Tielens 2008; Chown *et al.* 2024). There is a general consensus that these mid-IR features are caused by polycyclic aromatic hydrocarbon (PAH) molecules that cascade down to the ground state after being excited by ultraviolet (UV) photons from the interstellar radiation field (Leger & Puget 1984; Allamandola *et al.* 1985, 1989). Hence, these mid-IR features are commonly referred to as aromatic infrared bands (AIBs).

The band positions and shapes of the AIBs are fairly consistent among the various observed astronomical objects, although some variations exist. Based on these variations, the mid-IR spectra are grouped into class A through D, each of which represents an specific spectral type (Peeters et al. 2002; van Diedenhoven et al. 2004; Matsuura et al. 2014). Class A, the most commonly observed of the four, is characterized by features which peak at 6.22, 7.6 and 8.6 μ m. The 7.6 μ m feature also contains a shoulder at 7.8 μ m. A shift of these three class A features towards longer wavelengths results in class B, where the peaks are found at 6.24 - 6.28, 7.8 - 8.0 and $>8.62 \mu m$. Generally, the feature around $6.2 \mu m$ is broader for class B, compared to A. Class C contains a peak around 6.3 μ m and a broad feature ranging from 7.5 to 9.3 μ m with the peak at 8.22 μ m. Finally, class D is characterized by a feature at $6.24 \mu m$, and a broad feature ranging from 7.0 to 9.0 μm , peaking around 7.7 μm . Additionally, class D can contain a feature at 6.9 μ m. Numerous studies have demonstrated the very strong resemblance between the mid-IR emission of (ionic) homocyclic PAHs and astronomical objects (Bouwman et al. 2019, 2020, 2021; Allamandola et al. 1999; Hudgins & Sandford 1998a,b; Zhen et al. 2017, 2018; Palotás et al. 2020, 2021; Ricca et al. 2012; Mattioda et al. 2014; Mackie et al. 2015; Oomens et al. 2001, 2003; Banhatti et al. 2021; Gao et al. 2014). However, the experimental and computed IR emission features of cationic homocyclic PAHs around 6.2 μ m are consistently redshifted with respect to astronomical observations (Peeters et al. 2002; Hudgins et al. 2005).

Substitution of a carbon atom in the structure of a PAH cation by a nitrogen, forming a socalled polycyclic aromatic nitrogen heterocycle (PANH) cation, can introduce a spectral shift in the 6.2 μ m band (Hudgins et al. 2005; Ricca et al. 2021). Based on computational studies, it has been shown that this shift depends strongly on the position of the substituted carbon atom in the PAH structure and size and symmetry of the PAH. Substitution of an endoskeletal carbon atom in the PAH structure (commonly referred to as endo-PANH) results in a drastic change of the mid-IR emission spectrum with respect to its pure PAH congener, with most notably a blue-shift in the 6.2 μ m band, resulting in a better agreement with observational data (Hudgins et al. 2005; Ricca et al. 2021; Rap et al. 2022). Cationic endo-PANHs have, contrary to their cationic PAH congeners, a closed-shell electronic configuration. Recently, Ricca et al. (2021) estimated an upper limit for the fraction of cationic endo-PANH emission contributing to the astronomically observed 6.2 µm band to be about 12%, by assuming all of the 11.0 μ m emission is due to singly ionized endo-PANHs and combining their results with the reported values for NGC 7023 (Boersma et al. 2013). Exoskeletal PANHs (exo-PANHs), where a peripheral CH-group has been substituted by a nitrogen, do not show a significant change in the mid-IR range compared to their pure, cationic PAH analogue (Hudgins et al. 2005; Ricca et al. 2021; Schleier et al. 2024b). This makes exo-PANH cations challenging to distinguish from their PAH counterpart using IR spectra alone. Therefore, exo-PANHs are potential carriers of the observed mid IR features as well and potentially make up a large fraction of the PAH population.

Although nitrogen-bearing PAHs have been shown to be astronomically relevant (Hudgins et al. 2005; Ricca et al. 2021; McGuire et al. 2021), the number of experimental works studying (spectral) characteristics of (ionic) PANHs is limited. Moreover, the works that do report on



Figure 4.1: Molecular structure of cationic 1,5,9-triazacoronene (**TAC**^{•+}, C₂₁H₉N₃^{•+}).

nitrogen-bearing PAHs are often limited to species containing only a single nitrogen in the aromatic structure. For example, Ricca *et al.* (2021) studied PAHs up to 103 carbon atoms while mainly considering species containing a single nitrogen atom. Based on the interstellar abundances of elements, PAHs of astronomically relevant sizes could contain multiple nitrogen atoms as well, and thus spectroscopic studies of such species are needed.

The proton affinity (PA) of especially exo-PANHs is greatly enhanced relative to the analogous homocyclic PAHs. For example, the PA of naphthalene is determined to be 802.9 kJ mol⁻¹ (Hunter & Lias 1998), while its nitrogenated congeners quinoline and isoquinoline have a PA of 953.2 and 951.7 kJ mol⁻¹ (Hunter & Lias 1998), respectively. An exposed N-atom is a strong nucleophile that easily binds a proton, which are abundantly present in HII regions, resulting in the formation of very stable, closed-shell, protonated PANH species. Experimental and computational works have shown that protonated PANHs are potential carriers of the 6.2 μm band (Ricca et al. 2021; Alvaro Galué et al. 2010). Ricca et al. (2021) calculated the 6.2 μ m band position of protonated PANHs to span the 6.17 to 6.45 μ m range, depending on the size and symmetry of the protonated PANH. Singly dehydrogenated exo-PANH cations have been shown to be highly reactive. In a previous study by our group, we showed that after losing a hydrogen atom from the radical cation of 1,5,9-triazacoronene ($\mathbf{TAC}^{\bullet+}$, m/z 303, see Fig. 4.1 for the molecular structure), a species is formed that reacts very efficiently with background water vapor in the trap to form a very stable product ion (Schleier et al. 2024a). It is important to spectroscopically characterize exo-PANH radical cations, protonated exo-PANHs and even the reaction products formed in the reaction of exo-PANHs and water to determine their potential contribution to the interstellar mid-IR emission features.

In this work, we study the IR spectra of $\mathbf{TAC}^{\bullet+}$, protonated \mathbf{TAC} (\mathbf{TACH}^+ , m/z 304) and the product that forms when water adds to dehydrogenated $TAC^{\bullet+}$ $[TAC-H+H_2O]^+$ (m/z 320), using a combination of infrared multiple photon dissociation (IRMPD) spectroscopy at the Free Electron Laser for Infrared experiments (FELIX) and quantum chemical calculations. TAC is a threefold nitrogenated congener of coronene and its three available nitrogen sites allow for facile \mathbf{TACH}^+ formation and the formation of $[\mathbf{TAC} - \mathbf{H} + \mathbf{H_2O}]^+$ in the presence of H₂O. We first present the dissociation mass spectra that arise upon resonantly exciting TAC^{•+}, TACH⁺ and [TAC-H+H₂O]⁺ with IR radiation from FELIX. IR spectra are subsequently constructed from the frequency dependent dissociation mass spectra. The experimental spectra are analyzed using density functional theory (DFT) simulated spectra to characterize the vibrational normal modes. Next, we compare the experimental spectrum of $TAC^{\bullet+}$ to the coronene cation $(Cor^{\bullet+})$ to investigate the effect on the IR characteristics of triple nitrogen substitution in $\mathbf{Cor}^{\bullet+}$. Lastly, we put the ramification of nitrogen inclusion on the IR spectra in astrophysical context by comparing the TAC^{•+} and TACH⁺ IRMPD spectra to observed astronomical mid-IR spectra of each spectral type class.

62 4.2. Methods

4.2 Methods

4.2.1 Experimental

All IRMPD spectra were recorded on a commercially available Bruker Amazon Speed ETD 3D quadrupole ion trap mass spectrometer, which is modified to allow the radiation from the free electron laser for infrared experiments, FELIX, to be focused onto the ion cloud inside the trap. A detailed description of the system can be found in previous publications (Martens et al. 2016; Palotás et al. 2021) and a short description of the apparatus and the measurement procedures is provided here.

TAC was synthesized following the procedure described by Tan *et al.* (2016) and a detailed description is given in the appendix of **Chapter 5**. A glass direct insertion probe (DIP) was coated with the solid sample and subsequently introduced into an atmospheric pressure chemical ionization source (Bruker APCI II) which was connected to the mass spectrometer. The DIP is then gently heated to desorb the sample material. A flow of N_2 carries the desorbed molecules into a corona discharge region, where $\mathbf{TAC}^{\bullet+}$, \mathbf{TACH}^+ and $[\mathbf{TAC-H+H_2O}]^+$ are produced. These are subsequently guided through a spray shield into a glass capillary leading to the ion trap. The ions are mass-isolated using the mass spectrometer's tandem mass spectrometry features.

The trapped and mass isolated ions are exposed to five macropulses of mid-infrared (mid-IR) light originating from the free electron laser that operates at a 10 Hz repetition rate. Each macropulse has an average duration of 6 μ s with a maximum pulse energy of 75 mJ and a spectral full width at half maximum (FWHM) of about 5 – 10 cm⁻¹. The IR frequency is tuned from 730 to 1780 cm⁻¹ in steps of 5 cm⁻¹ and at each wavelength five mass spectra are averaged. The precursor ion absorbs multiple IR photons when the laser is tuned to a frequency that is resonant with a vibrational transition, which eventually leads to dissociation of the precursor ion. The resulting fragment ion signals and the remaining precursor ion signal are integrated and an IRMPD dissociation yield is calculated. The wavelength-dependent IRMPD yield is linearly corrected for free electron laser pulse energy, resulting in the IRMPD spectrum of our studied species.

4.2.2 Computational

Quantum chemical calculations of our species of interest were performed using the Gaussian16 (Frisch et al. 2016) suite of software. We employed the B3LYP functional (Becke 1993; Lee et al. 1988) in combination with the 6-311++G(d,p) basis set to calculate harmonic vibrational spectra. All computed normal mode frequencies are uniformly scaled with a factor of 0.9679 (Andersson & Uvdal 2005) to correct for anharmonicity effects. The stick spectrum is convolved with a 20 cm $^{-1}$ FWHM Gaussian function to facilitate comparison with the experimental IRMPD spectra.

4.3 Results

4.3.1 Mass spectrometry

Figure 4.2 shows the mass spectra obtained by irradiating the isolated $\mathbf{TAC}^{\bullet+}$, \mathbf{TACH}^{+} and $[\mathbf{TAC}-\mathbf{H}+\mathbf{H_2O}]^{+}$ ions. The top and bottom panel of each sub-figure represent mass spectra resulting from off- and on-resonance irradiation, respectively. From these figures it can be seen that clean isolation of the precursor ion is achieved and significant dissociation is induced when resonantly exciting the ions. Interestingly, the dissociation fragments that arise are significantly different for each of the precursor ions. Surprisingly, both $\mathbf{TAC}^{\bullet+}$ and

TACH⁺ show m/z 320 as a product after on-resonant irradiation. Likely, this product forms when H₂O binds with [**TAC-H**]⁺ (Schleier *et al.* 2024a). Section 4.3.4 will describe this in more detail. The mass spectra will not be analyzed further in this work, as it is outside of the scope of this study. The dissociation of **TAC**^{•+} using 630 nm radiation, however, has been described in much detail in a previous work by our group (Schleier *et al.* 2024a).

4.3.2 TAC^{•+} IRMPD spectroscopy

The IRMPD spectrum that is constructed by plotting the laser power-corrected dissociation yield of $\mathbf{TAC}^{\bullet+}$ as a function of IR frequency between 750 and 1750 cm⁻¹ is presented as the black trace in Fig. 4.3a. Shown together with the IRMPD is the scaled B3LYP/6-311++G(d,p) computed vibrational spectrum of $\mathbf{TAC}^{\bullet+}$, depicted by the blue vertical lines and the convolution in green. Neutral \mathbf{TAC} can be optimized at a C_{3h} symmetry and a molecular orbitals (MO) calculation of neutral \mathbf{TAC} , reveals that the highest occupied MO (HOMO), with E" orbital symmetry, is degenerate. The radical cation is therefore expected to undergo Jahn-Teller distortion. Indeed, optimizing the geometry of $\mathbf{TAC}^{\bullet+}$ at a C_{3h} symmetry, similar to neutral \mathbf{TAC} , resulted in an imaginary frequency. Optimization at the Jahn-Teller distorted C_s symmetry does converge with all real frequencies, and hence the $\mathbf{TAC}^{\bullet+}$ vibrational modes presented are calculated in this symmetry.

Six broad features at 803, 932, 1145, 1189, 1352 and 1553 cm⁻¹ can be distinguished from the IRMPD spectrum. The peak at 1352 cm⁻¹ is the broadest and also the most intense feature. According to our calculations (see Table 4.1 for computed modes), this most intense feature originates mainly from two modes which primarily consist of in-plane CH bending combined with some CC and CN stretching character at 1333 and 1346 cm⁻¹. The computed IR spectrum shows noticeable involvement of at least one of the nitrogen atoms in almost all vibrational modes, of which the most intense are the modes at 781 and 1542 cm⁻¹. This is in contrast with 3-azafluoranthene, where the nitrogen atom was shown to only be involved in a few specific modes (Schleier *et al.* 2024b).

The overall shape of the experimental spectrum is quite well replicated by the convolved calculated spectrum, but the relative intensities do not always match. This effect is sometimes observed, especially for systems with high dissociation thresholds, such as PAHs (Bouwman et al. 2019, 2020; Alvaro Galué et al. 2010; Bouwman et al. 2016; Schleier et al. 2024b; Oomens et al. 2000, 2001, 2003), and likely is the result of several effects, such as nonlinearities in the multiple-photon excitation in combination with the linear power correction applied (Oomens et al. 2001). Furthermore, there may be an apparent intensity cut-off below which the molecules cannot be excited to above the dissociation threshold with the employed laser powers (Munshi et al. 2017, 2020; Berden et al. 2019) (see section 4.3.3) and moreover, closely spaced bands in the spectrum may give rise to co-operative effects in the excitation that are absent in a (computed) absorption spectrum (Oomens et al. 2003; Parneix et al. 2013). In addition, there may be flaws in the computed spectra, which would require further computational investigation beyond the scope of this study. The scaled harmonic modes in the 1100 to 1400 cm⁻¹ spectral range do match quite well with the experimental spectrum. A plateau can be observed between 970 and 1100 cm⁻¹. Calculations show three harmonic modes at 983, 1003 and 1107 cm⁻¹ that do not correspond to an experimental IRMPD feature. In the observed spectrum, these features are likely blended, thus resulting in a featureless plateau. The IRMPD feature at 803 cm⁻¹ seems to consist mainly of two modes calculated at 781 and 813 cm⁻¹, albeit the feature is slightly blue-shifted with respect to the calculated 781 cm⁻¹ vibration. The first calculated mode originates from overall in-plane CC and CN stretching, and CH bending, while the latter originates mainly from out-of-plane CH bending and minor out-of-plane CC and CN stretching.

64 4.3. Results




Figure 4.2: Mass spectra of isolated $TAC^{\bullet+}$ (a), $TACH^+$ (b) and $[TAC-H+H_2O]^+$ (c) after exposure to off-resonance (top panel) and on-resonance (bottom panel) infrared radiation. The wavenumber of the IR radiation is depicted in each respective panel. The proposed geometry for the $[TAC-H+H_2O]^+$ species is discussed in section 4.3.4.

CHAPTER 4 65


Table 4.1: Experimentally measured band positions for TAC^{.+} shown together with the computed infrared intensity and symmetry. The computed modes are grouped to match experimental data, where possible.

Experimental	$Calculated^{a,b}$		
Pos.	Pos.	Int.	Carronnonatura
(cm^{-1})	$({\rm cm}^{-1})$	(km mol^{-1})	Symmetry
803	781	224	A'
	813	70	A''
932	931	33	A''
	983	20	A'
$-^c$	1003	44	A'
	1107	21	A'
1145	1128	56	A'
	1178	65	A'
1189	1191	46	A'
	1214	34	A'
	1289	39	A'
	1305	41	A'
	1333	147	A'
1352	1346	241	A'
	1389	54	A'
	1399	39	A'
	1429	26	A'
_	1467	41	A'
	1492	25	A'
1553	1512	27	A'
	1528	57	A'
	1534	71	A'
	1542	127	A'

^a The computed band positions are scaled with a factor of 0.9679. ^b Only computed modes with intensities \geq 20 km mol⁻¹ are listed.

^c A plateau is observed in the experimental data without resolved features.

66 4.3. Results






Figure 4.3: Experimental IRMPD (black) and simulated IR (blue vertical lines) spectra of $TAC^{\bullet+}$ (a) and $TACH^+$ (b). The B3LYP/6-311++G(d,p) computed spectrum (blue sticks) is uniformly scaled with a factor of 0.9679 and convolved with a 20 cm⁻¹ FWHM Gaussian function (green shaded area). The IRMPD intensity is represented by the left axis and the intensity of the computed spectra by the right axis. All spectra are normalized to their most intense peak. The horizontal grey, dotted line in (b) indicates an apparent threshold of 21 km mol⁻¹. Exciting modes with an intensity below line does not allow the molecule to reach the dissociation threshold.

4.3.3 TACH⁺ IRMPD spectroscopy

The APCI source on the mass spectrometer not only generated $\mathbf{TAC}^{\bullet +}$, but also protonated \mathbf{TAC} , \mathbf{TACH}^{+} . The black line in Fig. 4.3b corresponds to the \mathbf{TACH}^{+} IRMPD spectrum. Due to the high proton affinity of the nitrogen atom in the aromatic system (Ricca *et al.* 2021; Maclagan *et al.* 2015), it is expected that the proton will bind on one of the three available nitrogen atoms of \mathbf{TAC} . We optimized this geometry and calculated the vibrational spectrum and scaled them to account for anharmonicities. The calculated position and intensity of the \mathbf{TACH}^{+} modes are listed in Table 4.2 and depicted in Fig. 4.3b by the blue vertical sticks and their convolved spectral profile in green.

Features around 807, 1130, 1296, 1335, 1414, 1546 and 1597 cm⁻¹ can be identified in the IRMPD spectrum of **TACH**⁺, the most intense of these being at 1597 cm⁻¹ and 1296 cm⁻¹. According to our calculations the first feature mainly arises from three vibrational modes at 1593, 1595 and 1612 cm⁻¹. The modes at 1593 and 1612 cm⁻¹ originate mostly from in-plane NH and CH bending, with CC stretching and some CN stretching character. Contrarily, the 1595 cm⁻¹ mode involves barely any motion of the NH group and has mostly CC-stretching and in-plane CH bending mode character. The latter feature is calculated to consist of two modes at 1291 and 1299 cm⁻¹. The 1291 cm⁻¹ mode originates from in-plane CC and CN stretching, and CH and NH bending. This is comparable to the 1593 and 1612 cm⁻¹ calculated modes although the CC stretching motion is less pronounced. The 1299 cm⁻¹ mode consists mainly of in-plane CC and CN stretching, and CH bending. Furthermore, this mode involves barely any NH bending.

The **TACH**⁺ IRMPD displays no clear features between 820 and 1260 cm⁻¹ (see Fig. 4.7 in the appendix for a zoom-in), except for the 1130 cm⁻¹ feature, while modes are predicted to fall in this range. At these frequencies, the calculated intensities of the **TACH**⁺ vibrational normal modes are less than 21 km mol⁻¹, which apparently is not enough to drive the molecule to its dissociation threshold using IRMPD. This has been observed in earlier works as well and is not specific to PANHs (Munshi *et al.* 2017, 2020; Berden *et al.* 2019). **TAC**^{•+} does not seem to have a clear dissociation threshold, likely due to the higher intensity modes compared to **TACH**⁺. The intensity reduction in this wavenumber range of the IRMPD spectrum is also observed in an earlier work comparing IRMPD spectra of quinoline • and isoquinoline • to their protonated counterparts (Alvaro Galué *et al.* 2010), with these protonated species showing no distinct bands in the region from ~820 to ~1300 cm⁻¹. The calculated modes

of **TACH**⁺ in the 900 to 980 cm⁻¹ are dominated by out-of-plane CH bending. In-plane CC stretching and, CH and NH bending modes dominate the IR activity between 980 and 1000 cm⁻¹. Within the 1100 to 1260 cm⁻¹ range the calculated modes show mainly in-plane CH and NH bending character with minor CC stretching contribution.

Table 4.2: Experimentally measured band positions for **TACH**⁺ shown together with the computed infrared intensity and symmetry. The computed modes are grouped with respect to the matching experimental data, when possible.

Experimental	Calculated a,b		
$ \begin{array}{c} \text{Pos.} \\ \text{(cm}^{-1}) \end{array} $	$ \begin{array}{c} \text{Pos.} \\ \text{(cm}^{-1}) \end{array} $	Int. (km mol^{-1})	Symmetry
807	801 814	54 68	A" A"
1130	1126	22	A'
_	1155	20	A'
1296	1291 1299	54 43	$\begin{array}{c} A' \\ A' \end{array}$
1335	1326 1340 1357	27 109 35	A' A' A'
1414	1421 1432	36 28	$\begin{array}{c} A' \\ A' \end{array}$
_	1461	26	A'
1546	1540	36	A'
1597	1593 1595 1612	60 76 133	A' A' A'

^a The computed band positions are scaled with a factor of 0.9679.

Besides the range spanning 820 to 1260 cm⁻¹, the agreement between the computed (and scaled) normal modes and the IRMPD spectrum is generally good. Especially, the calculated frequencies match the measured IRMPD bands at 807, 1296, 1335, 1546 and 1597 cm⁻¹ well. A discrepancy can be observed for the calculated feature at around 1461 cm⁻¹, which has no distinct experimental counterpart. Also, the calculated spectrum shows a small feature at 762 cm⁻¹, which is not observed in the experimental spectrum, presumably because the dissociation threshold is not reached when exciting the molecule at this frequency.

4.3.4 [TAC-H+H₂O]⁺ IRMPD spectroscopy

While we attempted to record an IRMPD spectrum of $[\mathbf{TAC-H}]^+$ at m/z 302, we found that a peak at m/z 320 dominates the mass spectrum. This indicates that the loss of a hydrogen atom significantly increases the reactivity of the ion, causing it to react with background $\mathrm{H_2O}$ in the system to form $[\mathbf{TAC-H+H_2O}]^+$. Removing a hydrogen from the CH group that is located next to the nitrogen atom requires the least amount of energy (see Fig. 4.8

^b Only modes with intensities > 20 km mol⁻¹ are listed.

68 4.3. Results




Figure 4.4: Experimental IRMPD spectrum (black) and simulated IR spectra of four different isomers of $[TAC-H+H_2O]^+$. Also shown are the various structures of $[TAC-H+H_2O]^+$ isomers and their relative energies calculated using the B3LYP functional in combination with the 6-311++G(d,p) basis set. The hydrogen atoms that differ between the isomers are highlighted in cyan for clarity. The simulated spectrum in green indicates our tentative assignment that isomer 5 is likely the dominant isomer. The B3LYP/6-311++G(d,p) computed stick spectra are uniformly scaled with a factor of 0.9679. The calculated spectra are also convolved with a 20 cm⁻¹ FWHM Gaussian function. All spectra are normalized with respect to their most intense peak in the wavenumber range 750-1650 cm⁻¹.

in the appendix), and so the $\rm H_2O$ will likely bind to that site. Our previous work has shown that the resulting molecule does not dissociate after irradiation with intense light at 630 nm (Schleier *et al.* 2024a). However, the radiation from FELIX did induce dissociation when tuned to a vibrational resonance and, hence, an IRMPD spectrum could be recorded. The resulting spectrum is depicted as black lines in Fig. 4.4.

Many features can be seen in the IRMPD spectrum of which the more pronounced features are broad bands at 808, 1177, 1319, 1582 and 1718 cm⁻¹. The high wavenumber shoulder of the 1319 cm⁻¹ feature seems to contain subtle features at 1373, 1408, 1452 and 1487 cm⁻¹. Similarly, towards the lower wavenumber side of the 1582 cm⁻¹ feature, a peak can be observed at 1539 cm⁻¹. The broad 1177 cm⁻¹ feature seems to consist of multiple vibrational modes

A lead towards the most likely geometry of $[TAC-H+H_2O]^+$ can be found in the feature at 1718 cm⁻¹, which is indicative of the presence of a carbonyl (C=O) group (Francis 1951; Fuson *et al.* 1954). However, this does not exclude other functional groups, such as a

hydroxyl (OH) group, as an IRMPD spectrum can be comprised out of several isomers. A total of 16 plausible isomers were computationally investigated for comparison to the IRMPD spectrum. The geometries of the isomers are optimized and their respective vibrational normal modes are calculated, scaled and convolved with a Gaussian profile. The resulting structures of five plausible isomers, their relative energies and simulated IR spectra calculated using B3LYP/6-311++G(d,p) are shown in Fig. 4.4. The calculated spectra of the other isomers can be found in Fig. 4.9 in the appendix.

Isomer 1, 4 and 5 contain a C=O group and in isomers 4 and 5 the amide nitrogen is bound to a hydrogen. The remaining hydrogen is bound to one of the other nitrogen atoms. Isomer 1 has no hydrogen bound to the amide nitrogen, and both hydrogens are bound to the other two nitrogen atoms. Isomer 2 and 3 have an OH group bound to the bare carbon that forms upon hydrogen loss from TAC*. Both these isomers have no hydrogen bound to the nitrogen adjacent to the COH moiety and only differ, similar to isomer 4 and 5, by a hydrogen being bound to a different nitrogen atom relative to the OH group. Isomer 5 is found to be the lowest energy isomer and the calculated energies of the other isomers, with respect to this isomer, are 3.1, 18.2, 21.2 and 106.1 kJ/mol for isomer 4, 3, 2 and 1, respectively.

Comparing the calculated IR spectra of our isomers to the 750-1650 cm⁻¹ range of the IRMPD spectrum, does not allow for an unambiguous assignment of one specific isomer. Isomers **2**, **3** and **5** all match quite well with the experimental spectrum, albeit some intensity mismatches are noted. Only isomers **1**, **4** and **5** can account for the C=O stretching feature observed at 1718 cm⁻¹. The calculated C=O stretching modes for these isomers lay at 1673, 1724 and 1716 cm⁻¹, respectively. Despite having a carbonyl group, isomer **1** does not match the feature at 1718 cm⁻¹, thus indicating that the carbonyl group is likely neighboured by an NH group.

Based on this and the match in the 750-1650 cm⁻¹ range, we tentatively assign isomer 5 to be the most likely candidate for [TAC-H+H₂O]⁺. However, this does not rule out the possibility of several other isomers contributing to the [TAC-H+H₂O]⁺ IRMPD spectrum. Since isomer 4 and 5 are close in energy, we estimated the contribution of both isomers to the IRMPD spectrum at 300 K from the Boltzmann distribution based on their energies obtained from our DFT calculations. This showed a contribution of 22% and 78% isomer 4 and 5, respectively, thus indicating that isomer 4 is a potentially significant component in the IRMPD spectrum. However, this estimation should be treated with care as the energies were calculated using DFT. Perhaps additional measurements could narrow the number of possible contributing isomers. For example, since OH stretching is expected to result in a band around 3600 cm⁻¹ (Palotás et al. 2022), an IRMPD measurement in this range might help to confirm or exclude the contribution of the enol isomers.

4.4 Discussion and Astrophysical implication

4.4.1 Comparison of the TAC^{•+} IRMPD to coronene^{•+}

Comparing the IRMPD spectrum of **TAC**^{•+} to its non-nitrogen bearing analogue coronene ^{•+} (**Cor**^{•+}), measured by Oomens *et al.* (2001) in 2001 using the same IRMPD technique on a different ion trap mass spectrometer system gives direct insight into the spectral shift caused by the inclusion of three peripheral nitrogen atoms. Fig. 4.5 shows the IRMPD spectra of **Cor**^{•+} and **TAC**^{•+}.

From the comparison of the spectra in the spectral region between 1050 and 1700 cm⁻¹ it is clear that there is remarkable similarity in band positions between the experimental **TAC**^{•+} and **Cor**^{•+} spectra. All of the **TAC**^{•+} features in this region fall within about 3% of the position of the **Cor**^{•+} features. Some clear differences between the two spectra are




Figure 4.5: Experimental IRMPD of **TAC**^{•+} (black) and coronene^{•+} (green). Both spectra are normalized with respect to their most intense peak. The IRMPD spectrum of coronene^{•+} is taken from Oomens *et al.* (2001)

noted in the 750-1050 cm⁻¹ region. The **Cor**•+ spectrum only exhibits a single feature at 849 cm⁻¹, which is attributed to the out-of-plane CH bending mode (Oomens *et al.* 2001; Hudgins & Allamandola 1999), while **TAC**•+ exhibits two bands at 803 and 932 cm⁻¹. For the 932 cm⁻¹ **TAC**•+ feature no counterpart can be identified in the **Cor**•+ spectrum, as this feature is mainly inherent to a single out-of-plane CH bending mode neighbouring a nitrogen atom. Despite this, the spectra are thus remarkably similar. This is in agreement with earlier works predicting a minor impact on the overall IR spectrum after nitrogen atom incorporating on the periphery of a pure carbon PAH (Hudgins *et al.* 2005; Ricca *et al.* 2021).

4.4.2 Comparison of the TAC^{•+} and TACH⁺ IRMPD spectra to interstellar emission bands

In order to assess the interstellar relevance of $\mathbf{TAC}^{\bullet+}$ and \mathbf{TACH}^{+} , we compare the IRMPD spectra presented in this work to observed astronomical mid-IR spectra of each of the four spectral classes (Peeters *et al.* 2002; van Diedenhoven *et al.* 2004; Matsuura *et al.* 2014; Bouwman *et al.* 2019). Representative spectra of class A through D in the 6 to 10 μ m range are shown together with the experimental IRMPD spectra of $\mathbf{TAC}^{\bullet+}$ and \mathbf{TACH}^{+} in Fig. 4.6.

When comparing the IRMPD of $\mathbf{TAC}^{\bullet+}$ to the observed astronomical spectra, we find that $\mathbf{TAC}^{\bullet+}$ does not match well with many of the observed bands in the class A and B spectra, but may still contribute to some extent. Moreover, the modes at 8.41 and 8.73 μ m (1189 and 1145 cm⁻¹) can potentially contribute to the shoulders of the broad features in the class C and D spectra. This supports the hypothesis that cationic exo-PANHs can contribute to the AIBs as well. Thus, exo-PANH and PAH cations should both be considered to be contributors to interstellar IR emission bands.

Protonated PANHs have been proposed as possible carriers of the 6.2 μ m interstellar emission band (Ricca et al. 2021; Alvaro Galué et al. 2010). Ricca et al. (2021) calculated that these species can give rise to modes in the 6.17 to 6.45 μ m range, although, these calculations were performed on molecules containing $N_C \geq 51$. From our data we find that protonation of




Figure 4.6: a) Mid-IR spectra of astronomical representative sources of type A (IRAS 23133+6050; olive), type B (HD 44179; green), type C (IRAS 13416-6243; blue), and type D (IRAS 05110-6616; purple). All are normalized to the maximum intensity of the 7–8 μm complex and with continuum subtracted. The type D spectrum was recorded using Spitzer/IRS (Werner et al. 2004; Houck et al. 2004) and the other classes are obtained from ISO/SWS (Kessler et al. 1996; de Graauw et al. 1996) observations, as calibrated by Sloan et al. (2003). b) Experimental IRMPD of $\mathbf{TAC}^{\bullet+}$ (black) and \mathbf{TACH}^{+} (blue). Both spectra are normalized with respect to their most intense peak. The vertical grey, dashed and blue, dot-dashed lines represent the peaks of the most clear features in the IRMPD spectra of $\mathbf{TAC}^{\bullet+}$ and \mathbf{TACH}^{+} , respectively.

72 4.5. Conclusions

TAC results in an IRMPD spectrum with a feature that matches very well with the 6.2 μ m interstellar feature in all four classes and suggests medium-sized protonated exo-PANHs may contribute to the 6.2 μ m AIB feature. Furthermore, **TACH**⁺ has two intense bands at 7.49 and 7.72 μ m (1335 and 1296 cm⁻¹), which both match well with the 7.6 μ m feature of the class A spectrum. The match with the class A spectrum indicates that **TACH**⁺ could be a contributor to the AIBs, therefore suggesting that protonated exo-PANHs are possible carriers of the AIBs.

Based on the match/mismatch with the AIBs it seems that protonated exo-PANHs are more likely contributors than the cationic exo-PANHs. From a physical point of view this is also supported by the fact that exo-PANH radical cations are open-shell species, while protonated exo-PANHs are closed-shell species, making exo-PANH radical cations more reactive with other abundant atoms/molecules such as hydrogen as compared to protonated exo-PANHs. This possibly increases the fraction of interstellar protonated exo-PANH cations at the expense of cationic exo-PANHs.

4.4.3 Possible astronomical relevance of oxygen-functionalized PA(N)Hs

Weaker emission features between 5.7 and 5.9 μ m are observed towards various objects in the ISM (Tielens 2008; Chown et al. 2024). These features, however, are generally ignored due to their low intensity. The peak in the IRMPD spectrum of [TAC-H+H₂O]⁺ at 5.82 μ m (1718 cm⁻¹) shows that the carbonyl stretch of oxygen-functionalized exo-PANH cations is a possible carrier of the weak AIB features between 5.7 and 5.9 μ m. Previous studies on oxygen-functionalized PAHs concluded the C=O stretch to be at 6.01 μ m (1665 cm⁻¹) (Bernstein et al. 1999) and around 5.88 μ m (1700 cm⁻¹) (Pendleton & Allamandola 2002; Matrajt et al. 2005). The precise position of carbonyl stretching is thus influenced by the molecule and its chemical environment. However, it is also argued that these features could originate from a blend of CH stretching and in- and out-of-plane CH bending of pure PAHs (Boersma et al. 2008). Possibly, the further (IR) investigation of oxygen-functionalized PA(N)Hs might shed light on a clear origin of the 5.7–5.9 μ m AIB features, which could contain valuable information on the atomic and molecular environment of interstellar objects.

4.5 Conclusions

The gas-phase infrared spectra of 1,5,9-triazacoronene $^{\bullet+}$ (TAC $^{\bullet+}$), the protonated TAC (TACH $^+$) and the water-addition species [TAC $^-$ H $^+$ H $_2$ O] $^+$, obtained with IRMPD spectroscopy, are presented. TAC $^{\bullet+}$ shows six clear features at 803, 932, 1145, 1189, 1352 and 1553 cm $^{-1}$, with the 1352 cm $^{-1}$ feature being the broadest and also the most intense. The IRMPD of TACH $^+$ contains seven features at 807, 1130, 1296, 1335, 1414, 1546 and 1597 cm $^{-1}$. Of which the most intense feature is situated at 1597 cm $^{-1}$ (6.26 μ m), which is akin to the IR spectra of other protonated PANHs (Alvaro Galué *et al.* 2010; Hudgins *et al.* 2005; Ricca *et al.* 2021). This supports the argument of protonated PANHs being possible carriers of the interstellar 6.2 μ m band. [TAC $^-$ H $^+$ H $^-$ QO] $^+$ exhibits many features of which the most pronounced are peaks at 808, 1177, 1319, 1582 and 1718 cm $^{-1}$. From comparing the calculated IR spectra of 16 different isomers to the experimental IRMPD spectrum, the most likely candidate for [TAC $^-$ H $^+$ H $^-$ QO] $^+$ is determined to be a ketone.

We compared the experimental IRMPD spectrum of $\mathbf{TAC}^{\bullet+}$ to the IRMPD spectrum of coronene $^{\bullet+}$ ($\mathbf{Cor}^{\bullet+}$) measured by Oomens *et al.* (2001) In the 1050 to 1700 cm⁻¹ wavenumber region we find considerable overlap in band positions between the $\mathbf{TAC}^{\bullet+}$ and $\mathbf{Cor}^{\bullet+}$. Discrepancies only arise in the 750 to 1050 cm⁻¹ wavenumber region, where $\mathbf{TAC}^{\bullet+}$ has two features at 803 and 932 cm⁻¹ and $\mathbf{Cor}^{\bullet+}$ one at 849 cm⁻¹. Despite these features not

overlapping, the spectra are remarkably similar. This is in line with with the prediction from earlier works that a minor impact on the overall IR spectrum is expected after nitrogen atom incorporation on the periphery of a pure carbon PAH.

To put $\mathbf{TAC^{\bullet+}}$ and $\mathbf{TACH^{+}}$ into astronomical context, we compared their IRMPD spectra to observed astronomical mid-IR spectra of spectral type class A to D (Peeters *et al.* 2002; van Diedenhoven *et al.* 2004; Matsuura *et al.* 2014). The $\mathbf{TAC^{\bullet+}}$ IRMPD has a few features that have some overlap with the class C and D. This indicates the potential contribution of $\mathbf{TAC^{\bullet+}}$ to the class C and D spectra. Thus supporting the idea that cationic exo-PANHs can contribute to the AIBs as well. The IR spectra of exo-PANH cations generally do not match better with the AIBs than their unsubstituted PAH congener, due to the similarity of their IR spectra. So, both exo-PANH and PAH cations should be considered to be potential carriers of AIBs. $\mathbf{TACH^{+}}$ produces an IRMPD spectrum that matches well with the 6.2 μ m feature in all four classes. Furthermore, the two intense features at 7.49 and 7.72 μ m (1335 and 1296 cm⁻¹) have a considerable match with the 7.6 μ m feature of the class A spectrum. This match is indicative of $\mathbf{TACH^{+}}$ being a possible contributor to the AIBs, therefore suggesting the possibility of protonated exo-PANHs being carriers of the AIBs.

Acknowledgements

JB acknowledges the Netherlands Organisation for Scientific Research (Nederlandse Organisatie voor Wetenschappelijk Onderzoek, NWO) for a Vidi grant (grant number 723.016.006). This work was supported in part by NASA's Solar System Exploration Research Virtual Institute (SSERVI): Institute for Modeling Plasma, Atmosphere, and Cosmic Dust (IMPACT). DS acknowledges support by the PRIME programme of the German Academic Exchange Service (DAAD) with funds from the German Federal Ministry of Education and Research (BMBF). DS acknowledges a grant for computing time at the Paderborn Center for Parallel Computing PC^2 under the project PEPICON. DS gratefully acknowledges continued support by Prof. Tina Kasper and funding from DFG under project number KA 3871/3-2 and 270672969.

4.A Appendices

These appendices contain further information to chapter 4. First, we present a zoom-in of the experimental and simulated IR spectra of \mathbf{TACH}^+ between 820 and 1260 cm⁻¹. Next, we show the different energies of three possible isomers formed after losing a hydrogen atom from $\mathbf{TAC}^{\bullet+}$ from different positions. Then, we present the calculated IR spectra of the 16 $[\mathbf{TAC-H+H_2O}]^+$ isomers that we considered as potential products ions. Lastly, we show the coordinates of $\mathbf{TAC}^{\bullet+}$ and \mathbf{TACH}^+ used to calculate the IR spectra.

4.A.1 Experimental and simulated mid-IR spectra of TACH⁺ in the 800 – 1300 cm⁻¹ range

Here we present a zoomed-in version of the experimental and simulated IR spectra of **TACH**⁺ reported in chapter 4(see Fig. 4.7). Most notably, the **TACH**⁺ IRMPD spectrum displays no clear features between 820 and 1260 cm⁻¹, while modes are predicted to fall in this range.




Figure 4.7: Experimental IRMPD (black) and simulated IR (blue vertical lines) spectra of \mathbf{TACH}^+ . The B3LYP/6-311++G(d,p) computed spectrum (blue sticks) is uniformly scaled with a factor of 0.9679 and convolved with a 20 cm⁻¹ FWHM Gaussian function (green shaded area). The horizontal grey, dotted line indicates an apparent cut-off. Exciting modes with a lower intensity than this line does not lift the energy of the molecule over the dissociation threshold.

4.A.2 Energies of [TAC-H]⁺ isomers

Here we present the energies of three isomers that can form upon losing a hydrogen atom from three distinctly different positions in $\mathbf{TAC}^{\bullet +}$ (see Fig. 4.8). The isomer resulting from the loss of a hydrogen from the CH group adjacent to a nitrogen atom is found to be lowest in energy, according to our calculations at a B3LYP/6-311++G(d,p) level of theory.




Figure 4.8: Geometries and relative energies of three different $[TAC-H]^+$ isomers calculated at B3LYP/6-311++G(d,p) level of theory. The circled hydrogen atoms on $TAC^{\bullet+}$ depict the hydrogen that is removed and the arrows point toward the resulting isomer. The energies are depicted with respect to the lowest energetic isomer of $[TAC-H]^+$.

4.A.3 Calculated spectra of all considered $[TAC-H+H_2O]^+$ isomers

Here we present the calculated mid-IR spectra of the 16 isomers that we considered as $[TAC-H+H_2O]^+$ products (see Fig. 4.9). We propose that isomer 5 is the most likely candidate for $[TAC-H+H_2O]^+$, since it is the lowest in energy and its simulated spectrum matches well with the IRMPD spectrum of the $[TAC-H+H_2O]^+$ product ion. Other isomers such as 15 and 16 also provide a good match with the IRMPD spectrum, however, their energies are rather high with respect to isomer 5 and these isomers lack a feature at 1718 cm⁻¹. Therefore, we conclude that these are not the most likely candidate for $[TAC-H+H_2O]^+$. This does not necessarily mean that isomer 5 is the only contributor to the IRMPD and the other isomers may contribute to the experimental spectrum.

76 4.A. Appendices




Figure 4.9: Experimental IRMPD spectrum (black) compared to simulated IR spectra of all considered isomers of $[TAC-H+H_2O]^+$. Also shown are the structures of the various $[TAC-H+H_2O]^+$ isomers and their relative energies calculated using the B3LYP functional in combination with the 6-311++G(d,p) basis set. The simulated spectrum in green indicates our tentative assignment that isomer **5** is likely the dominant isomer. The B3LYP/6-311++G(d,p) computed stick spectra are uniformly scaled with a factor of 0.9679. The calculated spectra are also convolved with a 20 cm⁻¹ FWHM Gaussian function to facilitate comparison with the experimental spectrum. All spectra are normalized with respect to their most intense peak in the 750-1650 cm⁻¹ wavenumber range.




Figure 4.9 (Cont.)

78 4.A. Appendices

4.A.4 Coordinates of TAC^{•+} and TACH⁺

Here we presented the coordinates of $\mathbf{TAC}^{\bullet+}$ and \mathbf{TACH}^+ that were used to calculate the IR spectra presented in chapter 4.

Table 4.3: Cartesian coordinates of the molecular structure of $TAC^{\bullet+}$.

Atom	Coordinates		
	X	У	Z
С	2.343363	-2.831567	0.000000
\mathbf{C}	2.441080	-1.414550	0.000000
\mathbf{C}	1.227137	-0.700177	0.000000
\mathbf{C}	0.008865	-1.409349	0.000000
\mathbf{C}	0.032092	-2.816647	0.000000
\mathbf{C}	1.218829	0.711115	0.000000
\mathbf{C}	-1.218304	-0.710566	0.000000
\mathbf{C}	-1.227304	0.692024	0.000000
\mathbf{C}	0.000000	1.407314	0.000000
\mathbf{C}	-2.466785	1.384110	0.000000
\mathbf{C}	-3.655586	-0.598442	0.000000
\mathbf{C}	-2.455146	-1.411283	0.000000
\mathbf{C}	-2.430551	-2.821576	0.000000
\mathbf{C}	-1.223933	-3.510473	0.000000
Η	-1.202872	-4.594065	0.000000
Η	-3.362957	-3.374470	0.000000
Η	3.249407	-3.431837	0.000000
Η	-4.622094	-1.097693	0.000000
\mathbf{C}	3.671975	-0.664335	0.000000
\mathbf{C}	3.671439	0.705891	0.000000
\mathbf{C}	2.442357	1.443741	0.000000
Η	4.612702	-1.203561	0.000000
Η	4.596997	1.268094	0.000000
\mathbf{C}	1.301069	3.447027	0.000000
\mathbf{C}	0.011601	2.816789	0.000000
Η	1.359724	4.533148	0.000000
$^{\mathrm{C}}$	-2.442841	2.788733	0.000000
Η	-3.386345	3.321114	0.000000
$^{\mathrm{C}}$	-1.228657	3.492319	0.000000
Η	-1.247479	4.576922	0.000000
N	2.452632	2.803175	0.000000
N	-3.670011	0.704506	0.000000
N	1.200053	-3.507431	0.000000

 ${\bf Table~4.4:~Cartesian~coordinates~of~the~molecular~structure~of~{\bf TACH}^+.}$

Atom	(Coordinates	
	X	У	Z
С	-3.646949	0.587074	0.000000
\mathbf{C}	-2.456815	1.388059	0.000000
\mathbf{C}	-1.234483	0.691628	0.000000
$^{\mathrm{C}}$	-1.241235	-0.719561	0.000000
$^{\mathrm{C}}$	-2.475815	-1.404999	0.000000
$^{\mathrm{C}}$	0.000000	1.389690	0.000000
$^{\mathrm{C}}$	-0.022262	-1.432275	0.000000
$^{\mathrm{C}}$	1.208809	-0.734174	0.000000
\mathbf{C}	1.222632	0.676715	0.000000
$^{\mathrm{C}}$	2.418576	-1.466367	0.000000
$^{\mathrm{C}}$	1.269348	-3.465443	0.000000
$^{\mathrm{C}}$	-0.014760	-2.839075	0.000000
\mathbf{C}	-1.265952	-3.522988	0.000000
\mathbf{C}	-2.455549	-2.830283	0.000000
Η	-3.405499	-3.350499	0.000000
Η	-1.274228	-4.607283	0.000000
Η	-4.615172	1.083098	0.000000
Η	1.326281	-4.551696	0.000000
\mathbf{C}	-2.427831	2.808363	0.000000
\mathbf{C}	-1.235827	3.504159	0.000000
\mathbf{C}	-0.012550	2.795927	0.000000
Η	-3.362247	3.358308	0.000000
Η	-1.236558	4.588389	0.000000
\mathbf{C}	2.399553	2.771745	0.000000
\mathbf{C}	2.462926	1.376896	0.000000
Η	3.286046	3.393683	0.000000
\mathbf{C}	3.655109	-0.744409	0.000000
Η	4.572486	-1.320285	0.000000
\mathbf{C}	3.681432	0.625125	0.000000
Η	4.628918	1.151463	0.000000
N	1.221948	3.420766	0.000000
N	2.423324	-2.827044	0.000000
N	-3.663718	-0.724422	0.000000
H	1.238961	4.434882	0.000000