



Universiteit
Leiden
The Netherlands

Laser tonsil treatment under local anesthesia: a patient-friendly effective alternative?

Wong Chung, J.E.R.E.

Citation


Wong Chung, J. E. R. E. (2025, November 6). *Laser tonsil treatment under local anesthesia: a patient-friendly effective alternative?*. Retrieved from <https://hdl.handle.net/1887/4282135>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4282135>

Note: To cite this publication please use the final published version (if applicable).



CHAPTER 2

Tonsillotomy versus tonsillectomy in adults suffering from tonsil-related afflictions: a systematic review

Justin E.R.E. Wong Chung^{a,b}, Peter Paul G. van Benthem^a, Henk M. Blom^{a,b}

A Department of Otolaryngology Head and Neck Surgery, University Medical Center Leiden, Leiden, The Netherlands;

B Department of Otolaryngology Head and Neck Surgery, HagaZiekenhuis, the Hague, The Netherlands

Published in *Acta Oto-laryngologica*. 2018 May;
doi: <https://doi.org/10.1080/00016489.2017.1412500>



ABSTRACT

Objective: Tonsillotomy has emerged as an alternative for tonsillectomy in treating patients with tonsil-related afflictions. Tonsillotomy provides favorable outcomes in children, but treatment of choice in adults remains unclear. This systematic review sought to evaluate the current literature on the efficacy and adverse events of tonsillotomy compared to tonsillectomy in adults.

Methods: A Medline and Cochrane search was conducted for randomized clinical trials (RCTs) and cohort studies comparing tonsillotomy to tonsillectomy in adults. Risk of bias was assessed. Outcome measures were efficacy of the procedure in resolving the initial tonsil-related symptoms (tonsillitis, obstructive sleep apnea, tonsil stones, halitosis, dysphagia), procedure-related complications, recovery time, post-operative use of analgesics, patient satisfaction, and operating time.

Results: In total nine papers were included. These trials had a high risk of bias and the inter-comparability of results was poor. The reported studies found generally a similar efficacy for both interventions. With regard to pain, the use of analgesics, patient satisfaction and operation time, the results were generally in favor of tonsillotomy. Post-operative hemorrhages were more frequent after tonsillectomy.

Conclusion: Current evidence suggests an equal efficacy of tonsillotomy and tonsillectomy in adults and a preference for tonsillotomy in terms of pain, analgesics use, patient-satisfaction, operation time and post-operative complications.

INTRODUCTION

Classic tonsillectomy, the surgical removal of the palatine tonsils, is one of the most performed surgical interventions in the head-and-neck area. Within the United States alone, over half a million tonsillectomies are performed every year.

Tonsillectomies have been performed for over 3 millennia and even though different tonsillectomy methods have been developed over time, the traditional cold dissection tonsillectomy is still regarded as the gold standard.

The function of the palatine tonsils is related to the immune system, but their exact role in the immune response is under debate, especially since studies have shown that the function of the immune system was not compromised in early nor in late childhood in children that had undergone tonsillectomy compared to their age-matched healthy con-trols.¹

Indications for tonsillectomy vary between the pediatric and adult population. In the adult population, tonsillectomy is mainly performed for chronic or recurrent tonsillar infections rather than for tonsillar hypertrophy with obstructive symptoms. Other indications for tonsillectomy in both adults and children include established or suspected tonsil-related malignancies and dysphagia related to the tonsils. Tonsil-related symptoms, such as halitosis, dysphagia and detritus (tonsil stones) are rarely an indication for tonsillectomy. The median recovery-duration after tonsillectomy is around 10 d for adults and 7 d for children. Post-operative hemorrhage rates after adult tonsillectomy are reported to be around 5% and post-operative infection rates around 1–7%.²

Adult patients suffering from tonsillitis are reported to both miss around 9 days of work and use antibiotics for 6 weeks on a yearly basis. Altogether, these data underline the fact that the burden of tonsillectomy for patients is not to be underestimated.

An alternative surgical intervention for tonsillectomy is tonsillotomy. Although first described more than a century ago, tonsillotomy has only become a regular intervention in several areas around the world since its re-introduction in the 1990s. Tonsillotomy is defined as the partial, intracapsular removal of tonsil tissue, as opposed to the total, extra-capsular removal performed during tonsillectomy. Different synonyms are used in literature for the partial removal of tonsils, including tonsillotomy, partial tonsillectomy, tonsil ablation, intra-capsular tonsillectomy, radiofrequency-induced thermotherapy (RFITT) of the tonsils and subtotal tonsillectomy.³

Moreover, a wide variety of different surgical instruments have been used for tonsillotomy, including CO₂-Laser, diathermy, radiofrequency, microdebrider coblation, bipolar and cold-steel tonsillotomy. Tonsillotomy is mostly performed under general anesthesia, but it can also be executed under local anesthesia⁴

Tonsillotomy is currently mostly used in children with pediatric obstructive sleep apnea syndrome (pOSAS) and performed under general anesthesia. A recent systematic review by Wood et al. showed a comparable effect of tonsillotomy versus tonsillectomy on improving sleep disordered breathing, whereas tonsillotomy was associated with fewer post-operative complications.⁵ However, there is insufficient data to show that a single technique for surgical tonsillotomy is superior to others, or to indisputably determine whether tonsillotomy can replace tonsillectomy.⁶

Moreover, currently available study reports do not allow for a reliable estimation of the chance of infection of tonsil remnants after tonsillotomy, which might lead to recurrent tonsillitis, but studies on children have reported a median recurrent tonsillitis rate following tonsillotomy of 3.9%.⁶

In adult patients, tonsillotomy is also increasingly performed, but not yet as frequently as in children. A well-designed overview of current literature comparing the efficacy and safety of tonsillotomy and tonsillectomy in adults is currently lacking. Therefore, the aim of this literature review was to evaluate the current literature on the efficacy and adverse events of tonsillotomy compared to tonsillectomy in adults suffering from tonsil-related diseases and afflictions and identify the knowledge gaps.

MATERIALS AND METHODS

We performed a systematic review following the Cochrane guidelines to assess the efficacy and safety in tonsillotomy versus tonsillectomy in adults suffering from a tonsil-related disease. This study was exempt from institutional board approval as it is a review of previously published data.

Study groups (tonsillotomy)

Our predefined study group of interest consisted of adults or adolescent patients aged over 15 who had been treated with any technique of tonsillotomy for tonsil-related diseases. We used the system of Windfuhr and Werner to classify tonsillotomy interventions into two classes: tonsillotomy procedures in which only the protruding part of the tonsil, the part medial to the faucal pillars, is removed (class I), and tonsillotomy procedures in

which only the inner surface of the tonsil-capsule is preserved and approximately 90% of the tonsil is removed (class 2).³

Control group (tonsillectomy)

Our predefined control group consisted of adult patients with tonsil-related diseases who had undergone conventional tonsillectomy. Tonsillectomy was defined as the complete removal of the tonsil, including its complete capsule.

Tonsil-related diseases and afflictions

We included studies comparing the outcomes of tonsillotomy and tonsillectomy for the following tonsil-related diseases and afflictions: (recurrent) tonsillitis, peritonsillar abscess, obstructive symptoms including OSAS, dysphagia, halitosis and tonsil stones.

Outcome measures

Our predefined outcome measures of interest were efficacy of the procedure in terms of resolution of the initial tonsil-related symptoms that lead to surgery ([recurrent] tonsillitis, peritonsillar abscess, obstructive symptoms including OSAS, dysphagia, halitosis and tonsil stones), complications related to surgery, operating time, recovery time, post-operative pain and use of analgesics, and patient satisfaction.

Literature search

The predefined selection criteria were randomized controlled trials (RCT) and cohort studies comparing tonsillotomy to tonsillectomy in adult or adolescent (>15 y) patients, written in English, Dutch, French or German and published after 1960. We included studies with internal controls (one tonsil removed with tonsillotomy and the other with tonsillectomy in the same patient) and studies with external controls, which are patients undergoing classic tonsillectomies.

A three-step search strategy was executed. First, an initial limited search of MEDLINE and Cochrane collaboration databases was conducted, followed by an analysis of the wording used in the titles and abstracts, and of the index terms used to categorize the articles. Second, a search was performed using all the identified keywords and index terms across the MEDLINE and Cochrane databases. The following keywords and index terms were used: 'tonsillotomy', 'partial tonsillectomy', 'subtotal tonsillectomy', 'intracapsular tonsillectomy', 'hot tonsillectomy', 'radiofrequency induced thermotherapy tonsil', 'RFITT' and 'tonsil ablation'. Third, the reference lists of all the identified reports and articles were searched for additional studies.

Risk of bias

Prior to inclusion, all papers selected for retrieval were assessed by two independent reviewers (J. W. C. and H. B.) for methodological validity using the Cochrane Risk of Bias Tool. This tool addresses possible bias, more specifically selection bias, performance bias, detection bias, attribution bias and reporting bias. Any disagreements that arose between the reviewers were resolved through discussion between the two reviewers or in consultation with a third reviewer (P. P. v. B.).

RESULTS

Literature search

Our search and selection process are shown in **Figure 1**. The initial search performed on 1 April 2017 identified 512 articles. Of these, the majority focused on pediatric patients and had study designs other than RCT. Based on title and abstract, we further excluded papers that did not match the objective of this study (e.g. studies that only examined extracapsular tonsillectomies). Of a total of 20 articles eligible for full-text review, 3 studies were excluded because they only focused on extracapsular removal of the tonsils (tonsillectomy), and 7 studies were excluded for reporting on pediatric or pediatric and adult patients, and 1 study did not specify the age group. No additional articles were included after cross-reference checking of the included studies and reviews on tonsillotomies. This resulted in a total of nine RCT and cohort studies eligible for inclusion (**Table 1**).




FIGURE 1. SELECTION PROCESS OF ELIGIBLE STUDIES.

Table 1. Characteristics of included studies.

Reference	Study design	Publ. year	Country	Indication	Age (years)	Follow-up	Technique	Patients per tech	Blinding	Lost to follow-up
Bender B [13]	RCT	2014	Austria	Tonsillitis	18–65	6 months	1 ICTE (Class I) 2 Cold dissection tonsillectomy	1 50 2 54	Double blinded	10 patients (10%) 5 per group
Nemati S [14]	RCT	2010	Iran	Tonsillitis	15–65	12–24 months	1 RFITT (Class I) 2 Cold dissection tonsillectomy	1 38 2 24	Not stated	None
Arya A [10]	RCT	2003	United Kingdom	Tonsillitis	17–57	24 h	1 Coblation tonsillectomy (Class II) 2 Coblation tonsillectomy	1 14 2 14	Double blinded	None
Ericsson E [7]	RCT	2007	Sweden	Obstruction with or without recurrent tonsillitis	16–25	Until end of post-operative pain (up to 3 weeks)	1 RFITT (Class I) 2 Cold dissection tonsillectomy	1 32 2 44	Not blinded	None
Ericsson E [8]	RCT	2007	Sweden	Obstruction with or without recurrent tonsillitis	16–25	1 year	1 RFITT (Class I) 2 Cold dissection tonsillectomy	1 31 2 43	Not blinded	1 patient (1%)
Wireklint S [9]	RCT	2012	Sweden	Obstruction with or without recurrent tonsillitis	16–25	6 years	1 RFITT (Class I) 2 Cold dissection tonsillectomy	1 29 2 39	Not blinded	7 patients (10%)
Hall [11]	RCT	2004	Hawaii	Obstructive hypertrophy or recurrent tonsillitis	adult	14 d	1 Coblation tonsillectomy (Class II) 2 electrocautery tonsillectomy	1 28 2 28	Double blinded	4 patients (7%)
Lourijssen [4]	Prospective follow-up non-randomized cohort study	2016	Netherlands	Tonsillitis, halitosis, tonsilloliths, obstructing tonsillar hyperplasia or dysphagia	≥18	1 year	1 Outclinic CO ₂ -laser tonsillectomy (Class II) 2 Cold dissection	1 61 2 46	Not blinded	27 patients (25%)
Johnston [12]	Retrospective cohort study	2010	USA	Chronic tonsillitis, tonsillar hypertrophy	Teens (12–19), Adults (>19)	32 months	1 Powered intra-capsular tonsillectomy (PT) (Class II) 2 monopolar electrocautery tonsillectomy (MET)	1 117 adults 2 262 adults	Retrospective	None

ICTE: intra-capsular tonsillectomy; RFITT: radiofrequency-induced thermotherapy.

The nine included RCTs and cohort studies reported on a total of 770 (284 in RCTs and 486 in cohort studies) adult patients. The term used most frequently for describing tonsillotomy was *partial tonsillectomy*. **Table 2** shows the distribution of the terms used for tonsillotomy in the literature.

Table 2. Terminology use in current literature for describing tonsillotomy.

Term	Number of articles
Tonsillotomy	134
Partial tonsillectomy	35
Subtotal tonsillectomy	13
Intra-capsular tonsillectomy	81
Hot tonsillectomy	46
Radiofrequency-induced thermotherapy tonsil	3
RFITT	23
Tonsil ablation	83

The papers by Ericsson and Hultcrantz⁷, Ericsson and Ledin⁸, and Wireklint and Ericsson⁹ reported on the same cohort of patients in Linköping, Sweden, using a consecutively longer follow-up period. From here on, we will refer to this cohort as ‘the Swedish cohort’.

Of all 770 patients, 485 (63%) underwent a tonsillotomy intervention and 327 (42%) underwent tonsillectomy. In the 42 patients included in the studies by Arya et al.¹⁰ and Hall et al.¹¹, tonsillotomy was performed on one tonsil and tonsillectomy on the contralateral tonsil. We included these patients in both arms of our review, and outcome measures were included for the respective treatment.

The included papers studied different age ranges. The study by Bender et al. considered patients between 18 and 65 years as adults. Nemati et al. included all patients between 15 and 65 years, and Arya et al. reported on adults aged between 17 and 57. The Swedish cohort focused on ‘young adults’ aged 16 to 25^{7–9}, Hall et al. did not specify age other than ‘adults.’¹¹ Lourijssen et al. included all patients aged 18 and above and Johnston made a subdivision between teens (12–19 years old) and adults (>18 years old).^{4,12} We included results from the adult patient group of Johnston et al.

Indication for surgery

Primary indications for surgery were tonsillitis in five studies^{4,10,12–14}, obstructive symptoms with or without tonsillitis in five studies^{4,7–9,12}, obstructive symptoms or recurrent tonsillitis in two studies^{4,11}, and halitosis, tonsilloliths, and dysphagia in one study.⁴

Surgical technique

The surgical methods used for tonsillotomy were RFITT^{7-9,11,14}, coblation tonsillotomy^{10,11}, CO2 laser tonsillotomy⁴, power-assisted tonsillotomy¹², and scissor-assisted dissection of the tonsil part medial to the palatine arches followed by the removal of remnants with a microdebrider¹³ (**Table 1**). The surgical method for tonsillectomy was standard cold knife dissection in all studies except for the study by Arya et al.¹⁰, in which coblation was used for both tonsillotomy and tonsillectomy, and for the studies by Johnston et al. and Hall et al., which used electrosurgery^{11,12}.

The number of surgeons performing the surgical interventions was specified in five of nine studied populations. In the studies by Nemati et al. and Arya et al., operations were performed by a single surgeon^{10,14}. The Swedish cohort was operated on by three surgeons, and in the study by Hall et al., the interventions were performed by six surgeons, four of whom had no prior experience with coblation tonsillotomy.

RFITT was performed differently in the study by Nemati et al. and in the Swedish cohort. In the study by Nemati et al., the bipolar linear RFITT probe was introduced into the crypts of the tonsils at five up to nine locations, depending on the size of the tonsils, and the tonsil tissue was vaporized with the power set at 7 watts to perform complete cryptolysis. Afterwards, probable bleeding sites were coagulated.¹⁴ In the Swedish cohort, the RF probe was used to cut the tonsil parallel to the anterior pillar, followed by coagulation of bleeding vessels if necessary.⁸

Five studies performed a Class I tonsillotomy, and four studies performed a Class II tonsillotomy (Table 1). Arya et al. and Hall et al. removed all tonsil tissue, except the tonsillar capsule, during coblation tonsillotomy.^{10,11}

Follow-up

Follow-up varied between all the included studies. Follow-up was 1 day in the study by Arya et al.¹⁰, 14 days in the study by Hall et al.¹¹, 6 months in the study by Bender et al.¹³, 1 year in the study by Lourijssen et al.⁴, and 12 to 24 months in the study by Nemati et al.¹⁴ and 32 months in the study by Johnston et al.¹². In the consecutive papers on the Swedish cohort, follow-up was 3 weeks, 1 and 6 years.⁷⁻⁹

Risk of bias

Apart from the studies by Lourijssen et al.⁴ (prospective follow-up non-randomized cohort study) and Johnston et al. (retrospective study)¹², all included studies were RCTs. Risk of bias is summarized in **Table 3**. Studies by Bender et al.¹³ and Arya et al.¹⁰ had low risks of bias. The study by Nemati et al. had a medium risk of bias for blinding, the studies

on the Swedish cohort⁷⁻⁹, the study by Lourijsen et al.⁴, Johnston et al.¹², and the study by Hall et al.¹¹ had medium and high risk of bias. Possible bias will be further debated in the discussion section of this review.

Table 3. Risk of bias of included studies (Cochrane risk of bias tool).

Article	Bias	Authors' judgement		Support for judgement
Bender B [13]	Random sequence generation	Low risk		Randomization plan by Department of Medical statistics
	Allocation concealment	Low risk		Randomization plan by Department of Medical statistics
	Blinding? All outcomes	Low risk		Patients and examiners were blinded to the surgical procedure
	Incomplete outcome data addressed? All outcomes	Low risk		Per protocol analysis
Nemati S [14]	Selective reporting	Low risk		None
	Other sources of bias	Low risk		None
	Random sequence generation	Low risk		Block randomization
	Allocation concealment	Low risk		Block randomization
	Blinding? All outcomes	Medium risk		Not stated
	Incomplete outcome data addressed? All outcomes	Low risk		One case could not visit clinic but was questioned over the telephone
Arya A [10]	Selective reporting	Low risk		None
	Other sources of bias	Low risk		None
	Random sequence generation	Low risk		Sealed envelope allocation
	Allocation concealment	Low risk		Sealed envelope allocation
	Blinding? All outcomes	Low risk		Only operating surgeon would know type of intervention. PI was a different person.
	Incomplete outcome data addressed? All outcomes	Low risk		Complete data
Ericsson E [7] Ericsson E [8] Wireklint S [9] (Same cohort)	Selective reporting	Low risk		None
	Other sources of bias	Low risk		None
	Random sequence generation	Medium risk		Not stated
Hall D [11]	Allocation concealment	High risk		Invitation for participation and written consent to the randomized choice after randomization
	Blinding? All outcomes	High risk		Written consent to the randomized choice after randomization
	Incomplete outcome data addressed? All outcomes	Ericsson E [7]	Low risk	Complete data
		Ericsson E [8]	Medium risk	1 missing -> excluded
		Wireklint S [9]	Medium risk	7 missing -> excluded
	Selective reporting	Low risk		None
	Other sources of bias	Low risk		None
	Random sequence generation	Medium risk		Counterbalanced by order of presentation
	Allocation concealment	High risk		Invitation for participation after randomization
	Blinding? All outcomes	Medium risk		Patients and primary investigator (PI) were blinded except for 2 surgeries performed by the PI.
Johnston [12]	Incomplete outcome data addressed? All outcomes	Medium risk		Incomplete follow-up (4 patients) was reported but no statistical analysis was performed on missing data
	Selective reporting	Low risk		None
	Other sources of bias	Low risk		None
	Random sequence generation	High Risk		Retrospective chart review
	Allocation concealment	High risk		Retrospective chart review
	Blinding? All outcomes	High risk		Retrospective chart review
	Incomplete outcome data addressed? All outcomes	High risk		Incomplete follow-up (4 patients) was reported but no statistical
				Completeness of chart data not reported
Lourijssen [4]	Selective reporting	Low risk		None
	Other sources of bias	Low risk		None
	Random sequence generation	High risk		Non-randomized
	Allocation concealment	High risk		Freedom of choice
	Blinding? All outcomes	High risk		None
	Incomplete outcome data addressed? All outcomes	High risk		Incomplete follow-up (32%) was reported and addressed with best and worse cast scenarios.
	Selective reporting	Low risk		None
	Other sources of bias	Medium risk		Inclusion bias, most patients were redirected especially for the laser treatment and therefore highly motivated

OUTCOME MEASURES (TABLE 4)

Efficacy (primary outcome)

Efficacy of the surgical intervention in resolving the initial tonsil-related symptoms was reported in six of nine papers and was evaluated differently between studies. Eight of nine studies showed no difference in efficacy between tonsillotomy and tonsillectomy, while the study by Lourijzen et al. showed significantly better efficacy in favor of tonsillectomy. Bender et al.¹³ scored efficacy with the Tonsil and Adenoid Health Status Instrument (TAHSI). The TAHSI is a questionnaire for tonsil disease with a score ranging from 0 to 64, with a higher score indicating more severe tonsil-related morbidity. The TAHSI score in the study by Bender et al. did change significantly after surgery, and there were no differences between the two treatment groups.

Nemati et al.¹⁴ did not find a difference between the two treatment groups in controlling recurrent tonsillitis in 12 to 24 months follow-up. In the Swedish cohort, efficacy was studied using the Short Form 36 Health Survey to evaluate Health-Related Quality of Life (HRQoL) and the EuroQol five dimensions questionnaire (EQ-5D VAS) scales to evaluate the self-rated overall health. The HRQoL and EQ-5D VAS scales did improve significantly after both interventions. No significant difference was found between the treatment groups at 1-year and 6-year follow-up. A significant decrease in ENT infections was seen following both interventions at 1-year and 6-year follow-up, without any difference between treatment groups. Furthermore, the snoring intensity decreased in both treatment groups without a difference between treatments.

In the study by Lourijzen et al., 72.5% of patients were complaint-free 1 year after tonsilotomy compared to 97.2% of patients after tonsillectomy ($p < .001$).⁴

The study by Johnston et al. showed a difference in persistence of tonsillitis after both interventions in favor of tonsillectomy, but no statistical analyses had been performed between both groups.¹²

Complications

Complications were reported in all papers. Bender et al.¹³ reported post-tonsillectomy hemorrhage (PTH) in 16 of 54 patients (29.6%) after tonsillectomy and in 6 of 50 patients (12%) after tonsillotomy ($p = .03$). Furthermore, PTH was more severe and recurrent PTH was more frequent after tonsillectomy. In the Swedish cohort, six patients in the tonsillectomy group had a PTH (two primary, four secondary).⁷⁻⁹ Hall et al. found two minor PTHs on the coblation tonsillotomy side and one on the electrosurgery tonsillectomy side. None of the patients required transfusion or a return to the operating theatre.¹¹

There were no primary or secondary hemorrhages in the tonsillotomy group, but in one patient there were difficulties in maintaining intracapsular hemostasis during the tonsillotomy intervention and therefore surgery was converted to tonsillectomy. In the cohort studied by Arya et al., in which patients underwent tonsillotomy as well as tonsillectomy, PTH occurred in one patient, but the type of intervention causing the complication was not mentioned.¹⁰ No PTH occurred in the study by Nemati et al.¹⁴ Lourijsen et al. reported a significant difference in PTH in favor of tonsillotomy: three (3%) PTHs were seen after tonsillectomy, two requiring intervention, and two (1%) PTHs were seen after tonsillotomy, none requiring intervention.⁴ Johnston reported a PTH rate of 1.7% after tonsillotomy and 2.7% after tonsillectomy; no statistical analysis was performed.¹²

In addition, Lourijsen et al. reported four post-operative infections after tonsillectomy, none after tonsillotomy.⁴ Bender et al. reported more vomiting after tonsillectomy¹³, and Nemati et al. reported dysphagia in one patient after tonsillotomy caused by loose tonsil tissue in the throat post-operatively.¹⁴ No deaths occurred in any of the studies.

Pain

Post-operative pain or the use of pain medication was registered in all studies except in the study by Johnston et al.¹² Arya et al. did not find a difference in 24-hour post-operative pain scores between the sides operated with tonsillotomy and the tonsillectomy sides.¹⁰ This finding does not come unexpectedly since significant relief of post-operative pain is not expected within 24 hours. All other studies reported a difference in pain scores or analgesic medication use in favor of tonsillotomy. Bender et al. found that the tonsillotomy group used less and milder pain medication (only paracetamol and NSAID, instead of hydromorphone) than the tonsillectomy group.¹³ Hall et al. found less pain after tonsil ablation than after electrosurgical tonsillectomy.¹¹ Nemati et al. found a lower pain score on days 1, 3, 5, and 10 after tonsillotomy.¹⁴ The differences in pain scores were significant after day 1 and day 3. In the Swedish cohort, patients who had undergone tonsillotomy recorded less pain from the first day onwards, required less analgesics, and were pain-free 4 days earlier than the tonsillectomy group.⁷⁻⁹ In the study by Lourijsen et al., both overall pain scores (5.4 after tonsillotomy and 7.7 after tonsillectomy) and analgesic use (median of 5.4 days after tonsillotomy and 9.6 days after tonsillectomy) were significantly different in favor of tonsillotomy. The days until resumption of daily activities were significantly different (4.8 days after tonsillotomy and 9.6 days after tonsillectomy).⁴

Duration of surgery

Four studies reported on the duration of surgery. Bender et al. concluded that tonsillectomy prolonged the duration of surgery compared to microdebrider-assisted tonsil-

lotomy.¹³ Arya et al. concluded that the duration of coblation-assisted tonsillectomy was longer than the duration of coblation-assisted tonsillotomy.¹⁰ Hall et al. concluded that electrosurgical tonsillectomy took longer than coblation tonsillotomy¹¹, and Lourijsen et al. found a shorter operation time after CO2 laser tonsillotomy compared to cold dissection tonsillectomy.⁴

Table 4. Summary of outcomes.

Outcome reference	Efficacy	Complications	Operating time	Pain	Recovery time	Analgesics	Patient satisfaction
Bender B [13]	Median of 32.7 point improvement on TAHSI score, no significant differences between groups.	PTH after ICTE in 12% and after ECTE in 29.6% ($p = .03$).	Shorter for ICTE	Not stated	Not stated	Fewer and less intense analgesics in ICTE group ($p = .049$).	Not stated
Nemati S [14]	Significant decrease of tonsillitis, no significant differences between groups.	Remnant of tonsil tissue after RFTT causing dysphagia in one patient.	Shorter for RFTT	Lower mean post-operative pain day 1 & 3 after RFTT ($p < .001$). No significant differences on day 5 and 10.	1.8 d until return to normal diet after RFTT vs. 3.61 d after cold dissection (CD) ($p < .004$)	Not stated	Not stated
Arya A [10]	Not stated.	1 PTH, intervention not stated.	Tonsillectomy took 5–6 min, tonsillectomy took 15 min.	24-h VAS pain score: area under curve of 60.9 for tonsillectomy and 82.5 for tonsillectomy ($p = .13$).	Not stated	Not stated	Not stated
Ericsson E [7]	Not stated.	2 primary bleedings and 4 secondary bleedings after CD.	29.1 min for RFTT and 29.8 min for CD, difference was not significant.	First pain-free day after RFTT on day 8.6, after RFTT on day 12.8 ($p < .001$)	Days to normal eating: TT 7.9 TE 12.0 ($p < .001$). Days to normal activity: TT 6.1 TE 10.6 ($p < .001$)	Fewer analgesics in RFTT group for a shorter period of time. More NSAID use after TE.	Not stated
Ericsson E [8]	Significant: HRQoL improvement, decrease in snoring intensity and decrease in number of ENT-infections. No significant differences between groups.	None	See article by Ericsson et al. [7]	See article by Ericsson et al. [7]	See article by Ericsson et al. [7]	See article by Ericsson et al. [7]	Higher satisfaction after cold dissections tonsillectomy
Wierklint S [9]	Significant: HRQoL improvement, decrease in snoring intensity and decrease in number of ENT-infections. No significant differences between groups.	None	See article by Ericsson et al. [7]	See article by Ericsson et al. [7]	See article by Ericsson et al. [7]	See article by Ericsson et al. [7]	See article by Ericsson et al. [8]
Hall [11]	Not stated	3 minor PTH; 2 after tonsillectomy, 1 after tonsillectomy.	Shorter for Coblation tonsillectomy ($F = 13.271$, $p < .002$).	VAS score post-operatively significantly lower after tonsillectomy.	Not stated	Not stated	Patients choose coblation tonsillectomy over tonsillectomy ($p < .001$)
Johnston [12]	Recurring tonsillitis: 8.5% after PIT, 2.7% after MET (Statistical analysis not available).	PTH in 1.7% after PIT, 2.7% after MET. Dehydration was seen in 0.9% after PIT and 4.2% after MET. (Statistical analysis not available).	Not stated	Not stated	Not stated	Not stated	Not stated
Louijssen [4]	72.5% complaint free 1 year after TO, 97.2% complaint free 1 year after CD ($p < .001$).	3 PTH after tonsillectomy, two requiring intervention. 2 PTH after tonsillectomy, none requiring intervention. Four post-operative infections after tonsillectomy.	Shorter for tonsillectomy (17.1 min) compared to tonsillectomy (22.4 min) ($p = .002$)	Post-operative VAS pain score 5.4 after CD and 7.7 after TE ($p < .001$).	Resumption of daily activities on day 4.8 after TO and 9.6 after TE ($p < .001$). Subjective complete recovery on day 9.6 after TO and day 14.7 after TE ($p < .001$).	Analgesics use 5.4 d after TO and 9.9 d after CD ($p < .001$).	No significant difference on 2 week follow-up

TAHSI: tonsil and adenoid health status instrument; HRQoL: health-related quality of life.

DISCUSSION

We conducted a systematic review to assess the value of tonsillotomy versus conventional tonsillectomy for tonsil-related diseases in adults. We identified nine studies comparing the efficacy of tonsillotomy and tonsillectomy. In eight of the nine studies, tonsillotomy was found to be as effective as tonsillectomy, and in all nine studies, it was associated with less pain, a shorter period of recovery, and a shorter duration of surgery.

This systematic review confirms that tonsillotomy versus tonsillectomy in adults has been scarcely studied. Our search only revealed nine papers describing randomized controlled trials or cohort studies comparing intracapsular tonsillotomy with extracapsular tonsillectomy, with a combined population of 770 patients. These studies made use of a variety of tonsillotomy methods, inclusion criteria, and outcome measures, which led to poor inter-comparability and therefore no meta-analysis could be performed. Furthermore, some of the studies were of questionable quality. The study by Bender et al. was the only study of relatively high quality, with a clearly stated randomization and sufficient patients to meet predefined power. Unfortunately, this study had a relatively high percentage of lost-to-follow-up of 16%, and blinding of outcome assessment was not mentioned. A per-protocol analysis was performed, but an intention-to-treat analysis would have been preferred in this study.¹³ The study by Nemati et al. had a high dispersion in follow-up time (range 12–24 months) and reported sparse data on group differences. Therefore, possible bias cannot be evaluated thoroughly.¹⁴ Arya et al. chose a very different study design, in which patients were used as their own controls. The number of studied patients was small, with only 14 included patients, and follow-up for pain evaluation was as short as 24 hours. Therefore, no meaningful statements can be made on pain, symptom improvement, and post-operative bleeding. The study reported one complication, a PTH, but it did not state which type of intervention led to this complication. Moreover, patients being their own controls has its limitations in the context of the current interventions, since, for example, it may be hard to distinguish pain or discomfort from one side or the other.¹⁰

The study on the Swedish cohort reported on a more substantial number of patients ($n = 76$) with a total follow-up period of 6 years. Unfortunately, patients were not blinded to treatment, and randomization and allocation were not stated. Therefore, subsequent bias cannot be ruled out. Patients lost to follow-up were excluded from the analyses, which may potentially have led to classification bias.^{7–9} The study by Hall et al.¹¹ was, of all RCTs, most prone to bias. Randomization was based on order of presentation and was done before invitation to participate. Also, blinding was not performed correctly

since two patients were operated on by the primary investigator. Besides, there was no statistical correction for missing data during follow-up.

The study by Lourijssen et al. is limited by a non-randomized, non-blinded study design with a high loss-to-follow-up in both treatment groups.⁴ A selection bias of highly motivated patients is expected for the tonsillotomy treatment group since most of these were redirected especially to that specific hospital for outpatient laser treatment of their tonsils. The indication for surgery was significantly different between both groups (halitosis and tonsillolithiasis were predominantly treated with tonsillotomy), which makes the suspicion of selection bias stronger. Furthermore, CO2 laser tonsillotomy, an intervention with a clear learning curve, was not solely performed by an experienced laser surgeon but also by less experienced residents. Follow-up was 1 year, but tonsillitis has been shown to recur postoperatively predominantly after 1 year.¹⁵

The study by Johnston et al. has a retrospective study design with its inherent information bias and patient- and treatment-selection bias. This study sought to evaluate differences between teens and adults treated with powered intracapsular tonsillectomy or monopolar electrocautery tonsillotomy, and all analyses were based on this grouping. Therefore, no analysis was performed between both treatments, and thus nothing can be said about the statistical significance of the reported findings. The sizes of the groups were very disparate, and the indications for intervention varied greatly as well.¹²

In summary, the selected randomized controlled studies and cohort studies on tonsillotomy in adults are very diverse with regard to study population, surgical methods used, and outcomes measured. Thus, they cannot be compared to each other, and therefore the efficacy could not be evaluated by means of a meta-analysis. A better powered, well-designed randomized controlled trial needs to be undertaken to compare the efficacy and safety of tonsillotomy and tonsillectomy in adults and to allow for generalizable conclusions regarding the preferential surgical approach for tonsil surgery in adults. Especially, recovery and complication rates should be evaluated and put in light of socio-economic perspectives. Furthermore, confounding by indication should be assessed beforehand during patient inclusion.

Several procedural differences between tonsillotomy and tonsillectomy may explain the rather positive results that were reported in the few studies that we identified. Compared to tonsillectomy, tonsillotomy entails the removal of a smaller amount of tissue, while the major nerves and vessels running in the tonsillar bed to the tonsillar capsule are retained, and only the smaller nerves and vessels that radiate into the lymphoid outer part of the tonsil are affected. This may limit the risk of PTH and post-operative pain.

This assumption is supported by comparative studies of both interventions in pediatric populations, where this topic has been studied much more thoroughly. However, the anatomy of the palatine tonsillar region is strikingly different in children and adults. In children, the palatine tonsils lay relatively loose in the tonsillar bed and are only loosely connected by the vessel-nerve bundle between the tonsillar bed and tonsillar capsule. In adults, on the other hand, the palatine tonsils are often more rigidly connected to the tonsillar bed, and this connection is often enhanced by recurrent or chronic tonsillar infections. Therefore, tonsillectomy in adults is generally more time-consuming, has a longer post-operative recovery time, and is more frequently accompanied by PTH.¹⁶ Consequently, clinicians are generally much more reticent in performing tonsillectomy in adults than in performing the same operation in children.^{13,14}

Different methods for tonsillectomy and tonsillotomy have been described. The literature comprises reports on the use of CO₂ laser, coblation, shaver (microdebrider), diode laser, and radiofrequency. All these methods can be used for extracapsular tonsillectomy as well as for intracapsular tonsillotomy. However, regardless of the method used, tonsillotomy will result in less pain and lower post-operative bleeding rates. At present, conclusive evidence supporting the supremacy of any surgical technique is lacking.

Two recent Cochrane reviews on tonsillectomy could not find a difference in morbidity between cold knife dissection and diathermy tonsillectomy, nor could they find a difference in post-operative pain or in the speed and safety of recovery between coblation and other tonsillectomy interventions. Magdy et al. could not find a difference in tonsillar fossa healing when comparing coblation, dissection, and laser-assisted tonsillectomy, but monopolar cautery did show a slower healing process after 7 and 15 days.¹⁷ Coblation was associated with less thermal damage to surrounding tissue, which was presumed to be the result of the relatively low temperatures needed for sustaining the necessary plasma field. Currently, there is no sufficient evidence in favor of any method for tonsillectomy or tonsillotomy, and the choice of a surgical method is, at present, only based on the surgeon's preference and the availability of equipment.

Tonsillotomy as a surgical method for tonsil-related diseases and afflictions is associated with several limitations. First, tonsillotomy is contraindicated when tonsil-related malignancy is suspected. In these cases, the tonsil should be preserved for examination by a pathologist, and tumor spread needs to be prevented. Second, although there is currently no evidence to support transmission of Creutzfeldt-Jakob disease through inhalation of vaporized tissue, some have suggested that this may be a theoretical risk. This risk is thought to be extremely low, but guidelines indicate that there is no hard data available. Third, the possibility of post-operative tonsillitis caused by residual tissue

should be taken into account. Even though tonsillar remnants are also found after tonsillectomy, tonsil residues are always present in tonsillotomy and may (theoretically) lead to post-operative tonsillitis. No increase in post-operative tonsillitis rates was reported in the included studies, but there is no strong evidence that there is no difference in post-operative tonsillitis. Fourth, the possibility of regrowth of tonsils after tonsillotomy is often mentioned. Unfortunately, the risk of regrowth of tonsils in adults after tonsillotomy has not been reported in the literature and is thus a matter of 'expert' opinion. In adults, however, natural involution of the palatine tonsils is seen, and thus significant regrowth of tonsils in adults would be unexpected.^{7,18} In children, there are some data on regrowth: Doshi et al. reviewed 636 medical records of children that underwent tonsillotomy and concluded that there is a small risk of tonsillar regrowth and that regrowth is five times more likely at an age <5 years. Their findings were in concordance with the findings of other authors. Patients in the study on the Swedish cohort had surgery primarily for obstructive symptoms. At 6-year follow-up, no differences in snoring index or HRQL were found between the two treatments.⁹ This finding was in concordance with the studies by Hultcrantz et al. and Eviatar et al., which found similar effectiveness of tonsillotomy compared to tonsillectomy in children with pOSAS after 10 years of follow-up. Based on these data and on our own clinical experience with tonsillotomy, we expect the need for secondary tonsillotomy caused by regrowth in adults to be low.

As was mentioned before, different surgical methods for tonsillotomy have been put into practice, but the inter-comparability between studies is lacking. The less invasive nature of a tonsillotomy enables surgeons to perform tonsillotomies in an outpatient clinic setting using only local anesthesia without sedation,^{4,19,20} thereby reducing the morbidity of tonsil surgery. Tonsillotomy under local anesthesia can drastically reduce the costs of tonsil surgery since the need for general anesthesia and its associated specialty care disappears.

Other cost savings can possibly be found in the change in surgical tools, in the reduction of pain, and in the reduced risk of PTH. A decrease in PTH does not only lead to a significant decrease in morbidity and mortality, but also reduces the costs of in-hospital stay, readmission, and re-intervention. A reduction in pain leads to a decreased use of pain medication, earlier discharge from the hospital, and earlier return to work or school and study, which results in considerable socioeconomic gain.

CONCLUSION

Current evidence suggests that the efficacy of tonsillotomy and tonsillectomy in adults is equal and that tonsillotomy is preferable in terms of pain, use of analgesics, patient satisfaction, operation time, and post-operative complications. Post-operative hemorrhages were more frequent after tonsillectomy. This review only identified nine randomized controlled trials and cohort studies comparing the efficacy and safety of tonsillotomy and tonsillectomy in adults suffering from tonsil-related diseases and afflictions. An adequately powered, thorough, and well-designed randomized controlled trial should be performed to conclusively bridge the knowledge gap necessary to clinically decide which method should be used in which patient.

REFERENCES

1. Kaygusuz I, Alpay HC, Gödekmerdan A, et al. Evaluation of long-term impacts of tonsillectomy on immune functions of children: A follow-up study. *International Journal of Pediatric Otorhinolaryngology*. 2009;73(3):445-449. doi:10.1016/j.ijporl.2008.11.014
2. Bhattacharyya N, Kepnes LJ. Revisits and postoperative hemorrhage after adult tonsillectomy. *Laryngoscope*. 2014;124(7):1554-1556. doi:10.1002/lary.24541
3. Windfuhr JP, Savva K. [An update on tonsillotomy studies]. *HNO*. Published online September 2016. doi:10.1007/s00106-016-0237-4
4. Lourijsen ES, Wong Chung JERE, Koopman JP, Blom HM. Post-operative morbidity and 1-year outcomes in CO₂-laser tonsillotomy versus dissection tonsillectomy. *Acta Oto-Laryngologica*. 2016;136(10):983-990. doi:10.1080/00016489.2016.1183040
5. Wood JM, Cho M, Carney AS. Role of subtotal tonsillectomy ('tonsillotomy') in children with sleep disordered breathing. *The Journal of Laryngology and Otology*. 2014;128 Suppl 1:S3-7. doi:10.1017/S0022215113003058
6. Windfuhr JP, Savva K, Dahm JD, Werner JA. Tonsillotomy: facts and fiction. *European archives of oto-rhino-laryngology: official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS): affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2015;272(4):949-969. doi:10.1007/s00405-014-3010-x
7. Ericsson E, Hultcrantz E. Tonsil surgery in youths: good results with a less invasive method. *Laryngoscope*. 2007;117(4):654-661. doi:10.1097/mlg.0b013e318030ca69
8. Ericsson E, Ledin T, Hultcrantz E. Long-term improvement of quality of life as a result of tonsillotomy (with radiofrequency technique) and tonsillectomy in youths. *Laryngoscope*. 2007;117(7):1272-1279. doi:10.1097/MLG.0b013e31805559e1
9. Wireklint S, Ericsson E. Health-related quality of life after tonsillotomy versus tonsillectomy in young adults: 6 years postsurgery follow-up. *Eur Arch Otorhinolaryngol*. 2012;269(8):1951-1958. doi:10.1007/s00405-012-1990-y
10. Arya A, Donne AJ, Nigam A. Double-blind randomized controlled study of coblation tonsillotomy versus coblation tonsillectomy on postoperative pain. *Clin Otolaryngol Allied Sci*. 2003;28(6):503-506.
11. Hall DJ, Littlefield PD, Birkmire-Peters DP, Holtel MR. Radiofrequency ablation versus electrocautery in tonsillectomy. *Otolaryngology-Head and Neck Surgery: Official Journal of American Academy of Otolaryngology-Head and Neck Surgery*. 2004;130(3):300-305. doi:10.1016/j.otohns.2003.09.024
12. Johnston DR, Gaslin M, Boon M, Pribitkin E, Rosen D. Postoperative complications of powered intracapsular tonsillectomy and monopolar electrocautery tonsillectomy in teens versus adults. *Ann Otol Rhinol Laryngol*. 2010;119(7):485-489. doi:10.1177/000348941011900710
13. Bender B, Blassnigg EC, Bechthold J, et al. Microdebrider-assisted intracapsular tonsillectomy in adults with chronic or recurrent tonsillitis. *The Laryngoscope*. 2015;125(10):2284-2290. doi:10.1002/lary.25265
14. Nemati S, Banan R, Kousha A. Bipolar radiofrequency tonsillotomy compared with traditional cold dissection tonsillectomy in adults with recurrent tonsillitis. *Otolaryngol Head Neck Surg*. 2010;143(1):42-47. doi:10.1016/j.otohns.2010.03.031
15. Windfuhr JP, Werner JA. Tonsillotomy: it's time to clarify the facts. *Eur Arch Otorhinolaryngol*. 2013;270(12):2985-2996. doi:10.1007/s00405-013-2577-y
16. Windfuhr JP, Chen YS. Incidence of post-tonsillectomy hemorrhage in children and adults: a study of 4,848 patients. *Ear Nose Throat J*. 2002;81(9):626-628, 630, 632 passim.

17. Magdy EA, Elwany S, El-Daly AS, Abdel-Hadi M, Morshedy MA. *Coblation Tonsillectomy: A Prospective, Double-Blind, Randomised, Clinical and Histopathological Comparison with Dissection–Ligation, Monopolar Electrocautery and Laser Tonsillectomies.*; 2008. Accessed November 30, 2016. /core/journals/journal-of-laryngology-and-otology/article/coblation-tonsillectomy-a-prospective-double-blind-randomised-clinical-and-histopathological-comparison-with-dissectionligation-monopolar-electrocautery-and-laser-tonsillectomies/EFB5CAAD6E4351DE28747E975D5F2226/core-reader
18. Koempel JA, Solares CA, Koltai PJ. The evolution of tonsil surgery and rethinking the surgical approach to obstructive sleep-disordered breathing in children. *J Laryngol Otol.* 2006;120(12):993-1000. doi:10.1017/S0022215106002544
19. Remacle M, Keghian J, Lawson G, Jamart J. Carbon-dioxide laser-assisted tonsil ablation for adults with chronic tonsillitis: a 6-month follow-up study. *European Archives of Oto-Rhino-Laryngology.* 2003;260(8):456-459. doi:10.1007/s00405-003-0600-4
20. Andrews PJ, Latif A. Outpatient laser tonsillar ablation under local anaesthetic. *European Archives of Oto-Rhino-Laryngology and Head & Neck.* 2004;261(10):551-554. doi:10.1007/s00405-003-0718-4