



Universiteit
Leiden

The Netherlands

Lighting up dark exomoons: observational signatures of tidally induced volcanism in other worlds

Kleisioti, E.

Citation

Kleisioti, E. (2025, November 13). *Lighting up dark exomoons: observational signatures of tidally induced volcanism in other worlds*. Retrieved from <https://hdl.handle.net/1887/4282613>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4282613>

Note: To cite this publication please use the final published version (if applicable).

Lighting up dark exomoons

Observational signatures of tidally induced volcanism in other worlds

Proefschrift

ter verkrijging van
de graad van doctor aan de Universiteit Leiden,
op gezag van rector magnificus prof. dr. ir. H. Bijl,
volgens besluit van het college voor promoties
te verdedigen op donderdag 13 november 2025
klokke 10.00 uur

door

Evangelia Kleisioti

geboren te Athens, Greece
in 1994

Promotor:	Dr. M. A. Kenworthy	
Co-promotor:	Dr. D. Dirkx	TU Delft
Promotiecommissie:	Prof. dr. I.A.G. Snellen	
	Prof. dr. M.R. Hogerheijde	
	Dr. Y. Miguel	
	Prof.dr. C. Dominik	University of Amsterdam
	Dr. S.M. Cazaux	TU Delft

Printed by: Universitair Facilitair Bedrijf

Cover design: Illustrated by Phoebe Athanasopoulos.

An electronic copy of this thesis can be found at <https://openaccess.leidenuniv.nl>.

© Elina Kleisioti, 2025

To my parents

Contents

1	Introduction	1
1.1	Introduction	1
1.2	Detecting extrasolar worlds	3
1.2.1	Exoplanet detection methods	3
1.2.2	Exomoons and their detection methods	6
1.3	Tides	12
1.3.1	Tides and tidal heating: Basic concepts	14
1.3.2	Probing planetary interiors	16
1.4	This thesis	17
	References	20
2	Tidally Heated Exomoons around ϵ Eridani b: Observability and prospects for characterization	27
2.1	Introduction	28
2.2	Models	29
2.2.1	Interior model	30
2.2.2	Tidal model	32
2.2.3	Thermal equilibrium	34
2.3	Observability	36
2.4	Results	38
2.4.1	Interior structure dependence on orbital properties	38
2.4.2	Orbital and interior constraints from observations	41
2.5	Discussion	43
2.5.1	Longevity and mean motion resonance	44
2.5.2	Combination with other exomoon detection methods: Constraining the moon radius and semi-major axis	45
2.5.3	Larger moons	47
2.5.4	Future model validation	47
2.6	Conclusions	48
	References	49
3	The spectroastrometric detectability of nearby Solar System-like exomoons	53
3.1	Introduction	54
3.2	Methods	56
3.2.1	The centroid and the spectroastrometric signal	56
3.2.2	The spectroastrometric signal for closed Keplerian orbits	58
3.2.3	Sources of spectroastrometric noise	58
3.2.4	Consequences for the observation time allocation	61
3.2.5	Minimum required moon flux	61
3.2.6	Model scenario	62

3.3	Results	65
3.3.1	Minimum moon temperature for detectability	66
3.3.2	Eccentricity effects	66
3.3.3	Applicability to distant systems	66
3.4	Discussion	67
3.4.1	Comparison to previous work	69
3.4.2	Limitations on this formulation	69
3.4.3	Spectroastrometric capabilities of current-generation telescopes	71
3.4.4	Detectability of other moon-planet systems	71
3.4.5	Comparing against other moon detection methods	72
3.4.6	Synergies with other methods	73
3.5	Conclusion	73
	References	75
4	Direct detectability of tidally heated exomoons by photometric orbital modulation	79
4.1	Introduction	80
4.2	Photometric signal variability	81
4.2.1	Exo-Io modeled photometric light curves	81
4.2.2	Host modeled photometric light curves	83
4.3	Detectable moon signals in hosts' periodograms	84
4.4	Results	85
4.4.1	Tidally heated exomoons around a 2.5 h variable planet at 3 parsecs	87
4.4.2	Applicability to systems at larger distances	87
4.5	Discussion	88
4.5.1	Other sources of such signals and synergies with other exomoon detection methods	89
4.5.2	Hotspot time variability and spatial distribution on Io and exo-Ios	90
4.5.3	Possibility of hotspot detection around cooler planets	90
4.6	Conclusions	90
	References	92
5	Modeling the interiors of Trappist-1f, g, h: Tidal Heating, Subsurface Oceans, and Cryovolcanic Activity	99
5.1	Introduction	100
5.2	Methods	102
5.2.1	Heat transfer model	102
5.2.2	Tidal response model	103
5.2.3	Interior heat balance and structure calculation	104
5.2.4	Modelling observations of cryovolcanic activity	107
5.3	Results	110
5.3.1	Possible interior structures of the planets Trappist 1 - f,g,h	110
5.3.2	Observations of cryovolcanic activity	112
5.4	Discussion	113
5.4.1	Column density comparison with previous work	113

5.4.2	Water Mass Fraction comparison with previous work	115
5.4.3	Water content in the Trappist-1 system and other possible interior structures	116
5.4.4	Future work	117
5.5	Conclusions	118
	References	119
English summary		127
Nederlandstalige samenvatting		131
Ελληνική περίληψη		137
List of publications		143
Curriculum vitae		145
Acknowledgments		147

