



Universiteit  
Leiden

The Netherlands

## Translating preclinical insights into early psychopharmacology trials: application of the IB-Derisk analyser tool

Dijkstra, F.M.

### Citation

Dijkstra, F. M. (2025, December 4). *Translating preclinical insights into early psychopharmacology trials: application of the IB-Derisk analyser tool*. Retrieved from <https://hdl.handle.net/1887/4284607>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4284607>

**Note:** To cite this publication please use the final published version (if applicable).

CHAPTER VI

CENTRAL NERVOUS SYSTEM EFFECTS  
OF TAK-653, AN INVESTIGATIONAL  
ALPHA-AMINO-3-HYDROXY-5-METHYL-  
4-ISOXAZOLE RECEPTOR (AMPA)  
POSITIVE ALLOSTERIC MODULATOR IN  
HEALTHY VOLUNTEERS

Dijkstra, F.<sup>1,2</sup>, O'Donnell P.<sup>3,4</sup>, Klaassen E.<sup>1</sup>, Buhl D.<sup>3</sup>, Asgharnejad M.<sup>5</sup>,  
Rosen L.<sup>3</sup>, Zuiker R.G.J.A.<sup>1</sup>, van Gerven J.M.A.<sup>1,2</sup>, Jacobs G.E.<sup>1,2</sup>

*Transl Psychiatry*. 2022;12(1):408.  
doi:10.1038/s41398-022-02148-w

---

1 Centre for Human Drug Research (CHDR), Leiden, the Netherlands

2 Department of Psychiatry, Leiden University Medical Center (LUMC),  
Leiden, the Netherlands

3 Neuroscience Translational Medicine, Takeda Pharmaceutical  
Company Ltd, Cambridge, MA, USA

4 McLean Hospital, Department of Psychiatry, Harvard Medical  
School, Belmont, MA, USA

5 Neuroscience Therapeutic Area Unit, Takeda Pharmaceutical  
Company Ltd, Cambridge, MA, USA

---

## ABSTRACT

**AIMS** TAK-653 is a novel AMPA receptor positive allosteric modulator in clinical development for the treatment of major depressive disorder (MDD). This study aimed to measure the functional pharmacodynamic central nervous system (CNS) effects of TAK-653.

**METHODS** A randomised, double-blind, placebo-controlled, three-way crossover (placebo, TAK-653 0.5 mg and 6 mg) study with 24 healthy volunteers was performed. NeuroCart tests consisting of body sway (BS), saccadic peak velocity (SPV), smooth pursuit eye movements (SP), adaptive tracking (AT), Bowdle and Bond and Lader Visual Analogue Scales (B-VAS and BL-VAS) and Stroop test were performed pre-dose and 3.5 and 4 h post-dose. Data were analysed using a mixed model analysis of covariance with baseline as covariate.

**RESULTS** It was found that TAK-653 did not affect BS and subjective drug effects as measured by B-VAS and BL-VAS at either dose level. TAK-653 0.5 mg increased SPV (degrees/second) (19.49 [5.98, 32.99],  $P = 0.02$ ) and affected Stroop difference in reaction time between correct congruent and correct incongruent answers and number of correct responses in incongruent trials (22.0 [4.0, 40.0],  $P = 0.05$  and  $-0.3 [-0.5, -0.1]$ ,  $P = 0.02$ , respectively). TAK-653 6 mg improved AT (%) (1.68 [0.51, 2.84],  $P = 0.02$ ) and increased SPV (degrees/s) (15.40 [1.91, 28.90],  $P = 0.06$ ) and SP (%) (2.32 [0.37, 4.27],  $P = 0.05$ ).

**CONCLUSION** Based on these findings it can be concluded that TAK-653 demonstrated a psychostimulant-like pharmacodynamic profile on the NeuroCart consistent with previously reported increase of cortical excitability following Transcranial Magnetic Stimulation (TMS) of the human motor cortex.

## INTRODUCTION

Since ketamine, a N-methyl-D-aspartate (NMDA) receptor antagonist, has been shown to have rapid occurring antidepressant effects,<sup>1-3</sup> there is growing interest in the NMDA receptor as potential novel target for the pharmacological treatment of depressive disorders. Studies into the mechanisms underlying the antidepressant effects of NMDA receptor antagonism have demonstrated an important role for alpha-amino-3-hydroxy-5-methyl-4-isoxazole (AMPA) receptor-mediated signalling.<sup>4-7</sup> Blocking NMDA receptors and thereby indirectly stimulating AMPA receptors, leads to a shift towards predominantly stimulatory glutamate-mediated neurotransmission.<sup>4-7</sup> This is believed to affect molecular processes implicated in the pathophysiology of (chronic) mood disorders related to synaptic plasticity and/or cellular resilience,<sup>8</sup> including the enhanced production of brain-derived neurotrophic factor and triggering of the mammalian target of rapamycin (mTOR) signalling.<sup>6</sup> The importance of AMPA receptor mediated signalling is further supported by the finding that the preclinical antidepressant-like effects of ketamine and related compounds are opposed by AMPA receptor antagonists.<sup>9</sup> These findings support the development of novel antidepressants that target AMPA receptors. The novel AMPA receptor positive allosteric modulator (PAM) TAK-653 (9-[4-(cyclohexyloxy)phenyl]-7-methyl-3,4-dihydropyrazino[2,1-c][1,2,4]thiadiazine 2,2-dioxide) is an investigational potential therapeutic compound in clinical development for major depressive disorder. As full functional agonism of AMPA receptors is associated with potential untoward central nervous system (CNS) stimulation, AMPA receptor PAMs have been proposed as an alternative pharmacological strategy for glutamatergic modulation.<sup>10</sup> In initial healthy volunteer studies with oral doses of TAK-653 0.3 mg to 18 mg, the compound was well tolerated and, in contrast to ketamine, did not cause dissociative adverse effects.<sup>11</sup> Maximum plasma concentrations were attained within 1.25 hours to 5 hours after dosing, the terminal half-life varied from 33.1 hours to 47.8 hours and cerebrospinal fluid concentrations were suggestive of rapid brain penetration.<sup>11</sup> These pharmacokinetic (PK) and safety profiles in healthy volunteers were promising for further clinical development, but the pharmacodynamic (PD) properties of TAK-653 had not been systematically assessed.

In early phases of clinical drug development a full characterisation of both the PK and PD properties of innovative compounds is crucial to rationally guide drug development.<sup>12,13</sup> The question-based clinical development (QBCD) concept has previously been proposed as a conceptual framework for characterising drugs in early clinical development. Specific to CNS drug development, QBCD allows for systematic investigation of crucial issues such as blood-brain barrier (BBB) penetration, intended target engagement and off-target effects.<sup>12</sup> By explicitly incorporating methodologies to address these issues when designing early-phase CNS studies, findings may support go/no-go decisions in subsequent development phases.<sup>12</sup>

In order to characterise TAK-653's PD profile, we applied transcranial magnetic stimulation (TMS) as a potential biomarker for cortical excitability and we performed a test battery of extensively validated, drug-sensitive neurophysiological and neurocognitive CNS tests, the NeuroCart.<sup>14</sup> Based on its *in vitro* profile and preclinical effects, TAK-653 was hypothesised to yield stimulatory CNS effects in healthy clinical populations. As TAK-653 was the first AMPA receptor PAM to be tested using the NeuroCart it was decided to compare the NeuroCart profile of TAK-653 to the profiles of both excitatory or CNS-stimulant (e.g. dopamine releasers)<sup>15</sup> and inhibitory or CNS-depressant (e.g. GABA<sub>A</sub>-agonists) compounds.<sup>16-21</sup> Although the mechanism of action of these compounds differs from TAK-653's mechanism of action, their PD profiles were expected to be relevant for the 'pharmacological benchmarking' of TAK-653's functional PD effects in healthy humans.


Our previous paper reported the effects of TAK-653 0.5 mg and 6 mg on TMS motor evoked potentials following stimulation of the motor cortex.<sup>22</sup> It was observed that TAK-653 increased the amplitude of motor evoked potentials, indicative of increased AMPA receptor-mediated cortical excitability.<sup>22</sup> In the current paper, the NeuroCart data will be presented.

## METHODS

### STUDY DESIGN AND PARTICIPANTS

The study was a randomised, double-blind, placebo-controlled, three-period cross-over study (Figure 1). A fourth, open-label period was conducted with ketamine for assay sensitivity, but the TMS and NeuroCart results were equivocal for reasons discussed previously,<sup>22</sup> and will not be discussed here. During the three-period crossover phase, each treatment period was one day in duration and separated by a wash-out period of 10 to 15 days. During treatment days, volunteers received one oral dose of either placebo, TAK-653 0.5 mg or TAK-653 6 mg; all treatments had the same appearance to guarantee blinding of volunteers and research staff. At 12 to 16 days after the last study visit, a telephone call was made to volunteers as part of the follow-up procedure.

**Figure 1** Study design and the timing of NeuroCart assessments on treatment days



TAK-653 dose levels of 0.5 mg and 6 mg were selected as the lower dose was expected to have minimum to no effects based on the preclinical data and the higher dose to fall into the pharmacological active range.<sup>22</sup> Both dose levels were well tolerated in the previous healthy volunteer study.<sup>11</sup>

Twenty-four healthy volunteers were included in this study. To assess eligibility, volunteers were screened using the following procedures: a review of their medical and psychiatric history, a physical examination, measurement of vital signs, an electrocardiogram (ECG), blood chemistry and haematology laboratory assessments, and urinalysis. Volunteers who had a clinically significant previous or current psychiatric disorder according to Diagnostic and Statistical Manual-5 were excluded. In addition, volunteers who had a history of drinking an average of two or more alcoholic drinks per day were excluded.

During treatment days, volunteers reported at the research facility in the morning. Before a test dose was given, safety assessments were performed, including: adverse event (AE) occurrence, a physical examination, a suicidality assessment using the Columbia suicide severity rating scale,<sup>23</sup> measurement of vital signs, an ECG, laboratory assessments and urinalysis. The same safety assessments and AE recording were performed at set times after dosing.

During the study, volunteers were instructed to restrict the use of substances that could alter brain activity, including concomitant medication, alcoholic beverages, caffeinated products and nicotine-containing products. First, volunteers were instructed not to use concomitant medication starting from the seven days before the first test dose through to the end of the study. Second, they were instructed not to consume alcoholic beverages seven days before the screening visit and each treatment day. Third, they were instructed not to consume caffeinated products 24 hours before the screening visit and each treatment day. Outside of these restrictions, volunteers could consume up to six servings of caffeinated products a day. Finally, they were instructed not to use nicotine-containing products 48 h before the screening visit and each treatment day. Otherwise, volunteers could use up to five nicotine-containing products per day.

### PHARMACOKINETIC ASSESSMENTS

The PK sample collection times were aligned with timings of the TMS assessments, as these were the assessments of primary interest in this study; PK samples were collected before drug administration, and 0.5 h and 2.5 h after administration. Based on initial healthy volunteer studies, it was expected that the mean maximal plasma concentration ( $C_{max}$ ) would be reached at 2.5 h post-dose and due to the relatively long terminal half-life ( $t_{1/2}$ ) TAK-653 of 33.1 to 47.8 hours, plasma levels were then expected to remain stable over a few hours.<sup>11</sup> A validated high-performance liquid chromatography with tandem mass spectrometry assay with a lower limit of quantification of

0.1 ng/mL and coefficient of variation between 1.41% and 5.22% was used to measure TAK-653's plasma concentrations.

#### FUNCTIONAL PHARMACODYNAMIC NEUROCARD ASSESSMENTS

NeuroCard tests that have been shown sensitive to CNS depressant and/or CNS stimulant compounds were selected for this study.<sup>14</sup> These were: body sway, smooth pursuit eye movements, saccadic eye movements, adaptive tracking test, Stroop coloured word test, and Bond and Lader and Bowdle Visual Analogue Scales (VAS). In Table 1 the effects of different CNS depressant and CNS stimulant compounds on the NeuroCard tests performed in this study are summarised. The tests were performed twice prior to dosing as well as at the time of expected maximum plasma concentrations, namely 3.5 hours and 4 hours post-dose (Figure 1). During all tests, lighting conditions were standardised and volunteers were comfortably seated in front of a computer screen, except for body sway measurements, for which volunteers were standing.

**Table 1** Summary of effects of CNS depressant and CNS stimulant compounds on selected NeuroCard tests

Test	CNS depressant		CNS stimulant			
	Diazepam 10 mg	Benzo- diazepines (dose unspecified)	Modafinil 200 mg	Dexam- phetamine 20 mg	Methyl- phenidate (average 20 mg)	Caffeine (60 mg)
Body sway (%)	+ <sup>1</sup>	NR <sup>3</sup>	-	-	-	NR
Smooth pursuit (%)	- <sup>2</sup>	NR	NR	NR	+	NR
Saccadic peak velocity (deg/s)	NR	-	+	+	NR	+
Adaptive tracking (%)	NR	-	+	+	+	+
Stroop coloured word test	NR	-	NR	NR	NR	NR
Bond and Lader VAS alertness	NR	-	+	+	+	+
Bowdle VAS Feeling high	No effect	No effect	No effect	+	No effect	No effect

1. + indicates improvement or increase; 2. - indicates deterioration or decrease; 3. NR: not reported

#### BODY SWAY.

Body sway measurements are used to assess postural stability and are often used in pharmacologic studies.<sup>24-26</sup> Measurements of movements in the anteroposterior direction were performed as in previously published studies,<sup>26,27</sup> with a string similar to the Wright ataximeter<sup>28</sup> attached to the waist of participants. Volunteers were instructed to stand comfortably on a firm surface with their feet slightly apart and eyes closed for 2 minutes. In previous studies, CNS-stimulant compounds demonstrated reductions in body sway; for example, modafinil (200 mg), dexamphetamine (20 mg) and clinical doses (average 20 mg) of methylphenidate reduced body sway by

approximately 35%<sup>29</sup>, 19.4%<sup>15</sup> and 36.8%, respectively.<sup>30</sup> Conversely, CNS-depressant compounds such as benzodiazepines are associated with increased body sway; for example, diazepam (10 mg) increased body sway by 119%.<sup>31</sup>

#### SMOOTH PURSUIT EYE MOVEMENTS.

The computerised smooth pursuit measurement was performed as described previously<sup>32</sup> and used in many studies to assess drug effects.<sup>14</sup> During this test, participants followed a light source with their eyes, that moved continuously in a horizontal direction on a screen placed 58 cm away. The outcome of smooth pursuit was defined as the percentage of time the participant's eyes were in smooth pursuit of the target for each stimulus velocity and frequency. In a previous study, the velocity of smooth pursuit eye movements was impaired by diazepam (10 mg).<sup>32</sup> Improvements in smooth pursuit of approximately 6% have been reported with CNS stimulants such as methylphenidate (average 20 mg).<sup>30</sup>

#### SACCADIC EYE MOVEMENTS.

The computerised measurement of saccadic eye movements was performed as described previously<sup>33</sup> and used in many pharmacological studies.<sup>14</sup> Briefly, to measure saccadic eye movements, participants were positioned identically to when performing the smooth pursuit measurement and instructed to follow a light source that jumped from side to side.<sup>33</sup> The parameter collected was saccadic peak velocity (SPV) in degrees/second (deg/s). Previous studies have demonstrated that CNS-stimulant compounds such as caffeine (60 mg), modafinil (200 mg) and dexamphetamine (20 mg) increased the average SPV by 11.6 deg/s,<sup>34</sup> 24.6 deg/s,<sup>29</sup> and 12.7 deg/s,<sup>15</sup> respectively. CNS-depressant compounds such as benzodiazepines have been shown to decrease SPV.<sup>35</sup>

#### ADAPTIVE TRACKING TEST

Adaptive tracking tests have been used in many pharmacological studies to evaluate visuomotor coordination and vigilance.<sup>14</sup> In this study, we used an adaptive tracking test according to specifications from Borland and Nicholson.<sup>36</sup> During the test, a circle moved randomly on a screen. Participants were given a joystick and instructed to use it to keep a dot within the moving circle. When an effort was successful, the speed of the moving circle increased. Conversely, the speed decreased if the participant was not able to maintain the dot within the circle, resulting in a constant and individually adapted challenge throughout the procedure. The outcome of the test is the average speed of the moving circle as a percentage of the maximum speed of the circle. In previous studies, CNS-stimulant compounds such as caffeine (60 mg), modafinil (200 mg), dexamphetamine (20 mg) and methylphenidate (average 20 mg) improved

average adaptive tracking by approximately 1.6%,<sup>34</sup> 1.8%,<sup>29</sup> 4.2%,<sup>15</sup> and 2.2%,<sup>30</sup> respectively. For CNS-depressant compounds, such as benzodiazepines, an impairment of adaptive tracking has been demonstrated.<sup>14</sup>

#### STROOP COLOURED WORD TEST.

The Stroop effect test involves identifying the colour of coloured words,<sup>37</sup> many CNS-active compounds have an effect on this test.<sup>38-40</sup> In this study, we used a computer-adapted version from the Psychology Software Tools website (<https://pstnet.com/products/e-prime/>), comprising two subtests as described in a previous publication.<sup>41</sup> In the first subtest, six coloured items were presented at random. The possible colours were green, red and blue, and each colour corresponded to a number key on the numpad section of the keyboard; green corresponded with 1, red with 2 and blue with 3. Participants were instructed to place the index, middle and ring fingers of their dominant hand on keys 1, 2 and 3. When a coloured item appeared on the screen, participants were to press the corresponding key as quickly as possible. In the second subtest, which immediately followed the first, 34 colour and word pairs were presented randomly. The words that were used were 'red', 'green' and 'blue', and the colour and word pairs were either congruent or incongruent matches. Again, the participants were asked to identify the correct colour as quickly as possible by pressing either keys 1, 2 or 3 on the numpad. Each item or word was shown for 4 seconds, and there was a 0.5 s pause after every response. Two parameters were derived from this test: Stroop 1 is the difference in reaction time between correct congruent and correct incongruent answers (ms) and Stroop 2 is the number of correct responses in incongruent trials. Previous studies demonstrated that benzodiazepines impair performance on this test.<sup>38</sup>

#### BOND AND LADER AND BOWDLE VISUAL ANALOGUE SCALES.

VAS, as originally described by Norris, have commonly been used to quantify the subjective effects of sedative agents.<sup>35,42</sup> Subjects were instructed to use the computer mouse to select their response to each VAS item. The Bond and Lader VAS involved collecting scores from 16 horizontal scales related to how a person feels. From these measurements, three main factors, namely 'alertness', 'mood' and 'calmness', were calculated as described in previous publications.<sup>34,43</sup> Benzodiazepines have consistently shown reductions on VAS alertness,<sup>44</sup> whereas variable but consistent increases are observed with caffeine,<sup>34</sup> dexamphetamine,<sup>15</sup> modafinil,<sup>29</sup> and methylphenidate.<sup>30</sup> Psychedelic effects were measured using the Bowdle VAS as previously described.<sup>45</sup> This scale consists of 13 items on which three summary scales (internal perception, external perception and 'feeling high') are calculated using log transformation as described in previous publications.<sup>46</sup> Dexamphetamine (20 mg) has been shown to increase the summary scale 'feeling high',<sup>15</sup> in contrast to the other CNS stimulants mentioned earlier.

#### STATISTICAL ANALYSIS

Analyses were performed using SAS software version 9.4 (SAS Institute, Cary, NC, USA). Residual Q-Q plots were produced for all NeuroCart parameters to check the assumption of normality of the error term in the mixed effects models. This was done by visual inspection and the Shapiro–Wilk test statistic. To assess the treatment effects, data for each parameter were analysed with a mixed model analysis of covariance. We defined treatment, time, period and treatment by time as fixed factors; subject, subject by treatment and subject by time as random factors; and the (average) baseline measurement per study period as a covariate. The Kenward–Roger approximation was used to estimate denominator degrees of freedom and the model parameters were estimated using the restricted maximum likelihood method. Individual treatment effects over the 4-hour post-dose time period for the different doses were reported with the least squares mean estimated difference, the two-sided 90% confidence interval (CI) and the *P*-value. Owing to the exploratory nature of this study, a 90% CI instead of 95% CI was deemed sufficient. Next to that, no correction for multiple comparisons was performed as due to the exploratory nature of this study, hypothesis testing was not used in the strict way, but to guide the direction of future research.

#### RESULTS

##### DEMOGRAPHICS


In total, 69 volunteers were screened, of which 24 healthy volunteers (23 male and 1 female of non-childbearing potential) between 18 and 55 years of age were included (Table 2). Subject disposition can be found in Figure 2. All participants completed the three study periods.

**Table 2** Demographic characteristics of study participants

Characteristic	Subjects enrolled (N = 24)
AGE (YEARS)	
mean (SD) <sup>1</sup>	27.9 (9.0)
SEX, N (%)	
Female	1 (4.2%)
Male	23 (95.8%)
WEIGHT (KG)	
mean (SD)	79.12 (10.81)
HEIGHT (CM)	
mean (SD)	181.98 (9.88)
BMI (KG/M <sup>2</sup> )	
mean (SD)	23.92 (2.85)

1. SD: standard deviation; 2. BMI: body mass index

**Figure 2** Subject disposition



**PHARMACOKINETIC ASSESSMENTS**

As reported in our previous publication, mean (SD) TAK-653 plasma levels for the 0.5 mg dose were 0.99 (0.94) ng/ml at 0.5 h post-dose and 4.19 (0.83) ng/ml at 2.5 h post-dose.<sup>22</sup> Plasma levels for the 6 mg dose were 2.57 (3.29) ng/ml at 0.5 h post-dose and 45.99 (8.84) ng/ml at 2.5 h post-dose.<sup>22</sup>

**FUNCTIONAL PHARMACODYNAMIC NEUROCARD ASSESSMENTS**

All NeuroCart parameters were normally distributed, except body sway measurements, which were log-normal distributed and therefore natural log transformation was applied for their analysis. For interpretation back transformation was applied. To calculate summary scores for VAS Bowdle, log transformation was performed as well as described in previous publications.<sup>46</sup> Results are summarised in Table 3. On smooth pursuit eye movements (%), a clear statistically significant improvement was observed with the TAK-653 6 mg dose (Figure 3). At the same dose level, a similar improvement was observed for adaptive tracking (%) (Figure 4). Both doses of TAK-653 increased SPV (deg/s) to a similar extent (Figure 5).


**Table 3** Least squares mean overall treatment effects and individual treatment effects of TAK-653 0.5 mg and 6 mg over the 4-hour post-dose period.

	Least squares mean			Contrasts (90% CI) p-value	
	Placebo	TAK-653 0.5 mg	TAK-653 6 mg	TAK-653 0.5 mg vs. placebo Estimate of difference, 90% CI <sup>1</sup> , p-value	TAK-653 6 mg vs. placebo Estimate of difference, 90% CI <sup>1</sup> , p-value
Body sway, log (mm) (N=24)	202.3	199.6	185.7	-1.3% (-13.4%, 12.4%) P = 0.86	-8.2% (-19.4%, 4.6%) P = 0.28
Smooth pursuit (%) (N=24)	44.5	44.8	46.8	0.26 (-1.69, 2.21) P = 0.82	2.32 (0.37, 4.27) P = 0.05*
Saccadic peak velocity (deg/s) (N=24)	475.5	495.0	490.9	19.49 (5.98, 32.99) P = 0.02*	15.40 (1.91, 28.90) P = 0.06*
Adaptive tracking (%) (N=24)	30.8	31.2	32.5	0.41 (-0.73, 1.56) P = 0.55	1.68 (0.51, 2.84) P = 0.02*
Stroop 1 <sup>3</sup> (ms) (N=24)	71.4	93.4	71.0	22.0 (4.0, 40.0) P = 0.05*	-0.5 (-18.3, 17.3) P = 0.96
Stroop 2 <sup>4</sup> (ms) (N=24)	16.7	16.4	16.6	-0.3 (-0.5, -0.1) P = 0.02*	-0.1 (-0.3, 0.2) P = 0.61
VAS <sup>2</sup> alertness (mm) (N=24)	49.9	50.5	50.7	0.65 (-0.38, 1.67) P = 0.30	0.77 (-0.24, 1.79) P = 0.21
VAS <sup>2</sup> calmness (mm) (N=24)	52.4	52.4	52.2	-0.03 (-1.65, 1.60) P = 0.98	-0.25 (-1.88, 1.38) P = 0.80
VAS <sup>2</sup> mood (mm) (N=24)	51.5	51.8	52.0	0.28 (-0.29, 0.85) P = 0.41	0.48 (-0.10, 1.05) P = 0.17
VAS <sup>2</sup> external, log (mm) (N=24)	0.35	0.35	0.35	-0.01 (-0.03, 0.01) P = 0.63	-0.01 (-0.03, 0.01) P = 0.58
VAS <sup>2</sup> internal, log (mm) (N=24)	0.35	0.35	0.35	< 0.00 (-0.01, 0.01) P = 0.92	< 0.00 (-0.01, 0.02) P = 0.76
VAS <sup>2</sup> 'feeling high', log (mm) (N=24)	0.37	0.35	0.34	-0.02 (-0.05, 0.01) P = 0.25	-0.03 (-0.06, -0.01) P = 0.05*


1. CI: confidence interval; 2. VAS: Visual Analogue Scale; 3. Stroop 1 is the difference in reaction time between correct congruent and correct incongruent answers; 4. Stroop 2 is the number of correct responses in incongruent trials.

\*Indicates a statistically significant effect.


**Figure 3** Smooth pursuit eye movements: change from baseline time effect profile of the least square (LS) mean 90% confidence interval (CI)



**Figure 4** Adaptive tracking: change from baseline time effect profile of the least square (LS) mean 90% confidence interval (CI)



**Figure 5** Saccadic peak velocity: change from baseline time effect profile of the least square (LS) mean 90% confidence interval (CI)



The VAS Bowdle subscale 'feeling high' remained stable under TAK-653 0.5 mg and 6 mg; however, average VAS-high increased with placebo, resulting in a statistically significant reduction with TAK-653 6 mg. A review of the raw data revealed that this effect was caused by one subject who indicated a 20 mm (large) increase in VAS-high after placebo. This entry was judged to be an artefact, given 'feeling high' does not occur spontaneously or under placebo and the subject did not have AEs indicating subjective drug effects such as 'feeling abnormal', 'feeling drunk' or 'feeling high'.

On the Stroop coloured word test TAK-653 0.5 mg increased the difference in reaction time between correct congruent and correct incongruent answers and decreased the number of correct responses in incongruent trials compared to placebo whereas TAK-653 6 mg did not affect any parameter of the Stroop coloured word test.

No significant effects were observed on body sway (%), other VAS Bowdle subscales (mm) or any of the Bond and Lader VAS subscales (mm) (Table 3).

#### SAFETY AND TOLERABILITY

For details on TAK-653's safety and tolerability in this study, please refer to previous reported results.<sup>22</sup> In summary, TAK-653 was well tolerated, no serious AEs were observed and there were no withdrawals related to an AE. The most frequently reported AEs after administration of TAK-653 were somnolence (TAK-653 0.5 mg: 3 of 24 subjects [12.5%], TAK-653 6 mg: 3 of 24 subjects [12.5%]), headache (TAK-653 0.5 mg: 1

of 24 subjects [4.2%], TAK-653 6 mg: 4 of 24 subjects [16.7%]) and nasopharyngitis (TAK-653 0.5 mg: 3 of 24 subjects [12.5%], TAK-653 6 mg 1 of 24 subjects [4.2%]). Of these AEs, somnolence and headache were reported after administration of placebo as well (2 of 24 subjects [8.3%] each). No clinically significant effects on vital signs, ECGs or laboratory measurements were observed. Of note is that no AEs of seizure, dissociative effects or euphoria were observed.

## DISCUSSION

Similar to CNS stimulant compounds TAK-653 increased SPV, SP and adaptive tracking at the time maximum plasma concentration was reached. These effects were more pronounced with 6 mg than with 0.5 mg TAK-653. TAK-653 increased SPV at both 0.5 mg and 6 mg, while smooth pursuit eye movements and adaptive tracking increased at 6 mg but not at 0.5 mg. The effects of TAK-653 on the NeuroCart tests contrasted with the effects of CNS-depressant compounds such as benzodiazepines, which have been shown to decrease smooth pursuit eye movements,<sup>32</sup> SPV,<sup>44</sup> and adaptive tracking.<sup>14</sup> The absence of an effect of TAK-653 on any of the VAS subscales supports the finding that TAK-653 is devoid of subjective mood-related derangements observed with other CNS-stimulant compounds such as dexamphetamine.<sup>15</sup> When comparing the acute pharmacodynamic NeuroCart profile of TAK-653 to known profiles of CNS stimulant and CNS depressant compounds, TAK-653's profile is suggestive of stimulatory CNS effects. This is consistent with the TMS-EMG (electromyography) findings demonstrating increased cortical excitability with the 6 mg dose.<sup>22</sup> While one could argue that the observed effects on the NeuroCart tests are due to TMS itself, this can be ruled out for this study as a placebo arm was included and the effects of the different doses of TAK-653 on the NeuroCart were observed compared to placebo.

Compared to clinical doses of psychostimulants previously characterised using Neurocart test, TAK-653's stimulatory CNS effects appear more limited. TAK-653 increased SPV by 15.5 to 19.5 deg/s, which is larger than caffeine 60 mg (11.6 deg/s)<sup>34</sup> and dexamphetamine 20 mg (12.7 deg/s),<sup>15</sup> but smaller than modafinil 200 mg (24.6 deg/s).<sup>29</sup> Increases in smooth pursuit eye movements represented only roughly one-third of those induced by methylphenidate.<sup>30</sup> The increase in adaptive tracking of 1.6% with TAK-653 6 mg was comparable to modafinil (1.8%),<sup>29</sup> caffeine (1.6%),<sup>34</sup> and methylphenidate (2.2%),<sup>30</sup> but smaller than dexamphetamine (4.2%).<sup>15</sup> Next to that, decreases in BS have been observed for other CNS stimulant compounds, but no effect of both dose levels TAK-653 on BS was observed. Although direct comparisons should be made for an unequivocal interpretation of our findings, TAK-653 seems to have a novel stimulatory CNS profile that is generally more subtle than clinical doses of

known psychostimulants, and distinguishes itself by a relatively large stimulatory effect on saccadic peak velocity but devoid of any subjective mood-related derangement such as dysphoria, anxiety or feeling high.

Although TAK-653 demonstrated psychostimulant effects, its impact on different aspects of cognition was less consistent. The Stroop test was included as it can be helpful in understanding complex attention, perception and elements of executive function.<sup>47</sup> TAK-653 0.5 mg but not 6 mg increased the difference in reaction time between correct congruent and correct incongruent answers. Similarly, TAK-653 0.5 mg but not 6 mg decreased the number of correct responses in incongruent trials. It cannot be excluded that the lower dose may affect aspects of cognitive functioning, which are obscured at a higher dose. The overall pharmacodynamic profile however, provides no reason to assume a bell-shaped dose-response curve. Therefore, the Stroop results are currently best considered as a potential type I error of a less robust test with multiple complex endpoints.

The PK results of this study were in line with results from initial healthy volunteer studies, as mean maximal plasma concentrations for 0.5 and 6 mg TAK-653 were comparable to those observed at similar dose levels.<sup>11</sup> Therefore, although the area under the curve from time 0 to infinity ( $AUC_{\infty}$ ) was not determined in the current study, this was expected to correspond to the  $AUC_{\infty}$  observed in initial healthy volunteer studies with 406 h\*ng/ml and 3167 h\*ng/ml for TAK-653 0.5 and 6 mg, respectively. These data support dose and concentration dependence since more pronounced effects were observed with the 6 mg dose compared to the 0.5 mg dose.

A limitation of this study is that full dose/concentration-response characterisation was precluded by safety concerns. Given both AMPA receptor PAMs and TMS are associated with an increased albeit very limited risk of convulsions,<sup>10</sup> 6 mg was selected as the highest dose as it was expected to yield a mean maximum plasma concentrations well below those at which partial seizures were observed in primates (Takeda internal data).

Taken together, the PD profile of TAK-653 was characterised in this study according to recommendations by the 'QBDD' framework.<sup>12</sup> As hypothesised based on its mechanism of action of AMPA receptor PAM, TAK-653 demonstrated an acute functional PD profile of CNS stimulatory effects on the NeuroCart. This confirms BBB penetration and, moreover, target engagement that is consistent with the previously reported TMS results of increased cortical excitability.<sup>22</sup> No undesired pharmacological effects associated with AMPA receptor stimulation, such as seizures or euphoria, off-target effects or unexpected AEs, were observed in this acute dosing study. The insights obtained in this study, can be used to design future studies in both healthy individuals and selected patient populations that are hypothesised to benefit from AMPA receptor-mediated stimulatory CNS effects.

REFERENCES

- 1 Berman RM, Cappiello A, Anand A, Oren DA, Heninger GR, Charney DS, Krystal JH. Antidepressant effects of ketamine in depressed patients. *Biol Psychiatry*. 2000;47(4):351-354.
- 2 Murrough JW, Iosifescu DV, Chang LC, Al Jurdi RK, Green CE, Perez AM, et al. Antidepressant efficacy of ketamine in treatment-resistant major depression: a two-site randomized controlled trial. *Am J Psychiatry*. 2013;170(10):1134-1142. doi:10.1176/appi.ajp.2013.13030392
- 3 Zarate Jr. CA, Singh JB, Carlson PJ, Brutsche NE, Ameli R, Luckenbaugh DA, et al. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch Gen Psychiatry*. 2006;63(8):856-864. doi:10.1001/archpsyc.63.8.856
- 4 Collo G, Cavalleri L, Chiamulera C, Merlo Pich E. (2R,6R)-Hydroxynorketamine promotes dendrite outgrowth in human inducible pluripotent stem cell-derived neurons through AMPA receptor with timing and exposure compatible with ketamine infusion pharmacokinetics in humans. *Neuroreport*. 2018;29(16):1425-1430. doi:10.1097/WNR.000000000000131
- 5 Zanos P, Moaddel R, Morris PJ, Georgiou P, Fischell J, Elmer GI, et al. NMDAR inhibition-independent antidepressant actions of ketamine metabolites. *Nature*. 2016;533(7604):481-486. doi:10.1038/nature17998
- 6 Li N, Lee B, Liu RJ, Banasr M, Dwyer JM, Iwata M, et al. mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science* (80-). 2010;329(5994):959-964. doi:10.1126/science.1190287
- 7 Shaffer CL, Dutra JK, Tseng WC, Weber ML, Bogart LJ, Hales K, et al. Pharmacological evaluation of clinically relevant concentrations of (2R,6R)-hydroxynorketamine. *Neuropharmacology*. 2019;153:73-81. doi:10.1016/j.neuropharm.2019.04.019
- 8 Duman RS, Li N, Liu RJ, Duric V, Aghajanian G. Signaling pathways underlying the rapid antidepressant actions of ketamine. *Neuropharmacology*. 2012;62(1):35-41. doi:https://doi.org/10.1016/j.neuropharm.2011.08.044
- 9 Maeng S, Zarate Jr. CA, Du J, Schloesser RJ, McCammon J, Chen G, et al. Cellular mechanisms underlying the antidepressant effects of ketamine: role of alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptors. *Biol Psychiatry*. 2008;63(4):349-352. doi:10.1016/j.biopsych.2007.05.028
- 10 Leo A, Giovannini G, Russo E, Meletti S. The role of AMPA receptors and their antagonists in status epilepticus. *Epilepsia*. 2018;59(6):1098-1108. doi:10.1111/epi.14082
- 11 Asgharnejad Xu, Dong, Iadevaia, Dornier A, Murthy V, Ratti E, et al. Pharmacokinetic and Pharmacodynamic Properties of the Investigational AMPA Receptor Positive Allosteric Modulator TAK-653 After Single and Multiple Rising Doses in Healthy Volunteers. *Neuropsychopharmacology*. 43, S157 (2018).
- 12 Cohen AF, Burggraaf J, van Gerven JM, Moerland M, Groeneveld GJ. The use of biomarkers in human pharmacology (Phase I) studies. *Annu Rev Pharmacol Toxicol*. 2015;55:55-74. doi:10.1146/annurev-pharmtox-011613-135918
- 13 Morgan P, Van Der Graaf PH, Arrowsmith J, Feltner DE, Drummond KS, Wegner CD et al. Can the flow of medicines be improved? Fundamental pharmacokinetic and pharmacological principles toward improving Phase II survival. *Drug Discov Today*. 2012;17(9-10):419-424. doi:10.1016/j.drudis.2011.12.020
- 14 Groeneveld GJ, Hay JL, Van Gerven JM. Measuring blood-brain barrier penetration using the NeuroCart, a CNS test battery. *Drug Discov Today Technol*. 2016;20:27-34. doi:http://dx.doi.org/10.1016/j.dtttec.2016.07.004
- 15 Recourt K, van der Aart J, Jacobs G, de Kam M, Drevets W, van Nueten L, et al. Characterisation of the pharmacodynamic effects of the P2X7 receptor antagonist JNJ-54175446 using an oral dexamphetamine challenge model in healthy males in a randomised, double-blind, placebo-controlled, multiple ascending dose trial. *J Psychopharmacol*. 2020;34(9):1030-1042. doi:10.1177/0269881120914206
- 16 Chen X, Broeyer F, de Kam M, Baas J, Cohen A, van Gerven J. Pharmacodynamic response profiles of anxiolytic and sedative drugs. *Br J Clin Pharmacol*. 2017;83(5):1028-1038. doi:10.1111/bcp.13204
- 17 Chen X, Jacobs G, de Kam ML, et al. AZD6280, a novel partial  $\gamma$ -aminobutyric acid A receptor modulator, demonstrates a pharmacodynamically selective effect profile in healthy male volunteers. *J Clin Psychopharmacol*. 2015;35(1):22-33. doi:10.1097/JCP.0000000000000251
- 18 Chen X, Jacobs G, de Kam, Jaeger J, Lappalainen J, Maruff P, et al. The central nervous system effects of the partial GABA-A $\alpha$ 2,3-selective receptor modulator AZD7325 in comparison with lorazepam in healthy males. *Br J Clin Pharmacol*. 2014;78(6):1298-1314. doi:10.1111/bcp.12413
- 19 de Haas SL, de Visser SJ, van der Post JP, de Smet M, Schoemaker RC, Rijnbeek B, et al. Pharmacodynamic and pharmacokinetic effects of TPA023, a GABA $\alpha$ 2,3 subtype-selective agonist, compared to lorazepam and placebo in healthy volunteers. *J Psychopharmacol*. 2006;21(4):374-383. doi:10.1177/0269881106072343
- 20 de Haas SL, Franson KL, Schmitt JA, Cohen AF, Fau JB, Dubruc C, van Gerven JMA. The pharmacokinetic and pharmacodynamic effects of SL65,1498, a GABA-A $\alpha$ 2,3 selective agonist, in comparison with lorazepam in healthy volunteers. *J Psychopharmacol*. 2009;23(6):625-632. doi:10.1177/0269881108092595
- 21 de Haas SL, de Visser SJ, van der Post JP, Schoemaker RC, van Dyck K, Murphy MG, et al. Pharmacodynamic and pharmacokinetic effects of MK-0343, a GABA(A)  $\alpha$ 2,3 subtype selective agonist, compared to lorazepam and placebo in healthy male volunteers. *J Psychopharmacol*. 2008;22(1):24-32. doi:10.1177/0269881107082108
- 22 O'Donnell P, Dijkstra FM, Damar U, Quanhong L, de Goede A, Xu L, et al. Transcranial magnetic stimulation as a translational biomarker for AMPA receptor modulation. *Transl Psychiatry*. 2021;11(1):325. doi:10.1038/s41398-021-01451-2
- 23 Posner K, Brown GK, Stanley B, Posner, Brent DA, Yershova KV, et al. The Columbia-Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *Am J Psychiatry*. 2011;168(12):1266-1277. doi:10.1176/appi.ajp.2011.10.11704
- 24 van Steveninck AL, Schoemaker HC, Pieters MS, Kroon R, Breimer DD, Cohen AF. A comparison of the sensitivities of adaptive tracking, eye movement analysis and visual analog lines to the effects of incremental doses of temazepam in healthy volunteers. *Clin Pharmacol Ther*. 1991;50(2):172-180.
- 25 van Steveninck AL, van Berckel BN, Schoemaker RC, Breimer DD, van Gerven JM, Cohen AF. The sensitivity of pharmacodynamic tests for the central nervous system effects of drugs on the effects of sleep deprivation. *J Psychopharmacol*. 1999;13(1):10-17. doi:10.1177/026988119901301012
- 26 van Steveninck AL, Wallnofer AE, Schoemaker RC, Pieters MS, Danhof M, van Gerven JMA, et al. A study of the effects of long-term use on individual sensitivity to temazepam and lorazepam in a clinical population. *Br J Clin Pharmacol*. 1997;44(3):267-275. doi:10.1046/j.1365-2125.1997.t011-00580.x
- 27 van Steveninck AL, Gieschke R, Schoemaker HC, Pieters MS, Kroon JM, Breimer DD, et al. Pharmacodynamic interactions of diazepam and intravenous alcohol at pseudo steady state. *J Psychopharmacol*. 1993;11(4):471-478. doi:10.1007/bf02244655
- 28 Wright BM. A simple mechanical ataxia-meter. *J Physiol*. 1971;218 Suppl:27p-28p.
- 29 Baakman AC, Zuiker R, van Gerven JMA, Gross N, Yang R, Fetell M, et al. Central nervous system effects of the histamine-3 receptor antagonist CEP-26401, in comparison with modafinil and donepezil, after a single dose in a cross-over study in healthy volunteers. *Br J Clin Pharmacol*. 2019;85(5):970-985. doi:10.1111/bcp.13885
- 30 Klein RH. Minimally invasive methodology for pharmacological research involving children. Published online 2013.
- 31 van Steveninck AL, Gieschke R, Schoemaker RC, Roncari G, Tuk B, Pieters MS, et al. Pharmacokinetic and pharmacodynamic interactions of bretazenil and diazepam with alcohol. *Br J Clin Pharmacol*. 1996;41(6):565-573. doi:10.1046/j.1365-2125.1996.38514.x
- 32 Bittencourt PR, Wade P, Smith AT, Richens A. Benzodiazepines impair smooth pursuit eye movements. *Br J Clin Pharmacol*. 1983;15(2):259-262. doi:10.1111/j.1365-2125.1983.tb01495.x
- 33 Baloh RW, Sills AW, Kumlley WE, Honrubia V. Quantitative measurement of saccade amplitude, duration, and velocity. *Neurology*. 1975;25(11):1065-1070. doi:10.1212/wnl.25.11.1065
- 34 Wilhelmus MM, Hay JL, Zuiker RG, Okkerse P, Perdrieu C, Sauser J, et al. Effects of a single, oral 60 mg caffeine dose on attention in healthy adult subjects. *J Psychopharmacol*. 2017;31(2):222-232. doi:10.1177/0269881116668593
- 35 de Visser SJ, van der Post J, Pieters MS, Cohen AF, van Gerven JM. Biomarkers for the effects of antipsychotic drugs in healthy volunteers. *Br J Clin Pharmacol*. 2001;51(2):119-132.
- 36 Borland RG, Nicholson AN. Visual motor co-ordination and dynamic visual acuity. *Br J Clin Pharmacol*. 1984;18 Suppl 1(Suppl 1):69S-72S. https://www.ncbi.nlm.nih.gov/pubmed/6525331 https://www.ncbi.nlm.nih.gov/pmc/PMC1463351/
- 37 Stroop JR. Studies of interference in serial verbal reactions. *J Exp Psychol*. 1935;18(6):643-662. doi:10.1037/h0054651
- 38 Griffiths AN, Jones DM, Richens A. Zopiclone produces effects on human performance similar to flurazepam, lormetazepam and triazolam. *Br J Clin Pharmacol*. 1986;21(6):647-653. doi:10.1111/j.1365-2125.1986.tb05229.x
- 39 Bohnen NI, Kaufer DI, Hendrickson R, Ivancio LS, Lopresti BJ, Koeppe RA, et al. Degree of inhibition of cortical acetylcholinesterase activity and cognitive effects by donepezil treatment in Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 2005;76(3):315-319. doi:10.1136/jnnp.2004.038729
- 40 Rose AK, Duka T. The influence of alcohol on basic motoric and cognitive disinhibition. *Alcohol Alcohol*. 2007;42(6):544-551. doi:10.1093/alcac/agm073
- 41 Liem-Moolenaar M, te Beek ET, de Kam ML, Franson KL, Kahn RS, Hijman R, et al. Central nervous system effects of haloperidol on THC in healthy male volunteers. *J Psychopharmacol*. 2010;24(11):1697-1708. doi:10.1177/0269881109358200
- 42 Norris H. The action of sedatives on brain stem oculomotor systems in man. *Neuropharmacology*. 1971;10(21):181-191.
- 43 Hoefer P, de Haas S, Winkler J, Schoemaker RC, Chioffi E, van Gerven J, Dingemans J, et al. Orexin receptor antagonism, a new sleep-promoting paradigm: an ascending single-dose study with almorexant. *Clin Pharmacol Ther*. 2010;87(5):593-600. doi:10.1038/clpt.2010.19
- 44 De Visser SJ. A question based approach to drug development. 2003;PhD.
- 45 Bowdle AT, Radant AD, Cowley DS, Kharasch ED, Strassman RJ, Roy-Byrne PP. Psychedelic effects of ketamine in healthy volunteers relationship to steady-state plasma concentrations. *Anesthesiology*. 1998;88(1):82-88. doi:10.1097/0000542-199801000-00015
- 46 Zuurman L, Roy C, Schoemaker RC, Hazekamp A, den Hartigh J, Bender J, et al. Effect of intrapulmonary tetrahydrocannabinol administration in humans. *J Psychopharmacol*. 2008;22(7):707-716. doi:10.1177/0269881108089581
- 47 Gerstadt CL, Hong YJ, Diamond A. The relationship between cognition and action: performance of children 3 1/2-7 years old on a Stroop-like day-night test. *Cognition*. 1994;53(2):129-153. doi:10.1016/0010-0277(94)90068-x

## APPENDIX

### IB-DERISK ANALYSER OVERVIEW OF TAK-653

TAK-653 is an alpha-amino-3-hydroxy-5-methyl-4-isoxazole receptor (AMPA) positive allosteric modulator (PAM) in clinical development for the treatment of major depressive disorder (MDD). When sorted on  $C_{max}$  values in the IB-Derisk analyser tool, TAK-653 demonstrated an overall favourable profile, with desired green pharmacological effects at lower exposures followed by a range of acceptable yellow side effects, and only at higher exposures unacceptable orange and severe red side effects emerged.<sup>1</sup> It was noticeable however, that at dose levels above the NOAEL, the dose-response curve for adverse effects was relatively steep. This indicates that severe side effects (red), such as tonic-clonic seizures, occurred at exposure levels following exposure levels where only mild side effects (yellow), such as tremors, had been observed. In this respect, the IB-Derisk analyser overview did not show the optimal pattern of side effects gradually increasing in severity, where mild (yellow) adverse effects are followed by more severe (orange) adverse effects and ultimately severe (red) adverse effects.

In line with the excitatory/stimulatory effects expected from an AMPA-PAM, in preclinical experiments in rats, monkeys and mice convulsions were observed at exposure levels at least a factor 10 higher than levels associated with desired pharmacological effects. In rabbits, which were used for reproductive and developmental toxicity studies, no convulsions were observed. Desired pharmacological effects consisted of positive results observed on *in vivo* behavioural experiments testing memory and antidepressant-like effects in rats.

A dose range between 0.3 and 18 mg was tested in a first-in-human (FIH) study with TAK-653.<sup>2</sup> The exposure level associated with the highest dose of 18 mg was still a factor 13 below the values associated with convulsions in preclinical experiments, while it fell in the range associated with desired pharmacological effects. As expected based on the preclinical data, the dose range tested in the FIH study was well tolerated, specifically no tremors or convulsions were observed.


Since the FIH study with TAK-653 was primarily geared towards characterising safety and PK, a second clinical study was set up to assess the pharmacodynamics of TAK-653. In rats TAK-653 had shown to increase mechanomyography (MMG) amplitude elicited by Transcranial Magnetic Stimulation (TMS) of the motor cortex. It was therefore decided to perform a comparable measurement in humans, namely motor evoked potential (MEP) amplitude after TMS stimulation of the motor cortex. Because of the theoretical increased risk of seizures by potentiating glutamatergic synapses by both TMS and AMPA receptor potentiation, dose selection for the clinical study was

performed with great caution.<sup>3</sup> Doses selected for the human study were 0.5 and 6 mg, the estimated  $C_{max}$  values of these doses were expected to be within the anticipated dose range to be used in a phase 2 study of TAK-653 to obtain maximum treatment response while maintaining sufficient margins to the observed convulsions in pre-clinical studies. The results of the study demonstrated that increases in the amplitude of MMG in rats and MEPS in humans arose at comparable exposure levels (Figure 1).<sup>4</sup> Thereby demonstrating that MMG or MEPS as elicited by TMS can be used as a translational biomarker for AMPA receptor modulation.<sup>4</sup>

In addition to measuring TMS effects in humans, the Neurocart was performed as well with the 0.5 and 6 mg dose.<sup>5</sup> In line with TAK-653's mechanism of action and the findings of the TMS assessments, TAK-653 demonstrated a psychostimulant-like profile on the NeuroCart without unwanted effects of for example euphoria often observed with other psychostimulant compounds such as amphetamine.<sup>5</sup> While some small psychostimulant effects consisting of increased saccadic peak velocity, were observed with the 0.5 mg dose, the 6 mg dose demonstrated a more pronounced psychostimulant profile consisting of improved adaptive tracking, increased saccadic peak velocity and improved smooth pursuit eye movements.<sup>5</sup>

Taken together, these studies demonstrate how the overview of the available pre-clinical and clinical data on a compound obtained with the IB-Derisk analyser tool, can be used to guide decisions for early phase clinical studies. TAK-653 demonstrated consistency in pharmacokinetics and exposure-effect relations across species, which was reassuring for the clinical study. Furthermore, desired pharmacological effects were observed at much lower exposure levels than undesired effects. Based on the overview obtained with the IB Derisk analyser tool, we were able to select dose levels to study TAK-653's pharmacodynamics that were expected to sort out an effect, but still had sufficient margins to avoid seizures.

**Figure 1** IB-Derisk analyser overview TAK-653



Abbreviations: MEP, motor evoked potential; NOEL, no observed adverse effect level; CNS, central nervous system

**REFERENCES**

- 1 van Gerven J, Cohen A. Integrating data from the Investigational Medicinal Product Dossier/investigator's brochure. A new tool for translational integration of preclinical effects. *Br J Clin Pharmacol.* 2018;84(7):1457-1466. doi:10.1111/bcp.13529
- 2 Asgharnejad M, Zicha S, Lin X, Cheng D, Iadevaia S, Dörner A, Murthy V RE. Pharmacokinetic and pharmacodynamic properties of the investigational AMPA receptor positive allosteric modulator TAK-653 after single and multiple rising doses in healthy volunteers. *Poster Present Am Coll Neuropsychopharmacol 57th Annu Meet 9-13 Dec, 2018; Hollywood, Florida.* Published online 2018.
- 3 Rossi S, Hallett M, Rossini PM, Pascual-Leone A. Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clin Neurophysiol.* 2009;120(12):2008-2039. doi:10.1016/j.clinph.2009.08.016
- 4 O'Donnell P, Dijkstra FM, Damar U, et al. Transcranial magnetic stimulation as a translational biomarker for AMPA receptor modulation. *Transl Psychiatry.* 2021;11(1):325. doi:10.1038/s41398-021-01451-2
- 5 Dijkstra F, O'Donnell P, Klaassen E, et al. Central nervous system effects of TAK-653, an investigational alpha-amino-3-hydroxy-5-methyl-4-isoxazole receptor (AMPA) positive allosteric modulator in healthy volunteers. *Transl Psychiatry.* 2022;12(1):408. doi:10.1038/s41398-022-02148-w