



Universiteit  
Leiden

The Netherlands

## **Braver together: an exploration into the effectiveness of blended group cognitive behavior therapy as early intervention for socially anxious adolescents**

Velthuisen S.L.M.

### **Citation**

*Braver together: an exploration into the effectiveness of blended group cognitive behavior therapy as early intervention for socially anxious adolescents.* (2025, December 17). *Braver together: an exploration into the effectiveness of blended group cognitive behavior therapy as early intervention for socially anxious adolescents.* Retrieved from <https://hdl.handle.net/1887/4285450>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4285450>

**Note:** To cite this publication please use the final published version (if applicable).



# Chapter 4

---

## Does Cognitive Behavior Therapy Change Socially Anxious Adolescents' Behavior During a Public Speaking Task?

---

Based on:

Velthuisen, S. L. M., van den Bos, E., Miers, A. C., Chen, J., Westenberg, P. M. (in press). Does cognitive behavior therapy change socially anxious adolescents' behavior during a public speaking task?  
*Research on Child and Adolescent Psychopathology.*

## Abstract

Public speaking is one of the most commonly feared situations by socially anxious adolescents, often prompting behavioral anxiety markers including gaze avoidance and speech disruptions. While the potential adverse social consequences of behavioral anxiety markers in public speaking contexts have been established, research into how these markers might alter through cognitive behavior therapy is still in its infancy. In this preliminary study, we investigated changes in gaze behavior and speech disruptions from before to after 12 weeks of disorder-specific group cognitive behavior therapy among 41 adolescents aged 11-17 years ( $M = 14.46$ , 48.78% girls) with social anxiety disorder. Participants spoke for five minutes in front of a pre-recorded classroom audience while wearing an eye-tracker, before and after the *Skills for Academic and Social Success* program. Following treatment, we found an increase in frequency of gaze towards the faces of the audience while speaking, with greater changes among older participants. There were no changes in speech disruptions at the group level. We conclude that therapy may have a positive effect on gaze behavior, and discuss the clinical implications and opportunities for future research in this emerging field of study.

## Keywords

Social anxiety disorder; adolescents; public speaking; CBT; eye-tracking; speech disruptions

## Introduction

Social anxiety disorder (SAD) is the highly debilitating and persistent fear of being scrutinized by others, particularly during social interactions. This leads to suffering during and avoidance of fear-inducing social situations. The disorder typically has its onset in adolescence and, in youth, the fear is also present in the company of peers (American Psychiatric Association, 2022). SAD is thought to develop along a continuum, stemming from a combination of genetic and environmental influences, including parental and peer interactions and negative life events (Spence & Rapee, 2016). Cognitive distortions and safety behaviors resulting in poor social performance are involved in the maintenance of SAD (Hodson et al., 2008). Studies have shown that poor social skills can lead to detrimental social outcomes including disinterest in the short term, and lower acceptance and exclusion by peers in the long term (Miers & Masia Warner, 2023).

Yet, very little research has investigated the effect of treatment on behavioral markers of anxiety in a socially anxious population. Instead, treatment effectiveness is often regarded through the lens of remission rates or general symptom-reduction. While this remains the core purpose of treatment, insight into specific behavioral changes may further clarify the extent of remission and inform long-term prognoses. To fill this gap, the aim of this study was to investigate potential changes in socially anxious adolescents' behaviors during public speaking before and after cognitive behavior therapy (CBT). A public speaking situation was employed because it is one of the most commonly feared situations by socially anxious adolescents (Rao et al., 2007) and lends itself to the study of behavioral markers of anxiety. Moreover, public speaking tasks have been shown to elicit stress responses in clinical and non-clinical populations of adolescents with social anxiety (Asbrand et al., 2019; Inderbitzen-Nolan et al., 2007; Miers et al., 2011). Public speaking tasks have also been shown to trigger negative post-event processing in the days following the task (Asbrand et al., 2019), which according to theory can contribute to the maintenance of the disorder (Clark & Wells, 1995).

The Clark and Wells (1995) model of the development and maintenance of SAD in adults describes how self-focused attention causes socially anxious individuals to base their self-evaluations on negative assumptions about how others view them. This leads to negative expectations, causing them to use overt and covert safety behaviors. An overt safety behavior would be to avoid social situations in general (Hodson et al., 2008) whereas covert safety behaviors include avoidance of eye contact, decreasing the tone of the voice, speaking less, and standing on the outskirts of a group (Rapee & Heimberg, 1997; Wells et al., 1995). This theoretical model is applicable to socially anxious adolescents too (Hodson et al., 2008). Socially anxious adolescents may use

these behaviors with the implicit aim of reducing the chance of negative feedback, yet their very presence elicits negative feedback or exclusion from the group, resulting in a self-fulfilling prophecy (Leigh et al., 2021; Leigh & Clark, 2018). Thus, the disorder may in turn worsen and the situation deteriorates further. Reversing this negative spiral through treatment would be highly beneficial (Leigh et al., 2021).

Most studies on safety behaviors have used self-report measures (Piccirillo et al., 2016). Indeed, not all safety behaviors are observable (e.g., rehearsing what to say before giving a speech). However, reliance on the participants' awareness and relayed impression of their anxiety behaviors are limitations of self-report. Therefore, we concentrate on two of the most commonly studied observable behaviors in SAD during a behavior assessment task: eye-gaze avoidance and restricted speech. Although there is theoretical support for CBT improving behavioral anxiety markers, there is very little empirical evidence. Because of the scarcity of prior research, in particular with adolescent samples, we will discuss both adolescent and adult studies below.

## **Behavioral Anxiety Markers During Public Speaking**

### ***Eye Gaze***

Several public speaking studies using eye-tracking technology have provided evidence that adults with SAD look less at the faces of an audience, compared to non-anxious controls (Chen et al., 2015; Chen et al., 2016; Kim et al., 2018). Chen and colleagues (2021) administered a public speaking task, which also included a viewing task, to a community sample of young adults. Participants gave a one-minute speech in front of a pre-recorded audience and were subsequently instructed to simply look at the audience for the same duration of time. Participants with high levels of social anxiety showed avoidance of the audience's faces during public speaking but not while only viewing the audience. The authors concluded that avoidance is therefore not a default response, but rather occurs only when there is risk of negative judgement (see also Alden & Taylor 2004). However, the finding may not generalize to all SAD populations, as adults with SAD have been reported to look less at pictures of faces when there was no risk of being negatively evaluated (for review see Chen et al., 2020). These two findings together suggest that avoiding faces may generalize to non-evaluative situations over time and with increasing symptoms, thereby becoming avoiding faces may become a habit. Kitt and colleagues (2025) used the same speaking and viewing task with a sample of adolescents with and without anxiety disorders. In contrast to Chen and colleagues (2021), they did not find a difference in dwell time between the two groups during the speaking task. However, their population was not only younger, but also a mixed anxiety group including specific phobias; and gaze avoidance as a safety behavior may be particularly relevant to SAD.



### ***Disruptions while Speaking***

Speech duration, in the form of disruptions, appears to be different among socially anxious populations compared to non-socially anxious individuals during public speaking tasks. For example, in a study on speech disruptions in socially anxious adults versus healthy controls, Hofmann et al. (1997) found that the SAD group had significantly more speech disruptions with both filled (e.g., “uhm”, “er”) and silent pauses. They explained these results as socially anxious individuals shifting their attentional resources to other cognitive processes, including their own emotions and presence. Similarly, Levitan et al. (2012) found that adults with SAD were rated worse on intonation and speech fluency than control participants. In an adolescent study, speech disruptions were observed in nearly a third of adolescents with high levels of social anxiety, compared to 20% in the low anxiety group (Blöte et al., 2015).

In summary, evidence to date points to a link between SAD and both eye-gaze avoidance and speech disruptions. Previous research has shown that anxious behavior of socially anxious adolescents during a public speaking task can have negative academic and social consequences including poorer grades (Blöte et al., 2007) and more negative treatment by class peers, including not listening attentively while the person is speaking and ridiculing the speaker (Blöte & Westenberg, 2007). Negative social experiences may exacerbate symptoms of SAD and cumulate to increased impairment (Asbrand et al., 2019; Rapee & Spence, 2004); reducing anxious behavior such as eye gaze avoidance and frequent speech disruptions via CBT could therefore help reverse the negative spiral.

### **Changes in Anxiety Markers after Cognitive Behavior Therapy**

Despite CBT being one of the best-documented forms of psychotherapy to treat anxious youth (James et al., 2020), there are scarcely any studies on the potential effects of CBT on specific behavioral markers of SAD. Apart from a study demonstrating positive effects of CBT on global observer ratings of socially anxious adolescents' performance on behavioral assessment tasks (Herbert et al., 2009), we found only one prior study with an adolescent population, and three studies with adults on the effect of CBT on specific behaviors. These are summarized below.

Olivares-Olivares et al. (2019) studied the effect of treatment on socially anxious adolescents' eye-contact during an interview; duration of eye contact was obtained by studying video recordings of the interviews. They found that participants who received group CBT, the Spanish *Intervention in Adolescents with Generalized Social Phobia* program, with an added focus on social skills training, maintained a significantly longer duration of eye contact with the experimenter than participants who received the same

therapy without the social skills training. As such, CBT that incorporates social skills training could potentially improve eye gaze during a public speaking task.

To the best of our knowledge, no studies have investigated the effect of CBT on disruptions among socially anxious adolescents. Beidel et al. (2014) studied total speech duration in socially anxious adults. They conducted an impromptu speech task before and after Social Effectiveness Therapy (SET) – a 12-week CBT program combining exposure tasks with social skills training. They found SET led to improvements in total speech time, and in observer ratings of social skills. Two studies offering only the exposure ingredient of CBT compared the effects of 90 minutes of in vivo and virtual reality exposure therapy to a waitlist control condition. Anderson and colleagues (2013) asked participants to speak for ten minutes on up to three predetermined topics. Kampmann and colleagues (2016) asked their participants to perform a five-minute impromptu speech. Although the relative effectiveness of in vivo and virtual exposure varied between the studies, both showed a positive effect of exposure therapy on speech duration.

### ***Potential Moderators of Changes in Anxiety Markers after CBT***

Psychotherapy effectiveness research focusing on remission rates in youth with SAD has shown that higher initial clinician rated impairment scores are associated with less improvement at post-treatment (Compton et al., 2014; Wergeland et al., 2016). In the study by Kim et al. (2018) the time spent looking at a virtual audience was negatively related to state anxiety. Thus, it can be reasoned that eye gaze will increase when anxiety levels decrease. If individuals with higher initial anxiety levels experience a smaller improvement with treatment, behavioral outcomes such as eye-gaze and speech disruptions may be moderated in a similar way.

In the dynamic and continuously changing phase of adolescence it is relevant to investigate age as a potential moderator of treatment's influence on anxiety markers, along with gender. It has been suggested that safety behaviors may differ between younger and older youth (Leigh & Clark, 2018). In addition, public speaking behavior was related to both age and gender in a community sample of 9 to 16-year-olds: independent observer ratings of expressive and confident behaviors were positively related to age and higher in girls, whereas boys were rated higher on agitated behaviors, which included not looking at the audience (Blöte et al., 2015). Furthermore, an increase in eye-contact during adolescence has been found in peer conversation studies. A study comparing children of different ages to adults found that adult male dyads made 40% more eye-contact than 10-year-old boys, while adult female dyads made 50% more eye-contact than 10-year-old girls (Levine & Sutton-Smith, 1973). Likewise, Van Beek et al. (2006) showed that 16-year-old girls made more eye-contact than 13-year-old



girls or boys of either age. As relevant behaviors are still developing in adolescence, and given these findings, we include age and gender as exploratory moderators of a change in anxiety markers with treatment.

### The Present Study

In summary, to date there is a gap in empirical evidence investigating the potential influence of CBT on behavioral anxiety markers among socially anxious adolescents, despite the social benefits of improving these behaviors. We therefore aimed to study the extent to which CBT could alter socially anxious adolescents' eye-gaze behavior and speech disruptions during a public speaking task. In line with the finding that disorder-specific treatment programs have higher effect sizes than generic ones (Reynolds et al., 2012) we chose a disorder-specific group CBT protocol designed specifically for socially anxious adolescents: the *Skills for Academic and Social Success* (SASS; Masia Warner et al., 2018) program. This 12-week group intervention, developed in 2001 by Masia Warner and colleagues, is typically carried out in schools and, in addition to cognitive restructuring, focuses on practicing social skills and conducting social exposure tasks. Research has shown SASS to be an effective intervention, with medium to large treatment effects ( $d = 0.69$  in Masia Warner et al., 2016;  $d = 0.76$ ; Miller et al., 2011) and remission rates of 50-67% (Masia et al., 2001; Masia Warner et al., 2007; Masia Warner et al., 2005).

Before and after CBT, we administered a public speaking task with a subsequent viewing phase, as developed by Chen et al. (2021). Although other social interactions may be more frequent in the lives of adolescents, a public speaking context has the advantage that the audience does not have to make an active contribution to the interaction, so that the behavior of the socially anxious adolescent can be observed in a standardized situation. Adolescents are regularly asked to perform speeches at school in front of their class; our public speaking exercise was intended to capture and study their behavior in a naturalistic situation.

We hypothesized that at post-treatment, participants would display more eye-gaze towards audience members' faces (Olivares-Olivares et al., 2019), and longer speech duration (Anderson et al., 2013; Beidel et al., 2014; Kampmann et al., 2016) in the form of less disruption (Hofmann et al., 1997) compared to pre-treatment. Moreover, we explored whether changes after treatment were moderated by initial diagnosis severity, age and gender. In addition, and in line with previous eye-tracking studies (Chen et al., 2021; Kitt et al., 2025) we used a task with both a speaking and viewing phase in order to establish whether gaze behavior changed in both phases.

## Method

### Participants

Adolescents aged 11-17 years were recruited from public secondary schools in and surrounding a moderately large Dutch city for participation in a treatment efficacy study (Chapter 2). All included participants had SAD as their primary concern, and reached a Clinical Severity Rating (CSR) score of at least four in the child or parent clinical interview (ADIS-C/P; Silverman et al., 2001). We excluded adolescents with diagnosed or strongly suspected autism spectrum disorder, adolescents with suicidal inclinations or other difficulties that required immediate attention, and adolescents with behavioral difficulties that would disrupt the group treatment.

In total, 41 adolescents with a mean age of 14.46 ( $SD = 1.47$ ) were recruited. The sample consisted of 20 girls (48.78%), 20 boys (48.78%) and 1 genderfluid participant (2.44%). The study was approved by the university's Psychology Research Ethics Committee. All participants and their parents gave informed consent. The speaking task was completed by all participants except for one at T1 because the task was too demanding, and one at T2 because the participant was not available (speaking task  $N = 40$  at T1;  $N = 38$  at T2). In addition to the speaking task, all participants completed the viewing task, with the exception of the first therapy group because it was not yet introduced into the protocol (viewing task  $N = 33$  at T1;  $N = 31$  at T2). Two participants dropped out after the first therapy session.

### Design

These data were collected as part of a single-arm efficacy study, studying within-subject changes from before to after treatment. Data collection took place at four time points: intake (T0), pre-test (T1), post-test (T2) and follow-up (T3). In the present study, we only use the data collected at pre-test (T1) and post-test (T2) which are the only two time points when the participants performed the public speaking task. These two time points took place in the week before the start of treatment and the week after the completion of the 12-week treatment. Data were collected by university staff and graduate students. Therapists were involved in data collection at pre-test, but not at post-test.

### Instruments

#### *Eye-tracking Glasses*

To track eye gaze behavior during the public speaking exercise, wearable eye-tracking glasses, Tobii Pro Glasses 2 (Tobii AB, Sweden), were used. The experience is much like wearing reading or sunglasses, and is therefore not likely to evoke unrealistic distractions. The eye tracker is equipped with four eye cameras which track people's eye

movements in relation to the external environment they're watching (field of view 90° 16:9, visual angle 82° horizontally and 52° vertically, resolution 1920 × 1080 pixels). It records eye gaze at a sampling frequency of 100 Hz and a scene video at 25 Hz. The speech audio was recorded by a microphone embedded in the eye-tracking glasses, allowing us to measure disruptions. The eye-tracker was controlled by Tobii Glasses controller software installed on a tablet through a wireless connection. The software was used for recording and calibration.

### ***Clinical Severity Rating of Social Anxiety***

To establish the severity level of social anxiety disorder, the Anxiety Disorders Interview Schedule for children and parents (ADIS-C/P) was used (Silverman & Albano, 1996). This semi-structured clinical interview was developed to diagnose childhood anxiety disorders in accordance with DSM-IV (Kerns et al., 2013; Silverman et al., 2001), and has excellent reliability (Silverman et al., 2001). A CSR score is set on a scale of 0 to 8 for each disorder. Severity scores of 4 or higher indicate interference at clinical levels. In the present analysis, only the CSR scores of social anxiety disorder from the child interview at intake were used.

## **Procedure**

### ***Treatment***

The treatment consisted of 12 weekly group CBT sessions, *Skills for Academic and Social Success* (SASS; see Masia Warner et al., 2018 for the protocol). Developed specifically for socially anxious adolescents, the protocol features cognitive restructuring and social exposure exercises in addition to social skills training. The protocol covers varying social situations, starting small with skills like initiating a conversation with your desk neighbor at school, and progresses through to more difficult tasks like making phone calls and public speaking tasks, to doing embarrassing things in front of strangers. As part of the program, the adolescents practice public speaking skills; this has the primary focus during one session but also recurs throughout in the form of short public speaking exercises in the group. Moreover, behaviors such as eye-contact, speaking volume, speech fluency, posture, and more, were practiced throughout the protocol through various social skill exercises. We conducted SASS in a clinical setting, which allowed us to expand the typical 45-minute session duration to 120 minutes and go deeper into the protocol's themes. Participants also had access to a supplemental blended element (mHealth) with a CBT-focus throughout the treatment period.

### ***Public Speaking Task***

The public speaking task was based on the Leiden Public Speaking Task (LPST; Westenberg et al., 2009) using the same preparation time, total speech duration and pre-recorded audience. The LPST was modeled on a classroom presentation, which

Dutch students are regularly required to give in secondary school. It has previously been administered to 327 adolescents (Van den Bos et al., 2017). In the present study, participants performed the public speaking task in front of the pre-recorded audience while wearing eye-tracking glasses at pre- and post-test. The audience consisted of eight age peers and one teacher, seated in a classroom. The audience showed neutral affect because this is more realistic than angry faces yet still evokes a threat-like response due to the social nature of a face (Schmidtendorf et al., 2022; Yoon & Zinbarg, 2008). Two recordings of age peers were used: one audience for 12–14-year-old participants ( $N = 15$ ) and one for 15–17-year-old participants ( $N = 26$ ). The audience was projected life-size in front of the participant. Participants were aware that the audience was a video recording, and not a live audience via videocall. Virtual audiences have been shown to be similarly stress-evoking as real audiences (Kothgassner et al., 2016) and the video-recorded audiences and procedure used in this study have been shown to evoke stress responses in adolescents (Kitt et al., 2025; Van den Bos et al., 2017; Westenberg et al., 2009). While the therapy had a group format, all data collection including the public speaking task was conducted individually in a research facility with a projector and large screen onto which the audience was projected life-size. The participants were first informed of the public speaking task during the intake; thus, it was not impromptu. Participants were instructed not to wear eye-makeup in case this interferes with the validity of the gaze data; makeup remover was available in the laboratory in case participants had overlooked these instructions.

The public speaking task consisted of one minute of introducing themselves, followed by four minutes of a speech on a particular topic. When introduced to the task, the participants were given examples of what to talk about when introducing themselves for the first minute (e.g., “You can talk about who you live with, where you go to school, what you like to do in your spare time, and such.”). This instruction was the same at both time points. For the main part of the speech, the participants were assigned a general topic to talk about (e.g., “Tell us about your favorite movie and why you like it.”); the topic was different at post-test. Here they were given examples of what types of things they can talk about: e.g., themes, actors, specific scenes, etc. They were asked to keep talking for four minutes. The participants were then given five minutes to mentally prepare their speech. They were not allowed to use any written material during their speech.

After the five minutes of preparation time, the participant put on the eye-tracking glasses, and these were calibrated using Tobii’s default one-point calibration followed by a four-point calibration check. Once calibrated, they were asked to stand on a specified spot at a distance of about two meters from the projector screen, which showed an empty classroom. After a short moment, the image showed a class entering the

classroom and taking their seats. An auditory signal followed, at which the participant could start speaking. They introduced themselves and transitioned into their speech after they heard a second signal, a minute after the first signal. When a pause longer than 30 seconds fell, they were prompted via the intercom system to try to continue. This prompt was carried out a maximum of three times. Participants were videotaped during the speaking task, which they were aware of.

Following the speaking task, the participants removed the eye-tracking glasses and completed a questionnaire about their perception of the audience. Once this was finished, they put on the eye-tracking glasses once again, these were calibrated following the same procedure, and then they stood before the audience, now with a clipboard and pen in hand. They were then asked to first simply look at the audience members for one minute, and to then rate each audience member in turn as their number appeared above their head on screen, similar to the procedure used by Chen et al. (2021). Data on audience perception are not reported here. Participants were debriefed after each public speaking measurement. Participants did not receive compensation.

## **Data Preparation**

### ***Gaze Behavior***

The eye-tracking data were processed using Tobii Pro Lab software. Pictures of the video being displayed on the screen were used as reference images, drawing areas of interest (AOIs) on the faces of the 9 audience members. In this study, the specific AOI lied within the outline of each audience member's face, excluding the person's hair. We used the Tobii I-VT (Velocity-Threshold Identification) Attention gaze filter, which has been designed for the use of eye-tracking glasses in dynamic situations. The attention filter identifies fixations using a velocity threshold of 100°/s and a minimum fixation duration of 60 milliseconds (ms). Adjacent fixations are merged when the time between fixations is no more than 75 ms and the distance between fixations is no more than 0.5°, based on the average data from both eyes (Olsen, 2012). The fixations that had been registered relative to the scene video were automatically mapped onto the reference images for the five minutes between the start and end signal of the speaking task and the one minute between the start and end signal of the viewing task. Subsequently, these mappings were checked by a human observer. The software calculated fixation-based parameters: the number of fixations on each AOI and the duration of fixations on each AOI in milliseconds. Fixation counts and fixation time were cumulated across AOIs and divided by duration of the task in minutes to obtain comparable data for the 5-minute speaking task and the 1-minute looking task.

The eye-tracker also coded each gaze sample as valid or invalid for each eye. Behaviors that would lead to the eye-tracker temporarily losing contact with the pupil include the

participant blinking, closing their eyes, or moving their eyes in an extreme direction (e.g., keeping their head still but moving their gaze to their feet; turning their eyes up to the ceiling in thought). Raw data were exported from Tobii Pro Lab and imported in Matlab. A Matlab script was used to select the gaze samples pertaining to the 5-minute speaking task and the 1-minute looking task and compute the percentage of valid gaze samples (adding gaze samples for the left and right eye). The mean percentages of valid gaze samples were 84 ( $SD = 13$ ) at T1 and 85 ( $SD = 12$ ) at T2 for the speaking task and 90 ( $SD = 13$ ) at T1 and 92 ( $SD = 8$ ) at T2 for the viewing task. Seven data segments had less than 70% valid gaze samples. Since excluding these from the analyses did not affect the pattern of the results, the results reported are based on all available data.

### ***Disruptions while Speaking***

Although short silences and filled pauses commonly occur in spoken language, silences of more than two seconds have been shown to be rare in speech samples from a variety of contexts, including public speaking (Grosman et al., 2018). In the present study, participants were sometimes silent for a long time (Max = 54 s), suggesting a breakdown of speech performance. Likewise, some participants addressed the experimenter with a question or comment about the task (e.g., “How much time is left?”, “I don’t know what to say anymore”). Therefore, silences and filled pauses (i.e., silences interspersed with utterances that don’t contribute to the meaning of the sentence such as, “uhm”, “right”) of more than two seconds and interactions with the experimenter were manually coded as disruptions in Tobii Pro Lab, by marking their start and end times in the recording. Number and duration of disruptions were totaled and divided by the total duration of the speech to obtain the number of disruptions per minute and the duration of disruptions in milliseconds per minute.

### ***Statistical Analysis***

A power calculation of the required sample size to detect change in behavioral anxiety markers between pre- and post-test was computed prior to data collection. We based our calculations on the means and standard deviations reported by Beidel et al. (2000), who measured social skills in a behavioral assessment before and after Social Effectiveness Therapy for Children (SET-C), a treatment that was an inspiration for the SASS protocol (Fisher et al., 2004). With a desired power of .95 and an alpha level of .05, the computation showed that 47 participants would be needed.

We analyzed the differences in gaze behavior and speech disruptions between pre- and post-test, and moderation by age, gender, and initial CSR score. Gaze behavior and speech disruptions were both measured in number and in duration. Linear mixed omnibus models were run to prevent inflation of the alpha level by multiple testing. The omnibus models included time point, age, gender, initial CSR score, task (for gaze



behavior only) and interactions of time point with each of the other variables as predictors. Significant omnibus analyses were followed up by separate analyses per dependent variable including time point and those moderators that were significant in the omnibus model. Models including gender as a predictor were tested using data from male and female participants only. Models that did not include gender were tested using data from all participants. Effect sizes were interpreted using Cohen's (1988) *d* standards (small = 0.2, medium = 0.5, and large = 0.8). Analyses were conducted in RStudio version 4.3.1 using base packages (R Core Team, 2013), *effsize* package (version 0.8.1; Torchiano, 2020), the *lme4* package (version 1.1-35.3; Bates et al., 2015), the *dplyr* package (version 1.1.3; Wickham et al., 2023) and the *ggplot2* package (version 4.3.3; Wickham, 2016).

## Results

See Table 1 for the means and standard deviations of the total sample and for girls and boys separated, and Table 2 for an overview of the main results. The dependent variables included in the analyses were square root transformed to better approximate the normal distribution.

**Table 1**  
*Means and standard deviations*

	Total sample		Girls		Boys	
	T1	T2	T1	T2	T1	T2
	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)
<b>Eye-gaze</b>						
<i>Number of fixations</i>						
Speaking	9.07 (6.53)	11.8 (9.85)	9.31 (5.91)	10.3 (8.30)	8.85 (7.22)	13.1 (11.1)
Viewing	18.5 (10.5)	19.0 (15.4)	15.1 (11.2)	17.0 (17.8)	21.4 (9.31)	19.8 (13.8)
<i>Duration of fixations</i>						
Speaking	3.91 (2.89)	4.85 (4.15)	3.32 (2.24)	3.56 (2.73)	4.48 (3.35)	5.95 (4.85)
Viewing	11.89 (7.21)	12.39 (9.83)	8.35 (6.75)	10.08 (11.17)	15.21 (6.10)	14.16 (8.60)
<b>Disruptions while speaking</b>						
<i>Number of disruptions</i>						
	1.26 (0.75)	1.19 (0.77)	1.06 (0.59)	0.99 (0.63)	1.35 (0.79)	1.27 (0.77)
<i>Duration of disruptions</i>						
	10.37 (7.34)	10.37 (8.22)	8.59 (6.45)	10.29 (9.16)	11.81 (8.04)	10.19 (7.71)

*Note.* Reported data are untransformed; duration reported in seconds.

## Eye Gaze

The eye-tracking variables were highly correlated ( $r = .88, p < .001$ ). Omnibus testing of both eye-gaze variables revealed a main effect of time ( $t(70.58) = -2.68, p = .009$ ), qualified by an interaction with age ( $t(70.02) = 2.58, p = .012$ ). Moderation by gender and initial CSR score were non-significant (respectively  $t(75.89) = 0.33, ns$ ;  $t(82.97) = 0.12, ns$ ). However, there was a main effect of gender, with boys looking more at the audience member's faces than girls ( $t(46.99) = -2.10, p = .041$ ). Additionally, there was a main effect of task, with participants looking significantly more at the audience members' faces during the viewing task than the speaking task ( $t(68.60) = 5.13, p < .001$ ). Follow up analyses per dependent variable were run to clarify the interaction between time and age (Table 2).

**Table 2**

*Effect of treatment from pre- to post-test*

Interaction	Estimate	S.E.	Df	t-value	p
<b>Eye-Gaze</b>					
<i>Number of fixations</i>					
Time point	-4.85	1.79	97.66	-2.71	0.008 **
Age	-0.10	0.12	62.72	-0.81	0.42
Time point: Age	0.36	0.12	97.45	2.96	0.004 **
Task viewing	1.29	0.26	99.38	4.90	<0.001 ***
Time point: Task viewing	-0.44	0.37	95.95	-1.20	0.233
<i>Duration of fixations</i>					
Time point	-106.94	46.91	98.58	-2.28	0.025 *
Age	-3.12	2.97	68.30	-1.05	0.297
Time point: Age	7.83	3.21	98.36	2.44	0.016 *
Task viewing	43.80	6.88	100.50	6.37	<0.001 ***
Time point: Task viewing	-4.87	9.72	96.69	-0.50	0.618

### ***Number of Fixations on Faces***

The number of fixations on audience members' faces showed a significant main effect of time with a large effect size ( $t(97.66) = -2.71, p = .008$ ; *Cohen's d* = -2.87, 95% CI [-3.18, -2.56]), qualified by a significant interaction with age with a small effect size ( $t(97.45) = 2.96, p = .004$ ; *d* = 0.22, 95% CI [-0.00, 0.43]). Older participants showed a larger increase in the number of fixations on faces, see Figure 1. The main effect of task was significant with a large effect size ( $t(99.38) = 4.90, p < .001$ ; *d* = -0.75, 95% CI [-0.97, -0.54]). While the interaction between task and time was not significant ( $t(95.95) = -1.20, ns$ ), its effect size was medium (*d* = -0.44, 95% CI [-0.66, -0.22]),

warranting follow-up analyses of the interaction between task and time. Separate analyses by task showed that there was a significant increase in number of fixations between pre-test and post-test during the speaking task ( $t(37.71) = 2.10, p = .042$ ) with a small effect size,  $d = 0.31$ , 95% CI [-0.14, 0.77]; there was no significant interaction during the viewing task ( $t(30.76) = -0.53, ns$ ).


### ***Duration of Fixations on Faces***

Similar to the number of fixations, duration of fixations on audience members' faces showed a significant main effect of time with a large effect size ( $t(98.58) = -2.28, p = .025; d = -2.70$ , 95% CI [-3.00, -2.40]), qualified by a significant interaction with age ( $t(98.36) = 2.44, p = .016; d = 0.20$ , 95% CI [-0.02, 0.41]). Older participants showed a larger increase in the duration of fixations on faces, see Figure 2. The main effect of task was significant with a large effect size ( $t(100.50) = 6.37, p < .001; d = -1.21$ , 95% CI [-1.55, -0.88]); however, the interaction between time and task was not significant and had a small effect size ( $t(96.69) = -0.50, ns; d = -0.12$ , 95% CI [-0.34, 0.09]).

### ***Disruptions while Speaking***


The variables measuring the number and duration of disruptions were highly correlated ( $r = .79, p = .001$ ). Contrary to expectation, omnibus testing of both disruption variables did not show an effect of time point ( $t(19.37) = 0.50, ns$ ). Interactions between time and all moderating variables were non-significant: age ( $t(19.21) = 1.39, ns$ ), gender ( $t(20.06) = 1.09, ns$ ), and initial CSR score ( $t(20.89) = 1.68, ns$ ).

**Figure 1**  
*Age versus change score of number of fixations on audience members' faces, showing greater change among older adolescents*



**Figure 2**

*Age versus change score of duration of fixation on audience members' faces, showing greater change among older adolescents*



## Discussion

The present study investigated the extent to which disorder-specific group CBT for socially anxious adolescents improved behavioral markers of anxiety, specifically eye-gaze and speech disruptions, during public speaking. We also tested whether changes were moderated by age, gender, and initial severity of social anxiety. Participants completed a public speaking and viewing task while wearing an eye-tracker, before and after 12 weeks of CBT. We found that subsequent to treatment, participants displayed a greater degree of eye-gaze towards the audience's faces, both in number and in duration of fixations. The change was greater in older participants. We did not observe a difference in speech disruptions after CBT. Due to the small sample size of the present study, findings should be viewed as preliminary.

The significant increase in eye-gaze at post-test could indicate an influence of treatment. These results are in line with the only other adolescent study on the effects of CBT on these specific anxiety markers; Olivares-Olivares et al. (2019) found that CBT can have an effect on eye-contact during an interview, and that social skills training seems to be a particular driver of this effect. Age moderated the change of gaze behavior; indicating that gaze behavior improved more in older participants. Previous developmental studies demonstrated that older adolescents showed more expressive and confident behavior during public speaking than younger ones (Blöte et al., 2015) and that 16-year-old girls made more eye-contact during a conversation than 13-year-olds (Van Beek et al., 2006). This suggests that older adolescents may be developmentally more responsive to fine-tuning these social behaviors. Although boys in this study looked more at the faces of the audience members than girls (averaged over speaking and viewing tasks), gender did not moderate the change in gaze behavior (or any other outcome).

In the present study, an increase in number of fixations following CBT was observed in the speaking task but not during the viewing task. A change occurring in the speaking task only could support the notion that socially anxious individuals avoid eye-contact in socially evaluative situations, such as a speech, but not during non-threatening situations such as simply looking at an audience (Chen et al., 2021). However, given that the interaction between time and task was not significant, we cannot rule out that gaze behavior changed in the viewing task as well. Further research with a larger sample size is needed to clarify whether adolescents with SAD, like adults (Chen et al., 2020), avoid looking at faces in non-evaluative situations.

In line with two other studies comparing speaking and viewing (Chen et al., 2021; Kitt et al., 2025), participants looked more frequently and for longer at faces during the viewing task than during the speaking task. The mean fixation duration during



the viewing task (see Table 1) was also comparable to the results of Chen et al. (2021), obtained with a community sample of young adult females (9.3 - 10.5s), and the results of Kitt et al. (2025), obtained with an adolescent sample consisting of participants with an anxiety disorder and control participants (12.97s). However, the mean fixation duration during the speaking task was lower in the present study (Chen et al., 2021: 6.5 - 10.1s; Kitt et al., 2025: 6.08s). This might be related to our participants being exclusively socially anxious adolescents as opposed to an adult community sample or a mixed anxiety population. However, there was no significant relation with social anxiety symptoms in Kitt et al.'s study (2025). Rather, the difference may be related to the duration of the speaking task: four minutes in the present study versus one minute in the other two studies. Our more demanding task poses a higher risk of running out of things to say. During disruptions, participants tended to look down or turn towards the control room when addressing the experimenter. Hence, lower mean numbers and durations of fixations on the audience may be representative of a longer speech task.

While different behavioral factors involved in giving a speech were practiced during the treatment, this practice did not seem to have a substantial effect on speech disruptions. This was contrary to our expectations and previous treatment studies among adult samples (Anderson et al., 2013; Beidel et al., 2014; Kampmann et al., 2016). In these studies, participants were allowed to terminate their speech after a minimum duration (Beidel et al., 2014: 3 out of 10 min) or when they wanted to stop (Kampmann et al., 2016). In the present study, participants were encouraged to continue if they did not speak for the full five minutes and this may have obscured the effect. Moreover, there are other aspects involved in speech which may have been affected (e.g., volume, speed, intonation, frequency of pausing for less than two seconds) but these were not possible to address within the scope of this study.

Behavioral markers such as speech disruptions are likely caused by individuals being preoccupied: shifting their attention from the speech at hand to other cognitive processes, including their own emotions and presence (Hofmann et al., 1997). In the present study we did not find a change in speech disruptions following CBT. However, due to a lack of comparison to healthy controls, we cannot draw conclusions as to how typical the speech disruptions were at either time point. Moreover, generalization of cognitive restructuring and social skills learned during CBT to actual social situations could take more time than included in our study. The scope of this study covered the time points immediately before and after 12 weeks of CBT. However, as consistently observed in previous research using *SASS*, additional improvements are often made in the period following treatment (Masia Warner et al., 2007). Thus, the inclusion of follow-up measurements using a public speaking task should be considered, as additional benefits in behavior may evolve with more time.

There were no relations between age or gender and speech disruptions. Effects of age and gender need to be studied in greater detail as there may be nuanced differences in safety behaviors (Leigh & Clark, 2018), not captured here. Initial anxiety level did not moderate changes in gaze behavior or speech disruptions. Although in contrast with some studies (e.g., Compton et al., 2014; Wergeland et al., 2016), this result was in line with research by Butler and colleagues (2021) with socially anxious adults, finding that initial severity of diagnosis was not a predictor in symptom change across treatment; thereby indicating that treatment can have an equal effect across severity levels.

### **Clinical Implications**

Specific aspects of social skills, including public speaking, can have an influence on adolescents' school and peer functioning (Blöte et al., 2007; Blöte & Westenberg, 2007). The finding of a significant improvement in gaze behavior during public speaking after CBT has clinical relevance, as increased eye gaze can potentially reverse the negative spiral by reducing negative responses from others, improving social interactions and consequently leading to a reduction in impairment. The inclusion of public speaking training in treatment for socially anxious adolescents is a significant driver of symptom reduction (García-López et al., 2002). While public speaking skills were a recurring theme within our chosen treatment program, there are numerous other themes and skills covered throughout the twelve weeks of SASS. Thus, to see greater change in speech duration, more specific public speaking training within CBT, may be needed. It has been suggested that, generally, more emphasis may need to be placed on positive peer feedback in the group setting (Miers & Masia Warner, 2023). Blöte et al. (2019) suggested socially anxious adolescents who were poor performers at public speaking would benefit from specific social skill training paired with cognitive restructuring, whereas socially anxious adolescents with high quality performances may benefit more from video-feedback interventions. Combining the two suggestions, video feedback paired with structured positive peer response, may be highly beneficial. Future research could specifically test this hypothesis.

### **Limitations and Directions for Future Research**

This study extends current literature by studying changes in eye-gaze over the course of CBT using eye-tracking technology, thus combining methodological strengths from previous research that has either studied the effect of CBT (Olivares-Olivares et al., 2019) or used precise eye-tracking technology but only at one time point (Kitt et al., 2025). Moreover, the use of the material and procedure from the Leiden Public Speaking Task (Westenberg et al., 2009) enabled the comparison of results across other studies that have used the same format (e.g., Chen et al., 2021; Kitt et al., 2025).

While this study provides novel insights, it also has a number of limitations which could be addressed in future research. In addition to replication with a larger, more varied sample to increase the power to detect small effect sizes and to increase generalizability, a primary recommendation for future research is to include a control condition. This could be done in two ways: first, inclusion of a non-anxious control group to delineate typical public speaking behavior from what is unique to socially anxious adolescents (see Kitt et al., 2025). While we observed an improvement in number of fixations in the speaking task, we did not observe an effect in the viewing task. Previous studies have suggested that socially anxious individuals show similar gaze patterns to non-anxious individuals during tasks where they are not judged (Chen et al., 2021); however, as our study did not include a non-anxious control group, we cannot be certain to what extent participants' gaze behavior was on par with age peers. Second, use of an active control condition, employed within a randomized design and with a larger sample size, would be necessary to draw stronger conclusions about the observed effects of CBT rather than potential habituation to the public speaking task.

Moreover, this study utilized a video of an audience and not a live audience. This has the advantage of allowing for standardization but the disadvantage of lower ecological validity. A strength is that our audience was comprised of 'real' people rather than computer-animated people. However, it was nonetheless a video and the participants were aware of this; for example, there was no deception involved to suggest it was a videocall. Evidence shows that public speaking tasks elicit a stress response in adolescents (Miers et al., 2011; Westenberg et al., 2009) and it was observed to be demanding for many of our participants. It is not clear to what extent our participants would have behaved differently in front of a live audience. Future research should investigate the effects of treatment on behavioral markers in front of a live audience or during peer interaction. Such approaches would reduce standardization but may improve validity and generalizability of results.

There are numerous potential behavioral markers of SAD which may be present during a public speaking task, but which were not captured in this research (e.g., posture, facial expressions, volume). Future studies should code for other improvements in speech, both specific and global, following treatment using a coding scheme such as the Speech Performance Observation Scale for Youth (Blöte et al., 2015). Moreover, pubertal developmental level has been shown to be important in relation to (neuroendocrine) stress responses in this population (van den Bos et al., 2017); future studies should include this factor in addition to age. Furthermore, future studies may test the effect of CBT on more social skills beyond public speaking.

## **Conclusion**

The current study explored, for the first time, the potential effect of CBT for socially anxious adolescents on specific behavioral markers of anxiety during a public speaking task. Our findings reflect the notion that socially anxious individuals avoid eye-gaze in socially evaluative situations. Moreover, our results support that gaze behavior may be altered through general CBT treatment. The study also highlights the importance of including moderators such as age and possibly gender in adolescent treatment studies. Due to the preliminary nature of this study, results should be interpreted with caution. Whilst replication with a larger sample is required, the behavioral improvement in eye gaze demonstrates the potential for bringing about more rewarding social situations – particularly with same-age peers – thereby yielding social gains for this socially vulnerable adolescent population. Future research should include randomized control groups for a greater understanding of adolescents' typical behavior during public speaking tasks and to investigate the effect of treatment-related behavioral changes.

## References

- Alden, L. E., & Taylor, C. T. (2004). Interpersonal processes in social phobia. *Clinical Psychology Review*, 24(7), 857–882. <https://doi.org/10.1016/j.cpr.2004.07.006>
- American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders* (5th ed., text rev.). <https://doi.org/10.1176/appi.books.9780890425787>
- Anderson, P. L., Price, M., Edwards, S. M., Obasaju, M. A., Schmertz, S. K., Zimand, E., & Calamaras, M. R. (2013). Virtual reality exposure therapy for social anxiety disorder: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 81(5), 751.
- Asbrand, J., Schmitz, J., Krämer, M., Nitschke, K., Heinrichs, N., & Tuschen-Caffier, B. (2019). Effects of group-based CBT on post-event processing in children with Social Anxiety Disorder following an experimental social stressor. *Journal of Abnormal Child Psychology*, 47(12), 1945–1956. <https://doi.org/10.1007/s10802-019-00558-x>
- Bates D., Mächler, M., Bolker, B., Walker, S. (2015). Fitting linear mixed-effects models using lme4.” *Journal of Statistical Software*, 67(1), 1–48. doi:10.18637/jss.v067.i01.
- Beidel, D. C., Alfano, C. A., Kofler, M. J., Rao, P. A., Scharfstein, L., & Wong Sarver, N. (2014). The impact of social skills training for social anxiety disorder: A randomized controlled trial. *Journal of Anxiety Disorders*, 28(8), 908–918. <https://doi.org/10.1016/j.janxdis.2014.09.016>
- Beidel, D. C., Turner, S. M., & Morris, T. L. (2000). Behavioral treatment of childhood social phobia. *Journal of Consulting and Clinical Psychology*, 68(6), 1072–1080.
- Blöte, A. W., Miers, A. C., Heyne, D. A., Clark, D. M. & Westenberg, P. M. (2014), The relation between social anxiety and audience perception: Examining Clark and Wells' (1995) model among adolescents. *Behavioural and Cognitive Psychotherapy*, 42(5): 555–567.
- Blöte, A. W., Miers, A. C., Van den Bos, E., & Westenberg, P. M. (2019). The role of performance quality in adolescents' self-evaluation and rumination after a speech: Is it contingent on social anxiety level? *Behavioural and Cognitive Psychotherapy*, 47, 148–163. <https://doi.org/10.1017/S1352465818000310>
- Blöte, A. W., Pongjitt, A., Miers, A. C., Van Beek, Y., & Westenberg, P. M. (2015). The Speech Performance Observation Scale for Youth (SPOSY): Assessing social performance characteristics related to social anxiety. *Journal of Experimental Psychopathology*, 6(2), 168–184.
- Blöte, A. W., Kint, M. J., & Westenberg, P. M. (2007). Peer behavior toward socially anxious adolescents: Classroom observations. *Behaviour Research and Therapy*, 45(11), 2773–2779.
- Blöte, A. W., & Westenberg, P. M. (2007). Socially anxious adolescents' perception of treatment by classmates. *Behaviour Research and Therapy*, 45(2), 189–198.
- Butler, R. M., O'Day, E. B., Swee, M. B., Horenstein, A., & Heimberg, R. G. (2021). Cognitive Behavioral Therapy for social anxiety disorder: Predictors of treatment outcome in a quasi-naturalistic setting. *Behavior Therapy*, 52(2), 465–477. <https://doi.org/10.1016/j.beth.2020.06.002>
- Chen, J., van den Bos, E., & Westenberg, P. M. (2020). A systematic review of visual avoidance of faces in socially anxious individuals: Influence of severity, type of social situation, and development. *Journal of Anxiety Disorders*, 70, 102193–102193. <https://doi.org/10.1016/j.janxdis.2020.102193>
- Chen, J., van den Bos, E., Velthuisen, S. L. M., & Westenberg, P. M. (2021). Visual avoidance of faces in socially anxious individuals: The moderating effect of type of social situation. *Journal of Experimental Psychopathology*, 12(1), 2043808721989628.
- Chen, N. T. M., Clarke, P. J., MacLeod, C., Hickie, I. B., & Guastella, A. J. (2016). Aberrant gaze patterns in social anxiety disorder: An eye movement assessment during public speaking. *Journal of Experimental Psychopathology*, 7(1), 1–17.

- Chen, N. T. M., Thomas, L. M., Clarke, P. J. F., Hickie, I. B., & Guastella, A. J. (2015). Hyperscanning and avoidance in social anxiety disorder: The visual scanpath during public speaking. *Psychiatry Research*, 225(3), 667-672.
- Clark, D. M., & Wells, A. (1995). A cognitive model of social phobia. In R. G. Heimberg, M. Liebowitz, D. Hope, & F. Schneier (Eds.), *Social phobia: Diagnosis, assessment, and treatment* (pp. 69-93). Guilford Press.
- Compton, S. N., Peris, T. S., Almirall, D., Birmaher, B., Sherrill, J., Kendall, P. C., ... & Albano, A. M. (2014). Predictors and moderators of treatment response in childhood anxiety disorders: Results from the CAMS trial. *Journal of Consulting and Clinical Psychology*, 82(2), 212.
- García-López, L. J., Olivares, J., Turner, S. M., Beidel, D. C., Albano, A. M., & Sánchez-Meca, J. (2002). Results at long-term among three psychological treatments for adolescents with generalized social phobia (II): Clinical significance and effect size. *Psicología Conductual*, 10(2), 371-388.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Lawrence Erlbaum Associates.
- Evans, R., Chiu, K., Clark, D. M., Waite, P., & Leigh, E. (2021). Safety behaviours in social anxiety: An examination across adolescence. *Behaviour Research and Therapy*, 144, 103931.
- Grosman, I., Simon, A. C., & Degand, L. (2018). Variation de la durée des pauses silencieuses: Impact de la syntaxe, du style de parole et des disfluences. *Language*, 211(3), 13-40. DOI: 10.3917/lang.211.0013
- Herbert, J. D., Gaudiano, B. A., Rheingold, A. A., Moitra, E., Myers, V. H., Dalrymple, K. L., & Brandt, L. L. (2009). Cognitive behavior therapy for generalized social anxiety disorder in adolescents: A randomized controlled trial. *Journal of Anxiety Disorders*, 23(2), 167-177.
- Hodson, K. J., McManus, F. V., Clark, D. M., & Doll, H. (2008). Can Clark and Wells' (1995) cognitive model of social phobia be applied to young people? *Behavioural and Cognitive Psychotherapy*, 36(4), 449-461. <https://doi.org/10.1017/S1352465808004487>
- Hofmann, S. G., Gerlach, A. L., Wender, A., & Roth, W. T. (1997). Speech disturbances and gaze behavior during public speaking in subtypes of social phobia. *Journal of Anxiety Disorders*, 11(6), 573-585.
- Inderbitzen-Nolan, H. M., Anderson, E. R., & Johnson, H. S. (2007). Subjective versus objective behavioral ratings following two analogue tasks: A comparison of socially phobic and non-anxious adolescents. *Journal of Anxiety Disorders*, 21(1), 76-90. <https://doi.org/10.1016/j.janxdis.2006.03.013>
- James, A. C., Reardon, T., Soler, A., James, G., & Creswell, C. (2020). Cognitive behavioural therapy for anxiety disorders in children and adolescents. *Cochrane Database of Systematic Reviews*, (11).
- Kampmann, I. L., Emmelkamp, P. M. G., Hartanto, D., Brinkman, W.-P., Zijlstra, B. J. H., & Morina, N. (2016). Exposure to virtual social interactions in the treatment of social anxiety disorder: A randomized controlled trial. *Behaviour Research and Therapy*, 77, 147-156. <https://doi.org/10.1016/j.brat.2015.12.016>
- Kerns, C., Comer, J., Pincus, D., & Hofmann, S. (2013). Evaluation of the proposed social anxiety disorder specifier change for DSM-5 in a treatment-seeking sample of anxious youth. *Depression and Anxiety*, 30(8), 709-715.
- Kim, H., Shin, J. E., Hong, Y. J., Shin, Y. B., Shin, Y. S., Han, K., ... & Choi, S. H. (2018). Aversive eye gaze during a speech in virtual environment in patients with social anxiety disorder. *Australian & New Zealand Journal of Psychiatry*, 52(3), 279-285.
- Kitt, E. R., Abend, R., Amelio, P., Galbraith, J., Poe, A. D., Gee, D. G., Pine, D. S., & Harrewijn, A. (2025). Using mobile eye-tracking to evaluate gaze behavior during a speech in pediatric anxiety disorders. *Journal of Affective Disorders*, 369, 462-466. <https://doi.org/10.1016/j.jad.2024.10.024>



- Kothgassner, O. D., Felnhofer, A., Hlavacs, H., Beutl, L., Palme, R., Kryspin-Exner, I., & Glenk, L. M. (2016). Salivary cortisol and cardiovascular reactivity to a public speaking task in a virtual and real-life environment. *Computers in Human Behavior*, 62, 124–135. <https://doi.org/10.1016/j.chb.2016.03.081>
- Leigh, E., Chiu, K., & Clark, D. M. (2021). Self-focused attention and safety behaviours maintain social anxiety in adolescents: An experimental study. *PLoS One*, 16(2), e0247703.
- Leigh, E., & Clark, D. M. (2018). Understanding social anxiety disorder in adolescents and improving treatment outcomes: Applying the cognitive model of Clark and Wells (1995). *Clinical Child and Family Psychology Review*, 21(3), 388–414. <https://doi.org/10.1007/s10567-018-0258-5>
- Levitán, M. N., Falcone, E. M., Placido, M., Krieger, S., Pinheiro, L., Crippa, J. A., ... & Nardi, A. E. (2012). Public speaking in social phobia: A pilot study of self-ratings and observers' ratings of social skills. *Journal of Clinical Psychology*, 68(4), 397–402.
- Masia, C., Klein, R. G., Storch, E. A., & Corda, B. (2001). School-based behavioral treatment for social anxiety disorder in adolescents: Results of a pilot study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40, 780–786.
- Masia Warner, C., Colognori, D., Brice, C., Herzig, K., Mufson, L., Lynch, C., Reiss, P. T., Petkova, E., Fox, J., Mocerri, D. C., Ryan, J., & Klein, R. G. (2016). Can school counselors deliver cognitive-behavioral treatment for social anxiety effectively? A randomized controlled trial. *Journal of Child Psychology and Psychiatry*, 57(11), 1229–1238. <https://doi.org/10.1111/jcpp.12550>
- Masia Warner, C., Colognori, D., & Lynch, C. (2018). *Helping students overcome social anxiety: Skills for Academic and Social Success (SASS)*. Guilford Publications.
- Masia Warner, C., Fisher, P. H., Shrout, P. E., Rathor, S., & Klein, R. G. (2007). Treating adolescents with social anxiety disorder in school: An attention control trial. *Journal of Child Psychology and Psychiatry*, 48(7), 676–686. <https://doi.org/10.1111/j.1469-7610.2007.01737.x>
- Masia Warner, C., Klein, R., Dent, H., Fisher, P., Alvir, J., Malbano, A., & Guardino, M. (2005). School-based intervention for adolescents with social anxiety disorder: Results of a controlled study. *Journal of Abnormal Child Psychology*, 33(6), 707–722. <https://doi.org/10.1007/s10802-005-7649-z>
- Miers, A. C., & Masia Warner, C. (2023). Adolescent social anxiety: Bridging cognitive and social behavioral perspectives through a developmental lens. In C. R. Martin, V. B. Patel & V. R. Preedy (Eds.), *Handbook of lifespan Cognitive Behavioral Therapy: Childhood, adolescence, pregnancy, adulthood and aging*, 89–100. Academic Press.
- Miller, L. D., Gold, S., Laye-Gindhu, A., Martinez, Y. J., Yu, C. M., & Waechtler, V. (2011). Transporting a school-based intervention for social anxiety in Canadian adolescents. *Canadian Journal of Behavioural Science*, 43(4), 287–296. <https://doi.org/10.1037/a0023174>
- Olivares-Olivares, P. J., Ortiz-González, P. F., & Olivares, J. (2019). Role of social skills training in adolescents with social anxiety disorder. *International Journal of Clinical and Health Psychology*, 19(1), 41–48. <https://doi.org/10.1016/j.ijchp.2018.11.002>
- Olsen, A. (2012). Tobii I-VT fixation filter - Algorithm description. *White Paper* (Retrieved from <http://www.tobii.com/eye-tracking-research/global/library/white-papers/the-tobii-i-vt-fixation-filter/>).
- Piccirillo, M. L., Dryman, M. T., & Heimberg, R. G. (2016). Safety behaviors in adults with social anxiety: Review and future directions. *Behavior Therapy*, 47(5), 675–687.
- R Core Team (2013). R: A language and environment for statistical computing. R foundation for statistical computing. <https://www.R-project.org>.
- Rao, P. A., Beidel, D. C., Turner, S. M., Ammerman, R. T., Crosby, L. E., & Sallee, F. R. (2007). Social anxiety disorder in childhood and adolescence: Descriptive psychopathology. *Behaviour Research and Therapy*, 45(6), 1181–1191.

- Rapee, R. M., & Heimberg, R. G. (1997). A cognitive-behavioral model of anxiety in social phobia. *Behaviour Research and Therapy*, 35(8), 741–756. [https://doi.org/10.1016/S0005-7967\(97\)00022-3](https://doi.org/10.1016/S0005-7967(97)00022-3)
- Rapee, R. M., & Spence, S. H. (2004). The etiology of social phobia: Empirical evidence and an initial model. *Clinical Psychology Review*, 24(7), 737–767. <https://doi.org/10.1016/j.cpr.2004.06.004>
- Reynolds, S., Wilson, C., Austin, J., & Hooper, L. (2012). Effects of psychotherapy for anxiety in children and adolescents: A meta-analytic review. *Clinical Psychology Review*, 32(4), 251–262. <https://doi.org/10.1016/j.cpr.2012.01.005>
- Schmidtendorf, S., Herwig, A., Wiedau, S., Asbrand, J., Tuschen-Caffier, B., & Heinrichs, N. (2022). Initial maintenance of attention to threat in children with social anxiety disorder? Findings from an eye-tracking experiment. *Cognitive Therapy and Research*, 46(1), 197–208. <https://doi.org/10.1007/s10608-021-10244-2>
- Silverman, W.K. & Albano, A.M. (1996). *Anxiety Disorders Interview Schedule for DSM- IV: Child and Parent versions*. The Psychological Corporation.
- Silverman, W. K., Saavedra, L. M., & Pina, A. A. (2001). Test-retest reliability of anxiety symptoms and diagnoses with the Anxiety Disorders Interview Schedule for DSM-IV: Child and parent versions. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40(8), 937–944. <https://doi.org/10.1097/00004583-200108000-00016>
- Spence, S. H., & Rapee, R. M. (2016). The etiology of social anxiety disorder: An evidence-based model. *Behaviour Research and Therapy*, 86, 50–67. <https://doi.org/10.1016/j.brat.2016.06.007>
- Tobii AB. Tobii Pro Lab (Version 1,241) [Computer software]. Danderyd, Sweden: Tobii AB.
- Torchiano, M. (2020). *effsize: Efficient Effect Size Computation*. doi:10.5281/zenodo.1480624, R package version 0.8.1, <https://CRAN.R-project.org/package=effsize>.
- Van Beek, Y., Van Dolderen, M. S., & Demon Dubas, J. J. (2006). Gender-specific development of nonverbal behaviours and mild depression in adolescence. *Journal of Child Psychology and Psychiatry*, 47(12), 1272–1283.
- van den Bos, E., Tops, M., & Westenberg, P. M. (2017). Social anxiety and the cortisol response to social evaluation in children and adolescents. *Psychoneuroendocrinology*, 78, 159–167. <https://doi.org/10.1016/j.psyneuen.2017.02.003>
- Wells, A., Clark, D. M., Salkovskis, P., Ludgate, J., Hackmann, A., & Gelder, M. (1995). Social phobia: The role of in-situation safety behaviors in maintaining anxiety and negative beliefs. *Behavior Therapy*, 26(1), 153–161.
- Wergeland, G. J. H., Fjermestad, K. W., Marin, C. E., Bjelland, I., Haugland, B. S. M., Silverman, W. K., ... & Heiervang, E. R. (2016). Predictors of treatment outcome in an effectiveness trial of cognitive behavioral therapy for children with anxiety disorders. *Behaviour Research and Therapy*, 76, 1–12.
- Westenberg, P. M., Bokhorst, C. L., Miers, A. C., Sumter, S. R., Kallen, V. L., van Pelt, J. & Blöte-Aanhane A. W. (2009), A prepared speech in front of a pre-recorded audience: Subjective, physiological, and neuroendocrine responses to the Leiden Public Speaking Task, *Biological Psychology* 82(2): 116–124.
- Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis* (2nd ed.) [PDF]. Springer International Publishing.
- Wickham, H., François, R., Henry, L., Müller, K., Vaughan, D. (2023). *dplyr: A Grammar of Data Manipulation*. R package version 1.1.4.
- Yoon, K. L., & Zinbarg, R. E. (2008). Interpreting neutral faces as threatening is a default mode for socially anxious individuals. *Journal of Abnormal Psychology*, 117(3), 680.

