



Universiteit  
Leiden  
The Netherlands

## Microbial footprints of tomato domestication

Sarango Flores, S.W.

### Citation

Sarango Flores, S. W. (2026, January 6). *Microbial footprints of tomato domestication*. NIOO-thesis. Retrieved from <https://hdl.handle.net/1887/4285898>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4285898>

**Note:** To cite this publication please use the final published version (if applicable).

# Chapter 3

---

## **Changes in taxonomic diversity and functional traits of the tomato root microbiome along a domestication trajectory**

**Stalin Sarango Flores<sup>1,2</sup>, Ben O. Oyserman<sup>1</sup>, Luisa M. Arias Giraldo<sup>1</sup>, Nejc Stopnisek<sup>3</sup>,  
Pieter van 't Hof<sup>4,5</sup>, Jos M. Raaijmakers<sup>1,2</sup>, Viviane Cordovez<sup>1</sup>**

<sup>1</sup> Department of Microbial Ecology, Netherlands Institute of Ecology, Wageningen, 6708PB, Netherlands

<sup>2</sup> Institute of Biology, Leiden University, Leiden, 2333BE, Netherlands

<sup>3</sup> Department of Microbiological Research, National Laboratory of Health, Environment and Food, Maribor, 2000, Slovenia

<sup>4</sup> Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito USFQ, Quito, 170901, Ecuador

<sup>5</sup> Instituto de Microbiología, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito USFQ, Quito, 170901, Ecuador

---

## Abstract

Domestication and breeding have substantially altered the genetic and phenotypic traits of multiple plant species. To date, however, the impact of domestication on the taxonomic and functional diversity of microorganisms colonizing plant tissues remains largely unexplored for the majority of plant species. Here, we examined the bacterial microbiome associated with roots of eight wild and domesticated tomato genotypes grown in three distinct soil management gradients, ranging from native and agricultural soils from the tomato's center of origin in Ecuador to a soil from a tomato greenhouse in The Netherlands. Our findings revealed a higher taxonomic diversity in Ecuadorean agricultural soils than in native and greenhouse soils. Soil was the primary factor governing tomato microbiome assembly, followed by the root compartment and plant genotype. Root microbiomes of tomato grown in native Ecuadorian soils had a higher abundance in Acidobacteriota, Proteobacteria, Bacteroidota, Chloroflexi, and Myxococcota, whereas Actinobacteriota, Cyanobacteria, Firmicutes, and Patescibacteria dominated the root microbiome of tomato grown in agricultural and greenhouse soils. Also, we found significant differences in the root microbiome of the different tomato genotypes, with enhanced abundance in Bacteroidota, Proteobacteria and Chloroflexi on roots of wild tomato genotypes, and Actinobacteriota, Firmicutes and Cyanobacteria on roots of domesticated tomato genotypes. Representative metagenome assembled genomes (MAGs) of bacterial taxa of wild tomato genotypes grown in native Ecuadorean soil were enriched in genes associated with motility and chemotaxis, carbon metabolism, and stress response, while MAGs from the microbiome of domesticated tomatoes grown in agricultural and greenhouse soils were enriched for genes involved in metabolism of nitrogen, iron, amino acids and vitamins. This study highlights the pivotal role of habitat domestication and genetic changes of tomato in microbiome assembly, with alterations in the abundance of functional microbial traits associated with plant growth and health.

## Introduction

Plant domestication represents an evolutionary process where wild ancestral species were adapted to human needs, ultimately leading to our modern-day crop cultivars (Purugganan & Fuller, 2009). For most crops, this process was targeted at desirable traits such as larger fruit size, palatability, nutritional quality, reproductive timing or stress resistance (Bergougnoux, 2014). While favoring certain alleles during this process of artificial selection and breeding for these desired traits, many modern crop cultivars show a reduced genetic diversity as other (overlooked) genes were lost (Fernie & Yan, 2019). This phenomenon is referred to as the “domestication syndrome” and was first described for cereals in the 1970s by Harlan et al. (1973). The domestication syndrome not only comprises genetic changes caused by human artificial selection but also unintentional effects arising from agricultural practices to grow the domesticated crops, also referred to as habitat domestication (Barnes et al., 2024; Fernie & Yan, 2019; Soldan et al., 2021). Thus, domesticated species are notably different from their wild relatives, with altered morphology, physiology and ecological interactions, including those affecting microbiome assembly (Barnes et al., 2024; Cordovez et al., 2019; Hassani et al., 2020; Martínez-Romero et al., 2020; Sarango Flores et al., 2023).

Plant-associated microbiomes play essential roles in plant growth and health due to their contribution to nutrient acquisition and stress tolerance, thereby supporting and expanding the host plant’s functional capabilities (Adedayo et al., 2023; Ling et al., 2022; Pérez-Rodríguez et al., 2020; Santhanam et al., 2015; Shivega & Aldrich-Wolfe, 2017; Zuluaga et al., 2021). Several factors influence the assembly of the plant microbiome, including abiotic factors (e.g. soil types, climatic conditions, agricultural management practices), plant genotype, and the diversity of local microbial pools in the surrounding habitat (Abdullaeva et al., 2022; Berg & Smalla, 2009; Hewitt et al., 2023; Ofek-Lalzar et al., 2014). Traits such as root architecture, exudate composition and nutritional requirements in domesticated plants significantly affect the composition and diversity of their microbiomes, which in turn may affect their functional benefits to the plant host (Abbamondi et al., 2016; Dennis et al., 2010; Gutierrez & Grillo, 2022; Soldan et al., 2021). Plants recruit specific microbial taxa by specific constituents in root exudates that act as chemical signals and orchestrate the surrounding microbial communities to take advantage of the outsourced functions (Barnes et al., 2024; Nakayasu et al., 2023; Walker et al., 2003; Wen et al., 2023). Furthermore, prevailing soil types, changing climatic conditions and agricultural management practices, also modulate plant microbiome assembly (Cao et al., 2024; Flemer et al., 2022; Wang et al., 2022; Xue et al., 2018). For instance, domesticated plants rely on fertilizers and pesticides and much less on their microbial alliances, which may also limit crop adaptability to future adverse environmental conditions such as drought. This network of

different factors driving microbiome assembly creates complex dynamics that can vary across plant genotypes and growth conditions (Mo et al., 2024; Pérez-Jaramillo et al., 2018; Soldan et al., 2021).

Tomato (*Solanum lycopersicum* L.) is an iconic example of plant domestication, which started in the center of origin in the Andean region in South America (Knapp & Peralta, 2016; Razifard et al., 2020). The domestication journey of tomato began with the wild tomato relative *S. pimpinellifolium*, which provided the early traits selected and crossed by indigenous people into the landrace *S. lycopersicum* var. *cerasiforme*. This variety eventually spread beyond its center of origin and continued its domestication process in Central America (Blanca et al., 2012, 2022; Razifard et al., 2020). This process has continued for thousands of years outside of the native tomato habitat, to meet regional phenotypic preferences and cultivation management practices worldwide, which in turn resulted in many modern cultivated tomato varieties of *S. lycopersicum* (Mata-Nicolás et al., 2020; Sarango Flores et al., 2023; Sim et al., 2011).

Examples of notable phenotypic changes in domesticated tomato cultivars include larger fruits with a reduced seed number, uniform shapes and colors, as well as specific metabolic pathways to prioritize sweetness (or reduce bitterness) that are linked to changes in secondary metabolite composition (Bai & Lindhout, 2007; Bergougnoux, 2014). As a consequence of the domestication syndrome, modern cultivated tomatoes suffered a genetic bottleneck which caused a narrowed gene pool compared to their closest relatives, making them more vulnerable to pests, diseases and abiotic stresses (Aflitos et al., 2014; Gao et al., 2019; Kahlon et al., 2020). This loss of genetic variation may also have reduced the modern tomato's ability to effectively select for and interact with the microbial communities surrounding the plants tissues, which in turn can reflect on plant performance (Chen et al., 2022; Huang et al., 2022; Malacrinò et al., 2022; Nerva et al., 2022; Yue et al., 2023). This was exemplified in our previous elaborate analysis of the microbiome of a tomato recombinant inbred line (RIL) population by Oyserman et al. (2022), which revealed significant differences in microbiome assembly across 100 tomato lines and identified specific genetic associations between loci in the tomato genome and in the microbiome metagenome. Additionally, work by Smulders et al. (2021) suggested that tomato wild relatives establish more beneficial interactions with their rhizosphere microbiome than domesticated cultivars.

Building on the findings of Chapter 2, which explored the rhizosphere microbiome of wild tomato *Solanum pimpinellifolium* in its native Andean habitats, we identified a conserved microbial signature dominated by Proteobacteria. While these results revealed how wild tomatoes interact with their native microbiome, they did not address how tomato domestication process and agricultural management may have impacted plant-microbiome associations. To further explore the taxonomic and functional changes in tomato rhizosphere and endosphere microbiomes, we cultivated eight tomato genotypes, representing different stages of the tomato domestication trajectory, in native and agricultural soils from the center of origin in the South of Ecuador, and in a contrasting greenhouse production soil from The Netherlands. In this chapter, we analyzed the bacterial community composition of the rhizosphere and root endosphere by 16S rRNA gene amplicon sequencing and by shotgun metagenomics. We hypothesized that habitat domestication has the largest impact on microbiome assembly, influencing both the taxonomic and functional diversity of the root microbiome.

## Results

### ***Soil and genotype drive the taxonomic diversity of the tomato rhizosphere microbiome***

No significant differences (Kruskal-Wallis test,  $p > 0.05$ ) were found in alpha diversity, represented by Shannon's diversity index, for the rhizosphere bacterial communities from the tomato genotypes grown on the different soil types (Figure 1). However, the beta diversity revealed significant differences among soil origin (PERMANOVA,  $p = 0.0001$ ), soil type (PERMANOVA,  $p = 0.0001$ ) and tomato domestication degree (PERMANOVA,  $p = 0.024$ ) (Figure 2a). Soil origin and type explained 64% and 39% of the total variability in the rhizobacterial community composition, respectively.




**Figure 1.** Shannon diversity index of the rhizosphere microbiome of tomato plants grown in Ecuadorian native and agricultural soils, and Dutch greenhouse soil. Boxplots show no significant difference (Kruskal-Wallis test,  $p < 0.05$ ,  $n = 10$ ) of Shannon diversity index between tomato genotypes. SHA: *S. habrochaetes*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solariño; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

More detailed analysis by soil origin showed significant effects of the eight different tomato genotypes, ranging from 37% of the total variation in Ceiba and Limones native soils from the center of origin, 37-38% in the Ceiba and Limones agricultural soils from the center of origin, and 44% in the Dutch greenhouse production soil (Figure 2b–f). Significant differences were not observed between the wild and intermediate groups but were significant between the four intermediate-wild and four domesticated tomato genotypes (Adonis test,  $p < 0.05$ , Supplementary Table S4).




**Figure 2.** Rhizosphere bacterial community structure of different tomato genotypes grown in Ecuadorian native and agricultural soils, and Dutch greenhouse soil. Principal Coordinate Analysis (PCoA) of rhizosphere bacterial communities of wild ( $n = 2$ ), intermediate ( $n = 2$ ) and domesticated ( $n = 4$ ) tomato genotypes (a) PCoA by soil type; (b) – (f) PCoA by tomato genotype per soil type. (Agr = agricultural soil, Nat = native soil).

Subsequent differential abundance analysis revealed that soil type impacted the differential enrichment and depletion of ASVs in the rhizosphere of tomato. We first compared ASV abundance in the rhizosphere of tomatoes grown in native and agricultural soils from the South Andes in Ecuador. In native soils, a total of 490 ASVs were significantly more abundant in wild-intermediate tomato genotypes (cluster C3) and 1028 ASVs in the four domesticated tomato genotypes (cluster C4); in the agricultural soils, 1279 ASVs were significantly more abundant in wild-intermediate tomato genotypes (cluster C1) and 1121 ASVs in domesticated tomato genotypes (cluster C2) (Figure 3a). We next compared the differential abundance in the rhizosphere of tomato between the Ecuadorean agricultural soils and the Dutch greenhouse soil. In the Ecuadorean agricultural soils, 1286 ASVs were significantly more abundant in wild-intermediate tomato genotypes (cluster C5), and 2296 ASVs in domesticated tomato genotypes (cluster C4). In the Dutch greenhouse soil, 1148 ASVs showed higher abundance in wild and intermediate tomatoes (cluster C3), but only 109 ASVs were more abundant in domesticated tomatoes (cluster C2) (Figure 3b).




**Figure 3.** Differential abundance of the tomato rhizosphere amplicon sequence variants (ASVs) in pairwise comparisons of different soil types. **(a)** Heatmap of ASVs in the rhizosphere of tomatoes grown in Ecuadorian native and agricultural soils; clusters C1 = 1279; C2 = 1121; C3 = 490; C4 = 1028 ASVs. **(b)** Heatmap of ASVs in the rhizosphere of tomatoes grown in Ecuadorian agricultural and Dutch greenhouse soils; clusters C1 = 178; C2 = 109; C3 = 1148; C4 = 2296; C5 = 1286 ASVs. Significant ASVs identified from the DESeq differential abundance analysis were hierarchically clustered using Ward's method as the clustering algorithm. SHA: *S. habrochaites*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

By grouping the significantly abundant rhizosphere ASVs from each cluster at phylum and genus level (Figure 4), we observed that the rhizosphere of tomato plants grown in soils from the center of origin, i.e. native and agricultural soils, had a higher relative abundance of Actinobacteriota, Bacteroidota, Cyanobacteria and Proteobacteria when compared with tomato plants grown in the Dutch greenhouse soil, which were more abundant in Chloroflexi, Firmicutes, Gemmatimonadota, Myxococcota, Patescibacteria and Planctomycetota (Figure 4a). At genus level, the rhizosphere of tomatoes grown in native soils from the center of origin had a higher relative abundance of *Brevundimonas*, *Cellvibrio*, *Dyadobacter*, *Ohtaekwangia*, *Pseudomonas*, *Rhizobacter* and *Sphingomonas*. In contrast, tomatoes grown in agricultural soils had an increased relative abundance of *Actinoplanes*, *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium*, *Massilia*, *Nostoc* and *Streptomyces*, whereas tomatoes grown in the Dutch greenhouse soil showed greater abundance of *Bacillus*, *Devosia*, *Edaphobaculum*, *Mesorhizobium*, *Paenibacillus* and *Sericytochromatia*.

The tomato domestication degree was also linked to changes in specific rhizobacterial taxa. For instance, in native soils, wild and intermediate tomato genotypes had higher relative abundances of *Chitinophaga*, *Dyadobacter*, *Ohtaekwangia* (Bacteroidota) as well as *Cellvibrio* and *Rhizobacter* (Proteobacteria). In contrast, *Nocardioides* (Actinobacteriota) were enriched in the rhizosphere of domesticated tomato genotypes grown in the Ecuadorean native soils. In the two Ecuadorean agricultural soils, wild and intermediate tomato genotypes showed higher relative abundance of *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium* (Proteobacteria) whereas domesticated tomatoes harbored more *Actinoplanes* (Actinobacteriota). In the Dutch greenhouse soil, differences among tomato genotypes were observed in *Bacillus* and *Paenibacillus* (Firmicutes), which were more abundant in wild and intermediate tomatoes, while *Devosia* (Proteobacteria), *Mesorhizobium* (Proteobacteria) and *Sericytochromatia* (Cyanobacteria) were more abundant in domesticated tomatoes. In all three soil types, *Pseudomonas* and *Massilia* had higher abundance in wild-intermediate tomatoes, whereas members of the genera *Sphingomonas* (Proteobacteria) and *Streptomyces* (Actinobacteriota) showed greater relative abundance in domesticated tomatoes in all five soils (Figure 4b).





**Figure 4.** Rhizosphere bacterial community composition of tomato genotypes grown in native and agricultural soils. **(a)** Relative abundance of bacterial phyla in rhizosphere of different tomato genotypes grown in Ecuadorian native and agricultural soils and Dutch greenhouse soil; “Other” category corresponds to grouped phyla with relative abundance < 0.01; **(b)** Relative abundance of most abundant genera found in rhizosphere of different tomato genotypes. ASVs with significant differential abundance between soils and tomato genotypes were grouped according to their phylum or genus level and plotted as stacked bar and bubble charts, respectively. SHA: *S. habrochaites*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

### ***Effects of domestication on functional traits of the tomato rhizosphere microbiome***

Following shotgun metagenomics of the rhizosphere samples, 127 MAGs (high quality bins completeness >70%, contamination <10%) were obtained. Of these, 12 MAGs with the highest quality were selected for functional analysis based on their taxonomic consistency with amplicon sequencing results. Thus, functions annotated by SEED Subsystems from selected MAGs related to habitat and the domestication degree of the tomato genotypes revealed significant shifts in microbial functionality. For example, MAGs taxonomically delineated as Bacteroidota (*Chitinophaga*, *Fluviicola*) or Proteobacteria (*Cellvibrio*, *Pseudomonas*, *Rhizobacter* and Rhizobiaceae) were more abundant in wild-intermediate tomatoes in native and agricultural soils, showed a higher abundance of genes associated with motility and chemotaxis, as well as metabolism of carbohydrates and stress response. On the other hand, MAGs taxonomically delineated as Micromonosporaceae, Bacillaceae, Cyanobacteria, *Sphingomonas* and *Streptomyces*, representative taxa of domesticated tomatoes grown in agricultural and greenhouse soils, exhibited a higher number of genes associated with iron acquisition and metabolism of nitrogen, amino acids and derivatives; and cofactors, vitamins, prosthetic groups, pigments (Figure 5).



### ***Tomato endosphere microbiome assembly is soil type dependent***

The bacterial communities detected in the root endosphere displayed a similar Shannon diversity index across tomato genotypes and soils, differing only among the eight tomato genotypes grown in native Ecuadorean soil (Kruskal-Wallis test,  $p = 0.0044$ ). A higher diversity index was observed for domesticated tomato genotypes compared to wild-intermediate tomato genotypes. Interestingly, a same yet statistically insignificant trend was observed for agricultural and greenhouse soils, i.e. a minor yet progressive increase of bacterial alpha diversity along the tomato domestication trajectory (Figure 6).




**Figure 6.** Shannon diversity index in tomato root endosphere grown in Ecuadorian native and agricultural soils and Dutch greenhouse soil. Different letters above boxplots show significant difference (Wilcoxon test,  $p < 0.05$ ,  $n = 10$ ) of Shannon diversity index between tomato genotypes. SHA: *S. habrochaites*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

Overall, PCoA of the root endosphere showed significant differences between soil origin and soil types (PERMANOVA,  $p = 0.0001$ ), but surprisingly there were no significant differences between tomato genotypes or domestication groups (PERMANOVA,  $p > 0.05$ ) (Figure 7a). When the tomato endosphere microbiome composition was analyzed per soil type (i.e. native, agricultural, greenhouse), significant differences were observed between tomato genotypes and domestication degree. This variation observed in the PCoA plot appeared to be primarily driven by the broader dispersion of wild tomato genotypes across the different soils. Although some dispersion is observed in domesticated tomato genotypes, wild genotypes exhibited greater variability. This variation likely contributed to the significant differences found between domesticated tomatoes and both intermediate and wild tomato genotypes (Adonis test,  $p < 0.05$ , Supplementary Table S4). Specifically, tomato genotype showed 21% of the total variation in

Ceiba agricultural and native soils and in Limones native soil, while 22% was observed in Limones agricultural and greenhouse soils. The domestication effect, i.e. variation in endosphere microbiome composition between wild, intermediate and domesticated tomato groups, showed 7% of the total variation per soil. This percentage of variation represents a marked decrease compared to the variation observed in the rhizosphere microbiome of 20% (Figure 7b–f).




**Figure 7.** Principal Coordinate Analysis (PCoA) of bacterial communities in tomato root endosphere grown in different soil types. (a) PCoA by soil type and (b) – (f) PCoA by tomato genotype in each soil.

Furthermore, the differential abundance analysis of bacterial ASVs by DESeq (Wald test) through pairwise comparison between native and agricultural soils revealed no significant differences between the tomato domestication groups. The observed ASV clusters (Figure 8a) also suggest that the differences were mainly attributed to the soil type in which the tomato genotypes were grown. Similar results were found when agricultural and greenhouse soils were compared (Figure 8b).




**Figure 8.** Differential abundance of the tomato root endosphere amplicon sequence variants (ASVs) in pairwise comparisons of different soil types. **(a)** Heatmap of ASVs in the endosphere of tomatoes grown in Ecuadorian native and agricultural soils; clusters C1 = 129; C2 = 304 ASVs. **(b)** Heatmap of ASVs in the endosphere of tomatoes grown in Ecuadorian agricultural soils and Dutch greenhouse soil; clusters C1 = 466; C2 = 602 ASVs. Significant ASVs identified from the DESeq differential abundance analysis were hierarchically clustered using Ward's method as the clustering algorithm. SHA: *S. habrochaites*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

The significantly different ASV clusters in the tomato root endosphere from native and agricultural Ecuadorean soils showed increased abundance of the phyla Actinobacteriota, Bacteroidota, Cyanobacteria, Firmicutes and Proteobacteria. In contrast, Actinobacteriota and Firmicutes were predominantly more abundant in the endosphere of tomato plants grown in Dutch greenhouse soil (Figure 9a). At genus level, the endosphere microbiome of tomato plants grown in native soils showed higher abundance of *Ammoniphilus*, *Bacillus*, *Domibacillus*, *Fictibacillus*, *Lysinibacillus*, *Pseudomonas* and *Rhizobacter*, than that of plants grown in agricultural and greenhouse soils. On the other hand, agricultural and greenhouse soils, i.e. managed soils, showed higher abundance of *Clostridium*, *Cohnella*, *Massilia*, *Paenibacillus*, *Streptomyces* and *Tumebacillus*. Particularly, greenhouse soil increased the abundance of *Tumebacillus*, *Paenibacillus* and *Cohnella* in tomato endosphere. Although soil was shown to be the main driver of the root endosphere microbiome composition, domestication appeared to impact the abundance of *Actinoplanes*, *Cohnella*, *Massilia* and *Streptomyces*, which had higher abundance in the root endosphere of domesticated than of wild-intermediate tomato genotypes, especially when grown in agricultural soil (Figure 9b). Considering the substantial interference of plant DNA in shotgun metagenomic sequencing of endosphere samples, no data on functional genes and traits of the endophytic tomato microbiome were obtained.




**Figure 9.** Root endosphere bacterial community composition at phylum and genus level of tomato genotypes in different soil types. **(a)** Relative abundance of bacterial phyla in root endosphere of different tomato genotypes grown in Ecuadorian native and agricultural soils and Dutch greenhouse soil; “Other” category corresponds to grouped phyla with relative abundance < 0.01; **(b)** Relative abundance of highest abundant genera found in root endosphere of different tomato genotypes. ASVs with significant differential abundance between soils and tomato genotypes were grouped according to their phylum or genus level and plotted as stacked bar and bubble charts, respectively. SHA: *S. habrochaites*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

## Discussion

Domestication not only evolved specific traits of wild plant species to favor human needs, it also led to substantial changes in their habitats, moving from environmentally harsh, native ecosystems to heavily managed agricultural soils. Additionally, human migration and extensive exchanges of plant germplasm worldwide have further contributed to the vast genetic variation seen in modern tomatoes (Blanca et al., 2022). Agricultural environments can also deplete putative plant growth-promoting rhizobacteria in modern crop accessions (Reid et al., 2024). In this study, we used tomato as a 'model' plant to decipher the impact of domestication on microbiome assembly. To this end, we selected different genotypes and soils, including wild and domesticated tomato genotypes and native and managed soils from the center of origin (Ecuador), as well as greenhouse soil from the center of production (Netherlands) (Supplementary material Tables S1, S2). In this study, we used the same two parental tomato lines (*S. pimpinellifolium* and *S. lycopersicum* cv. Moneymaker) and greenhouse tomato soil as used in our earlier study on microbiome assembly of a tomato RIL population with approximately 100 genotypes (Oyserman et al., 2022). It should be noted that the number of tomato genotypes ( $n = 8$ ) and soils ( $n = 5$ ) used in this study is limited to make solid conclusions on the impact of domestication on microbiome assembly. However, the results obtained can serve as a stepping stone for more elaborate screenings and to generate hypotheses that can be experimentally validated on larger sets of plant genotypes and soils.

Our findings revealed that selective filtering of the soil microbiome was stronger in the tomato rhizosphere than in the endosphere (Figures 2, 7). Significant variations in rhizosphere microbiome composition were observed per soil, with wild-intermediate tomato genotypes clustering separately from the domesticated tomatoes (Figure 2). This was further supported by the differential abundance analysis which revealed a similar clustering pattern of the ASVs and whose separation was driven by soil type and the degree of tomato domestication (Figure 3). Soil plays a predominant role in plant microbiome assembly due to its properties, especially nutrient content and the diversity of the microbial 'seed bank' (Cordovez et al., 2019; Ling et al., 2022; Philippot et al., 2024). Particularly, soil amendments such as organic matter, N, P and K, have been shown in several studies to considerably alter the tomato bacterial community assembly in the rhizosphere (Cheng et al., 2020; Dixon et al., 2024; Garcia et al., 2024; Naumova et al., 2022; Zhang et al., 2022). On the other hand, plant genotype can act as a microbiome filter due to the quantitative and/or qualitative differences in root exudation composition, which selects for and enhances the proliferation of specific taxa within the microbiome, as reported in previous studies (French et al., 2020; Poudel et al., 2019). Our results suggest that the differentiation in rhizosphere microbiome

assembly occurred later in the domestication trajectory, likely during breeding and genetic improvement. Based on this, further examination of the differences in tomato root exudate composition per tomato genotype and the effect of soil amendments on the rhizosphere microbiome will be needed to decipher genotype-specific signatures in rhizosphere microbiome assembly.

When summarizing the impact of tomato and habitat domestication on rhizosphere assembly (Figure 4), our results show that: i) Bacteroidota and Proteobacteria had a higher relative abundance in the rhizosphere microbiome of tomatoes grown in native Ecuadorian soils, ii) Actinobacteriota and Cyanobacteria were more abundant in the rhizosphere microbiome of tomatoes grown in Ecuadorean agricultural soils, and iii) Firmicutes were more abundant in the rhizosphere of tomatoes grown in the Dutch greenhouse soil. At genus level, *Chitinophaga*, *Dyadobacter*, *Fluviicola*, *Ohtaekwangia* (Bacteroidota), as well as *Brevundimonas*, *Cellvibrio* and *Rhizobacter* (Proteobacteria) were more abundant in wild-intermediate tomato genotypes, while *Lysobacter* (Proteobacteria) and *Nocardioides* (Actinobacteriota) were more abundant in domesticated tomato genotypes grown in native soil. In contrast, in agricultural soil, wild-intermediate tomatoes were more abundant in *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium* (Proteobacteria), while domesticated tomatoes showed increased genera such as *Actinoplanes*, *Nocardioides* (Actinobacteriota) and *Nostoc* (Cyanobacteria). For greenhouse soil, *Bacillus*, *Paenibacillus* (Firmicutes) were significantly more abundant in wild-intermediate tomatoes, while *Sericytochromatia* (Cyanobacteria) was more abundant in domesticated tomato genotypes. Interestingly, across all soils, *Pseudomonas* and *Massilia* (Proteobacteria) were more abundant in wild-intermediate tomato genotypes, while *Streptomyces* (Actinobacteriota) and *Sphingomonas* (Proteobacteria) progressively increased along the tomato domestication trajectory (Figure 4).

Several *Pseudomonas* species are known as plant growth-promoting rhizobacteria (PGPR) due their versatile abilities in root colonization, nutrient acquisition, anti-oxidant activities, pathogen suppression, and stress adaptation (Alattas et al., 2024; Mekureyaw et al., 2022; Sah et al., 2021; Zboralski & Filion, 2020). *Massilia* exhibits copiotrophic behavior and effective attachment to biological surfaces, including fungal hyphae, filamentous algae and Cyanobacteria (Ofek et al., 2012; Salomon et al., 2003; Scheublin et al., 2010), which can make it a successful root colonizer. In our earlier studies, *Massilia* together with *Rhizobium* was also abundant in successively planted wild tomato (Cordovez et al., 2021). Whether these beneficial traits are also prominent for the *Pseudomonas* and *Massilia* species we detected in wild-intermediate tomato genotypes remains to be further investigated.

Our results on the enrichment of *Streptomyces* in the rhizosphere of domesticated tomatoes are consistent with findings from other studies. First and foremost, our results confirm and extend the results of our earlier study on the microbiome of a tomato RIL population, conducted with the same Dutch greenhouse soil and the same parental lines *S. pimpinellifolium* and *S. lycopersicum* cv. Moneymaker (Oyserman et al., 2022). We observed similar patterns of abundance of specific rhizosphere bacteria, especially the significant abundance of *Streptomyces* in the rhizosphere of domesticated tomatoes in any soil type. In our previous study, the increased abundance of *Streptomyces* was associated with a ‘modern allele’ on chromosome 6 (Oyserman et al., 2022). Also in other studies, similar patterns were reported. For example, Allard (2016) observed higher abundance of members of the Streptomycetaceae family in domesticated tomato (*S. lycopersicum* BHN602) grown in greenhouse conditions. Similarly, Dixon et al. (2024) found that modern (ca. 2020) and traditional tomatoes (ca. 1900) had a greater relative abundance of Actinobacteriota than wild tomatoes, especially in P-fertilized soil. In addition, Chen et al. (2022) demonstrated differential recruitment of rhizobacteria based on tomato fruit color phenotype, with Actinobacteriota being more abundant in tomatoes with yellow fruit compared to those with red fruit. Furthermore, Actinobacteriota can be competitive for root exudates and micronutrients, such as iron, as well as for antibiotic production (Nazari et al., 2023; Oyserman et al., 2022; Pérez-Jaramillo et al., 2016, 2018; Zhao et al., 2018), which may facilitate the successful colonization of *Streptomyces* in domesticated tomato rhizosphere. The observed enrichment of *Sphingomonas* in the rhizosphere of domesticated tomatoes is also confirms and extends the study by Lee et al. (2019) who proposed *Sphingomonas* as one of the indicators of the rhizosphere of domesticated tomato genotypes sampled from different greenhouses. Interestingly, we also observed an increase in Cyanobacteria genera, such as *Nostoc* and *Sericytochromatia* in domesticated tomato genotypes grown in agricultural and in greenhouse soils, respectively. This suggests that these bacterial genera are likely responding to nutrient availability, particularly N and P (Xu et al., 2017), and possibly to the reduced use of organic fertilization in agricultural soils, as observed by Zou et al. (2024).

Our profiling of the microbial traits of key tomato rhizosphere bacteria, in relation to genotype and habitat domestication, provided a better understanding of the interactions between these bacteria, their different tomato hosts, and the soil types (Figure 5; Supplementary Table S6). Unfortunately, we did not find MAGs assigned as *Massilia* or the Oxalobacteraceae family, which were abundant taxa in the rhizosphere of wild tomatoes across the soil types tested. We found a higher number of genes associated with the SEED subsystems Motility and Chemotaxis in rhizosphere MAGs of wild tomato genotypes grown in native soil and taxonomically delineated as Bacteroidetes, Cellvibrionales, Pseudomonadales and Rhizobiales. This observation aligns with find-

ings by Sun et al. (2021), who found that the root microbiome of wild rice accessions exhibited a greater abundance of bacterial chemotaxis genes than the microbiomes of their domesticated counterparts. Moreover, abundance of these bacterial taxonomic lineages was associated by Yin et al. (2020) with plant protection against pathogens. They found a higher abundance of Bacteroidetes, Pseudomonadales and Rhizobiales in tomato cultivars that resisted bacterial canker caused by *Clavibacter michiganensis*. In addition, the higher number of genes associated with flagellar motility observed in these MAGs, particularly in *Pseudomonas*, *Cellvibrio* and Rhizobiaceae, can be linked to enhanced rhizosphere colonization and increased carbohydrate metabolism as revealed in other studies (Barajas et al., 2020; de Weert et al., 2002; Liu et al., 2024; Ramoneda et al., 2024; Zuluaga et al., 2021). Moreover, carbohydrate metabolism also showed a higher number of genes in MAGs associated with the microbiomes of the wild tomato genotypes, consistent with findings by Zboralski & Filion (2020). Additionally, the higher abundance of microbial functions associated with stress response found here may align with the observation that the wild tomato rhizosphere has a greater capacity to adapt to environmental challenges, such as oxidative or osmotic stress, as reported in several studies (Alzate Zuluaga et al., 2021; Anjum et al., 2025; Schmitz et al., 2022).

MAGs associated with the rhizosphere of domesticated tomato genotypes, especially in agricultural and greenhouse soils, exhibited a higher abundance of genes involved in amino acid metabolism, cofactors and vitamins, as well as genes related to N, P and Fe metabolism. Specifically, the increased abundance of genes related to ammonification, allantoin utilization, and siderophore biosynthesis, suggests that the microbiome associated with domesticated tomatoes may respond to fertilizer inputs with both macro- and micronutrients, as indicated in other studies (Adedayo et al., 2022; Dixon et al., 2024; Garcia et al., 2024; Smulders et al., 2021; Terra et al., 2021; Zhang et al., 2022).

With respect to the endosphere tomato microbiome, soil type was the primary determinant of the community assembly (Figures 8, 9). This effect was particularly pronounced in the phylum Firmicutes with *Bacillus* remaining consistent across soils, while *Ammoniphilus*, *Domibacillus*, *Fictibacillus* and *Lysinibacillus* showed a progressive decline in abundance from Ecuadorean native to Dutch greenhouse soil. Conversely, *Cohnella*, *Paenibacillus* and *Tumebacillus* exhibited a progressive increase in abundance from Ecuadorean native to Dutch greenhouse soil. Collectively, these results suggest that soil management influenced microbiome assembly of the tomato root endosphere. Other taxa exhibited genotype-specific differences for Ecuadorean native and agricultural soils. For example, *Brevundimonas*, *Pseudomonas*, *Rhizobacter* (Proteobacteria) were more abundant in the endosphere of wild-intermediate tomatoes grown in native soils, while *Actinoplanes*, *Streptomyces* (Actinobacteriota) and *Tolypotrix* (Cyanobacteria) were more abundant in the endosphere of domesticated tomatoes grown in agricultural soil.

Further investigation is needed to understand how root exudates and soil management practices influence the selective enrichment of these endosphere taxa and their ability to withstand plant defense responses, as well as to adapt to the specific environmental conditions associated with different tomato genotypes and soil types.

In conclusion, our results highlight the significant impact of the tomato domestication process on its microbiome, with habitat domestication playing a pronounced role in root microbiome assembly. Our results suggest that the major differentiation in rhizosphere microbiome assembly occurred later in the domestication process, i.e. during the breeding and genetic improvement stages. Additionally, we observed that wild tomatoes grown in native soils exert strong selective pressure on the rhizosphere microbiome, favoring bacteria enriched in genes associated with motility, chemotaxis and stress response. In contrast, domesticated tomatoes establish microbiome associations with bacteria that exhibit functional adaptations to agricultural and greenhouse environments, presumably fertilizer inputs. Our study emphasizes the importance of understanding microbiome shifts driven by both genotype and habitat domestication. These insights are essential for integrating microbiome management strategies into tomato cultivation and considering the microbiome as a complementary feature in breeding programs. Future research should focus on the identity and role of specific root exudates in microbial assembly for both wild and domesticated tomatoes. Additionally, the effects of soil amendments (organic and synthetic fertilizers) and other agricultural practices on tomato genotypes should be further explored. It is also important to assess the phenotypic response of tomatoes to rhizobacterial inoculation, specifically in terms of plant growth, resistance to (a)biotic stresses, and yield, with an emphasis on the genetic background of different tomato cultivars to maximize the efficiency of microbial interventions. Such studies could provide valuable insights to promote sustainable farming practices through the integration of agricultural techniques with microbial management.

## Materials and Methods

### *Greenhouse experiment*

Native and agricultural soils from the tomato's center of origin were collected in June 2021 from two locations in the province of Loja in Southern Ecuador (Ceiba 4°18'07.6"S, 80°13'16.7"W and Limones 4°23'09.2"S, 80°20'50.7"W, Zapotillo, Loja, Ecuador). Native soils were collected from a natural vegetation close to the agricultural locations. Agricultural soil from Ceiba was sampled from a field that was cultivated with onion (vegetative growth stage), whereas the agricultural soil in Limones was sampled from a local rice field (grain filling stage). All four soils were air dried at room

temperature and subsequently sieved (2 mm diameter sieve mesh) and shipped to the greenhouse facility at the Netherlands Institute of Ecology (NIOO-KNAW) in Wageningen, The Netherlands in compliance to the permit approved by the Ecuadorian Ministry of Environment. A Dutch greenhouse used for tomato seed production in South-Holland, (51°57'47"N, 4°12'16"E, collected in June 2017), provided the greenhouse soil from the center of production (Supplementary material Table S1). The five soils of interest were coded as follows: Ceiba native (CN), Ceiba agricultural (CA), Limones native (LN), Limones agricultural (LA) and, Dutch greenhouse soil (GH) (Figure 10).

A total of 8 tomato genotypes were selected and included in this experiment, and given the following abbreviations: two wild tomato species *S. peruvianum* (SPE), and *S. habrochaites* (SHA); two accessions of the closest ancestor *S. pimpinellifolium* (LPI and SPI); and four modern *S. lycopersicum* tomato varieties Solarino (SOL), Trovanzo (TRO), Kivu (KIV) and Moneymaker (MON) (Supplementary material Table S2).

To facilitate the analysis of soils and tomato genotypes, they were grouped in soil types and tomato domestication degree. Hence, Ceiba and Limones native soils were catalogued as “native soil”, and Ceiba and Limones agricultural soils were named as “agricultural soil”. Also, tomatoes *S. peruvianum* and *S. habrochaites* were considered as “wild”, the two *S. pimpinellifolium* accessions were named as “intermediate”, and varieties of *S. lycopersicum* (Solarino, Trovanzo, Kivu and Moneymaker) were termed as “domesticated” tomatoes (Figure 10).




**Figure 10.** Soil origin (left panel) and genetic diversity of tomato genotypes (right panel) used in the experiment. Genetic diversity of the eight selected tomato genotypes was based on DArT genotyping data (17,855 SNP). The scale bar indicates the Jaccard distance to measure the dissimilarity between tomato genotypes based on SNP presence/absence. Hierarchical clustering was performed by unweighted pair group method with arithmetic mean (UPGMA). Tomato genotypes were grouped according to the domestication trajectory in wild, intermediate, and domesticated tomatoes. SHA: *S. habrochaites*; SPE: *S. peruvianum*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

Prior to the greenhouse experiment, the soil samples were conditioned during seven days by adding 20% (v/w) of sterile demi-water to the soil bags. The bags were then kept under greenhouse conditions in the shade (25 °C day; 18 °C night). After the conditioning phase, 5% (w/w) of each soil was mixed with 95% moistened sterile fine sand (0.4–0.8 mm), and 500 g of the mixture was transferred into polyethylene pots (10 × 10 × 11 cm).

Tomato seeds were surface sterilized by adding 5 ml of 80% ethanol and shaking for 2 min. The ethanol was then removed, and 5 ml of 1.5% (v/v) sodium hypochlorite was added. The seeds were shaken for 10 min, after which the solution was discarded. To remove any residual solution, seeds were rinsed five times with 5 ml of sterile demi-water, vortexing for 2 min during each cycle and discarding the water after each rinse. Following the sterilization, seeds were placed on a wet filter paper in a Petri dish containing 5 ml of sterile demi-water and incubated at 25 °C for three days. One pre-germinated tomato seed (radicle length ~1 cm) was sown in the center of the pot. Pots were covered with a plastic film for three days to maintain the humidity until the cotyledons emerged. The experiment consisted of eight tomato genotypes (SPE, SHA, SPI, LPI,

SOL, TRO, KIV, MON) and five soils (four Ecuadorian soils: CA, LA, CN, LN, and one Dutch tomato greenhouse soil GH), with a total of 239 pots [5 soils  $\times$  8 tomatoes  $\times$  5 replicates + 24 control (8 tomatoes in sand  $\times$  3 replicates) + 15 bulk soil (5 soil-sand mix  $\times$  3 replicates)] randomly distributed.

Plants were grown under greenhouse conditions set at 25 °C/18 °C ( $\pm$ 1 °C) day/night; with 16 h light and 60% relative humidity. The watering regime was adjusted to the plant's requirements during the growth period, starting with 5 ml of sterile demi-water per day and gradually increasing to 100 ml per day. Additionally, a 50% Hoagland nutrient solution was applied twice per week throughout the growth period (Supplementary material Table S3).

### ***Rhizosphere sampling***

Plants were sampled at their 6<sup>th</sup> true leaf stage (25–35 days after planting the germinated seeds). The shoots were cut and the root was carefully removed from the pot. Loosely attached soil was shaken off, leaving the tightly attached soil (i.e. rhizosphere soil). A 5 g sample of each root system and 5 g of each bulk soil sample were collected in 15 ml-tubes, immediately frozen in liquid nitrogen and stored at -80 °C until further processing.

Before DNA extraction, samples were defrosted at 4 °C and 5 ml of sterile demi-water was added and vortexed at maximum speed to remove the soil particles attached to the roots (rhizosphere). The roots were transferred to a new 15 ml-tube. Both rhizosphere soil suspension and frozen bulk soil samples were freeze-dried to remove the excess water. The roots were surface sterilized by vortexing with 5 ml of 75% ethanol at medium speed ( $\sim$ 2000 rpm) for 2 min. The ethanol solution was discarded and 5 ml of 1.5% (v/v) sodium hypochlorite were added. The roots were vortexed at medium speed for 5 min and after that, the hypochlorite solution was removed, followed by addition of 5 ml of sterile demi-water. Tubes were vortexed for 2 min, after which the water was carefully discarded. This washing process was repeated five times. Finally, the surface-sterilized roots were freeze dried to remove any remaining water.

### ***Soil and root DNA isolation and sequencing***

The rhizosphere soil samples were prepared by weighing 0.5 g of freeze-dried soil. For roots, 30 mg of freeze-dried roots were transferred into 2 ml-microtubes and the roots were homogenized with three metal beads ( $\varnothing$  1/8 inch) in the TissueLyser (Qiagen) at maximum speed (30 Hz, 1800 oscillations per minute) for 2 min. The resulting root powder was used for endosphere DNA isolation.

The Qiagen DNeasy® PowerSoil® Pro Kit was used to extract genomic DNA from soil and roots according to the manufacture's protocol. DNA samples were sent to Genome Québec (Canada) for amplicon library preparation and subsequent sequencing of the V3-V4 regions of the 16S rRNA gene using the universal bacterial primers 341F (CCTACGGGNGGCWGCAG) and 805R (GACTACHVGGGTATCTAATCC). Paired-end sequence reads (300 bp length) were generated using the Illumina NovaSeq-qX Plus platform. In addition, 24 rhizosphere samples corresponding to two genotypes (LPI and MON) grown in three soils (LA, LN and GH) representing four replicates were subjected to library preparation and shotgun sequencing by Novogene Co, Ltd. (Canada) to provide pair-end reads of 150 bp by Illumina NovaSeq 6000.

### ***Amplicon data analysis***

The compressed sequence reads contained in FASTQ format files were processed by the DADA2 v1.16.0 pipeline (Callahan et al., 2016) in RStudio environment (RStudio Team, 2020) to obtain the ASV abundance and taxonomy tables. The modeling of error rates associated with the sequencing process was adjusted to appropriately process NovaSeq data using the DADA2 pipeline, as suggested by Holland-Moritz (2021). The SILVA 16S ribosomal RNA gene reference database (v138) (Quast et al., 2013) was used for bacterial taxonomy assignment. ASVs present in control samples, i.e., plants grown in pure sand, were filtered out from all the samples before the analysis. The statistical analyses were performed using R software version 4.3.1 (R Core Team, 2023). Packages such as *vegan* (Oksanen et al., 2020), *phyloseq* (McMurdie & Holmes, 2013), *DESeq2* (Love et al., 2014) and *ggplot2* (Wickham, 2016) were used for alpha diversity, beta diversity (Bray–Curtis distance, PERMANOVA with 9,999 permutations), and differential abundance analyses, while the *tidyverse* package (Wickham et al., 2019) was used for formatting and visualization. Significant differences in the Shannon diversity index were determined by Kruskal-Wallis test and Wilcoxon post hoc test, while differences of beta diversity between tomato groups in pairwise comparisons were evaluated using the Adonis test. The abundance data were normalized by CSS (Cumulative Sum Scaling) for Principal Coordinates Analysis (PCoA) which was performed with the *cmdscale* function from the *vegan* package and the Bray–Curtis distance calculated previously. Differential abundance analyses were performed using the model “Soil\_class + Tom\_gen”, specified in the design formula of the *DESeq* function, where *Soil\_class* refers to the soil type, i.e. Native, Agricultural, or Greenhouse and *Tom\_gen* represents the tomato genotype (eight genotypes). Pairwise contrasts were tested between Native vs. Agricultural soil, and Agricultural vs. Greenhouse soil. To account for varying sequencing depths, size factors were estimated using a geometric mean approach from the counts data. ASVs resulted from differential abundance analysis were visualized in a heatmap using the R package *pheatmap* (Kolde, 2018). Clusters

were identified through hierarchical clustering, using correlation as the distance metric and Ward's method for the clustering algorithm.

### ***Metagenome data analysis***

To gain knowledge about the potential functionality of specific members of the rhizosphere tomato microbiome, bins were produced to be annotated further. First co-assembly of all samples was done using Megahit (Li et al., 2015). Read mapping against contigs was performed using Bowtie2 (Langmead & Salzberg, 2012) and binning was done using MaxBin2 (Wu et al., 2016) and Metabat2 (Kang et al., 2019). Results from both binning approaches were processed with DAS Tool to obtain refined bins (Sieber et al., 2018). Bins were taxonomically annotated using GTDB-tk (v2.4.0) (Chaumeil et al., 2020).

The quality of the resulting bins was assessed using CheckM (Parks et al., 2015). Afterwards, 11 out of 127 bins with completeness higher than 70% and contamination less than 10%, were considered representative metagenome-assembled genomes (MAGs) of the bacteria associated with tomato and soils, consistent with the amplicon results (Supplementary Table S5). These MAGs were submitted to the RAST server (Rapid Annotation using Subsystems Technology) (Aziz et al., 2008) for annotation of functional genes (Supplementary Table S6).

### **Acknowledgements**

We thank Dr. Sonia Zapata and Juan Mosquera (Universidad San Francisco de Quito–USFQ) for their support with the research permit procedures. We would also like to express our gratitude to Junnior Lalangui, José Luis Solano and Pablo Arévalo for their significant assistance with the collection of native and agricultural soils in Loja, Ecuador.

### **Author contributions**

SSF: methodology, investigation, data curation; writing - original draft, review & editing; VC: supervision, writing - review & editing; data curation; BOO: review & editing; LMAG: metagenomic data analysis; review & editing; NS: conceptualization; methodology; supervision; data curation; JMR: conceptualization; supervision; funding acquisition, writing - review & editing; PVTH: supervision; review & editing. All authors contributed critically to the drafts and gave final approval for publication.

## **Funding**

This research was supported by the Dutch Research Council Grant/Award Number: 024.004.014, the Ecuadorian National Secretary of Higher Education, Science, Technology and Innovation–SENESCYT with scholarship number: CZ07-000440-2018. Universidad San Francisco de Quito–USFQ supported this work through Chancellor Grant and COCIBA research funds to the HUBI project ID: 10093 entitled “Going back to the roots: Root microbiomes of tomato relatives”. The funders had no role in the study design, data collection, analysis, decision to publish, or preparation of the manuscript. This work was performed, in part, in collaboration with the MiCRop Consortium program, Harnessing the second genome of plants (grant number 024.004.014), funded by the Dutch Research Council (NWO/OCW), using its network and infrastructure.

## **Plant research permit**

This study was carried out under the Genetic Resource Permit N° MAE-DNB-CM-2018-0085, issued by the Ministry of Environment of Ecuador to USFQ.

## Supplementary material

### Supplementary Figures




**Figure S1.** Bacterial community structure of tomato genotypes grown in native, agricultural and greenhouse soils. Principal Coordinate Analysis (PCoA) of rhizosphere bacterial communities showing distinct separation between soils from center of origin and the center of production, as well as differentiation of bacterial communities between tomato rhizosphere and endosphere.




**Figure S2.** Shannon diversity index in tomato (a) rhizosphere and (b) endosphere grown in native, agricultural and greenhouse soils. Different letters above boxplots show significant difference (Wilcoxon test  $p < 0.05$ ) of Shannon diversity index between tomato genotypes ( $n = 5$ ). SPE: *S. peruvianum*; SHA: *S. habrochaïtes*; SPI and LPI: *S. pimpinellifolium*; and tomato varieties of *S. lycopersicum*: SOL: Solarino; TRO: Trovanzo; KIV: Kivu; MON: Moneymaker.

Supplementary Tables

Table S1. Physicochemical properties of soils employed in the experiment.

| Soil         | pH   | OM (g/ kg) | N (mg/ kg) | P (mg/ kg) | K (mg/ kg) | Ca (mg/ kg) | Mg (mg/ kg) | Fe (mg/ kg) | Mn (mg/ kg) | Cu (mg/ kg) | Zn (mg/ kg) | CIC (cmol/kg) | Sand (%) | Silt (%) | Clay (%) | Texture    |
|--------------|------|------------|------------|------------|------------|-------------|-------------|-------------|-------------|-------------|-------------|---------------|----------|----------|----------|------------|
| Ceiba_ Agr   | 7,46 | 13,10      | 700,00     | 19,10      | 50,80      | 2957,90     | 473,00      | 78,90       | 6,48        | 4,28        | 2,30        | 15,62         | 38       | 36       | 26       | Loam       |
| Ceiba_ Nat   | 7,29 | 15,10      | 800,00     | 29,60      | 62,60      | 3266,50     | 555,70      | 29,60       | 5,48        | 2,56        | 2,29        | 16,01         | 24       | 50       | 26       | Silt loam  |
| Limones_ Agr | 7,19 | 6,20       | 300,00     | 66,80      | 160,30     | 2154,30     | 340,50      | 56,80       | 6,07        | 3,67        | 3,29        | 9,24          | 52       | 32       | 16       | Sandy loam |
| Limones_ Nat | 7,83 | 2,90       | 100,00     | 47,70      | 215,10     | 2709,40     | 345,30      | 50,50       | 2,97        | 2,17        | 2,06        | 7,58          | 68       | 18       | 14       | Sandy loam |
| Green-house  | 7,10 | 24,00      | 1040,00    | 15,60      | 309,00     | 156,31      | 196,00      | 2,01        | 1,88        | 0,04        | 0,39        | 7,90          | 85       | 8        | 3        | Loamy sand |

Table S2. Tomato accessions used for DArT genotyping.

| Code | Species                                    | Accession number | Origin       | Passport                                                                                                                                                            |
|------|--------------------------------------------|------------------|--------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| SPE  | <i>Solanum peruvianum</i>                  | CGN23955         | Perú         | <a href="https://cgngenis.wur.nl/accessiondetails/CGN23955">https://cgngenis.wur.nl/accessiondetails/CGN23955</a>                                                   |
| SHA  | <i>Solanum habrochaites</i>                | CGN15792         | Ecuador      | <a href="https://cgngenis.wur.nl/accessiondetails/CGN15792">https://cgngenis.wur.nl/accessiondetails/CGN15792</a>                                                   |
| LPI  | <i>Solanum pimpinellifolium</i>            | CGN14498         | NA           | <a href="https://cgngenis.wur.nl/accessiondetails/CGN14498">https://cgngenis.wur.nl/accessiondetails/CGN14498</a>                                                   |
| SPI  | <i>Solanum pimpinellifolium</i>            | CGN23957         | Perú         | <a href="https://cgngenis.wur.nl/accessiondetails/CGN23957">https://cgngenis.wur.nl/accessiondetails/CGN23957</a>                                                   |
| MON  | <i>Solanum lycopersicum</i> cv Money-maker | CGN14330         | Netherlands  | <a href="https://cgngenis.wur.nl/accessiondetails/CGN14330">https://cgngenis.wur.nl/accessiondetails/CGN14330</a>                                                   |
| KIV  | <i>Solanum lycopersicum</i> cv Kivu        | NA               | © Rijk Zwaan | <a href="https://www.rijkzwaanusa.com/tomato/KIVU-RZ-F1-72-629-prdSL11187-ctgCrops">https://www.rijkzwaanusa.com/tomato/KIVU-RZ-F1-72-629-prdSL11187-ctgCrops</a> . |
| TRO  | <i>Solanum lycopersicum</i> cv Trovanzo    | NA               | © Rijk Zwaan | <a href="https://rijkzwaan.it/pomodoro/TROVANZO-RZ-F1-72-762-prdSL11225-ctgCrops">https://rijkzwaan.it/pomodoro/TROVANZO-RZ-F1-72-762-prdSL11225-ctgCrops</a> .     |
| SOL  | <i>Solanum lycopersicum</i> cv Solarino    | NA               | © Rijk Zwaan | <a href="https://www.rijkzwaan.de/tomate/SOLARINO-F1-72-150-prdSL11078-ctgCrops">https://www.rijkzwaan.de/tomate/SOLARINO-F1-72-150-prdSL11078-ctgCrops</a> .tomato |

**Table S3.** Hoagland solution composition.

| Hoagland solution stock                                                                         | M (g/mol) | Molarity | g/l Stock | 1 liter 1/2X |
|-------------------------------------------------------------------------------------------------|-----------|----------|-----------|--------------|
| <i>Macroelements:</i>                                                                           |           |          |           |              |
| Calcium nitrate $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$                            | 236.16    | 0.5 M    | 118.08    | 5 ml         |
| Potassium nitrate $\text{KNO}_3$                                                                | 101.11    | 1.0 M    | 101.11    | 2.5 ml       |
| Potassium phosphate $\text{KH}_2\text{PO}_4$                                                    | 136.09    | 1.0 M    | 136.09    | 0.5 ml       |
| Magnesium sulfate $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$                                     | 246.47    | 0.5 M    | 123.24    | 2 ml         |
| <i>Microelements:</i>                                                                           |           |          |           | 1 ml         |
| Boric acid $\text{H}_3\text{BO}_3$                                                              | 61.83     | 46.3 mM  | 2.86      |              |
| Manganese chloride $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$                                    | 197.91    | 9.1 mM   | 1.81      |              |
| Zinc sulfate $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$                                          | 287.54    | 0.77 mM  | 0.22      |              |
| Copper sulfate $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$                                        | 249.68    | 0.32 mM  | 0.08      |              |
| Sodium molybdate $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$                            | 241.95    | 0.52 mM  | 0.126     |              |
| EDTA ferric salt $\text{C}_{10}\text{H}_{12}\text{FeN}_2\text{NaO}_8 \cdot 3\text{H}_2\text{O}$ | 421.1     | 98.6 mM  | 41.52     |              |

*Note.* For preparing 1000 ml of Hoagland solution stock: Dissolve the components (g/l Stock) in 1000 ml of sterile demi water and store at room temperature in the shade. Stock solution of microelements can be prepared in 500 ml in an amber bottle or covered with aluminum foil to protect it against light.

For preparing 1000 ml of Hoagland 1/2X concentration: Autoclave 989 ml of demi water and add the volume of each Hoagland component by filtration (Whatman cellulose syringe filter 0,2  $\mu\text{m}$ ).

**Table S4.** Pairwise analysis by Adonis test in tomato rhizosphere and endosphere.

| Plant Compartment | Soil        | Pairs | R <sup>2</sup> | p.adj   | sig. |
|-------------------|-------------|-------|----------------|---------|------|
| Rhizosphere       | Ceiba_Agr   | W × I | 0,0653         | 0,0615  |      |
| Rhizosphere       | Ceiba_Agr   | W × D | 0,1366         | 0,00015 | *    |
| Rhizosphere       | Ceiba_Agr   | I × D | 0,1283         | 0,00015 | *    |
| Rhizosphere       | Ceiba_Nat   | W × I | 0,0633         | 0,0742  |      |
| Rhizosphere       | Ceiba_Nat   | W × D | 0,1818         | 0,00015 | *    |
| Rhizosphere       | Ceiba_Nat   | I × D | 0,1737         | 0,00015 | *    |
| Rhizosphere       | Limones_Agr | W × I | 0,0698         | 0,0442  | *    |
| Rhizosphere       | Limones_Agr | W × D | 0,1693         | 0,00015 | *    |
| Rhizosphere       | Limones_Agr | I × D | 0,1491         | 0,00015 | *    |
| Rhizosphere       | Limones_Nat | W × I | 0,0662         | 0,0605  |      |
| Rhizosphere       | Limones_Nat | W × D | 0,1554         | 0,00015 | *    |
| Rhizosphere       | Limones_Nat | I × D | 0,1799         | 0,00015 | *    |
| Rhizosphere       | Greenhouse  | W × I | 0,0608         | 0,1914  |      |
| Rhizosphere       | Greenhouse  | W × D | 0,2344         | 0,00015 | *    |
| Rhizosphere       | Greenhouse  | I × D | 0,2104         | 0,00015 | *    |
| Endosphere        | Ceiba_Agr   | W × I | 0,0558         | 0,1893  |      |
| Endosphere        | Ceiba_Agr   | W × D | 0,0509         | 0,0123  | *    |
| Endosphere        | Ceiba_Agr   | I × D | 0,0403         | 0,12    |      |
| Endosphere        | Ceiba_Nat   | W × I | 0,064          | 0,0506  |      |
| Endosphere        | Ceiba_Nat   | W × D | 0,0468         | 0,0306  | *    |
| Endosphere        | Ceiba_Nat   | I × D | 0,0485         | 0,0306  | *    |
| Endosphere        | Limones_Agr | W × I | 0,0585         | 0,0966  |      |
| Endosphere        | Limones_Agr | W × D | 0,0631         | 0,0021  | *    |
| Endosphere        | Limones_Agr | I × D | 0,0458         | 0,0461  | *    |
| Endosphere        | Limones_Nat | W × I | 0,0583         | 0,1749  |      |
| Endosphere        | Limones_Nat | W × D | 0,0506         | 0,0663  |      |
| Endosphere        | Limones_Nat | I × D | 0,0381         | 0,1749  |      |
| Endosphere        | Greenhouse  | W × I | 0,0542         | 0,3477  |      |
| Endosphere        | Greenhouse  | W × D | 0,0545         | 0,0108  | *    |
| Endosphere        | Greenhouse  | I × D | 0,0459         | 0,0402  | *    |

Table S5. High quality-bins assembled from rhizosphere tomato grown in different soils

| Bin_Id  | Marker_lineage               | Completeness | Contamination | Kingdom  | Phylum           | Class               | Order              | Family              | Genus             | Species |
|---------|------------------------------|--------------|---------------|----------|------------------|---------------------|--------------------|---------------------|-------------------|---------|
| bin.555 | p_Bacteroidetes (UID2605)    | 99.87        | 0.48          | Bacteria | Bacteroidota     | Bacteroidia         | Sphingobacteriales | Sphingobacteriaceae |                   |         |
| bin.732 | o_Cytophagales (UID2936)     | 99.7         | 1.07          | Bacteria | Bacteroidota     | Bacteroidia         | Cytophagales       | Spirosomaceae       | Emticia           |         |
| bin.204 | p_Bacteroidetes (UID2605)    | 99.52        | 1.9           | Bacteria | Bacteroidota     | Bacteroidia         | AKYH767            | b-17BO              |                   |         |
| bin.895 | p_Bacteroidetes (UID2591)    | 99.51        | 2.4           | Bacteria | Bacteroidota     | Bacteroidia         | Chitinophagales    | Chitinophagaceae    |                   |         |
| bin.601 | o_Cytophagales (UID2936)     | 99.03        | 5.7           | Bacteria | Bacteroidota     | Bacteroidia         | Cytophagales       | Cyclobacteriaceae   |                   |         |
| bin.795 | f_Xanthomonadaceae (UID4214) | 98.85        | 6.98          | Bacteria | Proteobacteria   | Gammaproteobacteria | Xanthomonadales    | Rhodanobacteraceae  | Dyella            |         |
| bin.952 | k_Bacteria (UID2569)         | 98.76        | 0             | Bacteria | Bacteroidota     | Bacteroidia         | Flavobacteriales   | Crocinitomiacaeae   |                   |         |
| bin.745 | o_Actinomycetales (UID1808)  | 98.54        | 3.73          | Bacteria | Actinobacteriota | Actinobacteria      | Mycobacteriales    | Mycobacteriaceae    | Fluviicola        |         |
| bin.518 | o_Sphingomonadales (UID3310) | 98.3         | 4.66          | Bacteria | Proteobacteria   | Alphaproteobacteria | Sphingomonadales   | Sphingomonadaceae   | Mycolicibacterium |         |
| bin.714 | k_Bacteria (UID203)          | 98.28        | 9.87          | Bacteria | Proteobacteria   | Gammaproteobacteria | Legionellales      | Legionellaceae      | Legionella        |         |
| bin.747 | k_Bacteria (UID3187)         | 98.16        | 0.89          | Bacteria | Bdellovibrionota | Bdellovibrionota    | Bdellovibrionales  | Bdellovibrionaceae  | Ga0074139         |         |
| bin.224 | o_Actinomycetales (UID2012)  | 97.82        | 8.06          | Bacteria | Actinobacteriota | Actinobacteria      | Mycobacteriales    | Pseudonocardiaceae  | Pseudonocardia    |         |
| bin.572 | p_Bacteroidetes (UID2605)    | 97.6         | 1.94          | Bacteria | Bacteroidota     | Bacteroidia         | Sphingobacteriales | Sphingobacteriaceae |                   |         |

**Table S5.** High quality-bins assembled from rhizosphere tomato grown in different soils (*continued*)

| Bin_Id  | Marker_lineage                  | Completeness | Contamination | Kingdom  | Phylum            | Class               | Order                | Family                | Genus            | Species         |
|---------|---------------------------------|--------------|---------------|----------|-------------------|---------------------|----------------------|-----------------------|------------------|-----------------|
| bin.53  | k_Bacteria (UID2569)            | 97.57        | 1.4           | Bacteria | Bacteroidota      | Bacteroidia         | Flavobacteriales     | Crocinitomiacaceae    | Fluviicola       |                 |
| bin.643 | p_Cyanobacteria (UID2182)       | 97.56        | 1.66          | Bacteria | Cyanobacteria     | Cyanobacteriia      | Elainellales         | Elainellaceae         | Lepolyngbya_A    |                 |
| bin.761 | o_Sphingomonadales (UID3310)    | 97.4         | 6.14          | Bacteria | Proteobacteria    | Alphaproteobacteria | Sphingomonadales     | Sphingomonadaceae     | Sphingopyxis     |                 |
| bin.929 | o_Actinomycetales (UID1696)     | 97.25        | 2.54          | Bacteria | Actinobacteriota  | Actinobacteriia     | Mycobacteriales      | Mycobacteriaceae      |                  |                 |
| bin.918 | p_Bacteroidetes (UID2605)       | 96.87        | 1.22          | Bacteria | Bacteroidota      | Bacteroidia         | Sphingobacteriales   | Sphingobacteriaceae   | Mucilaginibacter |                 |
| bin.215 | k_Bacteria (UID3187)            | 96.37        | 2.68          | Bacteria | Bdellovibrionota  | Bdellovibrionia     | Bdellovibrionales    | Bdellovibrionaceae    |                  | 21-14-0-10-47-8 |
| bin.296 | c_Deltaproteobacteria (UID3216) | 96.13        | 6.45          | Bacteria | Myxococcota       | Polyangia           | Polyangiales         | Sandaracinaceae       |                  |                 |
| bin.111 | p_Actinobacteria (UID1454)      | 96.07        | 1.95          | Bacteria | Actinobacteriota  | Thermoleophilae     | Solirubrobacteriales | Solirubrobacteriaceae |                  |                 |
| bin.690 | k_Bacteria (UID2982)            | 95.95        | 5.41          | Bacteria | Verrucomicrobiota | Verrucomicrobiae    | Pedospherales        | AV2                   |                  |                 |
| bin.890 | c_Gammaproteobacteria (UID4202) | 95.86        | 6.02          | Bacteria | Proteobacteria    | Gammaproteobacteria | Nevskiales           | Nevskiaceae           | Fontimonas       |                 |
| bin.721 | o_Pseudomonadales (UID4488)     | 95.73        | 5.18          | Bacteria | Proteobacteria    | Gammaproteobacteria | Pseudomonadales      | Pseudomonadaceae      | Pseudomonas_E    |                 |
| bin.485 | p_Bacteroidetes (UID2591)       | 95.71        | 7.51          | Bacteria | Bacteroidota      | Bacteroidia         | Chitinophagales      | Chitinophagaceae      | Taibaella_B      |                 |
| bin.188 | o_Burkholderiales (UID4000)     | 95.53        | 7.18          | Bacteria | Proteobacteria    | Gammaproteobacteria | Burkholderiales      | Burkholderiaceae      | Hermiimonas      |                 |

**Table S5.** High quality-bins assembled from rhizosphere tomato grown in different soils (*continued*)

| Bin_Id  | Marker_lineage                  | Completeness | Contamination | Kingdom  | Phylum           | Class                | Order               | Family               | Genus           | Species                   |
|---------|---------------------------------|--------------|---------------|----------|------------------|----------------------|---------------------|----------------------|-----------------|---------------------------|
| bin.591 | k_Bacteria (UID3187)            | 95.5         | 1.8           | Bacteria | Bdellovibrionota | Bdellovibrionia      | Bdellovibrionales   | Bdellovibrionaceae   | Bdellovibrion_A |                           |
| bin.844 | o_Rhizobiales (UID3450)         | 95.41        | 2.21          | Bacteria | Proteobacteria   | Alphaproteobacteria  | Rhizobiales         | Rhizobiaceae         | Chelativorans   | Chelativorans sp000014245 |
| bin.257 | k_Bacteria (UID3187)            | 95.39        | 5.73          | Bacteria | Acidobacteriota  | Thermoanaerobaculia  | Gp7-AA8             | Gp7-AA8              | QHVT01          |                           |
| bin.324 | k_Bacteria (UID2982)            | 95.35        | 2.86          | Bacteria | Verrucomicrobota | Verrucomicrobiae     | Verrucomicrobiales  | Akkermansiaceae      | Haloferula      |                           |
| bin.633 | k_Bacteria (UID2569)            | 95.26        | 1.62          | Bacteria | Bacteroidota     | Bacteroidia          | Flavobacteriales    | Crocinitomiacaeae    | Fluviicola      |                           |
| bin.221 | k_Bacteria (UID3187)            | 94.74        | 1.26          | Bacteria | Bdellovibrionota | Bdellovibrionia      | Bdellovibrionales   | Bdellovibrionaceae   | 21-14-0-10-47-8 |                           |
| bin.441 | c_Deltaproteobacteria (UID3216) | 94.62        | 3.39          | Bacteria | Myxococota       | Myxococcia           | Myxococcales        | Myxococcaceae        |                 |                           |
| bin.177 | k_Bacteria (UID3187)            | 94.12        | 0.89          | Bacteria | Bdellovibrionota | Bdellovibrionia      | Bdellovibrionales   | Bdellovibrionaceae   | 21-14-0-10-47-8 |                           |
| bin.329 | o_Sphingomonadales (UID3310)    | 93.95        | 6.7           | Bacteria | Proteobacteria   | Alphaproteobacteria  | Sphingomonadales    | Sphingomonadaceae    | Sphingobium     |                           |
| bin.117 | k_Bacteria (UID3187)            | 93.9         | 2.41          | Bacteria | Bdellovibrionota | Bacteriovoracia      | Bacteriovorales     | Bacteriovoracaceae   | Bacteriovorax   |                           |
| bin.350 | k_Bacteria (UID3187)            | 93.87        | 9.54          | Bacteria | Acidobacteriota  | Vicinamibacteria     | Vicinamibacteriales | Vicinamibacteriaceae | Luteitalea      |                           |
| bin.838 | c_Betaproteobacteria (UID3959)  | 93.69        | 5.35          | Bacteria | Proteobacteria   | Gammaaproteobacteria | Burkholderiales     | SG8-41               |                 |                           |
| bin.664 | o_Burkholderiales (UID4002)     | 93.57        | 4.3           | Bacteria | Proteobacteria   | Gammaaproteobacteria | Burkholderiales     | Burkholderiaceae     | Hermiinimonas   |                           |

**Table S5.** High quality-bins assembled from rhizosphere tomato grown in different soils (*continued*)

| Bin_Id  | Marker_lineage                  | Completeness | Contamination | Kingdom  | Phylum           | Class               | Order           | Family           | Genus               | Species                      |
|---------|---------------------------------|--------------|---------------|----------|------------------|---------------------|-----------------|------------------|---------------------|------------------------------|
| bin.273 | k_Bacteria (UID2982)            | 93.34        | 5.14          | Bacteria | Verucomicrobiota | Verucomicrobiae     | Opitutales      | Opitutaceae      | Didemnitutus        |                              |
| bin.737 | c_Deltaproteobacteria (UID3216) | 93.12        | 3.25          | Bacteria | Myxococota       | Polyangia           | Polyangiales    |                  |                     |                              |
| bin.319 | c_Gammaproteobacteria (UID4444) | 92.98        | 2.39          | Bacteria | Proteobacteria   | Gammaproteobacteria | Pseudomonadales | Cellvibrionaceae | Cellvibrio          |                              |
| bin.464 | p_Bacteroidetes (UID2591)       | 92.61        | 6.03          | Bacteria | Bacteroidota     | Bacteroidia         | Chitinophagales | Chitinophagaceae | Taibaiella_B        |                              |
| bin.515 | o_Burkholderiales (UID4000)     | 92.5         | 5.06          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales | Burkholderiaceae | Rhizobacter         |                              |
| bin.302 | p_Bacteroidetes (UID2605)       | 92.49        | 1.67          | Bacteria | Bacteroidota     | Bacteroidia         | AKYH767         | b-17BO           |                     |                              |
| bin.861 | g_Burkholderia (UID4006)        | 92.23        | 0.31          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales | Burkholderiaceae | Paraburkholderia    | Paraburkholderia sp000148685 |
| bin.427 | f_Micrococaceae (UID1623)       | 92.16        | 4.38          | Bacteria | Actinobacteriota | Actinobacteria      | Actinomycetales | Micrococaceae    | Sinomonas           |                              |
| bin.647 | c_Gammaproteobacteria (UID4266) | 92.16        | 1.75          | Bacteria | Proteobacteria   | Gammaproteobacteria | Legionellales   | Legionellaceae   | Legionella_C        |                              |
| bin.606 | o_Burkholderiales (UID4000)     | 91.79        | 2.29          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales | Burkholderiaceae | Novitherbaspirillum |                              |
| bin.607 | c_Betaproteobacteria (UID3959)  | 91.71        | 5.08          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales | SG8-41           | PLOWO2-02-64-14     |                              |
| bin.7   | p_Bacteroidetes (UID2591)       | 91.71        | 3.2           | Bacteria | Bacteroidota     | Bacteroidia         | Chitinophagales | Chitinophagaceae | Flavisolibacter     |                              |
| bin.124 | o_Burkholderiales (UID4000)     | 91.49        | 6.51          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales | Burkholderiaceae | Novitherbaspirillum |                              |

**Table S5.** High quality-bins assembled from rhizosphere tomato grown in different soils (*continued*)

| Bin_Id  | Marker_lineage                  | Completeness | Contamination | Kingdom  | Phylum         | Class                | Order           | Family              | Genus              | Species                    |
|---------|---------------------------------|--------------|---------------|----------|----------------|----------------------|-----------------|---------------------|--------------------|----------------------------|
| bin.868 | p_Proteobacteria (UID3887)      | 91.48        | 8.95          | Bacteria | Proteobacteria | Gamma proteobacteria | Burkholderiales | Burkholderiaceae    | Lautropia          |                            |
| bin.373 | c_Alphaproteobacteria (UID3422) | 90.57        | 3.32          | Bacteria | Proteobacteria | Alphaproteobacteria  | Caulobacterales | Caulobacteriaceae   | Asticcacaulis      |                            |
| bin.548 | o_Burkholderiales (UID4000)     | 90.38        | 4.35          | Bacteria | Proteobacteria | Gamma proteobacteria | Burkholderiales | Burkholderiaceae    | Noviherbaspirillum |                            |
| bin.915 | c_Gammaproteobacteria (UID4445) | 90.04        | 5.96          | Bacteria | Proteobacteria | Gamma proteobacteria | Pseudomonadales | Pseudomonadaceae    | Pseudomonas_A      | Pseudomonas_A stutzeri_AI  |
| bin.808 | f_Micrococaceae (UID1623)       | 89.55        | 2.18          | Bacteria | Actinobacteria | Actinobacteria       | Actinomycetales | Actinomycetaceae    | Arthrobacter_D     | Arthrobacter_D sp001428435 |
| bin.472 | o_Burkholderiales (UID4002)     | 89.5         | 6.46          | Bacteria | Proteobacteria | Gamma proteobacteria | Burkholderiales | Burkholderiaceae    | Noviherbaspirillum |                            |
| bin.96  | f_Bacillaceae (UID830)          | 89.34        | 4.14          | Bacteria | Firmicutes     | Bacilli              | Bacillales      | Anoxybacillaceae    | Parageobacillus    |                            |
| bin.494 | g_Streptomyces (UID2052)        | 89.32        | 2.72          | Bacteria | Actinobacteria | Actinobacteria       | Streptomyces    | Streptomycetaceae   | Streptomyces       |                            |
| bin.734 | o_Rhizobiales (UID3450)         | 89.14        | 2.98          | Bacteria | Proteobacteria | Alphaproteobacteria  | Rhizobiales     | Rhizobiaceae        | UBA1059            |                            |
| bin.716 | c_Gammaproteobacteria (UID4266) | 89.1         | 0.81          | Bacteria | Proteobacteria | Gamma proteobacteria | Legionellales   | Legionellaceae      | Legionella         | Herbaspirillum aquaticum   |
| bin.610 | o_Burkholderiales (UID4002)     | 88.51        | 5.86          | Bacteria | Proteobacteria | Gamma proteobacteria | Burkholderiales | Burkholderiaceae    | Herbaspirillum     |                            |
| bin.827 | c_Betaproteobacteria (UID3888)  | 88.47        | 8.33          | Bacteria | Proteobacteria | Gamma proteobacteria | Burkholderiales | Methylobacteriaceae |                    |                            |
| bin.577 | c_Alphaproteobacteria (UID3422) | 88.27        | 3.8           | Bacteria | Proteobacteria | Alphaproteobacteria  | Caulobacterales | Caulobacteriaceae   | Asticcacaulis      |                            |

**Table S5.** High quality-bins assembled from rhizosphere tomato grown in different soils (*continued*)

| Bin_Id  | Marker_lineage                 | Completeness | Contamination | Kingdom  | Phylum            | Class                | Order              | Family              | Genus              | Species                     |
|---------|--------------------------------|--------------|---------------|----------|-------------------|----------------------|--------------------|---------------------|--------------------|-----------------------------|
| bin.628 | k_Bacteria (UID3187)           | 88.24        | 3.3           | Bacteria | Bdellovibrionota  | Bdellovibrionia      | Bdellovibrionales  | Bdellovibrionaceae  | Ga0074137          |                             |
| bin.499 | o_Burkholderiales (UID4000)    | 87.98        | 4.95          | Bacteria | Proteobacteria    | Gamma proteobacteria | Burkholderiales    | Burkholderiaceae    | Noviherbaspirillum |                             |
| bin.505 | k_Bacteria (UID3187)           | 87.89        | 5.13          | Bacteria | Acidobacteriota   | Blastocatellia       | Pyrimonadales      | Pyrimonadaceae      | OLB17              |                             |
| bin.248 | o_Burkholderiales (UID4001)    | 86.66        | 4.57          | Bacteria | Proteobacteria    | Gamma proteobacteria | Burkholderiales    | Burkholderiaceae    | Paraburkholderia_E |                             |
| bin.280 | k_Bacteria (UID2982)           | 86.36        | 1.5           | Bacteria | Verrucomicrobiota | Verrucomicrobiae     | Verrucomicrobiales | Akkermansiaceae     | Haloferula         |                             |
| bin.445 | c_Betaproteobacteria (UID3888) | 86.09        | 3.75          | Bacteria | Proteobacteria    | Gamma proteobacteria | Burkholderiales    | Methylobacteriaceae |                    |                             |
| bin.783 | p_Bacteroidetes (UID2591)      | 85.47        | 2.02          | Bacteria | Bacteroidota      | Bacteroidia          | Chitinophagales    | Chitinophagaceae    | Chitinophaga       |                             |
| bin.113 | o_Rhizobiales (UID3449)        | 85.45        | 4.99          | Bacteria | Proteobacteria    | Alphaproteobacteria  | Rhizobiales        | Rhizobiaceae        | Pseudorhizobium    |                             |
| bin.8   | k_Bacteria (UID2982)           | 84.5         | 1.36          | Bacteria | Verrucomicrobiota | Verrucomicrobiae     | Verrucomicrobiales | Akkermansiaceae     | Haloferula         |                             |
| bin.777 | k_Bacteria (UID203)            | 84.48        | 6.03          | Bacteria | Proteobacteria    | Alphaproteobacteria  | Rhizobiales        | Rhizobiaceae        | Shinella           |                             |
| bin.898 | p_Bacteroidetes (UID2591)      | 84.44        | 2.4           | Bacteria | Bacteroidota      | Bacteroidia          | Chitinophagales    | Chitinophagaceae    | Niastella          |                             |
| bin.625 | f_Micrococcaceae (UID1623)     | 84.19        | 3.11          | Bacteria | Actinobacteriota  | Actinobacteria       | Actinomycetales    | Micrococcaceae      | Arthrobacter_D     | Arthrobacter_D subterraneus |
| bin.780 | o_Actinomycetales (UID1663)    | 84.06        | 6.49          | Bacteria | Actinobacteriota  | Actinobacteria       | Actinomycetales    | Dermatophilaceae    | Pedococcus         |                             |

Table S5. High quality-bins assembled from rhizosphere tomato grown in different soils (continued)

| Bin_Id  | Marker_lineage                  | Completeness | Contamination | Kingdom  | Phylum            | Class                | Order              | Family            | Genus          | Species                  |
|---------|---------------------------------|--------------|---------------|----------|-------------------|----------------------|--------------------|-------------------|----------------|--------------------------|
| bin.854 | g_Ensifer (UID3566)             | 84           | 4.67          | Bacteria | Proteobacteria    | Alphaproteobacteria  | Rhizobiales        | Rhizobiaceae      | Sinorhizobium  | Sinorhizobium meliloti_A |
| bin.651 | c_Alphaproteobacteria (UID3305) | 83.77        | 7.47          | Bacteria | Proteobacteria    | Alphaproteobacteria  | Sphingomonadales   | Sphingomonadaceae | Sphingomonas_A |                          |
| bin.225 | o_Burkholderiales (UID4000)     | 83.32        | 8.76          | Bacteria | Proteobacteria    | Gammaaproteobacteria | Burkholderiales    | Burkholderiaceae  | Rhizobacter    |                          |
| bin.725 | k_Bacteria (UID2565)            | 83.26        | 1.7           | Bacteria | Planctomycetota   | Physcisphaerae       | Physcisphaerales   | SM1A02            |                |                          |
| bin.537 | k_Bacteria (UID2570)            | 83.07        | 3.67          | Bacteria | Bacteroidota      | Rhodothermiales      | Rhodothermiales    | Rhodothermaceae   |                |                          |
| bin.434 | c_Alphaproteobacteria (UID3305) | 82.63        | 5.25          | Bacteria | Proteobacteria    | Alphaproteobacteria  | Sphingomonadales   | Sphingomonadaceae | Sphingomonas_A |                          |
| bin.581 | o_Cytophagales (UID2936)        | 82.62        | 1.99          | Bacteria | Bacteroidota      | Bacteroidia          | Cytophagales       | Cytophagaceae     |                |                          |
| bin.95  | p_Cyanobacteria (UID2182)       | 82.23        | 0.79          | Bacteria | Cyanobacteria     | Cyanobacteria        | Neosynechococcales |                   |                |                          |
| bin.383 | c_Betaproteobacteria (UID3959)  | 82.01        | 1.03          | Bacteria | Proteobacteria    | Gammaaproteobacteria | Burkholderiales    | Nitrosomonadaceae | Nitrospira     |                          |
| bin.402 | k_Bacteria (UID1452)            | 81.91        | 3.72          | Bacteria | Chloroflexota     | Chloroflexia         | 54-19              |                   |                |                          |
| bin.147 | k_Bacteria (UID2982)            | 81.74        | 7.77          | Bacteria | Verrucomicrobiota | Verrucomicrobiae     | Opitutales         | Opitutaceae       | IMCC26134_A    |                          |
| bin.500 | o_Rhizobiales (UID3447)         | 81.47        | 4.95          | Bacteria | Proteobacteria    | Alphaproteobacteria  | Rhizobiales        | Andersenellaceae  | QKVK01         |                          |
| bin.713 | k_Bacteria (UID3187)            | 81.41        | 2.61          | Bacteria | Acidobacteriota   | Blastocatellia       | Pyrimonadales      | Pyrimonadaceae    | OLB17          |                          |

Table S5. High quality-bins assembled from rhizosphere tomato grown in different soils (continued)

| Bin_Id  | Marker_lineage                  | Completeness | Contamination | Kingdom  | Phylum          | Class               | Order               | Family              | Genus              | Species |
|---------|---------------------------------|--------------|---------------|----------|-----------------|---------------------|---------------------|---------------------|--------------------|---------|
| bin.634 | k_Bacteria (UID203)             | 81.4         | 3.45          | Bacteria | Patescibacteria | Paceibacteria       | UBA9983_A           | UBA2163             | C7867-001          |         |
| bin.14  | p_Euryarchaeota (UID3)          | 81.36        | 2.67          | NA       | NA              | NA                  | NA                  | NA                  | NA                 | NA      |
| bin.708 | k_Bacteria (UID2495)            | 81.29        | 8.79          | Bacteria | Gemmatimonadota | Gemmatimonadetes    | Gemmatimonadales    | Gemmatimonadaceae   |                    |         |
| bin.585 | p_Cyanobacteria (UID2182)       | 81.24        | 1.57          | Bacteria | Cyanobacteria   | Cyanobacteria       |                     |                     |                    |         |
| bin.35  | c_Alphaproteobacteria (UID3305) | 81.05        | 6.81          | Bacteria | Proteobacteria  | Alphaproteobacteria | Sphingomonadales    | Sphingomonadaceae   | Sphingomonas_F     |         |
| bin.202 | k_Bacteria (UID203)             | 80.88        | 5.33          | Bacteria | Patescibacteria | Saccharimonadia     | Saccharimonadales   | UBA4665             |                    |         |
| bin.843 | p_Actinobacteria (UID1454)      | 80.85        | 7.98          | Bacteria | Actinobacteria  | Thermoleophilina    | Solirubrobacterales | Solirubrobacterales |                    |         |
| bin.560 | o_Burkholderiales (UID4001)     | 80.7         | 0.91          | Bacteria | Proteobacteria  | Gammaproteobacteria | Burkholderiales     | Burkholderiaceae    | Paraburkholderia_E |         |
| bin.602 | p_Euryarchaeota (UID3)          | 80.49        | 3.64          | NA       | NA              | NA                  | NA                  | NA                  | NA                 | NA      |
| bin.84  | p_Actinobacteria (UID1454)      | 79.31        | 5.03          | Bacteria | Actinobacteria  | Thermoleophilina    | 20CM-4-69-9         |                     |                    |         |
| bin.759 | k_Bacteria (UID203)             | 79.28        | 0             | Bacteria | Patescibacteria | Paceibacteria       | UBA9983_A           | UBA1006             |                    |         |
| bin.462 | o_Sphingomonadales (UID3310)    | 79.04        | 6             | Bacteria | Proteobacteria  | Alphaproteobacteria | Sphingomonadales    | Sphingomonadaceae   | Sphingomonas       |         |
| bin.549 | k_Bacteria (UID1453)            | 78.89        | 4.91          | Bacteria | Cyanobacteria   | Sericytochromatia   |                     |                     |                    |         |

**Table S5.** High quality-bins assembled from rhizosphere tomato grown in different soils (*continued*)

| Bin_Id  | Marker_lineage                    | Completeness | Contamination | Kingdom  | Phylum            | Class               | Order              | Family             | Genus           | Species |
|---------|-----------------------------------|--------------|---------------|----------|-------------------|---------------------|--------------------|--------------------|-----------------|---------|
| bin.688 | k_Bacteria<br>(UID2495)           | 78.61        | 3.3           | Bacteria | Gemmatimonadota   | Gemmatimonadetes    | Gemmatimonadales   | GW/C2-71-9         |                 |         |
| bin.449 | k_Bacteria<br>(UID203)            | 77.01        | 1.72          | Bacteria | Paenibacteriota   | Paenibacteriae      | UBA9983_A          | UBA2163            | C7867-001       |         |
| bin.416 | k_Bacteria<br>(UID203)            | 76.72        | 0.86          | Bacteria | Bacteroidota      | Bacteroidia         | AKYH767            | Palsa-965          | GCA-2737665     |         |
| bin.604 | k_Bacteria<br>(UID3187)           | 75.33        | 2.94          | Bacteria | Bdellovibrionota  | Bdellovibrionia     | Bdellovibrionales  | Bdellovibrionaceae | 21-14-0-10-47-8 |         |
| bin.180 | k_Bacteria<br>(UID2982)           | 75.1         | 2.7           | Bacteria | Verrucomicrobiota | Verrucomicrobiae    | Chthoniobacterales | Terrimicrobiaceae  |                 |         |
| bin.355 | o_Burkholderiales<br>(UID4000)    | 74.96        | 7.97          | Bacteria | Proteobacteriota  | Gammaproteobacteria | Burkholderiales    | Burkholderiaceae   | Vitreoscilla_A  |         |
| bin.419 | k_Bacteria<br>(UID3187)           | 74.6         | 4.35          | Bacteria | Bdellovibrionota  | Bdellovibrionia     | Bdellovibrionales  | Bdellovibrionaceae |                 |         |
| bin.640 | k_Bacteria<br>(UID203)            | 74.57        | 9.48          | Bacteria | Actinobacteriota  | Actinobacteria      | Frankiales         | Frankiaceae        |                 |         |
| bin.769 | k_Bacteria<br>(UID203)            | 74.22        | 7.99          | Bacteria | Gemmatimonadota   | Gemmatimonadetes    | Gemmatimonadales   |                    |                 |         |
| bin.276 | k_Bacteria<br>(UID2982)           | 74.19        | 0.04          | Bacteria | Verrucomicrobiota | Verrucomicrobiae    | Pedospirales       |                    |                 |         |
| bin.652 | k_Bacteria<br>(UID203)            | 74.14        | 6.9           | Bacteria | Proteobacteriota  | Gammaproteobacteria | Xanthomonadales    | Rhodanobacteraceae | Dyella          |         |
| bin.368 | c_Betaproteobacteria<br>(UID3888) | 73.87        | 5.85          | Bacteria | Proteobacteriota  | Gammaproteobacteria | Burkholderiales    | Leciaceae          | Lecia           |         |
| bin.934 | k_Bacteria<br>(UID2495)           | 73.08        | 6.66          | Bacteria | Gemmatimonadota   | Gemmatimonadetes    | Gemmatimonadales   | GW/C2-71-9         |                 |         |

Table S5. High quality-bins assembled from rhizosphere tomato grown in different soils (continued)

| Bin_Id  | Marker_lineage              | Completeness | Contamination | Kingdom  | Phylum           | Class               | Order             | Family             | Genus           | Species |
|---------|-----------------------------|--------------|---------------|----------|------------------|---------------------|-------------------|--------------------|-----------------|---------|
| bin.770 | o_Rhizobiales<br>(UID3654)  | 72.31        | 3.92          | Bacteria | Proteobacteria   | Alphaproteobacteria | Rhizobiales       | Beijerinckiaceae   | Microvinga      |         |
| bin.110 | k_Bacteria<br>(UID203)      | 71.9         | 2.59          | Bacteria | Bacteroidota     | Bacteroidia         | AKYH767           | Palsa-965          | GCA-2737665     |         |
| bin.209 | k_Bacteria<br>(UID203)      | 71.9         | 9.48          | Bacteria | Proteobacteria   | Gammaproteobacteria | Xanthomonadales   | Rhodanobacteraceae | Dyella          |         |
| bin.803 | k_Bacteria<br>(UID2495)     | 71.52        | 0             | Bacteria | Patiscibacteria  | Paceibacteria       | UBA9983_A         | UBA5272            | UBA11704        |         |
| bin.163 | k_Bacteria<br>(UID3187)     | 70.86        | 4.97          | Bacteria | Bdellovibrionota | Bdellovibrionia     | Bdellovibrionales | UBA1609            |                 |         |
| bin.125 | k_Bacteria<br>(UID203)      | 70.33        | 1.75          | Bacteria | Bdellovibrionota | Bdellovibrionia     | Bdellovibrionales | Bdellovibrionaceae | UBA2316         |         |
| bin.297 | o_Cytophagales<br>(UID2936) | 70.2         | 3.92          | Bacteria | Bacteroidota     | Bacteroidia         | Cytophagales      | Cytophagaceae      |                 |         |
| bin.437 | k_Bacteria<br>(UID203)      | 70.19        | 1.72          | Bacteria | Patiscibacteria  | Paceibacteria       | UBA9983_A         | UBA2163            | 1-14-0-10-47-16 |         |
| bin.469 | k_Bacteria<br>(UID203)      | 70.17        | 3.45          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales   | Burkholderiaceae   | Vitreoscilla_A  |         |
| bin.848 | k_Bacteria<br>(UID203)      | 70           | 1.72          | Bacteria | Proteobacteria   | Gammaproteobacteria | Burkholderiales   | Burkholderiaceae   | Roseateles      |         |

**Table S6.** Annotated protein encoding genes by RAST server using the SEED Subsystem database from bacterial assembled bins from different tomato genotypes grown in different soils.

| bin_name             | Category            | Subcategory                          | Subsystem                            | Role                                                           | Features                      |
|----------------------|---------------------|--------------------------------------|--------------------------------------|----------------------------------------------------------------|-------------------------------|
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrifying reductase gene clusters | Nitrous oxide reductase maturation protein NosF (ATPase)       | fig 66666666.1425568.peg.2979 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrifying reductase gene clusters | Copper-containing nitrite reductase (EC 1.7.2.1)               | fig 66666666.1425568.peg.2169 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrifying reductase gene clusters | Nitrous oxide reductase maturation trans-membrane protein NosY | fig 66666666.1425568.peg.2978 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrifying reductase gene clusters | Nitrous oxide reductase maturation protein NosD                | fig 66666666.1425568.peg.2980 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrifying reductase gene clusters | Nitrous-oxide reductase (EC 1.7.99.6)                          | fig 66666666.1425568.peg.2982 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrification                      | Copper-containing nitrite reductase (EC 1.7.2.1)               | fig 66666666.1425568.peg.2169 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrification                      | Nitrous oxide reductase maturation protein NosF (ATPase)       | fig 66666666.1425568.peg.2979 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrification                      | Nitrous oxide reductase maturation protein NosD                | fig 66666666.1425568.peg.2980 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrification                      | Nitrous-oxide reductase (EC 1.7.99.6)                          | fig 66666666.1425568.peg.2982 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrification                      | Nitrous oxide reductase maturation trans-membrane protein NosY | fig 66666666.1425568.peg.2978 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Denitrification                      | Denitrification                      | Nitric-oxide reductase (EC 1.7.99.7), quinol-dependent         | fig 66666666.1425568.peg.1512 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Nitrogen Metabolism - no subcategory | Ammonia assimilation                 | Glutamine synthetase type III, GlnN (EC 6.3.1.2)               | fig 66666666.1425568.peg.4262 |
| Chitinophaga_bin/783 | Nitrogen Metabolism | Nitrogen Metabolism - no subcategory | Ammonia assimilation                 | Glutamate synthase [NADPH] large chain (EC 1.4.1.13)           | fig 66666666.1425568.peg.1009 |

**Table S6.** Annotated protein encoding genes by RAST server using the SEED Subsystem database from bacterial assembled bins from different tomato genotypes grown in different soils. (*continued*)

| bin_name            | Category             | Subcategory                           | Subsystem                            | Role                                                               | Features                                                       |
|---------------------|----------------------|---------------------------------------|--------------------------------------|--------------------------------------------------------------------|----------------------------------------------------------------|
| Chitinophaga_bin783 | Nitrogen Metabolism  | Nitrogen Metabolism - no subcategory  | Ammonia assimilation                 | Glutamate synthase [NADPH] small chain (EC 1.4.1.13)               | fig 66666666.1425568.peg.1008                                  |
| Chitinophaga_bin783 | Nitrogen Metabolism  | Nitrogen Metabolism - no subcategory  | Ammonia assimilation                 | Ammonium transporter                                               | fig 66666666.1425568.peg.4388                                  |
| Chitinophaga_bin783 | Nitrogen Metabolism  | Nitrogen Metabolism - no subcategory  | Ammonia assimilation                 | Ferredoxin-dependent glutamate synthase (EC 1.4.7.1)               | fig 66666666.1425568.peg.242                                   |
| Chitinophaga_bin783 | Nitrogen Metabolism  | Nitrogen Metabolism - no subcategory  | Nitrosative stress                   | Nitric-oxide reductase (EC 1.7.99.7), quinol-dependent             | fig 66666666.1425568.peg.1512                                  |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | FKBP-type peptidyl-prolyl cis-trans isomerase SlyD (EC 5.2.1.8)    | fig 66666666.1425568.peg.838                                   |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | Potassium-transporting ATPase C chain (EC 3.6.3.12) (TC 3.A.3.7.1) | fig 66666666.1425568.peg.2163                                  |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | Potassium efflux system KefA protein                               | fig 66666666.1425568.peg.3042                                  |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | Potassium-transporting ATPase B chain (EC 3.6.3.12) (TC 3.A.3.7.1) | fig 66666666.1425568.peg.2162                                  |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | Potassium-transporting ATPase A chain (EC 3.6.3.12) (TC 3.A.3.7.1) | fig 66666666.1425568.peg.1077                                  |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | Large-conductance mechanosensitive channel                         | fig 66666666.1425568.peg.1787                                  |
| Chitinophaga_bin783 | Potassium metabolism | Potassium metabolism - no subcategory | Potassium homeostasis                | POTASSIUM/PROTON ANTIPORTER                                        | fig 66666666.1425568.peg.398                                   |
| Chitinophaga_bin783 | Protein Metabolism   | Protein degradation                   | Serine endopeptidase (EC 3.4.21.-)   | ROSB                                                               | fig 66666666.1425568.peg.446,<br>fig 66666666.1425568.peg.1370 |
| Chitinophaga_bin783 | Nitrogen Metabolism  | Denitrification                       | Denitrifying reductase gene clusters | Nitrous oxide reductase maturation protein NosF (ATPase)           | fig 66666666.1425568.peg.2979                                  |

*Note.* The full Table S6 is available online at <https://doi.org/10.5281/zenodo.15725106>