



Universiteit  
Leiden  
The Netherlands

## Microbial footprints of tomato domestication

Sarango Flores, S.W.

### Citation

Sarango Flores, S. W. (2026, January 6). *Microbial footprints of tomato domestication*. NIOO-thesis. Retrieved from <https://hdl.handle.net/1887/4285898>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4285898>

**Note:** To cite this publication please use the final published version (if applicable).

# **References**

---



- Abbamondi, G. R., Tommonaro, G., Weyens, N., Thijs, S., Sillen, W., Gkorezis, P., Iodice, C., de Melo Rangel, W., Nicolaus, B., & Vangronsveld, J. (2016). Plant growth-promoting effects of rhizospheric and endophytic bacteria associated with different tomato cultivars and new tomato hybrids. *Chemical and Biological Technologies in Agriculture*, 3(1), 1–10. <https://doi.org/10.1186/s40538-015-0051-3>
- Abdullaeva, Y., Ratering, S., Ambika Manirajan, B., Rosado-Porto, D., Schnell, S., & Cardinale, M. (2022). Domestication impacts the wheat-associated microbiota and the rhizosphere colonization by seed- and soil-originated microbiomes, across different fields. *Frontiers in Plant Science*, 12, 1–19. <https://www.frontiersin.org/articles/10.3389/fpls.2021.806915>
- Adedayo, A. A., & Babalola, O. O. (2023). Fungi that promote plant growth in the rhizosphere boost crop growth. *Journal of Fungi*, 9(2), 239. <https://doi.org/10.3390/jof9020239>
- Adedayo, A. A., Fadiji, A. E., & Babalola, O. O. (2022). The effects of plant health status on the community structure and metabolic pathways of rhizosphere microbial communities associated with *Solanum lycopersicum*. *Horticulturae*, 8(5), 404. <https://doi.org/10.3390/horticulturae8050404>
- Adedayo, A. A., Fadiji, A. E., & Babalola, O. O. (2023). Unraveling the functional genes present in rhizosphere microbiomes of *Solanum lycopersicum*. *PeerJ*, 11, 1–30. <https://doi.org/10.7717/peerj.15432>
- Aflitos, S., Schijlen, E., Jong, H. de, Ridder, D. de, Smit, S., Finkers, R., Wang, J., Zhang, G., Li, N., Mao, L., Bakker, F., Dirks, R., Breit, T., Gravendeel, B., Huits, H., Struss, D., Swanson-Wagner, R., Leeuwen, H. van, Ham, R. C. H. J. van, ... Peters, S. (2014). Exploring genetic variation in the tomato (*Solanum* section *Lycopersicon*) clade by whole-genome sequencing. *The Plant Journal*, 80(1), 136–148. <https://doi.org/10.1111/tpj.12616>
- Agaras, B. C., Grossi, C. E. M., & Ulloa, R. M. (2023). Unveiling the secrets of calcium-dependent proteins in Plant Growth-Promoting Rhizobacteria: An abundance of discoveries awaits. *Plants*, 12(19), 1–27. <https://doi.org/10.3390/plants12193398>
- Aghdam, M. S., Asghari, M., Farmani, B., Mohayjeji, M., & Moradbeygi, H. (2012). Impact of postharvest brassinosteroids treatment on PAL activity in tomato fruit in response to chilling stress. *Scientia Horticulturae*, 144, 116–120. <https://doi.org/10.1016/j.scienta.2012.07.008>
- Agrawal, S., Heiss, C., Zuckerman, D. M., So, J. M. T., Semeijn, K., Naran, R., Azadi, P., & Hoiczzyk, E. (2020). Structure and function of a novel osmoregulated periplasmic fiber-forming high-molecular-weight carbohydrate of *Myxococcus xanthus*. *bioRxiv*, 2020. <https://doi.org/10.1101/2020.07.26.208595>
- Ahmed, E., & Holmström, S. J. M. (2014). Siderophores in environmental research: Roles and applications. *Microbial Biotechnology*, 7(3), 196–208. <https://doi.org/10.1111/1751-7915.12117>
- Alattas, H., Glick, B. R., Murphy, D. V., & Scott, C. (2024). Harnessing *Pseudomonas* spp. For sustainable plant crop protection. *Frontiers in Microbiology*, 15, 1–21. <https://doi.org/10.3389/fmicb.2024.1485197>
- Al-Kaabi, F. K. (2004). *Insect Control Using Chitinolytic Soil Actinomycetes as Biocontrol Agents*. [https://scholarworks.uaeu.ac.ae/all\\_theses/594](https://scholarworks.uaeu.ac.ae/all_theses/594)
- Allard, S. M., Walsh, C. S., Wallis, A. E., Ottesen, A. R., Brown, E. W., & Micallef, S. A. (2016). *Solanum lycopersicum* (tomato) hosts robust phyllosphere and rhizosphere bacterial communities when grown in soil amended with various organic and synthetic fertilizers. *Science of The Total Environment*, 573, 555–563. <https://doi.org/10.1016/j.scitotenv.2016.08.157>
- Alsharif, W., Saad, M. M., & Hirt, H. (2020). Desert microbes for boosting sustainable agriculture in extreme environments. *Frontiers in Microbiology*, 11, 1–18. <https://doi.org/10.3389/fmicb.2020.01666>
- Alzate Zuluaga, M. Y., Milani, K. M. L., Miras-Moreno, M. B., Lucini, L., Valentinuzzi, F., Mimmo, T., Pii, Y., Cesco, S., Rodrigues, E. P., & de Oliveira, A. L. M. (2021). The adaptive metabolomic

- profile and functional activity of tomato rhizosphere are revealed upon PGPB inoculation under saline stress. *Environmental and Experimental Botany*, *189*, 1–14. <https://doi.org/10.1016/j.envexpbot.2021.104552>
- Amelia-Yap, Z. H., Azman, A. S., AbuBakar, S., & Low, V. L. (2022). *Streptomyces* derivatives as an insecticide: Current perspectives, challenges and future research needs for mosquito control. *Acta Tropica*, *229*, 1–18. <https://doi.org/10.1016/j.actatropica.2022.106381>
- Amelia-Yap, Z. H., Low, V. L., Saeung, A., Ng, F. L., Chen, C. D., Hassandarvish, P., Tan, G. Y. A., AbuBakar, S., & Azman, A. S. (2023). Insecticidal activities of *Streptomyces* sp. KSF103 ethyl acetate extract against medically important mosquitoes and non-target organisms. *Scientific Reports*, *13*(4), 1–14. <https://doi.org/10.1038/s41598-022-25387-9>
- Anjum, S., Sarwar, M., Mukhtar, A., Hussain, T., Liaquat, M., Batool, A., Alam, M. W., Ali, Q., & Ashraf, M. I. (2025). Impact of rhizobacterial inoculation in plant growth medium to mitigate lead stress in tomato (*Solanum lycopersicum* L.). *Scientific Reports*, *15*(1), 1–9. <https://doi.org/10.1038/s41598-024-84648-x>
- Aparicio, M. A., Lucena, C., García, M. J., Ruiz-Castilla, F. J., Jiménez-Adrián, P., López-Berges, M. S., Prieto, P., Alcántara, E., Pérez-Vicente, R., Ramos, J., & Romera, F. J. (2023). The nonpathogenic strain of *Fusarium oxysporum* FO12 induces Fe deficiency responses in cucumber (*Cucumis sativus* L.) plants. *Planta*, *257*(3), 1–17. <https://doi.org/10.1007/s00425-023-04079-2>
- Assis, R. de A. B., Sagawa, C. H. D., Zaini, P. A., Saxe, H. J., Wilmarth, P. A., Phinney, B. S., Salemi, M., Moreira, L. M., & Dandekar, A. M. (2021). A secreted chorismate mutase from *Xanthomonas arboricola* pv. *juglandis* attenuates virulence and walnut blight symptoms. *International Journal of Molecular Sciences*, *22*, 1–19. <https://doi.org/10.3390/ijms221910374>
- Attia, M. A. (2018). In vitro and in vivo characterization of three *Cellvibrio japonicus* glycoside hydrolase family 5 members reveals potent xyloglucan backbone-cleaving functions. *Biotechnol. Biofuels*, *11*(45), 1–16. <https://doi.org/10.1186/s13068-018-1039-6>
- Attia, M. S., Abdelaziz, A. M., Al-Askar, A. A., Arishi, A. A., Abdelhakim, A. M., & Hashem, A. H. (2022). Plant growth-promoting fungi as biocontrol tool against *Fusarium* wilt disease of tomato plant. *Journal of Fungi*, *8*(775), 1–19. <https://doi.org/10.3390/jof8080775>
- Audenaert, K., Pattery, T., Cornelis, P., & Höfte, M. (2002). Induction of systemic resistance to *Botrytis cinerea* in tomato by *Pseudomonas aeruginosa* 7NSK2: Role of salicylic acid, pyochelin, and pyocyanin. *Molecular Plant-Microbe Interactions*, *15*(11), 1147–1156. <https://doi.org/10.1094/MPMI.2002.15.11.1147>
- Avalos, M., Garbeva, P., Vader, L., Wezel, G. P. van, Dickschat, J. S., & Ulanova, D. (2022). Biosynthesis, evolution and ecology of microbial terpenoids. *Natural Product Reports*, *39*(2), 249–272. <https://doi.org/10.1039/D1NP00047K>
- Ayangbenro, A. S., & Babalola, O. O. (2021). Reclamation of arid and semi-arid soils: The role of plant growth-promoting archaea and bacteria. *Current Plant Biology*, *25*, 1–12. <https://doi.org/10.1016/j.cpb.2020.100173>
- Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., Formsma, K., Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman, A. L., Overbeek, R. A., McNeil, L. K., Paarmann, D., Paczian, T., Parrello, B., ... Zagnitko, O. (2008). The RAST Server: Rapid annotations using subsystems technology. *BMC Genomics*, *9*(75), 1–15. <https://doi.org/10.1186/1471-2164-9-75>
- Azzi, L., Deluche, C., Gévaudant, F., Frangne, N., Delmas, F., Hernould, M., & Chevalier, C. (2015). Fruit growth-related genes in tomato. *Journal of Experimental Botany*, *66*(4), 1075–1086. <https://doi.org/10.1093/jxb/eru527>

- Badri, D. V., Zolla, G., Bakker, M. G., Manter, D. K., & Vivanco, J. M. (2013). Potential impact of soil microbiomes on the leaf metabolome and on herbivore feeding behavior. *New Phytologist*, *198*(1), 264–273. <https://doi.org/10.1111/nph.12124>
- Bai, Y., & Lindhout, P. (2007). Domestication and breeding of tomatoes: What have we gained and what can we gain in the future? *Annals of Botany*, *100*(5), 1085–1094. <https://doi.org/10.1093/aob/mcm150>
- Balakrishnan, S., Santhanam, P., & Srinivasan, M. (2017). Larvicidal potency of marine actinobacteria isolated from mangrove environment against *Aedes aegypti* and *Anopheles stephensi*. *Journal of Parasitic Diseases: Official Organ of the Indian Society for Parasitology*, *41*(2), 387–394. <https://doi.org/10.1007/s12639-016-0812-3>
- Bandyopadhyay, P., Bhuyan, S. K., Yadava, P. K., Varma, A., & Tuteja, N. (2017). Emergence of plant and rhizospheric microbiota as stable interactomes. *Protoplasma*, *254*(2), 617–626. <https://doi.org/10.1007/s00709-016-1003-x>
- Bankevich, A. (2012). SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of computational biology: a journal of computational molecular cell biology*, *19*(5), 455–477. <https://doi.org/10.1089/cmb.2012.0021>
- Barajas, H. R., Martínez-Sánchez, S., Romero, M. F., Hernández-Álvarez, C., Servín-González, L., Peimbert, M., Cruz-Ortega, R., García-Oliva, F., & Alcaraz, L. D. (2020). Testing the two-step model of plant root microbiome acquisition under multiple plant species and soil sources. *Frontiers in Microbiology*, *11*, 1–17. <https://doi.org/10.3389/fmicb.2020.542742>
- Barber, C. C., & Zhang, W. (2021). Small molecule natural products in human nasal/oral microbiota. *Journal of Industrial Microbiology & Biotechnology*, *48*(3–4), 1–12. <https://doi.org/10.1093/jimb/kuab010>
- Barendse, G. W. M., & van der Weerden, G. M. (1997). The Solanaceae germplasm bank at the Botanical Garden of Nijmegen. *Botanic Gardens Conservation News*, *2*(9), 31–33.
- Barnawal, D., Singh, R., & Singh, R. P. (2019). Chapter Six - Role of Plant Growth Promoting Rhizobacteria in Drought Tolerance: Regulating Growth Hormones and Osmolytes. In A. K. Singh, A. Kumar, & P. K. Singh (Eds.), *PGPR Amelioration in Sustainable Agriculture* (pp. 107–128). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-815879-1.00006-9>
- Barnes, C. J., Bahram, M., Nicolaisen, M., Gilbert, M. T. P., & Vestergård, M. (2024). Microbiome selection and evolution within wild and domesticated plants. *Trends in Microbiology*, *33*(4), 447–458. <https://doi.org/10.1016/j.tim.2024.11.011>
- Barone, A., Chiusano, M. L., Ercolano, M. R., Giuliano, G., Grandillo, S., & Frusciante, L. (2008). Structural and functional genomics of tomato. *International Journal of Plant Genomics*, *2008*, 820274, 1–12. <https://doi.org/10.1155/2008/820274>
- Beauregard, P. B., Chai, Y., Vlamakis, H., Losick, R., & Kolter, R. (2013). *Bacillus subtilis* biofilm induction by plant polysaccharides. *Proc. Natl Acad. Sci. USA*, *110*(17), 1621–1630. <https://doi.org/10.1073/pnas.1218984110>
- Beilsmith, K. (2019). Genome-wide association studies on the phyllosphere microbiome: Embracing complexity in host-microbe interactions. *The Plant Journal*, *97*(1), 164–181. <https://doi.org/10.1111/tpj.14170>
- Benaud, N., Edwards, R. J., Amos, T. G., D'Agostino, P. M., Gutiérrez-Chávez, C., Montgomery, K., Nicetic, I., & Ferrari, B. C. (2021). Antarctic desert soil bacteria exhibit high novel natural product potential, evaluated through long-read genome sequencing and comparative genomics. *Environmental Microbiology*, *23*(7), 3646–3664. <https://doi.org/10.1111/1462-2920.15300>
- Bentley, S. D. (2002). Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature*, *417*, 141–147. <https://doi.org/10.1038/417141a>

- Berendsen, R. L., Pieterse, C. M. J., & Bakker, P. A. H. M. (2012). The rhizosphere microbiome and plant health. *Trends in Plant Science*, *17*(8), 478–486. <https://doi.org/10.1016/j.tplants.2012.04.001>
- Berg, G., & Raaijmakers, J. M. (2018). Saving seed microbiomes. *The ISME Journal*, *12*(5), 1167–1170. <https://doi.org/10.1038/s41396-017-0028-2>
- Berg, G., & Smalla, K. (2009). Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiology Ecology*, *68*(1), 1–13. <https://doi.org/10.1111/j.1574-6941.2009.00654.x>
- Berg, G., Grube, M., Schloter, M., & Smalla, K. (2014). Unraveling the plant microbiome: Looking back and future perspectives. *Frontiers in Microbiology*, *5*(148), 1–7. <https://doi.org/10.3389/fmicb.2014.00148>
- Bergelson, J., Brachi, B., Roux, F., & Vaillau, F. (2021). Assessing the potential to harness the microbiome through plant genetics. *Curr. Opin. Biotechnol.*, *70*, 167–173. <https://doi.org/10.1016/j.copbio.2021.05.007>
- Bergelson, J., Mittelstrass, J., & Horton, M. W. (2019). Characterizing both bacteria and fungi improves understanding of the *Arabidopsis* root microbiome. *Scientific Reports*, *9*(24), 1–11. <https://doi.org/10.1038/s41598-018-37208-z>
- Bergna, A., Cernava, T., Rändler, M., Grosch, R., Zachow, C., & Berg, G. (2018). Tomato seeds preferably transmit plant beneficial endophytes. *Phytobiomes Journal*, *2*(4), 183–193. <https://doi.org/10.1094/PBIOMES-06-18-0029-R>
- Bergougnoux, V. (2014). The history of tomato: From domestication to biopharming. *Biotechnology Advances*, *32*(1), 170–189. <https://doi.org/10.1016/j.biotechadv.2013.11.003>
- Berry, J. P., Gantar, M., Perez, M. H., Berry, G., & Noriega, F. G. (2008). Cyanobacterial toxins as allelochemicals with potential applications as algacides, herbicides and insecticides. *Marine Drugs*, *6*(2), 117–146. <https://doi.org/10.3390/md6020117>
- Bharti, N., & Barnawal, D. (2019). Chapter Five - Amelioration of Salinity Stress by PGPR: ACC Deaminase and ROS Scavenging Enzymes Activity. In A. K. Singh, A. Kumar, & P. K. Singh (Eds.), *PGPR Amelioration in Sustainable Agriculture* (pp. 85–106). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-815879-1.00005-7>
- Blanca, J., & Cañizares, J. (2021, December 3). El tomate no es natural, es un tesoro creado por el ingenio humano. *El País*. <https://elpais.com/ciencia/2021-12-03/el-tomate-no-es-natural-es-un-tesoro-creado-por-el-ingenio-humano.html>
- Blanca, J., Cañizares, J., Cordero, L., Pascual, L., Díez, M. J., & Nuez, F. (2012). Variation revealed by SNP genotyping and morphology provides insight into the origin of the tomato. *PLOS ONE*, *7*(10), 1–17. <https://doi.org/10.1371/journal.pone.0048198>
- Blanca, J., Montero-Pau, J., Sauvage, C., Bauchet, G., Illa, E., Díez, M. J., Francis, D., Causse, M., van der Knaap, E., & Cañizares, J. (2015). Genomic variation in tomato, from wild ancestors to contemporary breeding accessions. *BMC Genomics*, *16*(257), 1–19. <https://doi.org/10.1186/s12864-015-1444-1>
- Blanca, J., Sanchez-Matarredona, D., Ziarsolo, P., Montero-Pau, J., van der Knaap, E., Díez, M. J., & Cañizares, J. (2022). Haplotype analyses reveal novel insights into tomato history and domestication driven by long-distance migrations and latitudinal adaptations. *Horticulture Research*, *9*, 1–14. <https://doi.org/10.1093/hr/uhac030>
- Blin, K. (2021). antiSMASH 6.0: Improving cluster detection and comparison capabilities. *Nucleic Acids Res.*, *49*, 29–35. <https://doi.org/10.1093/nar/gkab335>
- Blin, K., Shaw, S., Augustijn, H. E., Reitz, Z. L., Biermann, F., Alanjary, M., Fetter, A., Terlouw, B. R., Metcalf, W. W., Helfrich, E. J. N., van Wezel, G. P., Medema, M. H., & Weber, T. (2023). an-

- tiSMASH 7.0: New and improved predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids Research*, *51*, 46–50. <https://doi.org/10.1093/nar/gkad344>
- Blundell, R., Schmidt, J. E., Igwe, A., Cheung, A. L., Vannette, R. L., Gaudin, A. C. M., & Casteel, C. L. (2020). Organic management promotes natural pest control through altered plant resistance to insects. *Nature Plants*, *6*(5), 483–491. <https://doi.org/10.1038/s41477-020-0656-9>
- Bontemps-Gallo, S., Bohin, J.-P., & Lacroix, J.-M. (2017). Osmoregulated Periplasmic Glucans. *EcoSal Plus*, *7*(2), 1–17. <https://doi.org/10.1128/ecosalplus.esp-0001-2017>
- Bowles, D. (2018, December 29). Mexican X Part IV: You Say “Tomato,” I Say You’re Missing a Syllable, Bro! *Medium*. <https://davidbowles.medium.com/mexican-x-part-iv-you-say-tomato-i-say-youre-missing-a-syllable-bro-1f002f4f110c>
- Bretting, P. K. (2018). 2017 Frank Meyer medal for plant genetic resources lecture: Stewards of our agricultural future. *Crop Science*, *58*(6), 2233–2240. <https://doi.org/10.2135/cropsci2018.05.0334>
- Brilli, F., Pollastri, S., Raio, A., Baraldi, R., Neri, L., Bartolini, P., Podda, A., Loreto, F., Maserti, B. E., & Balestrini, R. (2019). Root colonization by *Pseudomonas chlororaphis* primes tomato (*Lycopersicon esculentum*) plants for enhanced tolerance to water stress. *Journal of Plant Physiology*, *232*, 82–93. <https://doi.org/10.1016/j.jplph.2018.10.029>
- Broman, K. W. (2019). R/qtl2: Software for mapping quantitative trait loci with high-dimensional data and multiparent populations. *Genetics*, *211*(2), 495–502. <https://doi.org/10.1534/genetics.118.301595>
- Brooks, C., Nekrasov, V., Lippman, Z. B., & Van Eck, J. (2014). Efficient gene editing in tomato in the first generation using the clustered regularly interspaced short palindromic repeats/CRISPR-associated9 system. *Plant Physiology*, *166*(3), 1292–1297. <https://doi.org/10.1104/pp.114.247577>
- Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, *12*(1), 59–60. <https://doi.org/10.1038/nmeth.3176>
- Busby, P. E. (2017). Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLOS Biology*, *15*(3), 1–14. <https://doi.org/10.1371/journal.pbio.2001793>
- Busch, A., & Waksman, G. (2012). Chaperone–usher pathways: Diversity and pilus assembly mechanism. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1592), 1112–1122. <https://doi.org/10.1098/rstb.2011.0206>
- Caballero-Mellado, J., Onofre-Lemus, J., Estrada-de los Santos, P., & Martínez-Aguilar, L. (2007). The tomato rhizosphere, an environment rich in nitrogen-fixing *Burkholderia* species with capabilities of interest for agriculture and bioremediation. *Applied and Environmental Microbiology*, *73*(16), 5308–5319. <https://doi.org/10.1128/AEM.00324-07>
- Cai, F., Pang, G., Miao, Y., Li, R., Li, R., Shen, Q., & Chen, W. (2017). The nutrient preference of plants influences their rhizosphere microbiome. *Applied Soil Ecology*, *110*, 146–150. <https://doi.org/10.1016/j.apsoil.2016.11.006>
- Caicedo, A. L., & Schaal, B. A. (2004). Population structure and phylogeography of *Solanum pimpinellifolium* inferred from a nuclear gene. *Molecular Ecology*, *13*(7), 1871–1882. <https://doi.org/10.1111/j.1365-294X.2004.02191.x>
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583. <https://doi.org/10.1038/nmeth.3869>
- Calvo-Polanco, M., Sánchez-Romera, B., Aroca, R., Asins, M. J., Declerck, S., Dodd, I. C., Martínez-Andújar, C., Albacete, A., & Ruiz-Lozano, J. M. (2016). Exploring the use of recombinant inbred lines in combination with beneficial microbial inoculants (AM fungus and PGPR) to improve drought stress tolerance in tomato. *Environmental and Experimental Botany*, *131*, 47–57. <https://doi.org/10.1016/j.envexpbot.2016.06.015>

- Canarini, A., Kaiser, C., Merchant, A., Richter, A., & Wanek, W. (2019). Root exudation of primary metabolites: Mechanisms and their roles in plant responses to environmental stimuli. *Frontiers in Plant Science*, *10*(157), 1–19. <https://doi.org/10.3389/fpls.2019.00157>
- Cao, T., Luo, Y., Shi, M., Tian, X., & Kuzyakov, Y. (2024). Microbial interactions for nutrient acquisition in soil: Miners, scavengers, and carriers. *Soil Biology and Biochemistry*, *188*, 1–18. <https://doi.org/10.1016/j.soilbio.2023.109215>
- Caramante, M., Roupheal, Y., & Corrado, G. (2021). The genetic diversity and structure of tomato landraces from the Campania region (Southern Italy) uncovers a distinct population identity. *Agronomy*, *11*(3), 1–12. <https://doi.org/10.3390/agronomy11030564>
- Carmona, S. L., Burbano-David, D., Gómez, M. R., Lopez, W., Ceballos, N., Castaño-Zapata, J., Simbaqueba, J., & Soto-Suárez, M. (2020). Characterization of pathogenic and nonpathogenic *Fusarium oxysporum* isolates associated with commercial tomato crops in the Andean region of Colombia. *Pathogens*, *9*(70), 1–23. <https://doi.org/10.3390/pathogens9010070>
- Carrillo, J., Ingwell, L. L., Li, X., & Kaplan, I. (2019). Domesticated tomatoes are more vulnerable to negative plant–soil feedbacks than their wild relatives. *Journal of Ecology*, *107*(4), 1753–1766. <https://doi.org/10.1111/1365-2745.13157>
- Carrión, V. J. (2019). Pathogen-induced activation of disease-suppressive functions in the endophytic root microbiome. *Science*, *366*(6465), 606–612. <https://doi.org/10.1126/science.aaw9285>
- Carro, L., Veyisoglu, A., Guven, K., Schumann, P., Klenk, H.-P., & Sahin, N. (2022). Genomic analysis of a novel heavy metal resistant isolate from a black sea contaminated sediment with the potential to degrade alkanes: *Plantactinospora alkalitolerans* sp. nov. *Diversity*, *14*(11), 1–14. <https://doi.org/10.3390/d14110947>
- Casas, A. (2001). Silvicultura y domesticación de plantas en Mesoamérica. In B. Rendón Aguilar, S. Rebolgar Domínguez, J. Caballero Nieto, & M. Á. Martínez Alfaro (Eds.), *Plantas, cultura y sociedad: Estudio sobre la relación entre seres humanos y plantas en los albores del siglo XXI* (1. ed, pp. 123–157). Coloquio “Plantas, Cultura y Sociedad: Estudio sobre la Relación entre Seres Humanos y Plantas en los Albores del Siglo XXI,” México, D.F. Universidad Autónoma Metropolitana, Unidad Iztapalapa: Secretaría de Medio Ambiente, Recursos Naturales y Pesca.
- Cebolla-Cornejo, J., Roselló, S., & Nuez, F. (2013). Phenotypic and genetic diversity of Spanish tomato landraces. *Scientia Horticulturae*, *162*, 150–164. <https://doi.org/10.1016/j.scienta.2013.07.044>
- Cernava, T. (2024). Coming of age for Microbiome gene breeding in plants. *Nature Communications*, *15*(1), 1–3. <https://doi.org/10.1038/s41467-024-50700-7>
- Cesaro, P., Massa, N., Cantamessa, S., Todeschini, V., Bona, E., Berta, G., Barbato, R., & Lingua, G. (2020). Tomato responses to *Funneliformis mosseae* during the early stages of arbuscular mycorrhizal symbiosis. *Mycorrhiza*, *30*(5), 601–610. <https://doi.org/10.1007/s00572-020-00973-9>
- Challacombe, J. E., Hesse, C. N., Bramer, L. M., McCue, L. A., Lipton, M., Purvine, S., Nicora, C., Gallegos-Graves, L. V., Porrás-Alfaro, A., & Kuske, C. R. (2019). Genomes and secretomes of Ascomycota fungi reveal diverse functions in plant biomass decomposition and pathogenesis. *BMC Genomics*, *20*(1), 1–27. <https://doi.org/10.1186/s12864-019-6358-x>
- Chandrasekaran, M., Boopathi, T., & Manivannan, P. (2021). Comprehensive assessment of ameliorative effects of AMF in alleviating abiotic stress in tomato plants. *Journal of Fungi*, *7*(4), 1–21. <https://doi.org/10.3390/jof7040303>
- Chaparro, J. M. (2013). Root exudation of phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and correlate with soil microbial functions. *PLoS One*, *8*(2), 1–10. <https://doi.org/10.1371/journal.pone.0055731>

- Chaudhary, P., Sharma, A., Singh, B., & Nagpal, A. K. (2018). Bioactivities of phytochemicals present in tomato. *Journal of Food Science and Technology*, *55*(8), 2833–2849. <https://doi.org/10.1007/s13197-018-3221-z>
- Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2020). GTDB-Tk: A toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics*, *36*(6), 1925–1927. <https://doi.org/10.1093/bioinformatics/btz848>
- Cheema, D. S., & Dhaliwal, M. S. (2004). Hybrid tomato breeding. *Journal of New Seeds*, *6*, 1–14. [https://doi.org/10.1300/J153v06n02\\_01](https://doi.org/10.1300/J153v06n02_01)
- Chen, S., Sun, Y., Wei, Y., Li, H., & Yang, S. (2022). Different rhizosphere soil microbes are recruited by tomatoes with different fruit color phenotypes. *BMC Microbiology*, *22*(210), 1–15. <https://doi.org/10.1186/s12866-022-02620-z>
- Chen, Y. H., Gols, R., & Benrey, B. (2015). Crop domestication and its impact on naturally selected trophic interactions. *Annual Review of Entomology*, *60*(1), 35–58. <https://doi.org/10.1146/annurev-ento-010814-020601>
- Cheng, Z., Lei, S., Li, Y., Huang, W., Ma, R., Xiong, J., Zhang, T., Jin, L., Haq, H. ul, Xu, X., & Tian, B. (2020). Revealing the variation and stability of bacterial communities in tomato rhizosphere microbiota. *Microorganisms*, *8*(2), 1–15. <https://doi.org/10.3390/microorganisms8020170>
- Chepsergon, J., & Moleleki, L. N. (2023). Rhizosphere bacterial interactions and impact on plant health. *Current Opinion in Microbiology*, *73*, 1–10. <https://doi.org/10.1016/j.mib.2023.102297>
- Chewning, S. S. (2019). Root-associated streptomyces isolates harboring melC genes demonstrate enhanced plant colonization. *Phytobiomes Journal*, *3*(3), 165–176. <https://doi.org/10.1094/PBIOMES-01-19-0005-R>
- Chialva, M., Salvioli di Fossalunga, A., Daghino, S., Ghignone, S., Bagnaresi, P., Chiapello, M., Novero, M., Spadaro, D., Perotto, S., & Bonfante, P. (2018). Native soils with their microbiotas elicit a state of alert in tomato plants. *New Phytologist*, *220*(4), 1296–1308. <https://doi.org/10.1111/nph.15014>
- Chieb, M., & Gachomo, E. W. (2023). The role of plant growth promoting rhizobacteria in plant drought stress responses. *BMC Plant Biology*, *23*(407), 1–23. <https://doi.org/10.1186/s12870-023-04403-8>
- Choi, K., Choi, J., Lee, P. A., Roy, N., Khan, R., Lee, H. J., Weon, H. Y., Kong, H. G., & Lee, S.-W. (2020). Alteration of bacterial wilt resistance in tomato plant by microbiota transplant. *Frontiers in Plant Science*, *11*, 1–12. <https://doi.org/10.3389/fpls.2020.01186>
- Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2016). GenBank. *Nucleic Acids Research*, *44*(D1), D67–72. <https://doi.org/10.1093/nar/gkv1276>
- Cole, J. R. (2014). Ribosomal Database Project: Data and tools for high throughput rRNA analysis. *Nucleic Acids Research*, *42*(D1), D633–D642. <https://doi.org/10.1093/nar/gkt1244>
- Constante Tubay, G. B., Cañarte Bermudez, E. G., Navarrete Cedeño, J. B., Saldarriaga Lucas, V. A., Constante Tubay, G. B., Cañarte Bermudez, E. G., Navarrete Cedeño, J. B., & Saldarriaga Lucas, V. A. (2023). Situación actual de *Prodiplosis longifila* Gagné (Diptera: Cecidomyiidae) en zonas tomateras de Manabi, Ecuador. *Manglar*, *20*(1), 77–86. <https://doi.org/10.57188/manglar.2023.009>
- Cordero-Ramírez, J. D., López-Rivera, R., Calderón-Vázquez, C. L., Figueroa-López, A. M., Martínez-Álvarez, J. C., Leyva-Madrugal, K. Y., Cervantes-Gámez, R. G., & Maldonado-Mendoza, I. E. (2012). Microorganismos asociados a la rizosfera de jitomate en un agroecosistema del valle de Guasave, Sinaloa, México. *Revista mexicana de biodiversidad*, *83*(3), 712–730. <https://doi.org/10.7550/rmb.17897>
- Cordovez, V., Dini-Andreote, F., Carrión, V. J., & Raaijmakers, J. M. (2019). Ecology and evolution of plant microbiomes. *Annual Review of Microbiology*, *73*(1), 69–88. <https://doi.org/10.1146/annurev-micro-090817-062524>

- Cordovez, V., Rotoni, C., Dini-Andreote, F., Oyserman, B., Carrión, V. J., & Raaijmakers, J. M. (2021). Successive plant growth amplifies genotype-specific assembly of the tomato rhizosphere microbiome. *The Science of the Total Environment*, 772, 1–10. <https://doi.org/10.1016/j.scitotenv.2020.144825>
- Corpas, F. J., Gupta, D. K., & Palma, J. M. (2021). Tryptophan: A precursor of signaling molecules in higher plants. In D. K. Gupta & F. J. Corpas (Eds.), *Hormones and Plant Response* (pp. 273–289). Springer International Publishing. [https://doi.org/10.1007/978-3-030-77477-6\\_11](https://doi.org/10.1007/978-3-030-77477-6_11)
- Cosgrove, D. J. (2016). Catalysts of plant cell wall loosening. *F1000Res*, 5, 1–13. <https://doi.org/10.12688/f1000research.7180.1>
- Cosgrove, D. J. (2017). Microbial expansins. *Annual Review of Microbiology*, 71, 479–497. <https://doi.org/10.1146/annurev-micro-090816-093315>
- Costa, T. R. D., Patkowski, J. B., Macé, K., Christie, P. J., & Waksman, G. (2023). Structural and functional diversity of type IV secretion systems. *Nature Reviews Microbiology*, 22, 170–185. <https://doi.org/10.1038/s41579-023-00974-3>
- Craig, L., Forest, K. T., & Maier, B. (2019). Type IV pili: Dynamics, biophysics and functional consequences. *Nature Reviews Microbiology*, 17(7), 429–440. <https://doi.org/10.1038/s41579-019-0195-4>
- Crits-Christoph, A., Bhattacharya, N., Olm, M. R., Song, Y. S., & Banfield, J. F. (2021). Transporter genes in biosynthetic gene clusters predict metabolite characteristics and siderophore activity. *Genome research*, 31(2), 239–250. <https://doi.org/10.1101/gr.268169.120>
- Crowley, D. E. (2006). Microbial Siderophores in the Plant Rhizosphere. En L. L. Barton & J. Abadia (Eds.), *Iron Nutrition in Plants and Rhizospheric Microorganisms* (pp. 169–198). Springer Netherlands. [https://doi.org/10.1007/1-4020-4743-6\\_8](https://doi.org/10.1007/1-4020-4743-6_8)
- Dai, J., Wei, H., Tian, C., Damron, F. H., Zhou, J., & Qiu, D. (2015). An extracytoplasmic function sigma factor-dependent periplasmic glutathione peroxidase is involved in oxidative stress response of *Shewanella oneidensis*. *BMC Microbiology*, 15(34), 1–12. <https://doi.org/10.1186/s12866-015-0357-0>
- Daigham, G. E., Mahfouz, A. Y., Abdelaziz, A. M., Nofel, M. M., & Attia, M. S. (2023). Protective role of plant growth-promoting fungi *Aspergillus chevalieri* OP593083 and *Aspergillus egyptiacus* OP593080 as biocontrol approach against *Alternaria* leaf spot disease of *Vicia faba* plant. *Biomass Conversion and Biorefinery*, 14, 23073–23089. <https://doi.org/10.1007/s13399-023-04510-4>
- Dastogeer, K. M. G., Tumpa, F. H., Sultana, A., Akter, M. A., & Chakraborty, A. (2020). Plant microbiome—an account of the factors that shape community composition and diversity. *Current Plant Biology*, 23, 1–9. <https://doi.org/10.1016/j.cpb.2020.100161>
- de Lamo, F. J., Spijkers, S. B., & Takken, F. L. W. (2021). Protection to tomato wilt disease conferred by the nonpathogen *Fusarium oxysporum* Fo47 is more effective than that conferred by avirulent strains. *Phytopathology*, 111(2), 253–257. <https://doi.org/10.1094/PHYTO-04-20-0133-R>
- de Weert, S., Vermeiren, H., Mulders, I. H. M., Kuiper, I., Hendrickx, N., Bloembergen, G. V., Vanderleyden, J., De Mot, R., & Lugtenberg, B. J. J. (2002). Flagella-driven chemotaxis towards exudate components is an important trait for tomato root colonization by *Pseudomonas fluorescens*. *Molecular Plant-Microbe Interactions*, 15(11), 1173–1180. <https://doi.org/10.1094/MPMI.2002.15.11.1173>
- Deng, S. (2021). Genome wide association study reveals plant loci controlling heritability of the rhizosphere microbiome. *The ISME Journal*, 15(11), 3181–3194. <https://doi.org/10.1038/s41396-021-00993-z>
- Dennis, P. G., Miller, A. J., & Hirsch, P. R. (2010). Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiology Ecology*, 72(3), 313–327. <https://doi.org/10.1111/j.1574-6941.2010.00860.x>

- Desneux, N., Han, P., Mansour, R., Arnó, J., Brévault, T., Campos, M. R., Chailleux, A., Guedes, R. N. C., Karimi, J., Konan, K. A. J., Lavoit, A., Luna, M. G., Perez-Hedo, M., Urbaneja, A., Verheggen, F. J., Zappalà, L., Abbes, K., Ali, A., Bayram, Y., ... Biondi, A. (2022). Integrated pest management of *Tuta absoluta*: Practical implementations across different world regions. *Journal of Pest Science*, 95(1), 17–39. <https://doi.org/10.1007/s10340-021-01442-8>
- Díaz, E., Jiménez, J. I., & Nogales, J. (2013). Aerobic degradation of aromatic compounds. *Current Opinion in Biotechnology*, 24(3), 431–442. <https://doi.org/10.1016/j.copbio.2012.10.010>
- Díez, M. J., & Nuez, F. (2008). Tomato. In J. Prohens & F. Nuez (Eds.), *Vegetables II Fabaceae, Liliaceae, Solanaceae, and Umbelliferae* (Vol. 2, pp. 249–323). <https://link.springer.com/book/10.1007/978-0-387-74110-9>
- Distefano, M., Mauro, R. P., Page, D., Giuffrida, F., Bertin, N., & Leonardi, C. (2022). Aroma volatiles in tomato fruits: The role of genetic, preharvest and postharvest factors. *Agronomy*, 12(2), 1–27. <https://doi.org/10.3390/agronomy12020376>
- Dixon, M. M., Afkairin, A., Manter, D. K., & Vivanco, J. (2024). Rhizosphere microbiome co-occurrence network analysis across a tomato domestication gradient. *Microorganisms*, 12(9), 1–12. <https://doi.org/10.3390/microorganisms12091756>
- Doty, M., Roehr, J., Ahmed, R., & Dieterich, C. (2012). FLEXBAR—Flexible barcode and adapter processing for next-generation sequencing platforms. *Biology*, 1(3), 895–905 <https://doi.org/10.3390/biology1030895>
- Doebley, J. F., Gaut, B. S., & Smith, B. D. (2006). The molecular genetics of crop domestication. *Cell*, 127(7), 1309–1321. <https://doi.org/10.1016/j.cell.2006.12.006>
- Domínguez, M., Dugas, E., Benchouaia, M., Leduque, B., Jiménez-Gómez, J. M., Colot, V., & Quadrana, L. (2020). The impact of transposable elements on tomato diversity. *Nature Communications*, 11(1), 1–11. <https://doi.org/10.1038/s41467-020-17874-2>
- Dong, C.-J., Wang, L.-L., Li, Q., & Shang, Q.-M. (2019). Bacterial communities in the rhizosphere, phyllosphere and endosphere of tomato plants. *PLOS ONE*, 14(11), 1–17. <https://doi.org/10.1371/journal.pone.0223847>
- Doron, L., Sutter, M., & Kerfeld, C. A. (2023). Characterization of a novel aromatic substrate-processing microcompartment in Actinobacteria. *mBio*, 14(4), 1–20. <https://doi.org/10.1128/mbio.01216-23>
- Ducousso-Détréz, A., Fontaine, J., Lounès-Hadj Sahraoui, A., & Hijri, M. (2022). Diversity of phosphate chemical forms in soils and their contributions on soil microbial community structure changes. *Microorganisms*, 10(3), 1–22. <https://doi.org/10.3390/microorganisms10030609>
- Duque-Gamboa, D. N., Castillo-Cárdenas, M. F., Hernández, L. M., Guzmán, Y. C., Manzano, M. R., & Toro-Perea, N. (2018). Mitochondrial DNA suggests cryptic speciation in *Prodiplosis longifila* Gagné (Diptera: Cecidomyiidae) associated with geographic distance and host specialization. *Bulletin of Entomological Research*, 108(6), 739–749. <https://doi.org/10.1017/S0007485317001298>
- Egidi, E., Delgado-Baquerizo, M., Plett, J. M., Wang, J., Eldridge, D. J., Bardgett, R. D., Maestre, F. T., & Singh, B. K. (2019). A few Ascomycota taxa dominate soil fungal communities worldwide. *Nature Communications*, 10(1), 1–9. <https://doi.org/10.1038/s41467-019-10373-z>
- EPPO (European and Mediterranean Plant Protection Organization). (2017). *Pest risk analysis for Prodiplosis longifila*. EPPO. <https://gd.eppo.int/taxon/PRDILO>
- Ercolano, M. R., Di Donato, A., Sanseverino, W., Barbella, M., De Natale, A., & Frusciante, L. (2020). Complex migration history is revealed by genetic diversity of tomato samples collected in Italy during the eighteenth and nineteenth centuries. *Horticulture Research*, 7(1), 1–8. <https://doi.org/10.1038/s41438-020-0322-4>

- Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048. <https://doi.org/10.1093/bioinformatics/btw354>
- Fang, B.-Z., Salam, N., Han, M.-X., Jiao, J.-Y., Cheng, J., Wei, D.-Q., Xiao, M., & Li, W.-J. (2017). Insights on the effects of heat pretreatment, pH, and calcium salts on isolation of rare Actinobacteria from karstic caves. *Frontiers in Microbiology*, *8*, 1–9. <https://doi.org/10.3389/fmicb.2017.01535>
- FAO. (2020). *FAOSTAT Data, Crops and livestock products*. <https://www.fao.org/faostat/en/#data/QCL>
- FAO. (2021). *World Food and Agriculture – Statistical Yearbook 2021*. FAO. <https://doi.org/10.4060/cb4477en>
- Favela, A., O. Bohn, M., & D. Kent, A. (2021). Maize germplasm chronosequence shows crop breeding history impacts recruitment of the rhizosphere microbiome. *The ISME Journal*, *15*(8), 2454–2464. <https://doi.org/10.1038/s41396-021-00923-z>
- Feng, H., Fu, R., Hou, X., Lv, Y., Zhang, N., Liu, Y., Xu, Z., Miao, Y., Krell, T., Shen, Q., & Zhang, R. (2021). Chemotaxis of beneficial rhizobacteria to root exudates: The first step towards root–microbe rhizosphere interactions. *International Journal of Molecular Sciences*, *22*(13), 1–14. <https://doi.org/10.3390/ijms22136655>
- Fentik, D. A. (2017). review on genetics and breeding of tomato (*Lycopersicon esculentum* Mill). *Advances in Crop Science and Technology*, *05*(05), 1–6. <https://doi.org/10.4172/2329-8863.1000306>
- Fernie, A. R., & Yan, J. (2019). *De Novo* domestication: An alternative route toward new crops for the future. *Molecular Plant*, *12*(5), 615–631. <https://doi.org/10.1016/j.molp.2019.03.016>
- Finkel, O. M. (2020). A single bacterial genus maintains root growth in a complex microbiome. *Nature*, *587*, 103–108. <https://doi.org/10.1038/s41586-020-2778-7>
- Finkel, O. M., Castrillo, G., Herrera Paredes, S., Salas González, I., & Dangl, J. L. (2017). Understanding and exploiting plant beneficial microbes. *Current Opinion in Plant Biology*, *38*, 155–163. <https://doi.org/10.1016/j.pbi.2017.04.018>
- Fiodor, A., Singh, S., & Pranaw, K. (2021). The contrivance of plant growth promoting microbes to mitigate climate change impact in agriculture. *Microorganisms*, *9*, 1–36. <https://doi.org/10.3390/microorganisms9091841>
- Fischer, I., Camus-Kulandaivelu, L., Allal, F., & Stephan, W. (2011). Adaptation to drought in two wild tomato species: The evolution of the Asr gene family. *New Phytologist*, *190*(4), 1032–1044. <https://doi.org/10.1111/j.1469-8137.2011.03648.x>
- Flemer, B., Gulati, S., Bergna, A., Rändler, M., Cernava, T., Witzel, K., Berg, G., & Grosch, R. (2022). Biotic and abiotic stress factors induce microbiome shifts and enrichment of distinct beneficial bacteria in tomato roots. *Phytobiomes Journal*, *6*(4), 276–289. <https://doi.org/10.1094/PBIOMES-10-21-0067-R>
- Foolad, M. R., & Panthee, D. R. (2012). Marker-assisted selection in tomato breeding. *Critical Reviews in Plant Sciences*, *31*(2), 93–123. <https://doi.org/10.1080/07352689.2011.616057>
- Franza, T., & Expert, D. (2013). Role of iron homeostasis in the virulence of phytopathogenic bacteria: An ‘à la carte’ menu. *Molecular Plant Pathology*, *14*(4), 429–438. <https://doi.org/10.1111/mpp.12007>
- Freeman, K. R., Martin, A. P., Karki, D., Lynch, R. C., Mitter, M. S., Meyer, A. F., Longcore, J. E., Simmons, D. R., & Schmidt, S. K. (2009). Evidence that chytrids dominate fungal communities in high-elevation soils. *Proceedings of the National Academy of Sciences*, *106*(43), 18315–18320. <https://doi.org/10.1073/pnas.0907303106>
- French, E., Ghaste, M., Widhalm, J. R., & Iyer-Pascuzzi, A. S. (2019). Defense hormones modulate root microbiome diversity and composition in tomato. *bioRxiv*, 2019, <https://doi.org/10.1101/656769>

- French, E., Kaplan, I., & Enders, L. (2021). Foliar aphid herbivory alters the tomato rhizosphere microbiome, but initial soil community determines the legacy effects. *Frontiers in Sustainable Food Systems*, 5, 1–16. <https://doi.org/10.3389/fsufs.2021.629684>
- French, E., Tran, T., & Iyer-Pascuzzi, A. S. (2020). Tomato genotype modulates selection and responses to root microbiota. *Phytobiomes Journal*, 4(4), 314–326. <https://doi.org/10.1094/PBIOMES-02-20-0020-R>
- Friman, J., Pineda, A., Gershenzon, J., Dicke, M., & Loon, J. J. A. van. (2021). Differential effects of the rhizobacterium *Pseudomonas simiae* on above- and belowground chewing insect herbivores. *Journal of Applied Entomology*, 145(3), 250–260. <https://doi.org/10.1111/jen.12842>
- Frossard, S. M., Khan, A. A., Warrick, E. C., Gately, J. M., Hanson, A. D., Oldham, M. L., Sanders, D. A., & Csonka, L. N. (2012). Identification of a third osmoprotectant transport system, the OsmU system, in *Salmonella enterica*. *Journal of Bacteriology*, 194(15), 3861–3871. <https://doi.org/10.1128/JB.00495-12>
- Fuchs, J.-G., Moënne-Loccoz, Y., & Défago, G. (1997). Nonpathogenic *Fusarium oxysporum* strain Fo47 induces resistance to Fusarium wilt in tomato. *Plant Disease*, 81(5), 492–496. <https://doi.org/10.1094/PDIS.1997.81.5.492>
- Fuentes, A., Herrera, H., Charles, T. C., & Arriagada, C. (2020). Fungal and bacterial microbiome associated with the rhizosphere of native plants from the Atacama Desert. *Microorganisms*, 8(2), 1–14. <https://doi.org/10.3390/microorganisms8020209>
- Gahlot, D. K., Taheri, N., & MacIntyre, S. (2023). Diversity in genetic regulation of bacterial fimbriae assembled by the chaperone usher pathway. *International Journal of Molecular Sciences*, 24(1), 1–18. <https://doi.org/10.3390/ijms24010161>
- Gamalero, E., Lingua, G., Giusy Capri, F., Fusconi, A., Berta, G., & Lemanceau, P. (2004). Colonization pattern of primary tomato roots by *Pseudomonas fluorescens* A6RI characterized by dilution plating, flow cytometry, fluorescence, confocal and scanning electron microscopy. *FEMS Microbiology Ecology*, 48(1), 79–87. <https://doi.org/10.1016/j.femsec.2003.12.012>
- Ganal, M. W. (2013). Molecular Markers, Genetic Maps and Association Studies in Tomato. In B. E. Liedl, J. A. Labate, J. R. Stommel, A. Slade, & C. Kole (Eds.), *Genetics, Genomics, and Breeding of Tomato*. CRC Press.
- Gao, L., Gonda, I., Sun, H., Ma, Q., Bao, K., Tieman, D. M., Burzynski-Chang, E. A., Fish, T. L., Stromberg, K. A., Sacks, G. L., Thannhauser, T. W., Foolad, M. R., Diez, M. J., Blanca, J., Canizares, J., Xu, Y., van der Knaap, E., Huang, S., Klee, H. J., ... Fei, Z. (2019). The tomato pan-genome uncovers new genes and a rare allele regulating fruit flavor. *Nature Genetics*, 51(6), 1044–1051. <https://doi.org/10.1038/s41588-019-0410-2>
- Gao, S., Wu, H., Yu, X., Qian, L., & Gao, X. (2016). Swarming motility plays the major role in migration during tomato root colonization by *Bacillus subtilis* SWR01. *Biological Control*, 98, 11–17. <https://doi.org/10.1016/j.biocontrol.2016.03.011>
- García, J., Moravek, M., Fish, T., Thannhauser, T., Fei, Z., Sparks, J. P., Giovannoni, J., & Kao-Kniffin, J. (2024). Rhizosphere microbiomes derived from vermicompost alter gene expression and regulatory pathways in tomato (*Solanum lycopersicum*, L.). *Scientific Reports*, 14(21362), 1–12. <https://doi.org/10.1038/s41598-024-71792-7>
- García-Villaraco, A., Boukerma, L., Lucas, J. A., Gutierrez-Mañero, F. J., & Ramos-Solano, B. (2021). Tomato bio-protection induced by *Pseudomonas fluorescens* N21.4 involves ROS scavenging enzymes and PRs, without compromising plant growth. *Plants*, 10(2), 1–14. <https://doi.org/10.3390/plants10020331>

- Gardner, J. G. (2014). Systems biology defines the biological significance of redox-active proteins during cellulose degradation in an aerobic bacterium. *Molecular Microbiology*, *94*(5), 1121–1133. <https://doi.org/10.1111/mmi.12821>
- Garza Arizpe, M., & Molina Velázquez, M. (2008). Control climático en invernadero. In *Manual para la producción de tomate en invernadero en suelo en el estado de Nuevo León*. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación.
- Geibel, S., & Waksman, G. (2014). The molecular dissection of the chaperone–usher pathway. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, *1843*(8), 1559–1567. <https://doi.org/10.1016/j.bbamcr.2013.09.023>
- Geng, J., Liao, P., Tan, G. Y. A., Zhu, F.-Y., & Pradhan, N. (2023). Opportunities and challenges for n-alkane and n-alkene biosynthesis: A sustainable microbial biorefinery. *Biofuel Research Journal*, *10*(4), 1974–1988. <https://doi.org/10.18331/BRJ2023.10.4.4>
- Gherbawy, Y. A. M. H., & Abdelzaher, H. M. A. (1999). Isolation of fungi from tomato rhizosphere and evaluation of the effect of some fungicides and biological agents on the production of cellulase enzymes by *Nectria haematococca* and *Pythium ultimum* var. *ultimum*. *Czech Mycology*, *51*(2–3), 157–170.
- Giessen, T. W. (2022). Encapsulins. *Annual Review of Biochemistry*, *91*, 353–380. <https://doi.org/10.1146/annurev-biochem-040320-102858>
- Glick, B. R. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiological Research*, *169*(1), 30–39. <https://doi.org/10.1016/j.micres.2013.09.009>
- Goldstein, A. H. (1995). Recent progress in understanding the molecular genetics and biochemistry of calcium phosphate solubilization by gram negative bacteria. *Biological Agriculture & Horticulture*, *12*(2), 185–193. <https://doi.org/10.1080/01448765.1995.9754736>
- Gómez Expósito, R., de Bruijn, I., Postma, J., & Raaijmakers, J. M. (2017). Current insights into the role of rhizosphere bacteria in disease suppressive soils. *Frontiers in Microbiology*, *8*, 1–12. <https://www.frontiersin.org/articles/10.3389/fmicb.2017.02529>
- Gopal, M., & Gupta, A. (2016). Microbiome selection could spur next-generation plant breeding strategies. *Frontiers in Microbiology*, *7*, 1–10. <https://doi.org/10.3389/fmicb.2016.01971>
- Gran Diccionario Náhuatl. (2012). *Tomatl*. <https://gdn.iib.unam.mx/diccionario/tomatl/74196>
- Grandillo, S., Chetelat, R., Knapp, S., Spooner, D., Peralta, I., Cammareri, M., Perez, O., Termolino, P., Tripodi, P., Chiusano, M. L., Ercolano, M. R., Frusciantè, L., Monti, L., & Pignone, D. (2011). *Solanum* sect. *Lycopersicon*. In C. Kole (Ed.), *Wild Crop Relatives: Genomic and Breeding Resources: Vegetables* (pp. 129–215). Springer. [https://doi.org/10.1007/978-3-642-20450-0\\_9](https://doi.org/10.1007/978-3-642-20450-0_9)
- Grandillo, S., Termolino, P., & Knaap, E. van der. (2013). Molecular Mapping of Complex Traits in Tomato. In B. E. Liedl, J. A. Labate, J. R. Stommel, A. Slade, & C. Kole (Eds.), *Genetics, Genomics, and Breeding of Tomato*. CRC Press.
- Gruber, K. (2017). Agrobiodiversity: The living library. *Nature*, *544*, 8–10. <https://doi.org/10.1038/544S8a>
- Gu, S. (2020). Competition for iron drives phytopathogen control by natural rhizosphere microbiomes. *Nature Microbiology*, *5*, 1002–1010. <https://doi.org/10.1038/s41564-020-0719-8>
- Gu, Y., Banerjee, S., Dini-Andreote, F., Xu, Y., Shen, Q., Jousset, A., & Wei, Z. (2022). Small changes in rhizosphere microbiome composition predict disease outcomes earlier than pathogen density variations. *The ISME Journal*, *16*(10), 1–9. <https://doi.org/10.1038/s41396-022-01290-z>
- Guerry, P. (2007). *Campylobacter* flagella: Not just for motility. *Trends in Microbiology*, *15*(10), 456–461. <https://doi.org/10.1016/j.tim.2007.09.006>
- Gutierrez, A., & Grillo, M. A. (2022). Effects of domestication on plant-microbiome interactions. *Plant & Cell Physiology*, *63*(11), 1654–1666. <https://doi.org/10.1093/pcp/pcac108>

- Hale, M. G., Foy, C. L., & Shay, F. J. (1971). Factors affecting root exudation. *Advances in Agronomy*, 23, 89–109. [https://doi.org/10.1016/S0065-2113\(08\)60151-0](https://doi.org/10.1016/S0065-2113(08)60151-0)
- Halo, B. A., Al-Yahyai, R. A., & Al-Sadi, A. M. (2020). An endophytic *Talaromyces omanensis* enhances reproductive, physiological and anatomical characteristics of drought-stressed tomato. *Journal of Plant Physiology*, 249, 1–13. <https://doi.org/10.1016/j.jplph.2020.153163>
- Hanrahan-Tan, D. G., Lilje, O., & Henderson, L. (2023). Chytrids in soil environments: Unique adaptations and distributions. *Encyclopedia*, 3(2), 642–664. <https://doi.org/10.3390/encyclopedia3020046>
- Hao, T., Xie, Z., Wang, M., Liu, L., Zhang, Y., Wang, W., Zhang, Z., Zhao, X., Li, P., Guo, Z., Gao, S., Lou, C., Zhang, G., Merritt, J., Horsman, G. P., & Chen, Y. (2019). An anaerobic bacterium host system for heterologous expression of natural product biosynthetic gene clusters. *Nature Communications*, 10(1), 1–13. <https://doi.org/10.1038/s41467-019-11673-0>
- Harlan, J. R., de Wet, J. M. J., & Price, E. G. (1973). Comparative evolution of cereals. *Evolution*, 27(2), 311–325. <https://doi.org/10.2307/2406971>
- Hart, M., Ehret, D. L., Krumbein, A., Leung, C., Murch, S., Turi, C., & Franken, P. (2015). Inoculation with arbuscular mycorrhizal fungi improves the nutritional value of tomatoes. *Mycorrhiza*, 25(5), 359–376. <https://doi.org/10.1007/s00572-014-0617-0>
- Hassani, M. A., Özkurt, E., Franzenburg, S., & Stukenbrock, E. H. (2020). Ecological assembly processes of the bacterial and fungal microbiota of wild and domesticated wheat species. *Phytobiomes Journal*, 4(3), 217–224. <https://doi.org/10.1094/PBIOMES-01-20-0001-SC>
- Hayat, R., Ali, S., Amara, U., Khalid, R., & Ahmed, I. (2010). Soil beneficial bacteria and their role in plant growth promotion: A review. *Annals of Microbiology*, 60(4), 579–598. <https://doi.org/10.1007/s13213-010-0117-1>
- He, M.-Q., & Zhao, R.-L. (2021). Outline of Basidiomycota. In Ó. Zaragoza & A. Casadevall (Eds.), *Encyclopedia of Mycology* (pp. 310–319). Elsevier. <https://doi.org/10.1016/B978-0-12-819990-9.00065-2>
- Heaton, M. P., & Neuhaus, F. C. (1994). Role of the D-alanyl carrier protein in the biosynthesis of D-alanyl-lipoteichoic acid. *Journal of Bacteriology*, 176(3), 681–690. <https://doi.org/10.1128/jb.176.3.681-690.1994>
- Hernandez, L. M., Guzman, Y. C., Martínez-Arias, A., Manzano, M. R., & Selvaraj, J. J. (2015). The bud midge *Prodidiplosis longifila*: Damage characteristics, potential distribution and presence on a new crop host in Colombia. *SpringerPlus*, 4(1), 1–10. <https://doi.org/10.1186/s40064-015-0987-6>
- Herrou, J., Willett, J. W., Czyż, D. M., Babnigg, G., Kim, Y., & Crosson, S. (2017). Conserved ABC transport system regulated by the general stress response pathways of Alpha- and Gammaproteobacteria. *Journal of Bacteriology*, 199(5), 1–18. <https://doi.org/10.1128/JB.00746-16>
- Hewitt, S., Ghogare, R., Troxel, W., Tenic, E., Isaac, D., & Dhingra, A. (2023). Metatranscriptomic analysis of tomato rhizospheres reveals insight into plant-microbiome molecular response to biochar-amended organic soil. *Frontiers in Analytical Science*, 3, 1–15. <https://doi.org/10.3389/frans.2023.1205583>
- Holland-Moritz, H., Oliverio, A., Walsh, C., Gebert, M., & Fan, K. (2021). *Dada2 tutorial with NovaSeq dataset for Ernakovich Lab*. GitHub. [https://github.com/ErnakovichLab/dada2\\_ernakovichlab](https://github.com/ErnakovichLab/dada2_ernakovichlab)
- Holtsmark, I., Eijsink, V. G. H., & Brurberg, M. B. (2008). Bacteriocins from plant pathogenic bacteria. *FEMS Microbiology Letters*, 280(1), 1–7. <https://doi.org/10.1111/j.1574-6968.2007.01010.x>
- Horton, M. W. (2014). Genome-wide association study of *Arabidopsis thaliana* leaf microbial community. *Nature Communications*, 5, 1–7. <https://doi.org/10.1038/ncomms6320>
- Howe, H. F., & Smallwood, J. (1982). Ecology of Seed Dispersal. *Annual Review of Ecology and Systematics*, 13, 201–228.

- Hozzein, W. (2017, May). *Insecticidal Activities of Newly Isolated Actinomycetes Against the Rift Valley Fever Mosquito Aedes aegyptii*. Proceedings of International Academic Conferences, International Institute of Social and Economic Sciences, London. <https://ideas.repec.org/p/sek/iacpro/5107207.html>
- Hu, J., Wei, Z., Friman, V.-P., Gu, S., Wang, X., Eisenhauer, N., Yang, T., Ma, J., Shen, Q., Xu, Y., & Jousset, A. (2016). Probiotic diversity enhances rhizosphere microbiome function and plant disease suppression. *mBio*, 7(6), 1–8. <https://doi.org/10.1128/mBio.01790-16>
- Huang, J., Li, Y., Ma, Y., Li, Y., Jin, J., & Lian, T. (2022). The rhizospheric microbiome becomes more diverse with maize domestication and genetic improvement. *Journal of Integrative Agriculture*, 21(4), 1188–1202. [https://doi.org/10.1016/S2095-3119\(21\)63633-X](https://doi.org/10.1016/S2095-3119(21)63633-X)
- Huang, M. (2012). The major volatile organic compound emitted from *Arabidopsis thaliana* flowers, the sesquiterpene (E)- $\beta$ -caryophyllene, is a defense against a bacterial pathogen. *New Phytologist*, 193, 997–1008. <https://doi.org/10.1111/j.1469-8137.2011.04001.x>
- Hubbard, C. J., Li, B., McMinn, R., Brock, M. T., Maignien, L., Ewers, B. E., Kliebenstein, D., & Weinig, C. (2019). The effect of rhizosphere microbes outweighs host plant genetics in reducing insect herbivory. *Molecular Ecology*, 28(7), 1801–1811. <https://doi.org/10.1111/mec.14989>
- Huccetogullari, D., Luo, Z. W., & Lee, S. Y. (2019). Metabolic engineering of microorganisms for production of aromatic compounds. *Microbial Cell Factories*, 18(1), 1–29. <https://doi.org/10.1186/s12934-019-1090-4>
- Huerta-Cepas, J., Szklarczyk, D., Forslund, K., Cook, H., Heller, D., Walter, M. C., Rattei, T., Mende, D. R., Sunagawa, S., Kuhn, M., Jensen, L. J., von Mering, C., & Bork, P. (2016). eggNOG 4.5: A hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. *Nucleic Acids Research*, 44(D1), D286–293. <https://doi.org/10.1093/nar/gkv1248>
- Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal: Prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*, 11(119), 1–11. <https://doi.org/10.1186/1471-2105-11-119>
- Imlay, J. A. (2015). Transcription factors that defend Bacteria against reactive oxygen species. *Annual Review of Microbiology*, 69, 93–108. <https://doi.org/10.1146/annurev-micro-091014-104322>
- Islam, M. A., Islam, S., Akter, A., Rahman, M. H., & Nandwani, D. (2017). Effect of organic and inorganic fertilizers on soil properties and the growth, yield and quality of tomato in Mymensingh, Bangladesh. *Agriculture*, 7(3), 1–7. <https://doi.org/10.3390/agriculture7030018>
- Janahiraman, V., Anandham, R., Kwon, S. W., Sundaram, S., Karthik Pandi, V., Krishnamoorthy, R., Kim, K., Samaddar, S., & Sa, T. (2016). Control of wilt and rot pathogens of tomato by antagonistic pink pigmented facultative methylotrophic *Delftia lacustris* and *Bacillus* spp. *Frontiers in Plant Science*, 7, 1–11. <https://www.frontiersin.org/articles/10.3389/fpls.2016.01626>
- Jang, M.-S., Mouri, Y., Uchida, K., Aizawa, S.-I., Hayakawa, M., Fujita, N., Tezuka, T., & Ohnishi, Y. (2016). Genetic and transcriptional analyses of the flagellar gene cluster in *Actinoplanes missouriensis*. *Journal of Bacteriology*, 198(16), 1–9. <https://doi.org/10.1128/JB.00306-16>
- Jeanbille, M., Buée, M., Bach, C., Cébron, A., Frey-Klett, P., Turpault, M. P., & Uroz, S. (2016). Soil parameters drive the structure, diversity and metabolic potentials of the bacterial communities across temperate beech forest soil sequences. *Microbial Ecology*, 71(2), 482–493. <https://doi.org/10.1007/s00248-015-0669-5>
- Jenkins, J. A. (1948). The origin of the cultivated tomato. *Economic Botany*, 2(4), 379–392.
- Kahlon, P. S., Seta, S. M., Zander, G., Scheickl, D., Hückelhoven, R., Joosten, M. H. A. J., & Stam, R. (2020). Population studies of the wild tomato species *Solanum chilense* reveal geographically structured major gene-mediated pathogen resistance. *Proceedings of the Royal Society B: Biological Sciences*, 287(1941), 1–9. <https://doi.org/10.1098/rspb.2020.2723>

- Kalidasan, V., Azman, A., Joseph, N., Kumar, S., Awang Hamat, R., & Neela, V. K. (2018). Putative iron acquisition systems in *Stenotrophomonas maltophilia*. *Molecules*, *23*(8), 1–19. <https://doi.org/10.3390/molecules23082048>
- Kamilova, F. (2006). Organic acids, sugars, and L-tryptophane in exudates of vegetables growing on stone-wool and their effects on activities of rhizosphere bacteria. *MPMI*, *19*(3), 250–256. <https://doi.org/10.1094/MPMI-19-0250>
- Kandasamy, S., Weerasuriya, N., White, J. F., Patterson, G., & Lazarovits, G. (2021). Chapter 23—Soil’s physical and nutritional balance is essential for establishing a healthy microbiome. In J. White, A. Kumar, & S. Drobny (Eds.), *Microbiome Stimulants for Crops* (pp. 381–404). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-822122-8.00004-2>
- Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, *28*(1), 27–30. <https://doi.org/10.1093/nar/28.1.27>
- Kang, D. D., Froula, J., Egan, R., & Wang, Z. (2015). MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ*, *3*, 1–15. <https://doi.org/10.7717/peerj.1165>
- Kang, D. D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., & Wang, Z. (2019). MetaBAT 2: An adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ*, *7*, 1–13. <https://doi.org/10.7717/peerj.7359>
- Kannoja, P., Choudhary, K. K., Srivastava, A. K., & Singh, A. K. (2019). Chapter Four - PGPR Bioelicitors: Induced Systemic Resistance (ISR) and Proteomic Perspective on Biocontrol. In A. K. Singh, A. Kumar, & P. K. Singh (Eds.), *PGPR Amelioration in Sustainable Agriculture* (pp. 67–84). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-815879-1.00004-5>
- Karuppiyah, V., Natarajan, S., Gangatharan, M., Aldayel, M. F., Alsowayeh, N., & Thangavel, K. (2022). Development of siderophore-based rhizobacterial consortium for the mitigation of biotic and abiotic environmental stresses in tomatoes: An in vitro and in planta approach. *Journal of Applied Microbiology*, *133*(6), 3276–3287. <https://doi.org/10.1111/jam.15625>
- Kasotia, A., & Kumar Choudhary, D. (2014). Role of endophytic microbes in mitigation of abiotic stress in plants. In P. Ahmad & S. Rasool (Eds.), *Emerging Technologies and Management of Crop Stress Tolerance A Sustainable Approach* (Vol. 2, pp. 97–108).
- Kaur, T., Vasudev, A., Sohal, S. K., & Manhas, R. K. (2014). Insecticidal and growth inhibitory potential of *Streptomyces hydrogenans* DH16 on major pest of India, *Spodoptera litura* (Fab.) (Lepidoptera: Noctuidae). *BMC Microbiology*, *14*(227), 1–9. <https://doi.org/10.1186/s12866-014-0227-1>
- Kazerooni, E. A., Maharachchikumbura, S. S. N., Rethinasamy, V., Al-Mahrouqi, H., & Al-Sadi, A. M. (2017). Fungal diversity in tomato rhizosphere soil under conventional and desert farming systems. *Frontiers in Microbiology*, *8*, 1–8. <https://doi.org/10.3389/fmicb.2017.01462>
- Keoke, E. D., & Porterfield, K. M. (2009). *Encyclopedia of American Indian Contributions to the World: 15,000 Years of Inventions and Innovations*. Infobase Publishing.
- Khan, N. (2012). Exploring the natural variation for seedling traits and their link with seed dimensions in tomato. *PLoS One*, *7*(8), 1–14. <https://doi.org/10.1371/journal.pone.0043991>
- Kim, J.-S., Dungan, R. S., Kwon, S.-W., & Weon, H.-Y. (2006). The community composition of root-associated bacteria of the tomato plant. *World Journal of Microbiology and Biotechnology*, *22*(12), 1267–1273. <https://doi.org/10.1007/s11274-006-9171-z>
- Kim, K., Jang, Y.-J., Lee, S.-M., Oh, B.-T., Chae, J.-C., & Lee, K.-J. (2014). Alleviation of salt stress by *Enterobacter* sp. EJ01 in tomato and *Arabidopsis* is accompanied by up-regulation of conserved salinity responsive factors in plants. *Molecules and Cells*, *37*(2), 109–117. <https://doi.org/10.14348/molcells.2014.2239>

- Kimura, S., & Sinha, N. (2008). Tomato (*Solanum lycopersicum*): A model fruit-bearing crop. *Cold Spring Harbor Protocols*, 3(11), 1–10. <https://doi.org/10.1101/pdb.emo105>
- Kimura, T., Tezuka, T., Nakane, D., Nishizaka, T., Aizawa, S.-I., & Ohnishi, Y. (2019). Characterization of zoospore Type IV pili in *Actinoplanes missouriensis*. *Journal of Bacteriology*, 201(14), 1–12. <https://doi.org/10.1128/JB.00746-18>
- Klee, H. J., & Resende, M. F. R. (2020). Plant Domestication: Reconstructing the Route to Modern Tomatoes. *Current Biology*, 30(8), 359–361. <https://doi.org/10.1016/j.cub.2020.02.072>
- Knapp, S., & Peralta, I. E. (2016). The Tomato (*Solanum lycopersicum* L., Solanaceae) and Its Botanical Relatives. In M. Causse, J. Giovannoni, M. Bouzayen, & M. Zouine (Eds.), *The Tomato Genome* (pp. 7–21). Springer. [https://doi.org/10.1007/978-3-662-53389-5\\_2](https://doi.org/10.1007/978-3-662-53389-5_2)
- Knights, H. E., Jorriin, B., Haskett, T. L., & Poole, P. S. (2021). Deciphering bacterial mechanisms of root colonization. *Environmental Microbiology Reports*, 13(4), 428–444. <https://doi.org/10.1111/1758-2229.12934>
- Kodani, S., Lodato, M. A., Durrant, M. C., Picart, F., & Willey, J. M. (2005). SapT, a lanthionine-containing peptide involved in aerial hyphae formation in the streptomycetes. *Molecular Microbiology*, 58(5), 1368–1380. <https://doi.org/10.1111/j.1365-2958.2005.04921.x>
- Köhler, H.-R., & Triebskorn, R. (2013). Wildlife ecotoxicology of pesticides: Can we track effects to the population level and beyond? *Science*, 341(6147), 759–765. <https://doi.org/10.1126/science.1237591>
- Kolde, R. (2018). *pheatmap: Pretty Heatmaps* (Version 1.0.12) [Computer software]. <https://github.com/raivokolde/pheatmap>
- Kong, H. G., Song, G. C., Sim, H.-J., & Ryu, C.-M. (2021). Achieving similar root microbiota composition in neighbouring plants through airborne signalling. *The ISME Journal*, 15, 397–408. <https://doi.org/10.1038/s41396-020-00759-z>
- Kortbeek, R. W. J., Galland, M. D., Muras, A., van der Kloet, F. M., André, B., Heilijgers, M., van Hijum, S. A. F. T., Haring, M. A., Schuurink, R. C., & Bleeker, P. M. (2021). Natural variation in wild tomato trichomes; selecting metabolites that contribute to insect resistance using a random forest approach. *BMC Plant Biology*, 21(1), 1–19. <https://doi.org/10.1186/s12870-021-03070-x>
- Krügel, H., Krubasik, P., Weber, K., Saluz, H. P., & Sandmann, G. (1999). Functional analysis of genes from *Streptomyces griseus* involved in the synthesis of isorenieratene, a carotenoid with aromatic end groups, revealed a novel type of carotenoid desaturase. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, 1439(1), 57–64. [https://doi.org/10.1016/S1388-1981\(99\)00075-X](https://doi.org/10.1016/S1388-1981(99)00075-X)
- Kusstatscher, P., Wicaksono, W. A., Bergna, A., Cernava, T., Bergau, N., Tissier, A., Hause, B., & Berg, G. (2020). Trichomes form genotype-specific microbial hotspots in the phyllosphere of tomato. *Environmental Microbiome*, 15(1), 1–10. <https://doi.org/10.1186/s40793-020-00364-9>
- Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 9(4), 357–359. <https://doi.org/10.1038/nmeth.1923>
- Laslett, D., & Canback, B. (2004). ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Research*, 32(1), 11–16. <https://doi.org/10.1093/nar/gkh152>
- Lázaro, A. (2018). Tomato landraces: An analysis of diversity and preferences. *Plant Genetic Resources*, 16(4), 315–324. <https://doi.org/10.1017/S1479262117000351>
- Le, K. D., Kim, J., Nguyen, H. T., Yu, N. H., Park, A. R., Lee, C. W., & Kim, J.-C. (2021). *Streptomyces* sp. JCK-6131 protects plants against bacterial and fungal diseases via two mechanisms. *Frontiers in Plant Science*, 12, 1–13 <https://www.frontiersin.org/articles/10.3389/fpls.2021.726266>
- Lee Díaz, A. S., Rizaludin, M. S., Zweers, H., Raaijmakers, J. M., & Garbeva, P. (2022). Exploring the volatiles released from roots of wild and domesticated tomato plants under insect attack. *Molecules*, 27(5), 1–14. <https://doi.org/10.3390/molecules27051612>

- Lee, M. (2006). The Solanaceae: Foods and poisons. *The Journal of the Royal College of Physicians of Edinburgh*, 36(2), 162–169.
- Lee, S. A., Kim, Y., Kim, J. M., Chu, B., Joa, J.-H., Sang, M. K., Song, J., & Weon, H.-Y. (2019). A preliminary examination of bacterial, archaeal, and fungal communities inhabiting different rhizo-compartments of tomato plants under real-world environments. *Scientific Reports*, 9(9300), 1–15. <https://doi.org/10.1038/s41598-019-45660-8>
- Lee, S. A., Park, J., Chu, B., Kim, J. M., Joa, J.-H., Sang, M. K., Song, J., & Weon, H.-Y. (2016). Comparative analysis of bacterial diversity in the rhizosphere of tomato by culture-dependent and -independent approaches. *Journal of Microbiology*, 54(12), 823–831. <https://doi.org/10.1007/s12275-016-6410-3>
- Lee, S.-M., Kong, H. G., Song, G. C., & Ryu, C.-M. (2021). Disruption of Firmicutes and Actinobacteria abundance in tomato rhizosphere causes the incidence of bacterial wilt disease. *The ISME Journal*, 15(1), 330–347. <https://doi.org/10.1038/s41396-020-00785-x>
- Leo, J. C., Grin, I., & Linke, D. (2012). Type V secretion: Mechanism(s) of autotransport through the bacterial outer membrane. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1592), 1088–1101. <https://doi.org/10.1098/rstb.2011.0208>
- Leventis, G., Tsiknia, M., Feka, M., Ladikou, E. V., Papadakis, I. E., Chatzipavlidis, I., Papadopoulou, K., & Ehaliotis, C. (2021). Arbuscular mycorrhizal fungi enhance growth of tomato under normal and drought conditions, via different water regulation mechanisms. *Rhizosphere*, 19, 1–12. <https://doi.org/10.1016/j.rhisph.2021.100394>
- Levy, A., Salas Gonzalez, I., Mittelviehhaus, M., Clingenpeel, S., Herrera Paredes, S., Miao, J., Wang, K., Devescovi, G., Stillman, K., Monteiro, F., Rangel Alvarez, B., Lundberg, D. S., Lu, T.-Y., Lebeis, S., Jin, Z., McDonald, M., Klein, A. P., Felcher, M. E., Rio, T. G., ... Dangel, J. L. (2018). Genomic features of bacterial adaptation to plants. *Nature Genetics*, 50(1), 138–150. <https://doi.org/10.1038/s41588-017-0012-9>
- Lex, A., Gehlenborg, N., Strobel, H., Vuillemot, R., & Pfister, H. (2014). UpSet: Visualization of Intersecting Sets. *IEEE Transactions on Visualization and Computer Graphics*, 20(12), 1983–1992. <https://doi.org/10.1109/TVCG.2014.2346248>
- Li, C., Alam, K., Zhao, Y., Hao, J., Yang, Q., Zhang, Y., Li, R., & Li, A. (2021). Mining and biosynthesis of bioactive lanthipeptides from microorganisms. *Frontiers in Bioengineering and Biotechnology*, 9, 1–13. <https://doi.org/10.3389/fbioe.2021.692466>
- Li, D. (2016). MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced methodologies and community practices. *Methods*, 102, 3–11. <https://doi.org/10.1016/j.jmeth.2016.02.020>
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: An ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, 31(10), 1674–1676. <https://doi.org/10.1093/bioinformatics/btv033>
- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*, 2013, 1–3. <https://doi.org/10.48550/arXiv.1303.3997>
- Li, J., Wang, J., Liu, H., Macdonald, C. A., & Singh, B. K. (2022). Application of microbial inoculants significantly enhances crop productivity: A meta-analysis of studies from 2010 to 2020. *Journal of Sustainable Agriculture and Environment*, 1, 216–225. <https://doi.org/10.1002/sae2.12028>
- Li, M., Pommier, T., Yin, Y., Wang, J., Gu, S., Jousset, A., Keuskamp, J., Wang, H., Wei, Z., Xu, Y., Shen, Q., & Kowalchuk, G. A. (2022). Indirect reduction of *Ralstonia solanacearum* via pathogen helper inhibition. *The ISME Journal*, 16(3), 867–875. <https://doi.org/10.1038/s41396-021-01126-2>
- Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, 22(13), 1658–1659. <https://doi.org/10.1093/bioinformatics/btl158>

- Li, Y., & Chen, X. (2012). Sialic acid metabolism and sialyltransferases: Natural functions and applications. *Applied Microbiology and Biotechnology*, *94*(4), 887–905. <https://doi.org/10.1007/s00253-012-4040-1>
- Lima, G. P. P., Gómez, H. A. G., Seabra Junior, S., Maraschin, M., Tecchio, M. A., & Borges, C. V. (2022). Functional and nutraceutical compounds of tomatoes as affected by agronomic practices, postharvest management, and processing methods: A mini review. *Frontiers in Nutrition*, *9*, 1–7. <https://doi.org/10.3389/fnut.2022.868492>
- Lin, J. C.-Y., Chiang, B.-Y., Chou, C.-C., Chen, T.-C., Chen, Y.-J., Chen, Y.-J., & Lin, C.-H. (2015). Glutathionylspermidine in the modification of protein SH groups: The enzymology and its application to study protein glutathionylation. *Molecules*, *20*(1), 1452–1474. <https://doi.org/10.3390/molecules20011452>
- Lin, T., Zhu, G., Zhang, J., Xu, X., Yu, Q., Zheng, Z., Zhang, Z., Lun, Y., Li, S., Wang, X., Huang, Z., Li, J., Zhang, C., Wang, T., Zhang, Y., Wang, A., Zhang, Y., Lin, K., Li, C., ... Huang, S. (2014). Genomic analyses provide insights into the history of tomato breeding. *Nature Genetics*, *46*(11), 1220–1226. <https://doi.org/10.1038/ng.3117>
- Lin, Y.-P., Lu, C.-Y., & Lee, C.-R. (2020). The climatic association of population divergence and future extinction risk of *Solanum pimpinellifolium*. *AoB Plants*, *12*(2), 1–13. <https://doi.org/10.1093/aobpla/plaa012>
- Ling, H.-Q., Bauer, P., Bereczky, Z., Keller, B., & Ganai, M. (2002). The tomato *fer* gene encoding a bHLH protein controls iron-uptake responses in roots. *PNAS*, *99*(21), 1–6. <https://doi.org/10.1073/pnas.212448699>
- Ling, L., Han, X., Li, X., Zhang, X., Wang, H., Zhang, L., Cao, P., Wu, Y., Wang, X., Zhao, J., & Xiang, W. (2020). A *Streptomyces* sp. NEAU-HV9: isolation, identification, and potential as a biocontrol agent against *Ralstonia solanacearum* of tomato plants. *Microorganisms*, *8*(3), 1–15. <https://doi.org/10.3390/microorganisms8030351>
- Ling, N., Wang, T., & Kuzyakov, Y. (2022). Rhizosphere bacteriome structure and functions. *Nature Communications*, *13*(836), 1–13. <https://doi.org/10.1038/s41467-022-28448-9>
- Ling, S., Zhao, Y., Sun, S., Zheng, D., Sun, X., Zeng, R., Chen, D., & Song, Y. (2022). Enhanced anti-herbivore defense of tomato plants against *Spodoptera litura* by their rhizosphere bacteria. *BMC Plant Biology*, *22*(254), 1–10. <https://doi.org/10.1186/s12870-022-03644-3>
- Liu, S., Wang, H., Tian, P., Yao, X., Sun, H., Wang, Q., & Delgado-Baquerizo, M. (2020). Decoupled diversity patterns in bacteria and fungi across continental forest ecosystems. *Soil Biology and Biochemistry*, *144*, 1–9. <https://doi.org/10.1016/j.soilbio.2020.107763>
- Liu, Y., Xu, Z., Chen, L., Xun, W., Shu, X., Chen, Y., Sun, X., Wang, Z., Ren, Y., Shen, Q., & Zhang, R. (2024). Root colonization by beneficial rhizobacteria. *FEMS Microbiology Reviews*, *48*(1), 1–20. <https://doi.org/10.1093/femsre/fuad066>
- Liu, Z. (2016). Identification of a *Solanum pennellii* chromosome 4 fruit flavor and nutritional quality-associated metabolite QTL. *Frontiers in Plant Science*, *7*, 1–15. <https://doi.org/10.3389/fpls.2016.01671>
- Lloréns-Rico, V., Vieira-Silva, S., Gonçalves, P. J., Falony, G., & Raes, J. (2021). Benchmarking microbiome transformations favors experimental quantitative approaches to address compositionality and sampling depth biases. *Nature Communications*, *12*, 1–12. <https://doi.org/10.1038/s41467-021-23821-6>
- Long, J. (2022). De tomates y jitomates en el siglo XVI. *Estudios de Cultura Náhuatl*, *25*, 239–252.
- Longoria-Espinoza, R. M., Félix-Gastélum, R., Cordero-Ramírez, J. D., Longoria-Espinoza, R. M., Félix-Gastélum, R., & Cordero-Ramírez, J. D. (2020). Diversity of endophytic bacteria associated with

- tomato plants (*Solanum lycopersicum*). *Revista Mexicana de Fitopatología*, 38(2), 307–319. <https://doi.org/10.18781/r.mex.fit.2002-7>
- Loper, J. E., & Buyer, J. S. (1991). Siderophores in microbial interactions on plant surfaces. *Molecular Plant-Microbe Interactions*, 4(1), 5–13.
- Loper, J. E., & Henkels, M. D. (1999). Utilization of heterologous siderophores enhances levels of iron available to *Pseudomonas putida* in the rhizosphere. *Applied and Environmental Microbiology*, 65(12), 5357–5363.
- López Marín, L. M. P. R. de I. e I. por C. de V. A. (PRIICA). (2017). *Manual técnico del cultivo del tomate: Solanum lycopersicum*. Instituto Nacional de Innovación y Transferencia en Tecnología Agropecuaria (INTA). <https://repositorio.iica.int/handle/11324/3143>
- López, J. L., Fourie, A., Poppeliers, S. W. M., Pappas, N., Sánchez-Gil, J. J., de Jonge, R., & Dutilh, B. E. (2023). Growth rate is a dominant factor predicting the rhizosphere effect. *The ISME Journal*, 17(9), 1396–1405. <https://doi.org/10.1038/s41396-023-01453-6>
- López, S. M. Y., Pastorino, G. N., Fernández-González, A. J., Franco, M. E. E., Fernández-López, M., & Balatti, P. A. (2020). The endosphere bacteriome of diseased and healthy tomato plants. *Archives of Microbiology*, 202(10), 2629–2642. <https://doi.org/10.1007/s00203-020-01987-9>
- Lopez-Delacalle, M. (2019). Using tomato recombinant lines to improve plant tolerance to stress combination through a more efficient nitrogen metabolism. *Frontiers in Plant Science*, 10, 1–19. <https://doi.org/10.3389/fpls.2019.01702>
- Lorenz, K., & Cohen, B. A. (2012). Small- and large-effect quantitative trait locus interactions underlie variation in yeast sporulation efficiency. *Genetics*, 192(3), 1123–1132. <https://doi.org/10.1534/genetics.112.143107>
- Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12), 1–21. <https://doi.org/10.1186/s13059-014-0550-8>
- Lu, Y., Yang, Y., Liu, L., Yu, T., Zhao, J., Liu, L., & Li, C. (2021). Anti-adhesive effects of sialic acid and *Lactobacillus plantarum* on *Staphylococcus aureus* in vitro. *Journal of Food Safety*, 41(1), 1–11. <https://doi.org/10.1111/jfs.12875>
- Lucero, C. T., Lorda, G. S., Ludueña, L. M., Anzuay, M. S., & Taurian, T. (2020). Motility and biofilm production involved in the interaction of phosphate solubilizing endophytic strains with peanut, maize and soybean plants. *Rhizosphere*, 15, 1–9. <https://doi.org/10.1016/j.rhisph.2020.100228>
- Lugtenberg, B. J. J., Dekkers, L., & Bloemberg, G. V. (2001). Molecular determinants of rhizosphere colonization by *Pseudomonas*. *Annual Review of Phytopathology*, 39(1), 461–490. <https://doi.org/10.1146/annurev.phyto.39.1.461>
- Lundberg, D. S., Lebeis, S. L., Paredes, S. H., Yourstone, S., Gehring, J., Malfatti, S., Tremblay, J., Engelbrekton, A., Kunin, V., Rio, T. G. del, Edgar, R. C., Eickhorst, T., Ley, R. E., Hugenholtz, P., Tringe, S. G., & Dangl, J. L. (2012). Defining the core *Arabidopsis thaliana* root microbiome. *Nature*, 488(7409), 86–90. <https://doi.org/10.1038/nature11237>
- Lunn, J. E., Delorge, I., Figueroa, C. M., Dijck, P., & Stitt, M. (2014). Trehalose metabolism in plants. *The Plant Journal*, 79, 544–567. <https://doi.org/10.1111/tpj.12509>
- Macheroux, P., Schmid, J., Amrhein, N., & Schaller, A. (1999). A unique reaction in a common pathway: Mechanism and function of chorismate synthase in the shikimate pathway. *Planta*, 207(3), 325–334. <https://doi.org/10.1007/s004250050489>
- Malacrino, A., Abdelfattah, A., Berg, G., Benitez, M.-S., Bennett, A. E., Böttner, L., Xu, S., & Schena, L. (2022). Exploring microbiomes for plant disease management. *Biological Control*, 169, 1–7. <https://doi.org/10.1016/j.biocontrol.2022.104890>

- Marasco, R., Rolli, E., Ettoumi, B., Vigani, G., Mapelli, F., Borin, S., Abou-Hadid, A. F., El-Beairy, U. A., Sorlini, C., Cherif, A., Zocchi, G., & Daffonchio, D. (2012). A drought resistance-promoting microbiome is selected by root system under desert farming. *PLOS ONE*, *7*(10), 1–14. <https://doi.org/10.1371/journal.pone.0048479>
- Marçais, G. (2018). MUMmer4: A fast and versatile genome alignment system. *PLoS Computational Biology*, *14*, 1–14. <https://doi.org/10.1371/journal.pcbi.1005944>
- Marín, O., González, B., & Poupin, M. J. (2021). From microbial dynamics to functionality in the rhizosphere: A systematic review of the opportunities with synthetic microbial communities. *Frontiers in Plant Science*, *12*, 1–12. <https://doi.org/10.3389/fpls.2021.650609>
- Martínez-Romero, E., Aguirre-Noyola, J. L., Taco-Taype, N., Martínez-Romero, J., & Zuñiga-Dávila, D. (2020). Plant microbiota modified by plant domestication. *Systematic and Applied Microbiology*, *43*(5), 1–9. <https://doi.org/10.1016/j.syapm.2020.126106>
- Masella, A. P., Bartram, A. K., Truszkowski, J. M., Brown, D. G., & Neufeld, J. D. (2012). PANDA-seq: Paired-end assembler for illumina sequences. *BMC Bioinformatics*, *13*(31), 1–7. <https://doi.org/10.1186/1471-2105-13-31>
- Masood, S., Zhao, X. Q., & Shen, R. F. (2020). *Bacillus pumilus* promotes the growth and nitrogen uptake of tomato plants under nitrogen fertilization. *Scientia Horticulturae*, *272*, 1–9. <https://doi.org/10.1016/j.scienta.2020.109581>
- Massaretto, I. L., Albaladejo, I., Purgatto, E., Flores, F. B., Plasencia, F., Egea-Fernández, J. M., Bolarin, M. C., & Egea, I. (2018). Recovering tomato landraces to simultaneously improve fruit yield and nutritional quality against salt stress. *Frontiers in Plant Science*, *9*, 1–17. <https://www.frontiersin.org/articles/10.3389/fpls.2018.01778>
- Mata-Nicolás, E., Montero-Pau, J., Gimeno-Paez, E., Garcia-Carpintero, V., Ziarsolo, P., Menda, N., Mueller, L. A., Blanca, J., Cañizares, J., van der Knaap, E., & Díez, M. J. (2020). Exploiting the diversity of tomato: The development of a phenotypically and genetically detailed germplasm collection. *Horticulture Research*, *7*(1), 1–14. <https://doi.org/10.1038/s41438-020-0291-7>
- Matilla, M. A., Espinosa-Urgel, M., Rodríguez-Herva, J. J., Ramos, J. L., & Ramos-González, M. I. (2007). Genomic analysis reveals the major driving forces of bacterial life in the rhizosphere. *Genome Biology*, *8*(9), 1–13. <https://doi.org/10.1186/gb-2007-8-9-r179>
- Matos, R. C., Schwarzer, M., Gervais, H., Courtin, P., Joncour, P., Gillet, B., Ma, D., Bulteau, A.-L., Martino, M. E., Hughes, S., Chapot-Chartier, M.-P., & Leulier, F. (2017). D-Alanylation of teichoic acids contributes to *Lactobacillus plantarum*-mediated *Drosophila* growth during chronic undernutrition. *Nature Microbiology*, *2*(12), 1635–1647. <https://doi.org/10.1038/s41564-017-0038-x>
- Matson, P. A., Parton, W. J., Power, A. G., & Swift, M. J. (1997). Agricultural intensification and ecosystem properties. *Science*, *277*(5325), 504–509. <https://doi.org/10.1126/science.277.5325.504>
- Maul, J. E., Buyer, J. S., Lehman, R. M., Culman, S., Blackwood, C. B., Roberts, D. P., Zasada, I. A., & Teasdale, J. R. (2014). Microbial community structure and abundance in the rhizosphere and bulk soil of a tomato cropping system that includes cover crops. *Applied Soil Ecology*, *77*, 42–50. <https://doi.org/10.1016/j.apsoil.2014.01.002>
- Mayak, S., Tirosh, T., & Glick, B. R. (2004). Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiology and Biochemistry: PPB*, *42*(6), 565–572. <https://doi.org/10.1016/j.plaphy.2004.05.009>
- McCue, G. A. (1952). The history of the use of the tomato: An annotated bibliography. *Annals of the Missouri Botanical Garden*, *39*(4), 289–348. <https://doi.org/10.2307/2399094>
- McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE*, *8*(4), 1–11. <https://doi.org/10.1371/journal.pone.0061217>

- Mechan Llontop, M. E., Tian, L., Sharma, P., Hefflin, L., Bernal-Galeano, V., Haak, D. C., Clarke, C. R., & Vinatzer, B. A. (2021). Experimental evidence pointing to rain as a reservoir of tomato phyllosphere microbiota. *Phytobiomes Journal*, *5*(4), 382–399. <https://doi.org/10.1094/PBIOMES-04-21-0025-R>
- Medema, M. H., Kottmann, R., Yilmaz, P., Cummings, M., Biggins, J. B., Blin, K., de Bruijn, I., Chooi, Y. H., Claesen, J., Coates, R. C., Cruz-Morales, P., Duddela, S., Dusterhus, S., Edwards, D. J., Fewer, D. P., Garg, N., Geiger, C., Gomez-Escribano, J. P., Greule, A., ... Glöckner, F. O. (2015). Minimum information about a biosynthetic gene cluster. *Nature Chemical Biology*, *11*(9), 625–631. <https://doi.org/10.1038/nchembio.1890>
- Mediavilla, D. (2020, January 9). El viaje de una fruta diminuta nacida en los Andes que conquistó el mundo. *El País*. [https://elpais.com/elpais/2020/01/09/ciencia/1578590635\\_817541.html](https://elpais.com/elpais/2020/01/09/ciencia/1578590635_817541.html)
- Mekureyaw, M. F., Pandey, C., Hennessy, R. C., Nicolaisen, M. H., Liu, F., Nybroe, O., & Roitsch, T. (2022). The cytokinin-producing plant beneficial bacterium *Pseudomonas fluorescens* G20-18 primes tomato (*Solanum lycopersicum*) for enhanced drought stress responses. *Journal of Plant Physiology*, *270*, 1–15. <https://doi.org/10.1016/j.jplph.2022.153629>
- Mellerowicz, E. J., & Gorshkova, T. A. (2012). Tensional stress generation in gelatinous fibres: A review and possible mechanism based on cell-wall structure and composition. *Journal of Experimental Botany*, *63*(2), 551–565. <https://doi.org/10.1093/jxb/err339>
- Mendes, R., Kruijt, M., Bruijn, I. de, Dekkers, E., Voort, M. van der, Schneider, J. H. M., Piceno, Y. M., DeSantis, T. Z., Andersen, G. L., Bakker, P. A. H. M., & Raaijmakers, J. M. (2011). Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science*, *332*(6033), 1097–1100. <https://doi.org/10.1126/science.1203980>
- Meshram, S., & Adhikari, T. B. (2024). Microbiome-mediated strategies to manage major soil-borne diseases of tomato. *Plants*, *13*(3), 1–23. <https://doi.org/10.3390/plants13030364>
- Meuskens, I., Saragliadis, A., Leo, J. C., & Linke, D. (2019). Type V Secretion Systems: An overview of passenger domain functions. *Frontiers in Microbiology*, *10*, 1–19. <https://doi.org/10.3389/fmicb.2019.01163>
- Meyer, R. S., DuVal, A. E., & Jensen, H. R. (2012). Patterns and processes in crop domestication: An historical review and quantitative analysis of 203 global food crops. *New Phytologist*, *196*, 29–48. <https://doi.org/10.1111/j.1469-8137.2012.04253.x>
- Mikheenko, A., Prijbelski, A., Saveliev, V., Antipov, D., & Gurevich, A. (2018). Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics*, *34*(13), 142–150. <https://doi.org/10.1093/bioinformatics/bty266>
- Mimmo, T., Del Buono, D., Terzano, R., Tomasi, N., Vigani, G., Crecchio, C., Pinton, R., Zocchi, G., & Cesco, S. (2014). Rhizospheric organic compounds in the soil–microorganism–plant system: Their role in iron availability. *European Journal of Soil Science*, *65*(5), 629–642. <https://doi.org/10.1111/ejss.12158>
- Mishra, A. K., & Baek, K.-H. (2021). Salicylic acid biosynthesis and metabolism: A divergent pathway for plants and bacteria. *Biomolecules*, *11*(5), 1–16. <https://doi.org/10.3390/biom11050705>
- Mo, Y., Bier, R., Li, X., Daniels, M., Smith, A., Yu, L., & Kan, J. (2024). Agricultural practices influence soil microbiome assembly and interactions at different depths identified by machine learning. *Communications Biology*, *7*(1), 1–16. <https://doi.org/10.1038/s42003-024-07059-8>
- Mohammed, A. F., Oloyede, A. R., & Odeseye, A. O. (2020). Biological control of bacterial wilt of tomato caused by *Ralstonia solanacearum* using *Pseudomonas* species isolated from the rhizosphere of tomato plants. *Archives of Phytopathology and Plant Protection*, *53*(1–2), 1–16. <https://doi.org/10.1080/03235408.2020.1715756>

- Mohandas, S. (1988). Nitrogen fixation in tomato (*Lycopersicon esculentum* Mill 'Pusa Ruby'). *Plant and Soil*, 107(2), 219–225. <https://doi.org/10.1007/BF02370550>
- Morales Palacio, N., Morales Astudillo, R., Milian Garcia, Y., Artiles Valor, A., & Espinosa Lopez, G. (2014). Genetic characterization of *Solanum pimpinellifolium* and *Solanum lycopersicum* var. *cerasi-forme* tomato populations in Ecuador. *Gene Conserve*, 13(50), 21–45.
- Morcillo, R. (2021). Plant transcriptome reprogramming and bacterial extracellular metabolites underlying tomato drought resistance triggered by a beneficial soil bacteria. *Metabolites*, 11(6), 1–18. <https://doi.org/10.3390/metabo11060369>
- Morella, N. M., Weng, F. C.-H., Joubert, P. M., Metcalf, C. J. E., Lindow, S., & Koskella, B. (2020). Successive passaging of a plant-associated microbiome reveals robust habitat and host genotype-dependent selection. *Proceedings of the National Academy of Sciences*, 117(2), 1148–1159. <https://doi.org/10.1073/pnas.1908600116>
- Moreno, M. M., & Moreno, A. (2008). Effect of different biodegradable and polyethylene mulches on soil properties and production in a tomato crop. *Scientia Horticulturae*, 116(3), 256–263. <https://doi.org/10.1016/j.scienta.2008.01.007>
- Morsy, M., Cleckler, B., & Armuelles-Millican, H. (2020). Fungal endophytes promote tomato growth and enhance drought and salt tolerance. *Plants*, 9(7), 1–19. <https://doi.org/10.3390/plants9070877>
- Mueller, L. A., Solow, T. H., Taylor, N., Skwarecki, B., Buels, R., Binns, J., Lin, C., Wright, M. H., Ahrens, R., Wang, Y., Herbst, E. V., Keyder, E. R., Menda, N., Zamir, D., & Tanksley, S. D. (2005). The SOL genomics network. A comparative resource for solanaceae biology and beyond. *Plant Physiology*, 138(3), 1310–1317. <https://doi.org/10.1104/pp.105.060707>
- Müller, T., & Ruppel, S. (2014). Progress in cultivation-independent phyllosphere microbiology. *FEMS Microbiology Ecology*, 87(1), 2–17. <https://doi.org/10.1111/1574-6941.12198>
- Muñoz, P. A., Arismendi, M. J., Cárdenas, S. F., Cifuentes Bachmann, D. E., Venegas, F. A., & Sepúlveda-Chavera, G. F. (2020). Diversity of culturable bacteria isolated from ancestral crops of Arica and Parinacota Region, Atacama Desert. *Antonie Van Leeuwenhoek*, 113(12), 2123–2137. <https://doi.org/10.1007/s10482-020-01482-7>
- Murgia, M., Fiamma, M., Barac, A., Deligios, M., Mazzarello, V., Paglietti, B., Cappuccinelli, P., Al-Qahatani, A., Squartini, A., Rubino, S., & Al-Ahdal, M. N. (2018). Biodiversity of fungi in hot desert sands. *MicrobiologyOpen*, 8(1), 1–10. <https://doi.org/10.1002/mbo3.595>
- Nakayasu, M., Takamatsu, K., Yazaki, K., & Sugiyama, A. (2023). Plant specialized metabolites in the rhizosphere of tomatoes: Secretion and effects on microorganisms. *Bioscience, Biotechnology, and Biochemistry*, 87(1), 13–20. <https://doi.org/10.1093/bbb/zbac181>
- Nascimento, F. X., Rossi, M. J., & Glick, B. R. (2018). Ethylene and 1-aminocyclopropane-1-carboxylate (ACC) in plant–bacterial interactions. *Frontiers in Plant Science*, 9(114), 1–17. <https://doi.org/10.3389/fpls.2018.00114>
- Nassal, D., Spohn, M., Eltlbany, N., Jacquioid, S., Smalla, K., Marhan, S., & Kandeler, E. (2018). Effects of phosphorus-mobilizing bacteria on tomato growth and soil microbial activity. *Plant and Soil*, 427(1), 17–37. <https://doi.org/10.1007/s11104-017-3528-y>
- Nassar, M. M., Hafez, S. T., Nagaty, I. M., & Khalaf, S. A. (1999). The insecticidal activity of Cyanobacteria against four insects, two of medical importance and two agricultural pests with reference to the action on albino mice. *Journal of the Egyptian Society of Parasitology*, 29(3), 939–949.
- Natsiopoulou, D., Tziolias, A., Lagogiannis, I., Mantzoukas, S., & Eliopoulos, P. A. (2022). Growth-promoting and protective effect of *Trichoderma atrobrunneum* and *T. simmonsii* on tomato against soil-borne fungal pathogens. *Crops*, 2(3), 202–217. <https://doi.org/10.3390/crops2030015>

- Naumova, N., Baturina, O., Nechaeva, T., & Kabilov, M. (2022). Root and rhizosphere microbiome of tomato plants grown in the open field in the south of West Siberia under mineral fertilization. *Horticulturae*, 8(11), 1–19. <https://doi.org/10.3390/horticulturae8111051>
- Nazari, M. T., Schommer, V. A., Braun, J. C. A., dos Santos, L. F., Lopes, S. T., Simon, V., Machado, B. S., Ferrari, V., Colla, L. M., & Piccin, J. S. (2023). Using *Streptomyces* spp. As plant growth promoters and biocontrol agents. *Rhizosphere*, 27, 1–15. <https://doi.org/10.1016/j.rhisph.2023.100741>
- Nedeljković, M., Sastre, D. E., & Sundberg, E. J. (2021). Bacterial flagellar filament: A supramolecular multifunctional nanostructure. *International Journal of Molecular Sciences*, 22(14), 1–33. <https://doi.org/10.3390/ijms22147521>
- Nelson, C. E. (2017). Comprehensive functional characterization of the glycoside hydrolase family 3 enzymes from *Cellvibrio japonicus* reveals unique metabolic roles in biomass saccharification. *Environmental Microbiology*, 19(12), 5025–5039. <https://doi.org/10.1111/1462-2920.13959>
- Nerva, L., Sandrini, M., Moffa, L., Velasco, R., Balestrini, R., & Chitarra, W. (2022). Breeding toward improved ecological plant-microbiome interactions. *Trends in Plant Science*, 27(11), 1134–1143. <https://doi.org/10.1016/j.tplants.2022.06.004>
- Nicola, S., Tibaldi, G., & Fontana, E. (2009). Tomato production systems and their application to the tropics. *Acta Horticulturae*, 821, 27–34. <https://doi.org/10.17660/ActaHortic.2009.821.1>
- Nilsson, R. H., Larsson, K.-H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., Kennedy, P., Picard, K., Glöckner, F. O., Tedersoo, L., Saar, I., Kõljalg, U., & Abarenkov, K. (2019). The UNITE database for molecular identification of fungi: Handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Research*, 47, 259–264. <https://doi.org/10.1093/nar/kgy1022>
- Nozoye, T. (2018). The nicotianamine synthase gene is a useful candidate for improving the nutritional qualities and Fe-deficiency tolerance of various crops. *Frontiers in Plant Science*, 9(340), 1–7. <https://doi.org/10.3389/fpls.2018.00340>
- O'Brien, A. M., Jack, C. N., Friesen, M. L., & Frederickson, M. E. (2021). Whose trait is it anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. *Proceedings of the Royal Society B: Biological Sciences*, 288(1942), 1–10. <https://doi.org/10.1098/rspb.2020.2483>
- OECD. (2017). Tomato (*Solanum lycopersicum*). In Organisation for Economic Co-operation and Development (Ed.), *Safety Assessment of Transgenic Organisms in the Environment, Volume 7* (pp. 69–97). OECD Publishing. <https://www.oecd-ilibrary.org/content/publication/9789264279728-en>
- Ofek, M., Hadar, Y., & Minz, D. (2012). Ecology of root colonizing *Massilia* (Oxalobacteraceae). *PLOS ONE*, 7(7), 1–12. <https://doi.org/10.1371/journal.pone.0040117>
- Ofek-Lalzar, M., Sela, N., Goldman-Voronov, M., Green, S. J., Hadar, Y., & Minz, D. (2014). Niche and host-associated functional signatures of the root surface microbiome. *Nature Communications*, 5(1), 1–9. <https://doi.org/10.1038/ncomms5950>
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2020). *vegan: Community Ecology Package* (Version 2.5-7) [Computer software]. <https://CRAN.R-project.org/package=vegan>
- Olm, M. R. (2021). inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nature Biotechnology*, 39, 727–736. <https://doi.org/10.1038/s41587-020-00797-0>
- Ortega-Ortega, Y., Sarmiento-López, L. G., Baylón-Palomino, A., Vázquez-Lee, J., Maldonado-Bonilla, L. D., Flores-Olivas, A., & Valenzuela-Soto, J. H. (2024). *Enterobacter* sp. DBA51 produces ACC deaminase and promotes the growth of tomato (*Solanum lycopersicum* L.) and tobacco (*Nicotiana tabacum* L.) plants under greenhouse condition. *Current Research in Microbial Sciences*, 6, 1–9. <https://doi.org/10.1016/j.crmicr.2023.100207>

- Ottesen, A. R., Gorham, S., Reed, E., Newell, M. J., Ramachandran, P., Canida, T., Allard, M., Evans, P., Brown, E., & White, J. R. (2016). Using a control to better understand phyllosphere microbiota. *PLOS ONE*, *11*(9), 1–16. <https://doi.org/10.1371/journal.pone.0163482>
- Oyserman, B. O., Cordovez, V., Flores, S. S., Leite, M. F. A., Nijveen, H., Medema, M. H., & Raaijmakers, J. M. (2021). Extracting the GEMs: Genotype, environment, and microbiome interactions shaping host phenotypes. *Frontiers in Microbiology*, *11*, 1–8. <https://www.frontiersin.org/articles/10.3389/fmicb.2020.574053>
- Oyserman, B. O., Sarango Flores, S., Griffioen, T., Pan, X., van der Wijk, E., Pronk, L., Lokhorst, W., Nurfikari, A., Paulson, J. N., Movassagh, M., Stopnisek, N., Kupczok, A., Cordovez, V., Carrión, V. J., Ligterink, W., Snoek, B. L., Medema, M. H., & Raaijmakers, J. M. (2022). Disentangling the genetic basis of rhizosphere microbiome assembly in tomato. *Nature Communications*, *13*(1), 1–16. <https://doi.org/10.1038/s41467-022-30849-9>
- Oyserman, B. O., Medema, M. H., & Raaijmakers, J. M. (2018). Road MAPs to engineer host microbiomes. *Current Opinion in Microbiology*, *43*, 46–54. <https://doi.org/10.1016/j.mib.2017.11.023>
- Palleroni, N. J. (1976). Chemotaxis in *Actinoplanes*. *Archives of Microbiology*, *110*(1), 13–18. <https://doi.org/10.1007/BF00416963>
- Paradis, E., Blomberg, S., Bolker [aut, B., cph, Brown, J., Claramunt, S., Claude, J., Cuong, H. S., Desper, R., Didier, G., Durand, B., Duthel, J., Ewing, R. J., Gascuel, O., Guillaume, T., Heibl, C., Ives, A., Jones, B., Krah [aut, F., ... Vienne, D. de. (2023). *ape: Analyses of Phylogenetics and Evolution* (Version 5.7-1) [Computer software]. <https://cran.r-project.org/web/packages/ape/index.html>
- Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015). CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Research*, *25*(7), 1043–1055. <https://doi.org/10.1101/gr.186072.114>
- Parthasarathy, A., Cross, P. J., Dobson, R. C. J., Adams, L. E., Savka, M. A., & Hudson, A. O. (2018). A three-ring circus: Metabolism of the three proteogenic aromatic amino acids and their role in the health of plants and animals. *Frontiers in Molecular Biosciences*, *5*, 1–30. <https://doi.org/10.3389/fmolb.2018.00029>
- Patten, C. L., & Glick, B. R. (2002). Regulation of indoleacetic acid production in *Pseudomonas putida* GR12-2 by tryptophan and the stationary-phase sigma factor RpoS. *Canadian Journal of Microbiology*, *48*(7), 635–642. <https://doi.org/10.1139/w02-053>
- Paulson, J. N., Stine, O. C., Bravo, H. C., & Pop, M. (2013). Differential abundance analysis for microbial marker-gene surveys. *Nature Methods*, *10*(12), 1200–1202. <https://doi.org/10.1038/nmeth.2658>
- Peralta, I. E., & Spooner, D. M. (2000). Classification of wild tomatoes: A review. *Kurtziana*, *1*, 45–54.
- Peralta, I. E., Spooner, D. M., & Knapp, S. (2008). *Taxonomy of wild tomatoes and their relatives (Solanum sect. Lycopersicoides, sect. Juglandifolia, sect. Lycopersicon; Solanaceae)* (C. Anderson, Ed.). American Soc. of Plant Taxonomists.
- Pereira, P. A. A., Miranda, B. D., Attewell, J. R., Kmieciak, K. A., & Bliss, F. A. (1993). Selection for increased nodule number in common bean (*Phaseolus vulgaris* L.). *Plant and Soil*, *148*, 203–209. <https://doi.org/10.1007/BF00012858>
- Pérez-Jaramillo, J. E., Carrión, V. J., de Hollander, M., & Raaijmakers, J. M. (2018). The wild side of plant microbiomes. *Microbiome*, *6*(1), 1–6. <https://doi.org/10.1186/s40168-018-0519-z>
- Pérez-Jaramillo, J. E., Mendes, R., & Raaijmakers, J. M. (2016). Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Molecular Biology*, *90*(6), 635–644. <https://doi.org/10.1007/s11103-015-0337-7>
- Pérez-Rodríguez, M. M., Piccoli, P., Anzuay, M. S., Baraldi, R., Neri, L., Taurian, T., Lobato Ureche, M. A., Segura, D. M., & Cohen, A. C. (2020). Native bacteria isolated from roots and rhizosphere

- of *Solanum lycopersicum* L. increase tomato seedling growth under a reduced fertilization regime. *Scientific Reports*, 10(1), 1–14. <https://doi.org/10.1038/s41598-020-72507-4>
- Persmark, M., Expert, D., & Neilands, J. B. (1989). Isolation, characterization, and synthesis of chryso-bactin, a compound with siderophore activity from *Erwinia chrysanthemi*. *Journal of Biological Chemistry*, 264(6), 3187–3193. [https://doi.org/10.1016/S0021-9258\(18\)94049-4](https://doi.org/10.1016/S0021-9258(18)94049-4)
- Philippot, L., Chenu, C., Kappler, A., Rillig, M. C., & Fierer, N. (2024). The interplay between microbial communities and soil properties. *Nature Reviews Microbiology*, 22(4), 226–239. <https://doi.org/10.1038/s41579-023-00980-5>
- Philippot, L., Raaijmakers, J. M., Lemanceau, P., & van der Putten, W. H. (2013). Going back to the roots: The microbial ecology of the rhizosphere. *Nature Reviews Microbiology*, 11(11), 789–799. <https://doi.org/10.1038/nrmicro3109>
- Pieterse, C. M. J., Zamioudis, C., Berendsen, R. L., Weller, D. M., Van Wees, S. C. M., & Bakker, P. A. H. M. (2014). Induced systemic resistance by beneficial microbes. *Annual Review of Phytopathology*, 52(1), 347–375. <https://doi.org/10.1146/annurev-phyto-082712-102340>
- Pineda, A., Kaplan, I., & Bezemer, T. M. (2017). Steering soil microbiomes to suppress aboveground insect pests. *Trends in Plant Science*, 22(9), 770–778. <https://doi.org/10.1016/j.tplants.2017.07.002>
- Pineda, A., Kaplan, I., Hannula, S. E., Ghanem, W., & Bezemer, T. M. (2020). Conditioning the soil microbiome through plant–soil feedbacks suppresses an aboveground insect pest. *New Phytologist*, 226(2), 595–608. <https://doi.org/10.1111/nph.16385>
- Planet, P. J., Kachlany, S. C., Fine, D. H., DeSalle, R., & Figurski, D. H. (2003). The widespread colonization island of *Actinobacillus actinomycetemcomitans*. *Nature Genetics*, 34(2), 193–198. <https://doi.org/10.1038/ng1154>
- Poli, A., Lazzari, A., Prigione, V., Voyron, S., Spadaro, D., & Varese, G. C. (2016). Influence of plant genotype on the cultivable fungi associated to tomato rhizosphere and roots in different soils. *Fungal Biology*, 120(6), 862–872. <https://doi.org/10.1016/j.funbio.2016.03.008>
- Polizeli, M. L. T. M. (2005). Xylanases from fungi: Properties and industrial applications. *Applied Microbiology and Biotechnology*, 67, 577–591. <https://doi.org/10.1007/s00253-005-1904-7>
- Porcel, R., Zamarréño, Á. M., García-Mina, J. M., & Aroca, R. (2014). Involvement of plant endogenous ABA in *Bacillus megaterium* PGPR activity in tomato plants. *BMC Plant Biology*, 14(1), 1–12. <https://doi.org/10.1186/1471-2229-14-36>
- Porter, S. S., & Sachs, J. L. (2020). Agriculture and the disruption of plant–microbial symbiosis. *Trends in Ecology & Evolution*, 35(5), 426–439. <https://doi.org/10.1016/j.tree.2020.01.006>
- Porter, S. S., Bantay, R., Friel, C. A., Garoutte, A., Gdanetz, K., Ibarreta, K., Moore, B. M., Shetty, P., Siler, E., & Friesen, M. L. (2020). Beneficial microbes ameliorate abiotic and biotic sources of stress on plants. *Functional Ecology*, 34(10), 2075–2086. <https://doi.org/10.1111/1365-2435.13499>
- Poudel, R., Jumpponen, A., Kennelly, M. M., Rivard, C. L., Gomez-Montano, L., & Garrett, K. A. (2019). Rootstocks shape the rhizobiome: Rhizosphere and endosphere bacterial communities in the grafted tomato system. *Applied and Environmental Microbiology*, 85(2), 1–16. <https://doi.org/10.1128/AEM.01765-18>
- Poyart, C., Lamy, M. C., Boumaila, C., Fiedler, F., & Trieu-Cuot, P. (2001). Regulation of D-alanyl-lipoteichoic acid biosynthesis in *Streptococcus agalactiae* involves a novel two-component regulatory system. *Journal of Bacteriology*, 183(21), 6324–6334. <https://doi.org/10.1128/JB.183.21.6324-6334.2001>
- Prasad, M., Srinivasan, R., Chaudhary, M., Choudhary, M., & Jat, L. K. (2019). Chapter Seven - Plant Growth Promoting Rhizobacteria (PGPR) for Sustainable Agriculture: Perspectives and Challenges. In A. K. Singh, A. Kumar, & P. K. Singh (Eds.), *PGPR Amelioration in Sustainable Agriculture* (pp. 129–157). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-815879-1.00007-0>

- Priyadarshan, P. M. (2019). *Plant breeding: Classical to modern*. Springer.
- Purugganan, M. D. (2022). What is domestication? *Trends in Ecology & Evolution*, 37(8), 663–671. <https://doi.org/10.1016/j.tree.2022.04.006>
- Purugganan, M. D., & Fuller, D. Q. (2009). The nature of selection during plant domestication. *Nature*, 457(7231), 843–848. <https://doi.org/10.1038/nature07895>
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, 41(D1), D590–D596. <https://doi.org/10.1093/nar/gks1219>
- Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, 26(6), 841–842. <https://doi.org/10.1093/bioinformatics/btq033>
- R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Raaijmakers, J. M., & Kiers, E. T. (2022). Rewilding plant microbiomes. *Science*, 378(6620), 599–600. <https://doi.org/10.1126/science.abn6350>
- Radzki, W., Gutierrez Mañero, F. J., Algar, E., Lucas García, J. A., García-Villaraco, A., & Ramos Solano, B. (2013). Bacterial siderophores efficiently provide iron to iron-starved tomato plants in hydroponics culture. *Antonie van Leeuwenhoek*, 104(3), 321–330. <https://doi.org/10.1007/s10482-013-9954-9>
- Raivio, T. L., & Silhavy, T. J. (2001). Periplasmic stress and ECF sigma factors. *Annual Review of Microbiology*, 55, 591–624. <https://doi.org/10.1146/annurev.micro.55.1.591>
- Ramírez-Ojeda, G., Peralta, I. E., Rodríguez-Guzmán, E., Chávez-Servia, J. L., Sahagún-Castellanos, J., & Rodríguez-Pérez, J. E. (2021). Climatic diversity and ecological descriptors of wild tomato species (*Solanum* sect. *Lycopersicon*) and close related species (*Solanum* sect. *Juglandifolia* y sect. *Lycopersicoides*) in Latin America. *Plants*, 10(5), 1–17. <https://doi.org/10.3390/plants10050855>
- Ramírez-Rendón, D., Guzmán-Chávez, F., García-Ausencio, C., Rodríguez-Sanoja, R., & Sánchez, S. (2023). The untapped potential of actinobacterial lanthipeptides as therapeutic agents. *Molecular Biology Reports*, 50(12), 10605–10616. <https://doi.org/10.1007/s11033-023-08880-w>
- Ramón, G. (2008). Un espacio original: La región de los Andes Bajos. In *La nueva historia de Loja: La historia aborígen y colonial* (Vol. 1). Corporación Editora Nacional.
- Ramonedá, J., Fan, K., Lucas, J. M., Chu, H., Bissett, A., Strickland, M. S., & Fierer, N. (2024). Ecological relevance of flagellar motility in soil bacterial communities. *The ISME Journal*, 18(1), 1–14. <https://doi.org/10.1093/ismej/wrae067>
- Rashid, M. H.-O., & Chung, Y. R. (2017). Induction of systemic resistance against insect herbivores in plants by beneficial soil microbes. *Frontiers in Plant Science*, 8, 1–11. <https://doi.org/10.3389/fpls.2017.01816>
- Razali, R., Bougouffa, S., Morton, M. J. L., Lightfoot, D. J., Alam, I., Essack, M., Arold, S. T., Kamau, A. A., Schmöckel, S. M., Pailles, Y., Shahid, M., Michell, C. T., Al-Babili, S., Ho, Y. S., Tester, M., Bajic, V. B., & Negrão, S. (2018). The genome sequence of the wild tomato *solanum pimpinellifolium* provides insights into salinity tolerance. *Frontiers in Plant Science*, 9, 1–21. <https://doi.org/10.3389/fpls.2018.01402>
- Razifard, H., Ramos, A., Della Valle, A. L., Bodary, C., Goetz, E., Manser, E. J., Li, X., Zhang, L., Visa, S., Tieman, D., van der Knaap, E., & Caicedo, A. L. (2020). Genomic evidence for complex domestication history of the cultivated tomato in Latin America. *Molecular Biology and Evolution*, 37(4), 1118–1132. <https://doi.org/10.1093/molbev/msz297>
- Reid, T. E., Kavamura, V. N., Torres-Ballesteros, A., Smith, M. E., Abadie, M., Pawlett, M., Clark, I. M., Harris, J. A., & Mauchline, T. H. (2024). Agricultural intensification reduces selection of puta-

- tive plant growth-promoting rhizobacteria in wheat. *The ISME Journal*, 18(1), 1–14. <https://doi.org/10.1093/ismej/wrae131>
- Resendiz-Nava, C. N., Alonso-Onofre, F., Silva-Rojas, H. V., Rebollar-Alviter, A., Rivera-Pastrana, D. M., Stasiewicz, M. J., Nava, G. M., & Mercado-Silva, E. M. (2023). Tomato plant microbiota under conventional and organic fertilization regimes in a soilless culture system. *Microorganisms*, 11(7), 1–19. <https://doi.org/10.3390/microorganisms11071633>
- Reuscher, S. (2013). Genome-wide identification and expression analysis of aquaporins in tomato. *PLoS One*, 8(11), 1–13. <https://doi.org/10.1371/journal.pone.0079052>
- Rick, C. M. (1950). Pollination relations of *Lycopersicon esculentum* in native and foreign regions. *Evolution*, 4(2), 110–122. <https://doi.org/10.2307/2405388>
- Rick, C. M. (1978, January 8). The Tomato. *Scientific American*, 77–87. <https://doi.org/10.1038/scientificamerican0878-76>
- Rick, C. M. (1983). Genetic variability in tomato species. *Plant Molecular Biology Reporter*, 1(2), 81–87. <https://doi.org/10.1007/BF02680303>
- Ritchie, M. E. (2015). *limma* powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, 43(7), 1–13. <https://doi.org/10.1093/nar/gkv007>
- Rizaludin, M. S., Stopnisek, N., Raaijmakers, J. M., & Garbeva, P. (2021). The chemistry of stress: Understanding the “cry for help” of plant roots. *Metabolites*, 11(6), 1–14. <https://doi.org/10.3390/metabo11060357>
- Rodrigues, J. L. M., Pellizari, V. H., Mueller, R., Baek, K., Jesus, E. da C., Paula, F. S., Mirza, B., Hamaoui, G. S., Tsai, S. M., Feigl, B., Tiedje, J. M., Bohannan, B. J. M., & Nüsslein, K. (2013). Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proceedings of the National Academy of Sciences*, 110(3), 988–993. <https://doi.org/10.1073/pnas.1220608110>
- Rodrigues, M. H. P., & Furlong, E. B. (2022). Fungal diseases and natural defense mechanisms of tomatoes (*Solanum lycopersicum*): A review. *Physiological and Molecular Plant Pathology*, 122, 1–7 <https://doi.org/10.1016/j.pmpp.2022.101906>
- Rojas, C. M., Ham, J. H., Deng, W.-L., Doyle, J. J., & Collmer, A. (2002). HecA, a member of a class of adhesins produced by diverse pathogenic bacteria, contributes to the attachment, aggregation, epidermal cell killing, and virulence phenotypes of *Erwinia chrysanthemi* EC16 on *Nicotiana glauca* seedlings. *Proceedings of the National Academy of Sciences*, 99(20), 13142–13147. <https://doi.org/10.1073/pnas.202358699>
- Rolfe, S. A., Griffiths, J., & Ton, J. (2019). Crying out for help with root exudates: Adaptive mechanisms by which stressed plants assemble health-promoting soil microbiomes. *Current Opinion in Microbiology*, 49, 73–82. <https://doi.org/10.1016/j.mib.2019.10.003>
- Ronga, D., Caradonia, F., Francia, E., Morcia, C., Rizza, F., Badeck, F.-W., Ghizzoni, R., & Terzi, V. (2019). Interaction of tomato genotypes and arbuscular mycorrhizal fungi under reduced irrigation. *Horticulturae*, 5(4), 1–15. <https://doi.org/10.3390/horticulturae5040079>
- RStudio Team. (2020). *RStudio: Integrated Development for R* [Computer software]. RStudio, PBC. <http://www.rstudio.com/>
- Rudrappa, T., Czymmek, K. J., Paré, P. W., & Bais, H. P. (2008). Root-secreted malic acid recruits beneficial soil bacteria. *Plant Physiology*, 148(3), 1547–1556. <https://doi.org/10.1104/pp.108.127613>
- Runge, P., Ventura, F., Kemen, E., & Stam, R. (2022). Distinct phyllosphere microbiome of wild tomato species in central peru upon dysbiosis. *Microbial Ecology*, 85(1), 168–183. <https://doi.org/10.1007/s00248-021-01947-w>
- Saar-Dover, R., Bitler, A., Nezer, R., Shmuel-Galia, L., Firon, A., Shimoni, E., Trieu-Cuot, P., & Shai, Y. (2012). D-alanylation of lipoteichoic acids confers resistance to cationic peptides in group B *Streptococcus*

- tococcus* by increasing the cell wall density. *PLOS Pathogens*, 8(9), 1–13. <https://doi.org/10.1371/journal.ppat.1002891>
- Sade, N. (2009). Improving plant stress tolerance and yield production: Is the tonoplast aquaporin SITIP2; 2 a key to isohydric to anisohydric conversion? *New Phytologist*, 181(3), 651–661. <https://doi.org/10.1111/j.1469-8137.2008.02689.x>
- Sadrnia, M., Maksimava, N., Khromsova, E., Stanislavich, S., Owlia, P., & Arjomandzadegan, M. (2011). Study of the effect of bacterial 1-aminocyclopropane-1-carboxylate deaminase (ACC) deaminase on resistance to salt stress in tomato plant. *Analele Universității Din Oradea - Fascicula Biologie*, 18(2), 120–123.
- Safdarian, M., Askari, H., Shariati, J. V., & Nematzadeh, G. (2019). Transcriptional responses of wheat roots inoculated with *Arthrobacter nitroguajacolicus* to salt stress. *Scientific Reports*, 9(1792), 1–12. <https://doi.org/10.1038/s41598-018-38398-2>
- Sah, S., Krishnani, S., & Singh, R. (2021). *Pseudomonas* mediated nutritional and growth promotional activities for sustainable food security. *Current Research in Microbial Sciences*, 2, 1–12. <https://doi.org/10.1016/j.crmicr.2021.100084>
- Sakellaris, G., Kolisis, F. N., & Evangelopoulos, A. E. (1988). Presence of sialic acids in *Lactobacillus plantarum*. *Biochemical and Biophysical Research Communications*, 155(3), 1126–1132. [https://doi.org/10.1016/S0006-291X\(88\)81257-9](https://doi.org/10.1016/S0006-291X(88)81257-9)
- Saleem, M., Law, A. D., Sahib, M. R., Pervaiz, Z. H., & Zhang, Q. (2018). Impact of root system architecture on rhizosphere and root microbiome. *Rhizosphere*, 6, 47–51. <https://doi.org/10.1016/j.rhisph.2018.02.003>
- Salomon, P. S., Janson, S., & Granéli, E. (2003). Molecular identification of bacteria associated with filaments of *Nodularia spumigena* and their effect on the cyanobacterial growth. *Harmful Algae*, 2(4), 261–272. [https://doi.org/10.1016/S1568-9883\(03\)00045-3](https://doi.org/10.1016/S1568-9883(03)00045-3)
- Sandy, M., & Butler, A. (2011). Chrysobactin siderophores produced by *Dickeya chrysanthemi* EC16. *Journal of Natural Products*, 74(5), 1207–1212. <https://doi.org/10.1021/np200126z>
- Santhanam, R., Luu, V. T., Weinhold, A., Goldberg, J., Oh, Y., & Baldwin, I. T. (2015). Native root-associated bacteria rescue a plant from a sudden-wilt disease that emerged during continuous cropping. *Proceedings of the National Academy of Sciences*, 112(36), 5013–5020. <https://doi.org/10.1073/pnas.1505765112>
- Sarango Flores, S., Cordovez, V., Oyserman, B. O., Stopnisek, N., Raaijmakers, J. M., & Van 't Hof, P. (2023). The tomato's tale: Exploring taxonomy, biogeography, domestication and microbiome for enhanced resilience. *Phytobiomes Journal*, 8(1), 5–20 <https://doi.org/10.1094/PBIOMES-09-23-0091-MF>
- Sasse, J., Martinoia, E., & Northen, T. (2018). Feed your friends: do plant exudates shape the root microbiome? *Trends in Plant Science*, 23(1), 25–41. <https://doi.org/10.1016/j.tplants.2017.09.003>
- Scheublin, T. R., Sanders, I. R., Keel, C., & van der Meer, J. R. (2010). Characterisation of microbial communities colonising the hyphal surfaces of arbuscular mycorrhizal fungi. *The ISME Journal*, 4(6), 752–763. <https://doi.org/10.1038/ismej.2010.5>
- Schmidt, J. E., Vannette, R. L., Igwe, A., Blundell, R., Casteel, C. L., & Gaudin, A. C. M. (2019). Effects of agricultural management on rhizosphere microbial structure and function in processing tomato plants. *Applied and Environmental Microbiology*, 85(16), 1–18. <https://doi.org/10.1128/AEM.01064-19>
- Schmieder, R., & Edwards, R. (2011). Quality control and preprocessing of metagenomic datasets. *Bioinformatics*, 27(6), 863–864. <https://doi.org/10.1093/bioinformatics/btr026>
- Schmitz, L., Yan, Z., Schneijderberg, M., de Roij, M., Pijnenburg, R., Zheng, Q., Franken, C., Dechesne, A., Trindade, L. M., van Velzen, R., Bisseling, T., Geurts, R., & Cheng, X. (2022). Synthetic

- bacterial community derived from a desert rhizosphere confers salt stress resilience to tomato in the presence of a soil microbiome. *The ISME Journal*, 16(8), 1907–1920. <https://doi.org/10.1038/s41396-022-01238-3>
- Schneewind, O., & Missiakas, D. (2016). Lipoteichoic acid synthesis and function in Gram-Positive Bacteria. In O. Geiger (Ed.), *Biogenesis of Fatty Acids, Lipids and Membranes* (pp. 1–18). Springer International Publishing. [https://doi.org/10.1007/978-3-319-43676-0\\_17-1](https://doi.org/10.1007/978-3-319-43676-0_17-1)
- Schöner, T. A. (2016). Aryl polyenes, a highly abundant class of bacterial natural products, are functionally related to antioxidative carotenoids. *ChemBioChem*, 17(3), 247–253. <https://doi.org/10.1002/cbic.201500474>
- Schrempf, H., & Walter, S. (1995). The cellulolytic system of *Streptomyces reticuli*. *International Journal of Biological Macromolecules*, 17(6), 353–355. [https://doi.org/10.1016/0141-8130\(96\)81845-9](https://doi.org/10.1016/0141-8130(96)81845-9)
- Schwarz, B., & Bauer, P. (2020). FIT, a regulatory hub for iron deficiency and stress signaling in roots, and FIT-dependent and -independent gene signatures. *Journal of Experimental Botany*, 71(5), 1694–1705. <https://doi.org/10.1093/jxb/eraa012>
- Scott, J. W., Myers, J. R., Boches, P. S., Nichols, C. G., & Angell, F. F. (2013). Classical Genetics and Traditional Breeding. In B. E. Liedl, J. A. Labate, J. R. Stommel, A. Slade, & C. Kole (Eds.), *Genetics, Genomics, and Breeding of Tomato* (pp. 38–73). CRC Press.
- Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*, 30(14), 2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>
- Shabtai, I. A., Wilhelm, R. C., Schweizer, S. A., Höschen, C., Buckley, D. H., & Lehmann, J. (2023). Calcium promotes persistent soil organic matter by altering microbial transformation of plant litter. *Nature Communications*, 14(1), 1–13. <https://doi.org/10.1038/s41467-023-42291-6>
- Shaffer, M. (2020). DRAM for distilling microbial metabolism to automate the curation of microbiome function. *Nucleic Acids Research*, 48(16), 8883–8900. <https://doi.org/10.1093/nar/gkaa621>
- Sharanappa, C. H., Bheemanna, M., Prabhuraj, A., Harischandra, R. N., Nagaraj, M. N., Saroja, N. R., & Kariyanna, B. (2023). Toxic effect of cyanobacterial (blue–green algae) extracts as natural pesticides for the control of *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae). *Egyptian Journal of Biological Pest Control*, 33(77), 1–16. <https://doi.org/10.1186/s41938-023-00716-w>
- Sharma, M. K., Solanke, A. U., Jani, D., Singh, Y., & Sharma, A. K. (2009). A simple and efficient Agrobacterium-mediated procedure for transformation of tomato. *Journal of Biosciences*, 34(3), 423–433. <https://doi.org/10.1007/s12038-009-0049-8>
- Shen, W., & Ren, H. (2021). TaxonKit: A practical and efficient NCBI taxonomy toolkit. *Journal of Genetics and Genomics*, 48(9), 844–850. <https://doi.org/10.1016/j.jgg.2021.03.006>
- Shi, X., Chen, G., Tian, L., Peng, Z., Xie, W., Wu, Q., Wang, S., Zhou, X., & Zhang, Y. (2016). The salicylic acid-mediated release of plant volatiles affects the host choice of *Bemisia tabaci*. *International Journal of Molecular Sciences*, 17(7), 1–11. <https://doi.org/10.3390/ijms17071048>
- Shintu, P. V., & Jayaram, K. M. (2015). Phosphate solubilising bacteria (*Bacillus polymyxa*)—An effective approach to mitigate drought in tomato (*Lycopersicon esculentum* Mill.). *Tropical Plant Research*, 2(1), 17–22.
- Shivega, W. G., & Aldrich-Wolfe, L. (2017). Native plants fare better against an introduced competitor with native microbes and lower nitrogen availability. *AoB PLANTS*, 9(1), 1–14. <https://doi.org/10.1093/aobpla/plx004>
- Sideli, G. M. (2020). Quantitative phenotyping of shell suture strength in walnut (*Juglans regia* L.) enhances precision for detection of QTL and genome-wide association mapping. *PLoS One*, 15(4), 1–21. <https://doi.org/10.1371/journal.pone.0231144>

- Sieber, C. M. K., Probst, A. J., Sharrar, A., Thomas, B. C., Hess, M., Tringe, S. G., & Banfield, J. F. (2018). Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nature Microbiology*, *3*(7), 836–843. <https://doi.org/10.1038/s41564-018-0171-1>
- Sillo, F., Marino, G., Franchi, E., Haworth, M., Zampieri, E., Pietrini, I., Fusini, D., Mennone, C., Centritto, M., & Balestrini, R. (2022). Impact of irrigation water deficit on two tomato genotypes grown under open field conditions: From the root-associated microbiota to the stress responses. *Italian Journal of Agronomy*, *17*(3), 1–15. <https://doi.org/10.4081/ija.2022.2130>
- Silva, G. da C., Kitano, I. T., Ribeiro, I. A. de F., & Lacava, P. T. (2022). The potential use of Actinomycetes as microbial inoculants and biopesticides in agriculture. *Frontiers in Soil Science*, *2*, 1–20. <https://doi.org/10.3389/fsoil.2022.833181>
- Sim, S.-C., Robbins, M. D., Deynze, A. V., Michel, A. P., & Francis, D. M. (2011). Population structure and genetic differentiation associated with breeding history and selection in tomato (*Solanum lycopersicum* L.). *Heredity*, *106*(6), 927–935. <https://doi.org/10.1038/hdy.2010.139>
- Simonin, M., Briand, M., Chesneau, G., Rochefort, A., Marais, C., Sarniguet, A., & Barret, M. (2022). Seed microbiota revealed by a large-scale meta-analysis including 50 plant species. *New Phytologist*, *234*(4), 1448–1463. <https://doi.org/10.1111/nph.18037>
- Singh, M., Singh, D., Gupta, A., Pandey, K. D., Singh, P. K., & Kumar, A. (2019). Chapter Three - Plant Growth Promoting Rhizobacteria: Application in Biofertilizers and Biocontrol of Phytopathogens. In A. K. Singh, A. Kumar, & P. K. Singh (Eds.), *PGPR Amelioration in Sustainable Agriculture* (pp. 41–66). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-815879-1.00003-3>
- Singh, P. P., Kujur, A., Yadav, A., Kumar, A., Singh, S. K., & Prakash, B. (2019). Chapter Two—Mechanisms of Plant-Microbe Interactions and its Significance for Sustainable Agriculture. In A. K. Singh, A. Kumar, & P. K. Singh (Eds.), *PGPR Amelioration in Sustainable Agriculture* (pp. 17–39). Woodhead Publishing. <https://doi.org/10.1016/B978-0-12-815879-1.00002-1>
- Smith, K. P., Handelsman, J., & Goodman, R. M. (1999). Genetic basis in plants for interactions with disease-suppressive bacteria. *Proceedings of the National Academy of Sciences of the United States of America*, *96*(9), 4786–4790.
- Smith, N. W., Shorten, P. R., Altermann, E., Roy, N. C., & McNabb, W. C. (2019). The classification and evolution of bacterial cross-feeding. *Frontiers in Ecology and Evolution*, *7*, 1–15. <https://doi.org/10.3389/fevo.2019.00153>
- Smulders, L., Benítez, E., Moreno, B., López-García, Á., Pozo, M. J., Ferrero, V., de la Peña, E., & Alcalá Herrera, R. (2021). Tomato domestication affects potential functional molecular pathways of root-associated soil bacteria. *Plants*, *10*(9), 1–17. <https://doi.org/10.3390/plants10091942>
- Šnajdr, J., Dobiášová, P., Vetrovský, T., Valášková, V., Alawi, A., Boddy, L., & Baldrian, P. (2011). Saprotrophic basidiomycete mycelia and their interspecific interactions affect the spatial distribution of extracellular enzymes in soil. *FEMS Microbiology Ecology*, *78*(1), 80–90. <https://doi.org/10.1111/j.1574-6941.2011.01123.x>
- Soldan, R., Fusi, M., Cardinale, M., Daffonchio, D., & Preston, G. M. (2021). The effect of plant domestication on host control of the microbiota. *Communications Biology*, *4*(1), 1–9. <https://doi.org/10.1038/s42003-021-02467-6>
- Spaepen, S., & Vanderleyden, J. (2011). Auxin and plant-microbe interactions. *Cold Spring Harbor Perspectives in Biology*, *3*(4), 1–13. <https://doi.org/10.1101/cshperspect.a001438>
- Spaepen, S., Vanderleyden, J., & Remans, R. (2007). Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiology Reviews*, *31*(4), 425–448. <https://doi.org/10.1111/j.1574-6976.2007.00072.x>

- Spitz, O., Erenburg, I. N., Beer, T., Kanonenberg, K., Holland, I. B., & Schmitt, L. (2019). Type I Secretion Systems—One mechanism for all? *Microbiology Spectrum*, *7*(2), 1–9. <https://doi.org/10.1128/microbiolspec.psib-0003-2018>
- Steinegger, M., & Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nature Biotechnology*, *35*, 1026–1028. <https://doi.org/10.1038/nbt.3988>
- Sterken, M. G., Nijveen, H., van Zanten, M., Jiménez-Gómez, J. M., Geshnizjani, N., Willems, L. A. J., Rienstra, J., Hilhorst, H. W. M., Ligterink, W., & Snoek, B. L. (2023). Plasticity of maternal environment-dependent expression-QTLs of tomato seeds. *Theoretical and Applied Genetics*, *136*(28), 1–15. <https://doi.org/10.1007/s00122-023-04322-0>
- Stevens, M. A., & Rick, C. M. (1986). Genetics and breeding. In J. G. Atherton & J. Rudich (Eds.), *The Tomato Crop: A scientific basis for improvement* (pp. 35–109). Springer Netherlands. [https://doi.org/10.1007/978-94-009-3137-4\\_2](https://doi.org/10.1007/978-94-009-3137-4_2)
- Streit, W. R. (1996). Biotin and other water-soluble vitamins are key growth factors for alfalfa root colonization by *Rhizobium melioli* 1021. *MPMI*, *9*(5), 330–338. <https://doi.org/10.1094/MPMI-9-0330>
- Sugiura, H. (2020). Bacterial inducible expression of plant cell wall-binding protein YesO through conflict between *Glycine max* and saprophytic *Bacillus subtilis*. *Scientific Reports*, *10*(18691), 1–10. <https://doi.org/10.1038/s41598-020-75359-0>
- Sun, Y., Tian, L., Chang, J., Shi, S., Zhang, J., Xie, H., Cai, Y., Chen, D., Kuramae, E. E., van Veen, J. A., Li, W., Tran, L.-S. P., & Tian, C. (2021). Rice domestication influences the composition and function of the rhizosphere bacterial chemotaxis systems. *Plant and Soil*, *466*(1), 81–99. <https://doi.org/10.1007/s11104-021-05036-2>
- Swapnil, P., Meena, M., Singh, S. K., Dhuldhaj, U. P., Harish, & Marwal, A. (2021). Vital roles of carotenoids in plants and humans to deteriorate stress with its structure, biosynthesis, metabolic engineering and functional aspects. *Current Plant Biology*, *26*, 1–11. <https://doi.org/10.1016/j.cpb.2021.100203>
- Sweeney, C., Lakshmanan, V., & Bais, H. P. (2017). Interplant aboveground signaling prompts upregulation of auxin promoter and malate transporter as part of defensive response in the neighboring plants. *Frontiers in Plant Science*, *8*(595), 1–10. <https://doi.org/10.3389/fpls.2017.00595>
- Tabrett, A., & Horton, M. W. (2020). The influence of host genetics on the microbiome. *F1000Research*, *9*(84), 1–9. <https://doi.org/10.12688/f1000research.20835.1>
- Taffner, J., Bergna, A., Cernava, T., & Berg, G. (2020). Tomato-associated archaea show a cultivar-specific rhizosphere effect but an unspecific transmission by seeds. *Phytobiomes Journal*, *4*(2), 133–141. <https://doi.org/10.1094/PBIOMES-01-20-0017-R>
- Tamames, J., & Puente-Sánchez, F. (2019). SqueezeMeta, a highly portable, fully automatic metagenomic analysis pipeline. *Frontiers in Microbiology*, *9*, 1–10. <https://doi.org/10.3389/fmicb.2018.03349>
- Tank, N., & Saraf, M. (2010). Salinity-resistant plant growth promoting rhizobacteria ameliorates sodium chloride stress on tomato plants. *Journal of Plant Interactions*, *5*(1), 51–58. <https://doi.org/10.1080/17429140903125848>
- Tanksley, S. D., & McCouch, S. R. (1997). Seed banks and molecular maps: Unlocking genetic potential from the wild. *Science*, *277*(5329), 1063–1066. <https://doi.org/10.1126/science.277.5329.1063>
- Taylor, I. B. (1986). Biosystematics of the tomato. In J. G. Atherton & J. Rudich (Eds.), *The Tomato Crop: A scientific basis for improvement* (pp. 1–34). Springer Netherlands. [https://doi.org/10.1007/978-94-009-3137-4\\_1](https://doi.org/10.1007/978-94-009-3137-4_1)
- Tchagang, C. F., Xu, R., Doumbou, C. L., & Tambong, J. T. (2018). Genome analysis of two novel *Pseudomonas* strains exhibiting differential hypersensitivity reactions on tobacco seedlings reveals differences in nonflagellar T3SS organization and predicted effector proteins. *MicrobiologyOpen*, *7*(2), 1–13. <https://doi.org/10.1002/mbo3.553>

- Tchakounté, G. V. T., Berger, B., Patz, S., Becker, M., Fankem, H., Taffouo, V. D., & Ruppel, S. (2020). Selected rhizosphere bacteria help tomato plants cope with combined phosphorus and salt stresses. *Microorganisms*, 8(11), 1–16. <https://doi.org/10.3390/microorganisms8111844>
- Terra, L., Ratcliffe, N., Castro, H. C., Vicente, A. C. P., & Dyson, P. (2021). Biotechnological potential of *Streptomyces* siderophores as new antibiotics. *Current Medicinal Chemistry*, 28(7), 1407–1421. <https://doi.org/10.2174/0929867327666200510235512>
- The Tomato Genome Consortium. (2012). The tomato genome sequence provides insights into fleshy fruit evolution. *Nature*, 485(7400), 635–641. <https://doi.org/10.1038/nature11119>
- Thomas, P., Knox, O. G. G., Powell, J. R., Sindel, B., & Winter, G. (2023). The hydroponic rockwool root microbiome: under control or underutilised? *Microorganisms*, 11(4), 1–17. <https://doi.org/10.3390/microorganisms11040835>
- Tieman, D., Zhu, G., Resende, M. F. R., Lin, T., Nguyen, C., Bies, D., Rambla, J. L., Beltran, K. S. O., Taylor, M., Zhang, B., Ikeda, H., Liu, Z., Fisher, J., Zemach, I., Monforte, A., Zamir, D., Granell, A., Kirst, M., Huang, S., & Klee, H. (2017). A chemical genetic roadmap to improved tomato flavor. *Science*, 355(6323), 391–394. <https://doi.org/10.1126/science.aal1556>
- Timofeeva, A. M., Galyamova, M. R., & Sedykh, S. E. (2022). Bacterial siderophores: Classification, biosynthesis, perspectives of use in agriculture. *Plants*, 11(22), 1–22. <https://doi.org/10.3390/plants11223065>
- Toju, H., Okayasu, K., & Notaguchi, M. (2019). Leaf-associated microbiomes of grafted tomato plants. *Scientific Reports*, 9(1), 1–11. <https://doi.org/10.1038/s41598-018-38344-2>
- Toledo Cabrera, B. (2021). Effect of *Rhizobium* inoculation on tomato (*Solanum lycopersicum* L.) yield in protected crops. *Biology and Life Sciences Forum*, 3(52), 1–7. <https://doi.org/10.3390/IECAG2021-09993>
- Tracanna, V., Ossowicki, A., Petrus, M. L. C., Overduin, S., Terlouw, B. R., Lund, G., Robinson, S. L., Warris, S., Schijlen, E. G. W. M., van Wezel, G. P., Raaijmakers, J. M., Garbeva, P., & Medema, M. H. (2021). Dissecting disease-suppressive rhizosphere microbiomes by functional amplicon sequencing and 10× metagenomics. *mSystems*, 6(3), 1–15. <https://doi.org/10.1128/mSystems.01116-20>
- Tran, H., Ficke, A., Asimwe, T., Höfte, M., & Raaijmakers, J. M. (2007). Role of the cyclic lipopeptide massetolide A in biological control of *Phytophthora infestans* and in colonization of tomato plants by *Pseudomonas fluorescens*. *New Phytologist*, 175(4), 731–742. <https://doi.org/10.1111/j.1469-8137.2007.02138.x>
- Traving, C., & Schauer, R. (1998). Structure, function and metabolism of sialic acids. *Cellular and Molecular Life Sciences CMLS*, 54(12), 1330–1349. <https://doi.org/10.1007/s000180050258>
- Treangen, T. J., Sommer, D. D., Angly, F. E., Koren, S., & Pop, M. (2011). Next generation sequence assembly with AMOS. *Current Protocols in Bioinformatics*, 11, 1–31. <https://doi.org/10.1002/0471250953.bi1108s33>
- Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., & Singh, B. K. (2020). Plant-microbiome interactions: From community assembly to plant health. *Nature Reviews Microbiology*, 18(11), 607–621. <https://doi.org/10.1038/s41579-020-0412-1>
- Tronson, E., Kaplan, I., & Enders, L. (2022). Characterizing rhizosphere microbial communities associated with tolerance to aboveground herbivory in wild and domesticated tomatoes. *Frontiers in Microbiology*, 13, 1–17. <https://doi.org/10.3389/fmicb.2022.981987>
- Tsang, M.-J., & Bernhardt, T. G. (2015). A role for the FtsQLB complex in cytokinetic ring activation revealed by an ftsL allele that accelerates division. *Molecular Microbiology*, 95(6), 925–944. <https://doi.org/10.1111/mmi.12905>

- Turner, T. R., James, E. K., & Poole, P. S. (2013). The plant microbiome. *Genome Biology*, *14*(6), 1–10. <https://doi.org/10.1186/gb-2013-14-6-209>
- Tyagi, V., & Tyagi, S. (2016). Qualitative and quantitative variations of fungi in tomato rhizosphere in response to plant age. *Plant Archives*, *16*(2), 934–936.
- Uchida, K., Jang, M.-S., Ohnishi, Y., Horinouchi, S., Hayakawa, M., Fujita, N., & Aizawa, S.-I. (2011). Characterization of *Actinoplanes missouriensis* spore flagella. *Applied and Environmental Microbiology*, *77*(7), 2559–2562. <https://doi.org/10.1128/AEM.02061-10>
- Vacheron, J., Desbrosses, G., Bouffaud, M.-L., Touraine, B., Moëgne-Loccoz, Y., Muller, D., Legendre, L., Wisniewski-Dyé, F., & Prigent-Combaret, C. (2013). Plant growth-promoting rhizobacteria and root system functioning. *Frontiers in Plant Science*, *4*, 1–19. <https://doi.org/10.3389/fpls.2013.00356>
- Valarezo Cely, O., Cañarte Bermúdez, E., Navarrete Cedeño, J. B., & Arias de López, M. (2003). *Prodiplosis longifila* principal plaga de tomate en el Ecuador. INIAP, Estación Experimental Portoviejo, Ecuador. <http://repositorio.iniap.gob.ec/handle/41000/1324>
- Valdez, F. (2008). Inter-zonal Relationships in Ecuador. In H. Silverman & W. H. Isbell (Eds.), *The Handbook of South American Archaeology* (pp. 865–888). Springer. [https://doi.org/10.1007/978-0-387-74907-5\\_43](https://doi.org/10.1007/978-0-387-74907-5_43)
- van de Mortel, J. E., de Vos, R. C. H., Dekkers, E., Pineda, A., Guillod, L., Bouwmeester, K., van Loon, J. J. A., Dicke, M., & Raaijmakers, J. M. (2012). Metabolic and transcriptomic changes induced in *Arabidopsis* by the rhizobacterium *Pseudomonas fluorescens* SS101. *Plant Physiology*, *160*(4), 2173–2188. <https://doi.org/10.1104/pp.112.207324>
- van der Heijden, M. G. A., & Hartmann, M. (2016). Networking in the plant microbiome. *PLOS Biology*, *14*(2), 1–9. <https://doi.org/10.1371/journal.pbio.1002378>
- van Staden, A. D. P., van Zyl, W. F., Trindade, M., Dicks, L. M. T., & Smith, C. (2021). Therapeutic application of lantibiotics and other lanthipeptides: Old and new findings. *Applied and Environmental Microbiology*, *87*(14), 1–24. <https://doi.org/10.1128/AEM.00186-21>
- Vančura, V., & Hovadík, A. (1965). Root exudates of plants: II. Composition of root exudates of some vegetables. *Plant and Soil*, *22*(1), 21–32.
- Vargas, P., Bosmans, L., Van Calenberge, B., Van Kerckhove, S., Lievens, B., & Rediers, H. (2021). Bacterial community dynamics of tomato hydroponic greenhouses infested with hairy root disease. *FEMS Microbiology Ecology*, *97*(12), 1–10. <https://doi.org/10.1093/femsec/fiab153>
- Velasco-Cuervo, S. M., Espinosa, L. L., Duque-Gamboa, D. N., Castillo-Cárdenas, M. F., Hernández, L. M., Guzmán, Y. C., Manzano, M. R., & Toro-Perea, N. (2016). Barcoding, population structure, and demographic history of *Prodiplosis longifila* associated with the Andes. *Entomologia Experimentalis et Applicata*, *158*(2), 217–227. <https://doi.org/10.1111/eea.12397>
- Veloso, J., & Díaz, J. (2012). *Fusarium oxysporum* Fo47 confers protection to pepper plants against *Verticillium dahliae* and *Phytophthora capsici*, and induces the expression of defence genes. *Plant Pathology*, *61*(2), 281–288. <https://doi.org/10.1111/j.1365-3059.2011.02516.x>
- Vikram, S., Ramond, J.-B., Ortiz, M., Maggs-Kölling, G., Pelsler, K., & Cowan, D. A. (2023). Soil fungal diversity and assembly along a xeric stress gradient in the central Namib Desert. *Fungal Biology*, *127*(4), 997–1003. <https://doi.org/10.1016/j.funbio.2023.03.001>
- Vimr, E. R., Kalivoda, K. A., Deszo, E. L., & Steenbergen, S. M. (2004). Diversity of microbial sialic acid metabolism. *Microbiology and Molecular Biology Reviews*, *68*(1), 132–153. <https://doi.org/10.1128/MMBR.68.1.132-153.2004>
- Volpe, V., Chitarra, W., Cascone, P., Volpe, M. G., Bartolini, P., Moneti, G., Pieraccini, G., Di Serio, C., Maserti, B., Guerrieri, E., & Balestrini, R. (2018). The association with two different arbuscular

- mycorrhizal fungi differently affects water stress tolerance in tomato. *Frontiers in Plant Science*, 9, 1–16. <https://www.frontiersin.org/articles/10.3389/fpls.2018.01480>
- Voorrips, R. E., Verkerke, W., Finkers, R., Jongerius, R., & Kanne, J. (2000). Inheritance of taste components in tomato. *Acta Physiologiae Plantarum*, 22(3), 259–261. <https://doi.org/10.1007/s11738-000-0026-1>
- Vorholt, J. A., Vogel, C., Carlström, C. I., & Müller, D. B. (2017). Establishing causality: Opportunities of synthetic communities for plant microbiome research. *Cell Host & Microbe*, 22(2), 142–155. <https://doi.org/10.1016/j.chom.2017.07.004>
- Vosman, B. (2019). QTL mapping of insect resistance components of *Solanum galapagense*. *Theoretical and Applied Genetics*, 132(2), 531–541. <https://doi.org/10.1007/s00122-018-3239-7>
- Wagner, M. R. (2021). Microbe-dependent heterosis in maize. *PNAS*, 118(30), 1–8. <https://doi.org/10.1073/pnas.2021965118>
- Waheed, K., Nawaz, H., Hanif, M. A., & Rehman, R. (2019). Tomato. In M. Hanif, H. Nawaz, M. Khan, & H. Byrne (Eds.), *Medicinal Plants of South Asia Novel Sources for Drug Discovery* (1st ed., pp. 631–644). Elsevier.
- Walker, T. S., Bais, H. P., Grotewold, E., & Vivanco, J. M. (2003). Root exudation and rhizosphere biology. *Plant Physiology*, 132(1), 44–51. <https://doi.org/10.1104/pp.102.019661>
- Wallace, J. G., Kremling, K. A., Kovar, L. L., & Buckler, E. S. (2018). Quantitative genetics of the maize leaf microbiome. *Phytobiomes Journal*, 2(4), 208–224. <https://doi.org/10.1094/PBIOMES-02-18-0008-R>
- Wallden, K., Rivera-Calzada, A., & Waksman, G. (2010). Type IV secretion systems: Versatility and diversity in function. *Cellular Microbiology*, 12(9), 1203–1212. <https://doi.org/10.1111/j.1462-5822.2010.01499.x>
- Wallenstein, M. D. (2017). Managing and manipulating the rhizosphere microbiome for plant health: A systems approach. *Rhizosphere*, 3, 230–232. <https://doi.org/10.1016/j.rhisph.2017.04.004>
- Wang, N., Li, H., Wang, B., Ding, J., Liu, Y., Wei, Y., Li, J., & Ding, G.-C. (2022). Taxonomic and functional diversity of rhizosphere microbiome recruited from compost synergistically determined by plant species and compost. *Frontiers in Microbiology*, 12, 1–12. <https://doi.org/10.3389/fmicb.2021.798476>
- Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology*, 73(16), 5261–5267. <https://doi.org/10.1128/AEM.00062-07>
- Wang, W., & Lu, X. (2013). Microbial synthesis of alka(e)nes. *Frontiers in Bioengineering and Biotechnology*, 1, 1–5. <https://doi.org/10.3389/fbioe.2013.00010>
- Wang, X., Gao, L., Jiao, C., Stravoravdis, S., Hosmani, P. S., Saha, S., Zhang, J., Mainiero, S., Strickler, S. R., Catala, C., Martin, G. B., Mueller, L. A., Vrebalov, J., Giovannoni, J. J., Wu, S., & Fei, Z. (2020). Genome of *Solanum pimpinellifolium* provides insights into structural variants during tomato breeding. *Nature Communications*, 11(5817), 1–11. <https://doi.org/10.1038/s41467-020-19682-0>
- Wells, T. J., & Henderson, I. R. (2013). Chapter 16—Type 1 and 5 secretion systems and associated toxins. In M. S. Donnenberg (Ed.), *Escherichia coli (Second Edition)* (pp. 499–532). Academic Press. <https://doi.org/10.1016/B978-0-12-397048-0.00016-4>
- Wen, T., Xie, P., Liu, H., Liu, T., Zhao, M., Yang, S., Niu, G., Hale, L., Singh, B. K., Kowalchuk, G. A., Shen, Q., & Yuan, J. (2023). Tapping the rhizosphere metabolites for the prebiotic control of soil-borne bacterial wilt disease. *Nature Communications*, 14(1), 1–15. <https://doi.org/10.1038/s41467-023-40184-2>

- White, A. R., Borkum, M. I., Rivas-Ubach, A., Bilbao, A., Wendler, J. P., Colby, S. M., Köberl, M., & Jansson, C. (2017). From data to knowledge: The future of multi-omics data analysis for the rhizosphere. *Rhizosphere*, 3, 222–229. <https://doi.org/10.1016/j.rhisph.2017.05.001>
- Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., ... Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1–6. <https://doi.org/10.21105/joss.01686>
- Wickham, H., François, R., Henry, L., Müller, K., Vaughan, D., Software, P., & PBC. (2023). *dplyr: A Grammar of Data Manipulation* (Version 1.1.3) [Computer software]. <https://cran.r-project.org/web/packages/dplyr/index.html>
- Wille, L., Messmer, M. M., Studer, B., & Hohmann, P. (2019). Insights to plant–microbe interactions provide opportunities to improve resistance breeding against root diseases in grain legumes. *Plant, Cell & Environment*, 42(1), 20–40. <https://doi.org/10.1111/pce.13214>
- Wissuwa, M., Mazzola, M., & Picard, C. (2008). Novel approaches in plant breeding for rhizosphere-related traits. *Plant and Soil*, 321(1), 409–430. <https://doi.org/10.1007/s11104-008-9693-2>
- Wood, D. E., Lu, J., & Langmead, B. (2019). Improved metagenomic analysis with Kraken 2. *Genome Biol.*, 20(257), 1–13. <https://doi.org/10.1186/s13059-019-1891-0>
- Wood, S. (2000). *Tomatl*. Online Nahuatl Dictionary. <https://nahuatl.uoregon.edu/content/tomatl>
- Worsley, S. F. (2021). Investigating the role of root exudates in recruiting *Streptomyces* bacteria to the *Arabidopsis thaliana* microbiome. *Frontiers in Molecular Biosciences*, 8, 1–16. <https://doi.org/10.3389/fmolb.2021.686110>
- Wu, T., Chellemi, D. O., Graham, J. H., & Roskopf, E. N. (2008). Assessment of fungal communities in soil and tomato roots subjected to diverse land and crop management systems. *Soil Biology and Biochemistry*, 40(7), 1967–1970. <https://doi.org/10.1016/j.soilbio.2008.02.012>
- Wu, X., Bei, S., Zhou, X., Luo, Y., He, Z., Song, C., Yuan, H., Pivato, B., Liesack, W., & Peng, J. (2023). Metagenomic insights into genetic factors driving bacterial niche differentiation between bulk and rhizosphere soils. *Science of The Total Environment*, 891, 1–10. <https://doi.org/10.1016/j.scitotenv.2023.164221>
- Wu, Y.-W., Simmons, B. A., & Singer, S. W. (2016). MaxBin 2.0: An automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics*, 32(4), 605–607. <https://doi.org/10.1093/bioinformatics/btv638>
- Wubet, T., Christ, S., Schöning, I., Boch, S., Gawlich, M., Schnabel, B., Fischer, M., & Buscot, F. (2012). Differences in soil fungal communities between European beech (*Fagus sylvatica* L.) dominated forests are related to soil and understory vegetation. *PLOS ONE*, 7(10), 1–14. <https://doi.org/10.1371/journal.pone.0047500>
- Xu, H., Paerl, H. W., Zhu, G., Qin, B., Hall, N. S., & Zhu, M. (2017). Long-term nutrient trends and harmful cyanobacterial bloom potential in hypertrophic Lake Taihu, China. *Hydrobiologia*, 787(1), 229–242. <https://doi.org/10.1007/s10750-016-2967-4>
- Xu, L. (2021). Genome-resolved metagenomics reveals role of iron metabolism in drought-induced rhizosphere microbiome dynamics. *Nature Communications*, 12(3209), 1–17. <https://doi.org/10.1038/s41467-021-23553-7>
- Xue, P.-P., Carrillo, Y., Pino, V., Minasny, B., & McBratney, A. B. (2018). Soil properties drive microbial community structure in a large scale transect in South Eastern Australia. *Scientific Reports*, 8(1), 1–11. <https://doi.org/10.1038/s41598-018-30005-8>

- Yan, J., Smith, M. D., Glick, B. R., & Liang, Y. (2014). Effects of ACC deaminase containing rhizobacteria on plant growth and expression of Toc GTPases in tomato (*Solanum lycopersicum*) under salt stress. *Botany*, *92*(11), 775–781. <https://doi.org/10.1139/cjb-2014-0038>
- Ye, J. (2017). An InDel in the promoter of Al-ACTIVATED MALATE TRANSPORTER9 selected during tomato domestication determines fruit malate contents and aluminum tolerance. *Plant Cell*, *29*(9), 2249–2268. <https://doi.org/10.1105/tpc.17.00211>
- Ye, L., Zhao, X., Bao, E., Li, J., Zou, Z., & Cao, K. (2020). Bio-organic fertilizer with reduced rates of chemical fertilization improves soil fertility and enhances tomato yield and quality. *Scientific Reports*, *10*(177), 1–11. <https://doi.org/10.1038/s41598-019-56954-2>
- Yi, H.-S., Yang, J. W., Ghim, S.-Y., & Ryu, C.-M. (2011). A cry for help from leaf to root. *Plant Signaling & Behavior*, *6*(8), 1192–1194. <https://doi.org/10.4161/psb.6.8.15780>
- Yin, J., Yu, Y., Zhang, Z., Chen, L., & Ruan, L. (2020). Enrichment of potentially beneficial bacteria from the consistent microbial community confers canker resistance on tomato. *Microbiological Research*, *234*, 1–10. <https://doi.org/10.1016/j.micres.2020.126446>
- Yin, Y. (2012). dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Research*, *40*(1), 445–451. <https://doi.org/10.1093/nar/gks479>
- Yoo, S.-J., Shin, D. J., Won, H. Y., Song, J., & Sang, M. K. (2018). *Aspergillus terreus* JF27 promotes the growth of tomato plants and induces resistance against *Pseudomonas syringae* pv. Tomato. *Mycobiology*, *46*(2), 147–153. <https://doi.org/10.1080/12298093.2018.1475370>
- Yue, H., Yue, W., Jiao, S., Kim, H., Lee, Y.-H., Wei, G., Song, W., & Shu, D. (2023). Plant domestication shapes rhizosphere microbiome assembly and metabolic functions. *Microbiome*, *11*(70), 1–19. <https://doi.org/10.1186/s40168-023-01513-1>
- Zarrillo, S., Gaikwad, N., Lanaud, C., Powis, T., Viot, C., Lesur, I., Fouet, O., Argout, X., Guichoux, E., Salin, F., Solorzano, R. L., Bouchez, O., Vignes, H., Severts, P., Hurtado, J., Yepez, A., Grivetti, L., Blake, M., & Valdez, F. (2018). The use and domestication of *Theobroma cacao* during the mid-Holocene in the upper Amazon. *Nature Ecology & Evolution*, *2*(12), 1879–1888. <https://doi.org/10.1038/s41559-018-0697-x>
- Zboralski, A., & Filion, M. (2020). Genetic factors involved in rhizosphere colonization by phyto-beneficial *Pseudomonas* spp. *Computational and Structural Biotechnology Journal*, *18*, 3539–3554. <https://doi.org/10.1016/j.csbj.2020.11.025>
- Zechner, E. L., Lang, S., & Schildbach, J. F. (2012). Assembly and mechanisms of bacterial type IV secretion machines. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1592), 1073–1087. <https://doi.org/10.1098/rstb.2011.0207>
- Zeist, A. R., Silva, A. A., Resende, J. T. V. de, Maluf, W. R., Gabriel, A., Zanin, D. S., & Guerra, E. P. (2018). Tomato Breeding for Insect-Pest Resistance. In S. Tatu Nyaku & A. Danquah (Eds.), *Recent Advances in Tomato Breeding and Production* (pp. 27–46). IntechOpen. <https://doi.org/10.5772/intechopen.75978>
- Zhang, H., Jay Forman, H., & Choi, J. (2005).  $\gamma$ -glutamyl transpeptidase in glutathione biosynthesis. In H. Sies & L. Packer (Eds.), *Methods in Enzymology* (Vol. 401, pp. 468–483). Academic Press. [https://doi.org/10.1016/S0076-6879\(05\)01028-1](https://doi.org/10.1016/S0076-6879(05)01028-1)
- Zhang, J., Wang, P. C., Fang, L., Zhang, Q.-A., Yan, C. S., & Chen, J. Y. (2017). Isolation and characterization of phosphate-solubilizing bacteria from mushroom residues and their effect on tomato plant growth promotion. *Polish Journal of Microbiology*, *66*(1), 57–65. <https://doi.org/10.5604/17331331.1234993>
- Zhang, Z., Xiao, Y. S., Zhan, Y., Zhang, Z., Liu, Y., Wei, Y., Xu, T., & Li, J. (2022). Tomato microbiome under long-term organic and conventional farming. *iMeta*, *1*(3), 1–13. <https://doi.org/10.1002/imt.2.48>

- Zhao, K., Li, J., Zhang, X., Chen, Q., Liu, M., Ao, X., Gu, Y., Liao, D., Xu, K., Ma, M., Yu, X., Xiang, Q., Chen, J., Zhang, X., & Penttinen, P. (2018). Actinobacteria associated with *Glycyrrhiza inflata* Bat. Are diverse and have plant growth promoting and antimicrobial activity. *Scientific Reports*, 8(13661), 1–13. <https://doi.org/10.1038/s41598-018-32097-8>
- Zheng, Q., Hu, Y., Zhang, S., Noll, L., Böckle, T., Dietrich, M., Herbold, C. W., Eichorst, S. A., Woebken, D., Richter, A., & Wanek, W. (2019). Soil multifunctionality is affected by the soil environment and by microbial community composition and diversity. *Soil Biology & Biochemistry*, 136, 1–13. <https://doi.org/10.1016/j.soilbio.2019.107521>
- Zhong, S., Fei, Z., Chen, Y.-R., Zheng, Y., Huang, M., Vrebalov, J., McQuinn, R., Gapper, N., Liu, B., Xiang, J., Shao, Y., & Giovannoni, J. J. (2013). Single-base resolution methylomes of tomato fruit development reveal epigenome modifications associated with ripening. *Nature Biotechnology*, 31(2), 154–159. <https://doi.org/10.1038/nbt.2462>
- Zhou, F., & Pichersky, E. (2020). The complete functional characterisation of the terpene synthase family in tomato. *New Phytologist*, 226(5), 1341–1360. <https://doi.org/10.1111/nph.16431>
- Zhou, L., Song, C., Muñoz, C. Y., & Kuipers, O. P. (2021). *Bacillus cabrialesii* BH5 protects tomato plants against botrytis cinerea by production of specific antifungal compounds. *Frontiers in Microbiology*, 12, 1–12. <https://www.frontiersin.org/articles/10.3389/fmicb.2021.707609>
- Zhu, D., Si, H., Zhang, P., Geng, A., Zhang, W., Yang, B., Qian, W.-J., Gabriel, M., & Sun, J. (2018). Genomics and biochemistry investigation on the metabolic pathway of milled wood and alkali lignin-derived aromatic metabolites of *Comamonas serinivorans* SP-35. *Biotechnology for Biofuels*, 11(338), 1–15. <https://doi.org/10.1186/s13068-018-1341-3>
- Zou, X., Jiang, X., Guan, J., Huang, S., Chen, C., Zhou, T., Kuang, C., Ye, J., Liu, T., Cheng, J., Chen, S., & Yu, S. (2024). Both organic fertilizer and biochar applications enhanced soil nutrition but inhibited cyanobacterial community in paddy soils. *Frontiers in Environmental Science*, 12, 1–12. <https://doi.org/10.3389/fenvs.2024.1376147>
- Zuluaga, M. Y. A., Milani, K. M. L., Miras-Moreno, B., Lucini, L., Valentinuzzi, F., Mimmo, T., Pii, Y., Cesco, S., Rodrigues, E. P., & Oliveira, A. L. M. de. (2021). Inoculation with plant growth-promoting bacteria alters the rhizosphere functioning of tomato plants. *Applied Soil Ecology*, 158, 1–12. <https://doi.org/10.1016/j.apsoil.2020.103784>
- Zuriaga, E., Blanca, J. M., Cordero, L., Sifres, A., Blas-Cerdán, W. G., Morales, R., & Nuez, F. (2009). Genetic and bioclimatic variation in *Solanum pimpinellifolium*. *Genetic Resources and Crop Evolution*, 56(1), 39–51. <https://doi.org/10.1007/s10722-008-9340-z>