



## **Daily resilience: a systematic review of measures and associations with well-being and mental health in experience sampling studies**

Zietse, J.; Keijsers, L.; Hillegers, M. H. J.; Vreeker, A.; Harmelen, A.-L. van; Vries, L.P. de

### **Citation**

Zietse, J., Keijsers, L., Hillegers, M. H. J., Vreeker, A., Harmelen, A. -L. van, & Vries, L. P. de. (2025). Daily resilience: a systematic review of measures and associations with well-being and mental health in experience sampling studies. *Development And Psychopathology*, 1-26. doi:10.1017/S0954579425000197

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/4286371>

**Note:** To cite this publication please use the final published version (if applicable).

## Regular Article

# Daily resilience: A systematic review of measures and associations with well-being and mental health in experience sampling studies

Juul Zietse<sup>1</sup>, Loes Keijsers<sup>2</sup>, Manon H.J. Hillegers<sup>1</sup>, Annabel Vreeker<sup>1,2</sup>, Anne-Laura van Harmelen<sup>3</sup> and Lianne P. de Vries<sup>1,4</sup> 

<sup>1</sup>Department of Child and Adolescents Psychiatry/Psychology Erasmus MC Sophia Children's Hospital, Erasmus University Medical Center, Rotterdam, the Netherlands, <sup>2</sup>Department of Psychology, Education & Child Studies, Erasmus School of Social and Behavioural Sciences, Erasmus University Rotterdam, Rotterdam, the Netherlands, <sup>3</sup>Department of Child and Education Studies, Leiden University, Leiden, the Netherlands. and <sup>4</sup>Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

## Abstract

Resilience is the dynamic process of adapting to or recovering from stressors, maintaining positive mental health. While most studies have investigated resilience after major life events, less is known about resilience in everyday life. To understand how individuals recover from everyday stressors, and associations with other psychosocial variables, well-being and mental health, we conducted a systematic review of studies to daily resilience, i.e., recovery from daily stressors, using the experience sampling method (ESM). Out of 36 included studies, 11 studies investigated daily resilience in youth (10.9–24.7 years) and 25 in adult samples. Daily resilience was operationalized either with self-report items adapted from trait measures (17 studies) or in terms of affective recovery from daily stressors (20 studies). The self-reported ability to recover from daily stressors reflects subjective experiences of coping with stressors, whereas daily resilience as recovery from daily stressors captures the dynamic process, but is understudied in youth. Daily resilience was associated with psychosocial variables, including better sleep quality and greater optimism. Furthermore, individuals with mental health problems consistently showed longer recovery times after daily stressors. Overall, ESM studies highlight that daily resilience could help to identify individuals at-risk for mental health problems. The findings may facilitate timely interventions.

**Keywords:** resilience; daily life; experience sampling method; ecological momentary assessment; youth; adolescents; systematic review

(Received 2 April 2024; revised 3 March 2025; accepted 3 March 2025)

## Introduction

Everyone is exposed to stressors in their lives, which may range from daily hassles, such as a conflict with a neighbor, a poor test result, an angry bus driver, to major stressful life events, such as an earthquake, a divorce, or being bullied. How individuals respond to these stressors depends on both the person and context. If individuals are able to recover from or adapt to the stressor relatively quickly, they show resilience (Ioannidis et al., 2020; Rutter, 2012). Resilience can either be defined as the dynamic process of adaptation or recovery that unfolds after stressful or negative experiences or the positive outcome of functioning well after stress (Kalisch et al., 2017, 2019; Masten et al., 1990, 2021). If individuals are unable to recover or adapt to a stressor, they are at risk for later mental health problems.

From a developmental perspective, resilience early in life, during childhood and adolescence, is particularly important. That is, mental health problems during adolescence have a substantial impact on daily life and later development (Thapar et al., 2012) and

can lead to increased risks of psychopathology later in life (Clayborne et al., 2019; Kim-Cohen et al., 2003; Patel et al., 2007). Currently, around 13%–25% of adolescents experience depression or anxiety (WHO, 2021). This percentage is expected to increase due to heightened stress levels caused by a rapidly changing world, societal challenges, and major crises such as the COVID-19 pandemic and climate change (Daly, 2022; Lass-Hennemann et al., 2023; Poletti et al., 2023; Racine et al., 2021; Wiederhold, 2022). To prevent this increase, a proactive approach and focus on resilience across the lifespan is needed, rather than focusing only on risk factors of mental health problems (Pluess, 2024; Uhlhaas et al., 2023).

Most of the knowledge about resilience stems from research on the long-term responses to major life events (e.g., the loss of a loved one, an earthquake or natural disaster) or adversity in childhood (e.g., parental divorce or emotional or physical abuse) (Fritz et al., 2018; Ioannidis et al., 2020; Werner, 1995). Everyone experiences one or more of these major events in their life, and this generally causes high levels of stress (Kendler et al., 1999; Kessler, 1997; McLaughlin et al., 2020). However, besides major stressors, all of us encounter smaller stressors in their everyday lives, e.g., a fight with a friend or being late for school/work, which can lead to increased negative affect and lower positive affect. Here, we define the process in which individuals recover from these daily hassles

**Corresponding author:** Lianne P de Vries; Email: [lp.devries@erasmusmc.nl](mailto:lp.devries@erasmusmc.nl)

**Cite this article:** Zietse, J., Keijsers, L., Hillegers, M. H. J., Vreeker, A., van Harmelen, A-L, & de Vries, L. P. (2025). Daily resilience: A systematic review of measures and associations with well-being and mental health in experience sampling studies. *Development and Psychopathology*, 1–26, <https://doi.org/10.1017/S0954579425000197>

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



(Almeida, 2005; Montpetit et al., 2010; Seery & Quinton, 2016) in the short-term as “*daily resilience*.” In this structured review, we describe the importance of daily resilience, synthesize the existing body of research on daily resilience and describe the measures used and associations of daily resilience with psychosocial variables, mental health and well-being.

### Definitions and measurement of resilience to major life events or adversity

While our study focusses on daily resilience to smaller daily stressors, we first provide context by discussing the definitions and measurement of resilience to major life events or adversity for two reasons. First, the majority of existing resilience research has investigated recovery from major stressful life events or adversity. Second, these definitions influences how daily resilience can be defined and measured in the context of daily stressors. There is a broad variety of operationalizations and measurements of resilience to major life events or adversity (Anderson & Priebe, 2021; Bonanno, 2012; Schultze-Lutter et al., 2016; Southwick et al., 2014; Walsh et al., 2010). At the moment, research into both adult and youth resilience mostly includes measures of trait resilience using static questionnaires (Mesman et al., 2021). These measures are suitable to assess the self-reported trait aspects of resilience, but cannot capture the dynamic process of recovery from stress. There is thus a growing theoretical consensus that instead of a trait, resilience is either a process or outcome (Kalisch et al., 2017, 2019; van Harmelen, 2022). Resilience can be defined as the process of affective recovery from stressors or as the outcome of this process: individuals that recover quickly from the stress show good well-being and mental health, i.e., *resilient functioning*.

Importantly, within individuals, resilience is not stable over time. Individuals can experience resilience at certain times or in certain contexts, whereas they might be less able to adapt to or recover from stressors at other times or in other contexts (Cahill et al., 2023; Kalisch et al., 2017, 2019; van Harmelen, 2022). Different *resilience factors* can help individuals to be able to recover from major stressful experiences of adversity. According to Ann Masten, resilience after major stressful or traumatic life events is “*ordinary magic*,” such that ‘*resilience comes not from rare qualities, but from the everyday magic of the ordinary, normative human resources*’ (Masten, 2015). Resilience factors include personal and social skills, parental resilience, and social support from friends and family (Fritz et al., 2018; King et al., 2021; Mesman et al., 2021; van Harmelen et al., 2017).

### Measuring daily resilience

Major life events are relatively rare on a daily basis, making it difficult to capture and study the experience of such events and subsequent recovery in real time. Alternatively, smaller stressors and daily hassles occur every day and can be tracked in real-time daily life. We refer to the short-term recovery from these daily stressors as “*daily resilience*.” Since the 1980s, researchers have emphasized the importance of recovery from daily stressors for overall mental health. In fact, it has been argued that the impact of daily stressors on well-being and mental health could be as great or greater than that of major life events (Delongis et al., 1982; Kanner et al., 1981; Monroe, 1983). Recent research seems to confirm the relation between recovery from (multiple) daily stress and mental health, although the number of studies is limited (Charles et al., 2013; Parrish et al., 2011; Vaessen et al., 2017; Wright et al., 2019). For example, slower recovery from daily stressors puts adolescents


at risk for persistence or worsening of psychopathology (Vaessen et al., 2017). Moreover, daily stressors have been suggested to be the mechanism by which major life events influence mental health. Major life events can disrupt daily routines and the (perceived) ability to deal with daily challenges (Wright et al., 2019). In line with this idea, the response to and recovery from daily stressors have been found to mediate the association between major life events and mental health outcomes, while also having independent predictive effects (Eckenrode, 1984; Kanner et al., 1981; Schneiders et al., 2006; Wagner et al., 1988). Therefore, even though small daily stressors do not cause immediate mental health problems, recovery from such stressors are potentially related to future mental health. This may be due to the accumulation of poor short-term recovery and the resulting pile-up of stress. To fully understand this recovery from daily stressors, i.e., daily resilience, and to be able to support individuals in strengthening resilience, we need to identify resilience factors and mechanisms that protect against the development of mental health problems following daily stressors.

To be able to investigate daily resilience, we need valid measurements. The recovery from everyday stressors in real-time requires multiple measurements within or across days. In recent years, major progress has been made in the study of daily life, which has opened up new opportunities to trace the dynamic processes of daily resilience. Daily stressful events and responses to these events can be assessed using the experience sampling method (ESM) or daily diaries via the smartphones of participants (Hamaker & Wichers, 2017; van Roekel et al., 2019). ESM refers to structured self-report diary techniques assessing mood, symptoms, behavior, and context as they occur in daily life and real-time (Csikszentmihalyi & Larson, 1987; Myin-Germeys & Kuppens, 2022; de Vries et al., 2021). Participants receive multiple very short questionnaires on their phone per day, in which they answer, for instance, how they are feeling, where they are, and what they are doing. In daily diary studies, participants receive one questionnaire at the end of the day about their daily mood, behaviors, and events.

### The current study

Despite the increasing recognition of the importance of daily resilience for mental health and the increasing technical feasibility of measuring feelings, behavior, and context in real-time, a structured overview of existing empirical studies, the approaches to measuring daily resilience, and their associations with other variables is lacking. We aim to advance the understanding of daily resilience by systematically reviewing the increasingly popular approach of ESM and daily diary studies on daily resilience. We address two research questions:

1. What are existing approaches to measure daily resilience and what is the quality of these operationalizations? We include studies that measure perceived daily resilience directly via self-report items and studies that measure daily resilience as the recovery of positive or negative affect after experiencing daily stressors, in line with more dynamic accounts in the resilience literature.
2. How does daily resilience relate to psychosocial variables, mental health and well-being in youth and adult samples? Daily resilience during childhood and adolescence might be especially important for well-being and mental health, therefore we discuss the results separately for youth (<25 years) and adult samples (25+ years).



**Figure 1.** PRISMA 2020 flow diagram for new systematic reviews which included searches of databases and registers only. *from: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al., The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ 2021;372:n71. doi: 10.1136/bmj.n71.*

## Methods

This preregistered systematic review (PROSPERO: [https://www.crd.york.ac.uk/PROSPERO/display\\_record.php?RecordID=475693](https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=475693)) is conducted in accordance with the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guidelines (Moher et al., 2009). In October 2023, the medical library of the Erasmus MC Rotterdam conducted the search for relevant articles in the bibliographic databases Embase, Medline ALL, PsycINFO, Web of Science, Cochrane, and Google Scholar. Additional articles that were missed during this search were identified via reference lists of the selected articles. The search strategy included combinations of search terms related to (1) resilience (resilience, stress or affect recovery, adaptation or adjustment) and (2) experience sampling and daily diaries (Ecological Momentary Assessment, Experience Sampling, daily diary, momentary assessment or longitudinal sampling; see Supplementary Material for the full search strategy). A total of 3060 articles were identified across the records, which led to 2183 articles after duplicates removed (see Figure 1).

Two independent reviewers (LdV, JZ) screened the titles and abstracts of collected articles for eligibility. Articles were included if daily resilience was assessed in daily life in daily diary or ESM

studies. There was a strong level of agreement between the reviewers (89.1% interrater reliability, McHugh, 2012). Discrepancies between the reviewers were resolved by discussions.

The full-text papers of the 141 remaining articles were independently assessed by the two reviewers. Exclusion criteria were: (1) no measure of daily resilience to daily stressors, (2) only a static baseline measure of resilience, (3) only a measure of stress reactivity, i.e., assessing the affective response only at the same time point of the stressor instead of recovery, i.e., hours or days later, (4) review papers, or (5) descriptive planned studies or methodological papers not including a resilience measure. The full-text screening resulted in a strong mean interrater reliability of 89.5% and the inclusion of 36 studies.

Data on characteristics, design, and results of the 36 studies were extracted. To assess the quality of included articles, we used a checklist for the design and reporting for ESM studies (van Roekel et al., 2019). This checklist includes 25 items on recruitment, procedure, and materials (see Supplementary Material). Each item was rated as 1, 0.5 or 0, or cannot determine/not applicable, and scores were converted to percentages. Studies rated >80% were considered good quality, 60%–80% was considered fair quality and <60% was considered poor quality. The two reviewers had 94.0% consensus and discrepancies were resolved by discussions.

## Results

We examined the existing approaches to measure daily resilience and the associations with well-being and mental health. The 36 included studies assessed the micro-dynamics of daily resilience via two operationalizations. Seventeen studies used daily or momentary self-report items to assess the perceived ability of participants to recover from daily stressors, hereafter referred to as *self-reported ability to recover from daily stressors*. Twenty studies assessed daily resilience indirectly, in terms of short-term recovery of positive or negative affect after the experience of daily stressors, i.e., *affective recovery*. One study included both a daily self-report item and a measure of affective recovery in 68 adolescents and reported no relation between the measures ( $r = -0.01$  [95%CI:  $-0.21$ ,  $0.19$ ], Kuranova et al., 2021).

Table 1 provides an overview of the samples and characteristics of the 36 included studies, split by daily resilience operationalization (self-report vs affective recovery) and age group (youth, defined as the mean age of the sample below 25 years, and adults). Table 2, 3, and 4 provide the results of the studies. We discuss the findings of the studies separately for youth and adults to take the development of daily resilience into account. Table 3 includes the quality assessment.

### *Self-reported ability to recover from daily stressors*

#### *Study characteristics and quality*

Eight and nine studies respectively assessed self-reported ability to recover from daily stressors in youth samples and adults. Note that Harpøth et al., (2020, 2021) used data from the same participants. Sample sizes ranged from 43 to 248 participants, with an average of 96.4 ( $SD = 45.5$ ) participants. The mean age in the youth samples ranged from 18.1 to 24.7 years, with a mean age of 21.3 ( $SD = 2.5$ ) and in the adult samples from 29 to 46 years ( $M_{age} = 35.5$ ,  $SD = 6.3$ ). The proportion of males was on average 39.3% ( $SD = 22.7\%$ , range 0%–100%). Specific populations were included, such as patients with depression (Heininga & Oldehinkel, 2024; Hoorelbeke et al., 2019) or firefighters (Schwerdtfeger & Dick, 2019) (see Table 1).

Of the 17 studies, seven were of fair quality ( $>60\%$  (Harpøth et al., 2020; Heininga & Oldehinkel, 2024; Hoorelbeke et al., 2019; Jennings et al., 2023; LoSavio et al., 2011; Martinez-Corts et al., 2015; Reininghaus et al., 2023)). The other 10 studies showed poor quality (36%–56%), resulting in an average rating of 54.4% ( $SD = 10.6\%$ ). Only three studies did a power analysis to determine the needed sample size and showed sufficient power to detect effects (Hill et al., 2021; Kuranova et al., 2021; Martinez-Corts et al., 2015). Most studies did not provide full details of the methods used. For example, few studies reported the software used to collect ESM data or response window for the surveys (see Table 3).

Ten studies used daily diaries with one survey per day and lasted on average 19.5 days ( $SD = 25.7$ , range = 5–90 days). The compliance of daily diary or ESM studies gives important information about the study and data quality. The average compliance was reported for six daily diary studies and was on average 88.3% ( $SD = 7.9\%$ ). The seven ESM studies included between 5 to 16 prompts per day or students could choose how often to complete surveys for a maximum of 280 days (Brogly et al., 2024). The average compliance, based on five studies, was 66.0% ( $SD = 19.3\%$ ).

## *Instruments*

Different items or scales were used to assess daily or momentary self-reported ability to recover from daily stressors (see Table 2). The number of items used ranged from 1 to 21 ( $M = 4.9$ ,  $SD = 5.5$ ). The items were mostly adaptations of trait resilience questionnaire items (11 of the 17 studies, 64.7%), e.g., adding “today” or “in this moment” to the items. The other studies either used a self-developed scale, the Monitoring of Actual Resilience State (MARS) scale (Lohner & Aprea, 2021) or items with no source definition. The instruments varied in timescale. Ten studies used items to assess daily self-reported ability to recover (e.g., “Today I could handle what came my way” (Kuranova et al., 2021)). Four studies focused on momentary ability to recover (e.g., “Right now, I feel like I can deal with whatever comes”; Mertens et al., 2023)). Finally, three studies assessed self-reported ability to recover in response to a stressful or negative event (e.g., “I had difficulties to recover” (Reininghaus et al., 2023)). Most studies investigated the self-reported ability to recover from daily stressors in relation to general daily stressors, whereas Jennings et al. (2023) and Martinez-Corts et al. (2015) examined resilience in the context of work conflicts.

Ten of the twelve studies with multiple items reported the internal consistency of the items. The average Cronbach’s alpha of 0.81 ( $SD = .11$ ) indicated good consistency. Moreover, to assess if self-reported ability to recover from daily stressors changes within a person over time, an intraclass correlation coefficient (ICC) was reported in nine studies. The average ICC was 0.52 ( $SD = .15$ , range = .29–.80). This indicates that, on average, 52% of the variance in self-reported ability to recover from daily stressors was due to stable between-person differences and the remaining 48% of the variance in self-reported ability to recover from daily stressors was due to within-participant variability (and/or measurement error).

### *Results and associations of self-reported ability to recover from daily stressors in youth*

First, we assessed mean levels of self-reported ability to recover from daily stressors. Due to the use of different measurement scales, the means could not be directly compared. Therefore, scores were standardized to a range of 1–10 using the R *rescale* function. Mean levels then ranged from 2.5 (LoSavio et al., 2011) to 7.2 (Lohner & Aprea, 2021), with a mean of 5.7 ( $SD = 1.7$ ; see Table 4).

Self-reported ability to recover from daily stressors showed significant associations with different psychological and environmental measures in youth (see Table 4). Some studies investigated under which circumstances youth report more ability to recovery from daily stressors. At this within-person level, LoSavio et al. (2011) found that in 82 students from the USA self-reported ability to recover from daily stressors, defined as stress-related growth, was higher in moments with more positive affect or more negative affect. Similarly, USA students reported higher daily resilience on days with more hope, optimism, or stress (Sytine et al., 2019; Yang, 2020) and days with less somatic symptoms and neuroticism (Yang, 2020).

The other studies provided insight into which adolescents report higher daily ability to recover from daily stressors compared to others. At this between-person level, self-reported ability to recover from daily stressors was positively related to overall trait resilience in both young adults with different levels of risk for

**Table 1.** Descriptives of the 36 included studies, split by operationalization of daily resilience (self-report vs affective recovery) and age (youth vs adults)

Study	N	Self-reported ability to recover from daily stressors										
		Sample population specifics	Country	% male	Mean age	SD age	Range	Method	Survey platform	Duration (days)	Prompts per day	Compliance
<b>Youth samples (&lt; 25 years)</b>												
(Brogly et al., 2024)	94	students	Canada	0.77	–	–	–	ESM	Smartphone	max 280	min of 1	–
(Hill et al., 2021)	62	students who engage in competitive sports	NL	0.34	22	–	–	daily diary	Website	21	1	–
(Kuranova et al., 2021)	68	young adults with different levels of risk for psychosis	NL	0.24	24.7	4.2	–	daily diary	Smartphone	90	1	–
(Lohner & Aprea, 2021)	100	students	Germany	0.34	23.7	2.4	19–30	daily diary	Website	5	1	–
(LoSavio et al., 2011)	82	students	USA	0.18	18.1	0.5	–	daily diary	Website	7	1	93.40%
(Reininghaus et al., 2023)	92	youth at risk for psychological distress	Germany	0.27	21.7	2.5	14–25	ESM	Smartphone	6	8	–
(Sytine et al., 2019)	109	university students	USA	0.45	19	–	–	daily diary	Website	8	1	–
(Yang, 2020)	248	students	USA	0.28	19.7	3.5	–	daily diary	Website	7	1	78%
<b>Adult samples (&gt; 25 years)</b>												
(Harpøth et al., 2020)	72	women with borderline personality disorder	Denmark	0	29.3	8	18–59	daily diary	Website	21	1	96%
(Harpøth et al., 2021)	72	women with borderline personality disorder	Denmark	0	29.3	8	18–59	daily diary	Website	21	1	96%
(Heininga & Oldehinkel, 2024)	86	outpatients with depression	NL	0.44	34.1	12.7	18–64	ESM	Smartphone	28	5 (only 1 resilience)	81.20%
(Hoorelbeke et al., 2019)	85	adults with a history of depression	Belgium	0.38	45.5	13.1	23–65	ESM	Smartphone	7	6	83%
(Jennings et al., 2023)	128	employees	USA	0.38	42.5	11.9	–	daily diary	E-mail	10	3 (only 1 resilience)	82.70%
(Martinez-Corts et al., 2015)	113	employees	Spain	0.4	40	11	–	daily diary	Pen and paper	5	1	83.70%
(Mertens et al., 2023)	71	adults	NL	0.39	29.8	13.5	18+	ESM	Smartphone	7	10	35%
(Schwerdtfeger & Dick, 2019)	43	firefighters	Germany	1	32.7	6.9	–	ESM	Smartphone	1	16	66.70%
(Unger et al., 2023)	90	adults	NL	0.42	30	13.6	18–65	ESM	Smartphone	7	10	64.30%

**Table 1** (Continued)

Study	N	Sample population specifics	Country	Affect recovery after daily stressors					Duration (days)	Prompts per day	Compliance	
				% male	Mean age	SD age	Range	Method				
<b>Youth samples (&lt; 25 years)</b>												
(Bai & Repetti, 2018)	83	children (fifth grade)	USA	0.49	10.9	0.5	–	ESM	Pen and paper	5	5	98%
(Kuranova et al., 2020)	157	adolescents at risk for mental health	Belgium	0.34	17.5	4.0, 3.6	14–34	ESM	Undefined	6	10	–
(Kuranova et al., 2021)	68	young adults with different levels of risk for psychosis	NL	0.24	24.7	4.2	–	daily diary	Smartphone	90	1	–
(Lachowicz et al., 2024)	1031	adolescents	Belgium	0.34	13.8	1.9	11–19	ESM	Provided smartphone	6	10	43%
<b>Adults samples (&gt; 25 years)</b>												
(Ader et al., 2022)	921	individuals with a mental disorder vs at risk vs controls	Germany	0.51	36.5	11.42, 13.12, 12.56	–	ESM	Digital wristwatch - Pen and paper	6	10	75, 78, 82%
(Almeida et al., 2020) 1	115	adults	USA	0.25	41.2	11.6	19–63	ESM	Handheld computer	3	6	89.4%
(Almeida et al., 2020) 2	297	adults	USA	0.5	42.4	12.8	21–70	ESM	Handheld computer	2	Every 45 min	65.0%
(Bergeman & Deboeck, 2014)	783	midlife and older adults	USA	0.41	59.3	10	37–90	daily diary	Pen and paper	3 × 56	1	63.0%
(Congard et al., 2011)	49	students and employees	France	0.39	36	12	19–77	ESM	Pen and paper	40	3	–
(De Calheiros Velozo et al., 2023)	349	adults with depression, at risk and healthy controls	NL	0.23	32.1	12.8	16–65	ESM	Digital wristwatch - Pen and paper	6	10	72%
(Deboeck & Bergeman, 2013)	68	older adults	USA	0.25	79	6.2	–	daily diary	Undefined	56	1	–
(Ekas & Whitman, 2011)	46	mothers of child with autism	USA	0	41	6.6	29–61	daily diary	Pen and paper	15, 10, 20	1	85.60%
(Fleuren et al., 2023)	410	employees	19 countries	0.41	38	13.3	–	daily diary	Website	30	1	63%
(Hamilton et al., 2008)	89	women with fibromyalgia	USA	0	44.4	8.8	–	ESM	Pen and paper	30	3 (only 1 resilience)	98.3%
(Lee et al., 2022)	233	adults	USA	1	76.7	6.5	60–92	daily diary	Pen and paper	3 × 8	1	99%
(Leger et al., 2018)	1155	adults	USA	0.43	55		30–84	daily diary	Telephone interviews	8	1	–
(Marcusson-Clavertz et al., 2022) 1	234	adults	USA	0.34	46.9	10.9	25–65	ESM	Handheld computer	14	5	83%
(Marcusson-Clavertz et al., 2022) 2	117	adults	USA	0.28	44.5	13.8	–	ESM	Handheld computer	7	5	79%
(Marcusson-Clavertz et al., 2022) 3	172	adults	USA	0.49	49.5	16.9	20–80	ESM	Handheld computer	7	5	87%
(Montpetit et al., 2010)	42	older adults	USA	0.2	78.8	6.6	65–92	daily diary	Pen and paper	56	1	–

(Ong et al., 2006) 1a	27	older adults	USA	0.52	72.1	5.3	62–80 daily diary	Pen and paper	45	1	92%
(Ong et al., 2006) 1b	40	older adults	USA	0.5	75.5	6.3	60–85 daily diary	Pen and paper	30	1	96%
(Ong et al., 2006) 2	34	bereaved widows	USA	0	71.9	6.1	61–83 daily diary	Pen and paper	90	1	78%
(Scott et al., 2017)	199	adults	USA	0.33	47.2	10.7	25–65 ESM	Provided smartphones	14	5	85%
(Vaessen et al., 2019)	509	adults with psychosis, early stages, healthy controls	Belgium	0.52	32.2	–	– ESM	Pen and paper/ dedicated electronic device	6	10	75%

Note. NL = the Netherlands, USA = United States of America, ESM = experience sampling method.

psychosis (Kuranova et al., 2021) and German students (Lohner & Aprea, 2021). Furthermore, self-reported ability to recover from daily stressors was positively related to overall conscientiousness and agreeableness (Lohner & Aprea, 2021) and predicted mental health outcomes one year later (Kuranova et al., 2021). Overall, self-reported ability to recover from daily stressors in youth is related to both within-person factors such as positive affect and optimism, and between-person factors such as trait resilience and personality traits.

### *Results and associations of self-reported ability to recover from daily stressors in adults*

Similar to the findings in youth samples, the rescaled mean score for self-reported ability to recover from daily stressors varied from 4.3 (Harpøth et al., 2020, 2021) to 8.0 (Jennings et al., 2023), with a mean of 6.3 (SD = 1.4; see Table 4). Self-reported ability to recover from daily stressors in adults (aged 25 or older) was related to several psychological measures (see Table 4). At a within-person level, self-reported ability to recover from daily stressors was higher on days/moments with more positive affect or less negative affect in both women with borderline personality disorder (Harpøth et al., 2021) and full-time employees in the US (Jennings et al., 2023). Furthermore, self-reported ability to recover from daily stressors was related to an increase in positive affect and decrease in negative affect the next day in outpatients with depression (Heininga & Oldehinkel, 2024), more optimism and less work conflicts in Spanish employees (Martinez-Corts et al., 2015), more daily meaning in life and less depletion in US employees (Jennings et al., 2023), and lower heart rate variability in firefighters (Schwerdtfeger & Dick, 2019).

At a between person level, self-reported ability to recover from daily stressors was positively related to overall trait resilience in both firefighters and adults from the general population (Schwerdtfeger & Dick, 2019; Unger et al., 2023). Furthermore, self-reported ability to recover from daily stressors was consistently positively related higher levels of positive affect or lower levels of negative affect in various samples (Harpøth et al., 2020; Heininga & Oldehinkel, 2024; Hoorelbeke et al., 2019; Jennings et al., 2023). Similarly, self-reported ability to recover from daily stressors was consistently linked to the severity of mental health problems in (sub)clinical samples (Harpøth et al., 2020, 2021; Heininga & Oldehinkel, 2024; Hoorelbeke et al., 2019). For instance, in outpatients with depression, participants with higher self-reported ability to recover from daily stressors showed a better daily mood and fewer depressive symptoms at baseline and three and six months later (Heininga & Oldehinkel, 2024).

Overall, when ability to recover from daily stressors was assessed with self-reported questionnaires, the findings among adolescents and adults provided quite similar patterns. Self-reported ability to recover from daily stressors in adults is associated with various psychological measures across diverse samples, including within-person factors of positive and negative affect, and between-person factors such as trait resilience and severity of mental health problems.

### *Affective recovery after daily stressors*

#### *Study characteristics and quality*

Whereas 17 studies operationalized daily resilience with self-reported instruments to assess perceived ability to recover from daily stressors, 20 studies operationalized the underlying dynamics, in terms of the recovery of positive or negative affect after daily

**Table 2.** Items used for self-reported ability to recover from daily stressors

Youth samples (<25 year)							
Study	Timeframe	Items	Item or, if not all available, examples	Scale	Adapted from	alpha	ICC
Brogly et al., 2024	Daily	2	It was hard for me to snap back when something bad happened today.	1 (Strongly disagree) to 5 (Strongly agree)	Brief Resilience Scale (Smith et al., 2008)	–	–
			It did not take me long to recover from a stressful event today.				
Hill et al., 2021	Daily	6	I tend to bounce back quickly after hard times.	0 (strong disagreement) to 100 (strong agreement)	Brief Resilience Scale (Smith et al., 2008)	.84 (.75–.90)	–
			I have a hard time making it through stressful events.				
			It does not take me long to recover from a stressful event.				
			It is hard for me to snap back when something bad happens.				
			I usually come through difficult times with little trouble.				
			I tend to take a long time to get over setbacks in my life.				
Kuranova et al., 2021	Daily	1	Today I could handle what came my way	0 (Not at all) to 100 (Very much)	Undefined	N.A.	–
Lohner & Aprea, 2021	Daily	8	Today... I had support when I needed it.	1 (strongly disagree) to 100 (strongly agree)	Monitoring of Actual Resilience State (MARS) scale (self-developed)	0.75 (b: 0.79, w: 0.73)	0.29
			I could rely on myself to overcome challenges.				
			I did not give up in the face of adversities.				
			I dealt well with negative emotions.				
			My actions did not lead to a higher goal.				
			I had difficulties with recovering from stress.				
			I should have tried harder to achieve my goals.				
			I lacked something to overcome challenges.				
LoSavio et al., 2011	About negative event	21	I changed my priorities about what is important in life	1 (Not at all) to 6 (To a very great degree)	Posttraumatic Growth Inventory (Tedeschi & Calhoun, 1996)	.90 (within)	–
Reininghaus et al., 2023	About negative event	1	I had difficulties to recover	1 (not at all) to 7 (very much)	Undefined	N.A.	–
Sytine et al., 2019	Daily	6	I usually managed difficulties one way or another	1 (Strongly disagree) to 7 (Strongly agree)	Psychological Capital Questionnaire (Luthans et al., 2007)	.68–.82	0.58
			When I had a setback, I had trouble recovering from it, moving on				
Yang, 2020	Daily	3	Today, I got through difficult times because I've experienced difficulty before	1 (totally disagree) to 5 (totally agree)	Resilience scale (Luthans et al., 2007)	.66–.82	0.57
			Today, I felt I could handle many things at a time				
			Today, I took stressful things in my stride.				

**Table 2** (Continued)

Adult samples (>25 year)							
Study	Timeframe	Items	Item or, if not all available, examples	Scale	Adapted from	alpha	ICC
Harpøth et al., 2020; 2021	Daily	14	I quickly get over and recover from being startled  I get over my anger at someone reasonably quickly  My daily life is full of things that keep me interested  I enjoy dealing with new and unusual situations	1 (does not apply at all) to 4 (applies very strongly)	Ego-Resiliency 89 scale (Block & Kremen, 1996)	0.82	0.61
Heininga & Oldehinkel, 2024	Daily	1	To which extent are you able to handle today's challenges?	0 (Not at all) to 100 (very much)	Undefined	N.A.	0.4
Hoorelbeke et al., 2019	Momentary	1	To what extent do you feel resilient?	1 (not at all) to 100 (very much)	Undefined	N.A.	0.53
Jennings et al., 2023	Daily	3	Today at work, I managed difficulties one way or another  Today at work, I felt I could handle many things at a time  Today at work, I took stressful things in stride	1 (Strongly disagree) to 5 (Strongly agree)	Resilience scale (Luthans et al., 2007)	0.82	-
Martinez-Corts et al., 2015	Daily	3	Today, I got through difficult times because I've experienced difficulty before  Today, I felt I could handle many things at a time  Today, I took stressful things in my stride.	1 (totally disagree to 5 (totally agree)	Resilience scale (Luthans et al., 2007)	0.58	0.39
Mertens et al., 2023	Momentary	2	Right now, I feel like I can deal with whatever comes.  Right now, I feel like I can handle unpleasant situations	1 (Strongly disagree) to 7 (Strongly agree)	modified CD-RISC (Dong et al., 2013)	-	-
Mertens et al., 2023	About negative event	1	I could handle the event	1 (Strongly disagree) to 7 (Strongly agree)	Undefined	N.A.	-
Schwerdtfeger & Dick, 2019	Momentary	4	I am determined  I am cheerful  I am content  I am interested in the things I am doing	1 (not at all) to 7 (very much so)	Resilience scale RS-25 (Wagnild & Young, 1993)	0.91	0.8
Unger et al., 2023	Momentary	2	I feel like I can deal with whatever comes.  I feel like I can handle unpleasant situations.	1 (not at all) to 7 (very)	Connor-Davidson Resilience Scale (Connor & Davidson, 2003)	0.94	0.5

Note. N.A. = not applicable. Cronbach's alpha is reported for between-person measures. If the Cronbach's alpha is based on within-person measures, this is defined: B = between-person Cronbach's alpha, W = within-person Cronbach's alpha.

**Table 3.** Quality assessment of 36 included studies

Study	1. Recruitment method	2. A priori power analysis	3. Devices (or on paper)	4. Software (NA when paper)	5. Prompt design	6. Study duration	7. Response window (NA when paper)	8. Items per assessment	9. Assessments per day	10. Inclusion criteria	11. Instructions participants	12. Incentive structure	13. Monitoring scheme	14. Problems data collection	15. Adjustments to protocol	16. Questionnaire duration	17. Overall compliance	18. Reasons noncompliance	19. Time lag (NA for paper)	20. Patterns noncompliance	21. Exclusion participants	22. Compliance after exclusion	23. Scale construction	24. Current state	25. Psychometric properties	Quality %	
<b>Self-reported ability to recover from daily stressors</b>																											
<b>Youth</b>																											
Brogly et al., 2024	1	0	1	1	.5	1	.5	1	1	1	0	1	1	0	0	0	.5	0	0	0	0	NA	0	1	0	42%	
Hill et al., 2021	1	1	.5	0	1	1	1	1	1	1	1	0	0	0	0	0	5	0	0	0	0	1	0	0	.5	0	40%
Kuranova et al., 2021	1	1	1	0	1	1	0	.5	1	1	0	0	0	0	0	0	0	0	0	0	0	1	NA	1	1	0	42%
Lohner & Aprea, 2021	1	0	1	1	.5	1	0	1	1	.5	1	1	0	0	0	0	0	0	0	0	0	1	0	1	1	1	48%
LoSavio et al., 2011	1	0	0	0	1	1	1	1	1	0	1	1	1	0	0	0	1	0	0	0	0	1	1	1	1	1	60%
Reininghaus et al., 2023	1	0	1	1	1	1	0	1	1	1	0	1	1	0	0	0	1	0	0	0	1	NA	1	1	1	67%	
Sytine et al., 2019	.5	0	.5	0	1	1	0	1	1	0	1	1	0	0	0	0	0	0	0	0	0	1	1	1	1	1	36%
Yang, 2020	1	0	.5	0	1	1	1	.5	1	0	1	1	0	0	0	0	1	0	0	0	1	0	NA	1	1	0	46%
<b>Adults</b>																											
Harpøth et al., 2020	1	0	1	.5	1	1	1	1	1	1	1	.5	0	0	0	0	1	0	0	0	1	1	1	1	1	1	64%
Harpøth et al., 2021	1	0	0	1	0	1	0	1	1	1	1	1	0	0	0	0	1	0	0	0	1	1	0	1	1	1	56%
Heininga & Oldehinkel, 2024	1	0	.5	1	1	1	1	1	1	1	0	1	0	0	1	0	0	0	0	1	1	1	1	1	1	1	64%
Hoorelbeke et al., 2019	1	0	.5	1	1	1	1	1	1	1	1	1	0	0	0	0	0	1	0	1	1	1	1	1	1	1	68%
Jennings et al., 2023	1	0	1	1	1	1	.5	1	1	1	.5	1	0	0	0	0	0	1	0	1	1	1	1	1	1	1	60%
Martinez-Corts et al., 2015	1	1	1	NA	1	1	NA	1	1	1	1	1	0	0	0	NA	.5	0	NA	0	.5	NA	1	1	1	1	70%
Mertens et al., 2023	1	0	.5	1	1	1	1	.5	1	1	1	1	.5	1	0	0	0	0	0	1	1	0	1	0	1	0	52%
Schwerdtfeger & Dick, 2019	.5	0	.5	1	1	1	1	1	1	1	1	1	0	0	0	0	1	0	0	0	1	NA	1	1	1	1	54%
Unger et al., 2023	1	0	1	1	1	1	1	1	1	1	0	1	0	0	0	0	1	0	0	0	1	0	0	1	1	1	56%
<b>Affective recovery</b>																											
<b>Youth</b>																											
Bai & Repetti, 2018	1	0	1	NA	1	1	NA	1	1	1	1	1	0	0	0	0	1	0	NA	0	1	NA	1	1	1	1	71%
Kuranova et al., 2020	1	0	1	1	1	1	0	.5	1	1	0	0	0	0	0	0	0	0	0	0	1	0	1	1	1	1	44%

Kuranova et al., 2021	1	1	1	0	1	1	0	.5	1	1	0	0	0	0	0	0	0	0	1	NA	1	1	0	42%			
Lachowicz et al., 2024	1	0	1	1	1	1	0	.5	1	0	1	1	0	0	0	0	0	1	0	0	1	NA	1	1	1	54%	
<b>Adults</b>																											
Ader et al., 2022	1	0	.5	NA	1	1	1	1	1	1	.5	0	0	0	0	0	1	0	0	0	1	0	1	1	0	46%	
Almeida et al., 2020 1	1	0	1	1	1	1	0	1	1	1	1	0	0	0	0	0	1	0	0	0	1	NA	1	1	0	54%	
Almeida et al., 2020 2	0	0	.5	1	1	1	0	1	1	0	1	0	0	0	1	0	1	0	0	0	1	NA	1	1	0	46%	
Bergeman & Deboeck, 2014	1	0	1	NA	1	1	NA	1	1	0	1	1	0	0	0	NA	1	0	NA	0	1	0	1	1	1	62%	
Congard et al., 2011	1	0	1	NA	1	1	NA	1	1	0	1	1	0	0	0	NA	0	0	NA	0	0	0	1	1	1	52%	
De Calheiros Velozo et al., 2023	1	0	1	NA	1	1	0	.5	1	1	0	0	0	0	0	0	1	0	0	0	.5	0	1	1	1	42%	
Deboeck & Bergemann 2013	1	0	0	0	1	1	0	1	1	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	1	40%	
Ekas & Whitman, 2011	1	0	1	NA	1	1	NA	1	1	1	1	1	0	0	0	NA	1	0	NA	1	1	NA	1	1	1	75%	
Fleuren et al., 2023	1	0	.5	0	1	1	1	1	1	0	1	1	0	0	0	0	1	0	0	0	1	NA	1	1	1	54%	
Hamilton et al., 2008	1	0	1	1	1	1	1	.5	1	1	1	1	1	0	0	0	1	0	0	0	1	NA	1	1	1	67%	
Lee et al., 2022	1	0	1	NA	1	1	NA	1	1	1	1	1	0	0	0	0	NA	1	0	NA	1	1	NA	1	1	1	70%
Leger et al., 2018	1	1	1	1	1	1	NA	1	1	1	.5	0	1	0	0	NA	0	0	NA	0	0	0	1	1	1	57%	
Marcusson-Clavertz et al., 2022	1	0	.5	0	1	1	0	1	1	1	1	0	0	0	0	0	1	0	0	0	0	0	1	1	.5	40%	
Montpetit et al., 2010	1	0	1	NA	1	1	NA	1	1	.5	.5	0	0	0	0	0	0	0	0	0	1	1	0	1	1	48%	
Ong et al., 2006	1	0	1	NA	1	1	NA	1	1	.5	1	1	0	0	0	NA	1	0	NA	0	1	NA	1	1	1	65%	
Scott et al., 2017	1	0	1	0	1	1	.5	1	1	1	1	1	0	0	1	1	0	0	0	1	NA	1	1	1	67%		
Vaessen et al., 2019	1	0	1	0	1	1	0	1	1	1	0	0	0	0	0	0	1	0	0	0	1	NA	1	1	0	46%	

Note. N.A. = not applicable. Quality ratings are percentages of all non-N.A. ratings. <60% is poor quality, 60%-80% is fair quality, and >80% is good quality.

**Table 4.** Results of the studies assessing self-reported ability to recover from daily stressors and associations with other measures

Youth samples (<25 year)			
Study	Mean daily resilience (SD)	Rescaled mean 1–10	Associations and other results
Brogly et al., 2024	~2.5	~4.4	Step count (within-person): $B = -.001 (.03)$ , $p = .958$ GPS locations visited (within-person): $B = -.004 (.03)$ , $p = .861$ Floors walked (within-person)*: $B = -1.07 (.03)$ , $p < .001$ Campus proximity (within-person)*: $-0.09 (.04)$ , $p = .008$
Hill et al., 2021	54.76 (7.04)	5.9	Individual differences*: y-intercept: 20.23, slope = .57, CI .47–.71
Kuranova et al., 2021	52.18 (13.7)	5.7	Trait resilience*: $r = .35$ , $p < .03$ Psychopathological symptoms one year later*: $\beta = -0.22$ , $p < 0.001$
Lohner & Aprea, 2021	~68/70	7.2	Trait resilience*: $r = .09$ –.22, $p < .05$ Conscientiousness*: $r = .21$ –.29, $p < .05$ Agreeableness*: $r = .19$ –.21, $p < .05$ Neuroticism: $r = -.06$ –.16, $p = \text{ns}$ Extraversion: $r = .05$ –.11, $p = \text{ns}$ Openness: $r = .04$ –.10, $p = \text{ns}$ Satisfaction with life*: $r = .39$ , $p < .05$ Effect resilience journal intervention: $F (1,98) = 0.32$ , $p = 0.574$ , $\eta^2 = .003$
LoSavio et al., 2011	1.82 (1.03)	2.5	Positive affect (within-person): $r = .18$ Negative affect (within-person): $r = .09$ Daily core belief disruption (within-person): $r = .37$ Daily stress reactive rumination (within-person): $r = .27$ Daily resolution of day's worst event (within-person): $r = -.04$
Reininghaus et al., 2023	–	–	Intervention*: $B = 0.55$ , 95% CI 0.18–0.92, $d = 0.33$
Sytine et al., 2019	4.83 (.91)	6.7	Daily hope (within-person)*: $r = .56$ , $p < .01$ Daily optimism (within-person)*: $r = .6$ , $p < .01$ Daily self-efficacy (within-person)*: $r = .56$ , $p < .01$ Daily uplifts (within-person)*: $r = .35$ , $p < .01$ Daily savoring (within-person)*: $r = .31$ , $p < .01$ Daily demands (within-person)*: $r = -.21$ , $p < .01$ Psychological capital (within-person)*: $r = .83$ , $p < .01$
Yang, 2020	3.75 (0.92)	7.2	Daily somatic symptoms (within-person)*: $r = -.19$ , $p < .01$ Daily stress (within-person)*: $r = .01$ , $p < .05$ Daily negative affect (within-person): $r = -.12$ Neuroticism*: $r = -.17$ , $p < .01$ Aggression: $r = -.12$ Resilience $\times$ aggression $\rightarrow$ daily somatic symptoms*: $b = -0.07$ , 95% CI: [ −0.13, 0.00], $p = 0.04$
Adult samples (>25 year)			
Study	Mean daily resilience (SD)	Rescaled mean 1–10	Associations and other results
Harpøth et al., 2020; 2021	2.09 (.47), within: .36	4.3	Positive affect*: $r = .82$ , $p < .001$ Negative affect*: $r = -.28$ , $p < .001$ Quality of life*: $r = .63$ , $p < .001$ Number other mental disorders*: $\beta = 0.01$ , $SE = 0.03$ , $p = .843$ Positive affect same day*: $B = 0.39$ , $p < .001$ Positive affect day before*: $B = 0.41$ , $SE = 0.03$ , $p < .001$

(Continued)

**Table 4.** (Continued)

Harpoth et al., 2021	2.09 (.47), within: .36	4.3 Positive affect (within-person)*: $r = .16, p < .05$ Negative affect (within-person)*: $r = -.09, p < .05$ Quality of life (within-person)*: $r = .17, p < .05$ Borderline personality disorder symptoms severity (within-person)*: $r = -.07, p < .05$ Positive affect mediator*: resilience -> BPD $-0.05^*$ vs .02 not included Negative affect mediator*: resilience -> BPD $-0.07^*$ vs .02 not included
Heininga & Oldehinkel, 2024	53.07 (14.21)	5.8 Positive affect*: $r = .58, p < .001$ Negative affect*: $r = -.62, p < .001$ Confidence and purpose*: $r = .22, p < .05$ Overall mood*: $r = .74, p < .001$ Dysfunction in the daily life*: $r = -.43, p < .001$ Baseline depression*: $r = -.31; p = < .001$ Increase in positive affect next day (within-person)*: $b = .20; p = .002$ Decrease in negative affect next day (within-person)*: $b = -.15; p = .004$ Depressive symptoms*: $B = -.49, p < .001$ and $B = -.55, p < .001$ three and six months later General dysfunction*: $B = -.48, p < .001$ and $B = -.57, p < .001$ three and six months later Age: $r = -.18$ , Gender: $r = -.03$ , Rumination: $r = -.12$ , Self-management: $r = .06$ , Social support: $r = .20$ , Connectedness: $r = .12$ , Professional help: $r = .01$ , Having a caring community $r = .10$ , all $p = \text{ns}$
Hoorelbeke et al., 2019	50.26 (15.14)	5.5 Positive affect*: $r = .76, p < .001$ Positive appraisal*: $r = .64, p < .001$ Depressive complaints*: $r = -.30, p < .001$ Cognitive complaints*: $r = -.15, p < .001$ Rumination*: $r = -.07, p < .001$ Networks (within-person): resilience ? decreased rumination, cognitive complaints, depressive symptoms and increased positive appraisal over time.
Jennings et al., 2023	4.13 (0.64)	8 Positive affect*: $r = .51, p < .01$ , within person*: $r = .15, p < .01$ Negative affect*: $r = -.29, p < .01$ , within person*: $r = -.08, p < .05$ Insomnia*: $r = -.26, p < .01$ , within person: $r = .01, p = \text{ns}$ Daily work self-esteem*: $r = .70, p < .01$ , within person: $r = -.02, p = \text{ns}$ Daily work engagement*: $r = .71, p < .01$ , within person: $r = -.01, p = \text{ns}$ Daily depletion*: $r = -.42, p < .01$ , within person*: $r = -.19, p < .01$ Daily meaning in life*: $r = .62, p < .01$ , within person*: $r = .12, p < .01$
Martinez-Corts et al., 2015	3.83 (.51)	7.4 Task conflict*: $r = -.12, p < .05$ , within person: $r = -.17$ Work related relationship conflict*: $r = -.19, p < .01$ , within person: $r = -.14$ Optimism*: $r = .22, p < .01$ , within person: $r = .40$ Strain-based work conflict $\times$ resilience on interpersonal conflict*: $\gamma = 0.33, \text{SE} = 0.08, z = 3.85, p < .001$
Mertens et al., 2023	5.09	7.1 Spiritual vs religious individuals*: $B = 1.21, \text{SE} = 0.53, t = 2.27, p = .026$
Schwerdtfeger & Dick, 2019	5.16 (1.19)	7.2 Trait resilience*: $r = .58, p < .01$ Heart rate variability (within-person)*: $b = -.14, \text{CI: } -.12, -.03$ Stress situation $\times$ resilience on heart rate variability (within-person)*: $b = -.033, \text{SE} = 0.16, \text{CI: } -.64, -.07$
Unger et al., 2023	5 (1.44)	7 Trait resilience*: $r = .68, p < .001$ Momentary loneliness*: $r = -.66, p < .001$ Momentary self-esteem*: $r = .84, p < .001$

**Table 5.** Results of the studies that operationalized daily resilience as affective recovery after daily stressors and associations with other measures

Youth samples (<25 year)					
Study	Time scale recovery	Daily resilience operationalization	Recovery NA or PA (Mean, SD)	Stressors	Results / associations with daily recovery
Bai & Repetti, 2018	Within a day	association between daily school problems and bedtime negative and positive emotion	NA: -0.39 (0.28)  PA: -0.32 (0.20)	School problems: 5 academic problems, 5 peer problems: 1 (definitely false) to 4 (definitely true).	Depressive symptoms (CDI)*: $r = -.37^*$  Internalizing problems (parent report)*: $r = -.24^*$  Externalizing problems (parent report): $r = -.02$  Depressive symptoms (CDI)*: $r = -.31^*$  Internalizing problems (parent report)*: $r = -.13^*$  Externalizing problems (parent report): $r = -.03$
Kuranova et al., 2020	Within a day	multilevel models with negative affect as dependent variable and lagged variables of pleasantness of event as predictor (t, t-1, t-2, ... t-4) to assess amount of time until the effect of unpleasant events on negative/positive affect is no longer significantly different from the person-specific mean of negative/positive affect. Extract random slopes per individual per model to compute area under the curve with respect to baseline (AUCb)	NA: stable: lag 1 (~90 min), increase: lag 2 (~180 min)  PA: stable: lag 1 (~90 min), increase: lag 2 (~180 min)	Most important event and how pleasant/unpleasant this event was: (-3) very unpleasant to (+3) very pleasant	Group*: $B = 0.05^*$ , $p = 0.02$  Psychopathological symptoms: $\beta = 0.09$ , $p = 0.051$  Group: $B = -0.06$ , $p = 0.15$  Psychopathological symptoms: $\beta = -0.02$ , $p = 0.74$
Kuranova et al., 2021	Across multiple days	speed of affect recover: amount of time it takes until the effect of unpleasant events on negative/positive affect is no longer significantly different from the person-specific mean of negative/positive affect	NA: 5.11 (4.33)	Most important negative event and how unpleasant this event was: (0) Very unpleasant to (100) Neutral	Self-report daily resilience: $r = -0.01$ (CI: -0.21 -0.19)  Trait resilience: $r = -0.16$ (CI: -0.35 -0.04)  Psychopathology: $\beta = -0.01$ (CI: -0.13 - 0.11)
Lachowicz et al., 2024	Within a day	delineate the event interval, i.e. the time between stressor (t0) and the moment that negative affect had recovered to baseline values, using survival analysis	Valence: 138.95 (24.37) min	Activity stress: I would rather do something else + This is difficult for me + I can do this well: 1 (not at all) to 7 (very much)	Anxiety symptoms*: $\beta = .071$ , SE = .034, $p = .036$

					Cumulative stress*: $\beta=0.432$ , SE = 0.057, $p < .001$
					Anxiety symptoms (control for depr)*: $\beta=.111$ , SE = .051, $p = .029$
					Depression: $\beta=-0.048$ , SE = 0.044, $p = .275$
					Anxiety symptoms ~ 3 years later: $\beta=-0.001$ , SE = 0.003, $p = .661$
			Arousal: 133.49 (19.39) min		Anxiety symptoms: $\beta=.042$ , SE = .036, $p = .246$
					Cumulative stress*: $\beta=0.339$ , SE = 0.053, $p < .001$
					Anxiety symptoms (control for depr)*: $\beta=.105$ , SE = .048, $p = .029$
					Depression: $\beta=-0.077$ , SE = 0.039, $p = .050$
					Anxiety symptoms ~ 3 years later: $\beta=-0.00$ , SE = 0.002, $p = .885$
Adult samples (>25 year)					
Study	Time scale recovery	Daily resilience operationalization	Recovery NA or PA (Mean, SD)	Stressors	Results / associations with daily recovery
Ader et al., 2022	Within a day	multilevel models with negative affect as dependent variable and time since negative event ( $-1, t1, t2, \dots, t8$ ) as predictor. Estimate contrasts between $tj$ for $j=1, \dots, 8$ vs $t-1$ . No longer significant = recovery	PA: patients and at-risk: $t2$ (~180 min), HC: $t1$ (~90 min)	Event related stress: most important event + pleasantness. Activity related stress: this costs energy, I'm skilled at this, this is a challenge, and I prefer doing something else: (1) very pleasant/not at all to (7) very unpleasant/very much	Group: patients vs controls: $b = .01$ , $p = .89$ , at-risk vs controls: $b = .02$ , $p = .62$ , and patients vs at-risk: $b = .02$ , $p = .74$
Almeida et al., 2020 1	Within a day	NA score on a given stressor moment ( $n$ ) minus the NA score the subsequent non-stressor moment ( $n+1$ : 2 hour later)	NA: 0.07 (0.82)	Experience of a stressor + perceived stressfulness: (0) no or (1) yes AND (0) not at all to (6) very much	Negative affect*: $r = .22^*$ , and within-person $r = -.55^*$
					Perceived stress*: $r = .21^*$ and within-person $r = -.17^*$
Almeida et al., 2020 2	Within a day	NA score on a given stressor moment ( $n$ ) minus the NA score the subsequent non-stressor moment ( $n+1$ : 45 min later)	NA: 0.41 (0.54)	Experience of a stressor + perceived stressfulness: (0) no or (1) yes AND (1) not at all to (7) extremely	Negative affect*: $r = .07$ and within-person $r = -.12^*$

(Continued)

Table 5. (Continued)

Adult samples (>25 year)					
Study	Time scale recovery	Daily resilience operationalization	Recovery NA or PA (Mean, SD)	Stressors	Results / associations with daily recovery
					Perceived stress: $r = .01$ and within-person $r = .01$
					Physical activity: 0–10 min: $B = 0.14(0.17)$ , 0–60 min: $B = -0.05 (0.13)$ , 0–120 min: $B = -0.04 (0.11)$ (between-person) and 0–10 min: $B = -0.12 (0.24)$ , 0–60 min: $B = -0.17 (0.18)$ , 0–120 min: $B = -0.23 (0.17)$ (within-person)
Bergeman & Deboeck, 2014	Across multiple days	Reservoir model: dissipation of stress	Stress: wave 1: -0.655 (.627), wave 3: -0.719 (.686), wave 5: -0.776 (.720)	11 items Perceived Stress Scale: (0) strongly agree to (4) strongly disagree	Trait resilience*: $r = -.20^*$ , Ego resilience*: $r = -.15^*$
					Social Coping*: $r = -.14^*$ , Friend Support*: $r = -.17^*$ , Family Support*: $r = -.14^*$
					Environment Mastery*: $r = -.25^*$ , Self-Esteem*: $r = -.23^*$
					Depression*: $\beta = 5.61^*$ , SE = .65
					Self-reported health*: $\beta = 0.97^*$ , SE = .38
Congard et al., 2011	Across multiple days	attractor strength: the coefficient linking the speed of change in affect level to the distance between the current affect level and the homebase, taking into account events	NA: .35 (.51)	Rate the events that had happened to them since last survey: (-3) extremely negative impact to (+3) extremely positive impact	Anxiety: $B = .004$ , SE = .03, $p = .51$
					Positive affect (within-person)*: $B = .15^*$ , SE = .15, $p = <.01$
			PA: 54 (.48)		Anxiety: $B = .001$ , SE = .01, $p = .67$
					Negative affect (within-person)*: $B = .45^*$ , SE = .13, $p = <.01$
De Calheiros Velozo et al., 2023	Within a day	delineate the event interval, i.e. the time between stressor (t0) and the moment that negative affect had recovered to baseline values, using survival analysis	NA	Most important event and how pleasant/ unpleasant this event was: (-3) very unpleasant to (+3) very pleasant	Depression group*: SSD and RRS: 53 % and 49 % less likely to have recovered at any given time point compared to HC. There is a 50 % chance of HC not recovering after 56 min or below versus SSD and RRS groups at 89 min and 78 min.
Deboeck & Bergemann 2013	Across multiple days	Reservoir model: dissipation of negative affect after stress	NA: -1.54 (1.01)	10 items Perceived Stress Scale: (0) never to (4) very often	Neuroticism*: $B = .29^*$ [CI: 0.01, 0.56], $p = .04$ , $r^2 = .08$
Ekas & Whitman, 2011	Next day	lagged associations between daily stress and the next day's negative affect	NA	General stress: 16 items from the Small Life Events Scale (Zautra et al., 1986). Child related stress: 8 item Child-Related Stress measure: (1) not at all stressful to (7) extremely stressful	Days in which both child and life stress were above average (within-person)* Positive affect (within-person)*: buffers negative affect recovery

Fleuren et al., 2023	Across multiple days	change in slope after the event relative to before the event: discontinuous random coefficient growth modeling (RCGM)	NA	COVID-19 diagnosis of a close other: (0) no or (1) yes	Burnout*: $\gamma = -.01^*$ , $p < .05$ . High burnout → steeper negative slope in negative affect after a COVID-19 event.
			PA		Burnout*: $\gamma = .01^*$ , $p < .05$ . High burnout → more positive slope in positive affect after a COVID-19 event.
Hamilton et al., 2008	Next day	lagged associations between daily stress $\times$ sleep and the next day's negative affect and positive affect	NA	Inventory of Small Life Events: (1) not at all desirable to (7) extremely desirable	Sleep duration (within-person)*: interaction with the lagged effect of negative events ( $B = -.03^*$ ). More stress + less sleep → decreased PA, whereas more stress + more sleep → no effect PA.
			PA		Sleep duration (within-person)*: interaction with the lagged effect of negative events ( $B = -.03^*$ ). More stress + less sleep → increased NA, whereas more stress + more sleep → no effect NA.
Lee et al., 2022	Next day	the slope derived from regressing daily negative or positive affect on the presence of prior-day stressors	NA	Daily Inventory of Stressful Event: 7 items: (0) no or (1) yes	Optimism: $B = -.02$ , 95% CI: $-.24$ , $.21$
					Average stressor exposure: $B = -.37$ , 95% CI: $-.18$ , $.93$
			PA		Optimism: $B = -.24$ , 95% CI: $-.26$ , $.73$
					Average stressor exposure*: $3.88^*$ , 95% CI: $1.08$ , $6.69$
Leger et al., 2018	Next day	slope derived from regressing daily negative or positive affect on the presence of previous day stressors	NA: $-.012$ (.032)	Daily Inventory of Stressful Event: 7 items: (0) no or (1) yes	Individual differences in slope*: $.005^*$
					Chronic conditions*: $r = -.127^*$
					Chronic conditions + 10 years*: $r = -.096^*$
Marcusson-Clavertz et al., 2022 I	Within a day	Proximal recovery: difference between negative affect on the same assessment as a stressor and negative affect on the subsequent non-stressor assessment	proximal: $1.2$ (1.3), distal: $-.04$ (1.1)	Experience of a stressful event: (0) no or (1) yes	Sleep duration (within-person): $B = -.81$ , SE = $.52$ (prox), $B = -.50$ , SE = $.75$ (dis)
					Sleep quality (within-person): $B = -.55$ , SE = $.41$ (prox), $B = .34$ , SE = $.60$ (dis)
					Sleep latency (within-person): $B = 1.01$ , SE = $.53$ (prox), $B = -1.31$ , SE = $.76$ (dis)
Marcusson-Clavertz et al., 2022 II	Within a day		proximal: $1.4$ (1.5), distal: $-.01$ (1.6)	Experience of a stressful event: (0) no or (1) yes	Duration (within-person): $B = -.65$ , SE = $.69$ (prox), $B = 1.32$ , SE = $1.17$ (dis)
		Distal recovery: difference between the typical negative affect score and the negative affect score of the subsequent non-stressor assessment. (4-6 hour post stressor)			Quality (within-person): $B = -.78$ , SE = $.83$ (prox), $B = 1.75$ , SE = $1.33$ (dis)
Marcusson-Clavertz et al., 2022 III	Within a day		proximal: $1.4$ (2.1), distal: $-.06$ (1.1)	Experience of a stressful event: (0) no or (1) yes	Duration (within-person): $B = 0.26$ , SE = $.91$ (prox), $B = -1.36$ , SE = $1.13$ (dis)
					Quality (within-person): $B = 1.10$ , SE = $.87$ (prox), $B = 1.75$ , SE = $1.33$ (dis)

(Continued)

**Table 5.** (Continued)

Adult samples (>25 year)					
Study	Time scale recovery	Daily resilience operationalization	Recovery NA or PA (Mean, SD)	Stressors	Results / associations with daily recovery
					Latency (within-person)*: $B = -.89$ , $SE = .83$ (prox), $B = -2.58^*$ , $SE = 1.29$ (dis)
Montpetit et al., 2010	Across multiple days	Multilevel model: non-linear dampening between stress and negative affect	NA	10 items Perceived Stress Scale: (0) never to (4) very often	Trait resilience*: $B = -0.0032$ , $SE = .002$ , $t (38) = -2.09^*$ , $p = .04$
					Social support friends: $B = .013$ , $SE = .007$ , $t (38) = 1.83$ , $p = .07$
					Family Support: $B = -.011$ , $SE = .011$ , $t (38) = -1.07$ , $p = .29$ .
Ong et al., 2006 1a	Next day	lagged associations between daily stress and next day negative affect	NA	Most stressful event of the day + how stressful: (1) not very stressful to (5) very stressful	Positive affect mediator (within-person)*: stress → NA .08 vs .31* no PA
					Trait resilience*: $y = -.243^*$ , $t(925) = 3.46$ , $p = .01$
Ong et al., 2006 1b					Positive affect mediator (within-person)*: stress → NA .03 vs .27* no PA
					Trait resilience*: $y = .285^*$ , $t(984) = 3.47$ , $p < .01$
Ong et al., 2006 2					Positive affect mediator (within-person)*: stress → NA .06 vs .32* no PA
					Trait resilience*: $y = .269^*$ , $t (2388) = 4.38$ , $p < .05$ .
Scott et al., 2017	Within a day	multilevel model with age × stressor recency (how long ago the stressor was). Elevations in NA at post-stressor periods after the initial reactivity or peak response	NA	Experience of a stressful event + how long ago: yes/no, and 0-5, 5-10, 10-30, 30-60, 60 or more minutes ago	Age*: 10-30 min after stressor: $B = -.33^*$ , $SE = .11$
					Age*: 30-60 min: $B = -.32^*$ , $SE = .12$
					Age*: 60-150 min: $B = -.23^*$ , $SE = .11$
					Age: after 2.5-5 hour: age: $\beta = -0.04$ , $p = 0.44$ .
Vaessen et al., 2019	Within a day	multilevel models with negative affect as dependent variable and time since negative event (-1, t1, t2, ..., t8) as predictor. Estimate contrasts between $t_j$ for $j = 1, \dots, 8$ vs $t-1$ . No longer significant = recovery	NA: HC + CP: t1 (~90 min), EP: t4 (~270 min)	Most important event and how pleasant/ unpleasant this event was: (-3) very unpleasant to (+3) very pleasant	Group*: t1 – t3 was greater in EP compared to HV ( $B = 0.185$ ; $SE = 0.068$ ; $p = .007$ ) and CP ( $B = 0.228$ ; $SE = 0.072$ ; $p = .002$ )

Note.  $r$  = correlation, \* indicates significant correlations. Prox= proximal effects, Dis=distal effects. Unless indicated by *within-person*, between person correlations and associations are reported.

stressors. The average sample size was 289 participants ( $SD = 334$ , range = 27–1155). The mean age of the four youth samples was 16.7 ( $SD = 6.0$ ), ranging from 10.9 (Bai & Repetti, 2018) to 24.7 (Kuranova et al., 2021). The mean age of the adult samples was 52.4 years ( $SD = 16.5$ ), ranging from 32.1 to 79.0. The proportion of males included ranged from 0% to 100%, with a mean of 36% ( $SD = 21\%$ ). Different populations were included, from healthy participants to patients with mental health problems (see Table 1).

Seven studies were of fair quality ( $>60\%$  (Bai & Repetti, 2018; Bergeman & Deboeck, 2014; Bergh & Silverman, 2018; Ekas & Whitman, 2011; Hamilton et al., 2008; Lee et al., 2022; Ong et al., 2006; Scott et al., 2017). The other studies showed poor quality regarding design and reporting (40%–57%), resulting in an average rating of 54.4% ( $SD = 11.2\%$ , see Table 3). Only Leger et al. (2018) performed a power analysis and showed sufficient power to detect effects. Most studies did not include all details on the methods to be able to replicate the study (see Table 3). Nine studies (including 11 samples) were daily diary studies where participants completed one survey a day for 8–168 days ( $M = 55.6$  days,  $SD = 46.2$  days). The average compliance of these daily diaries, based on 10 samples, was 82.4% ( $SD = 14.9\%$ ). The 11 ESM studies included on average 7 surveys per day ( $SD = 2.7$ , range = 3–20) for 2–40 days ( $M = 10.9$  days,  $SD = 10.9$  days). The compliance was on average 79.4% ( $SD = 15.1\%$ ).

#### *Operationalization of affective recovery*

In this second type of studies, daily resilience was operationalized as the recovery of positive or negative affect after the experience of a stressor, i.e., affective recovery. Different operationalizations of affective recovery were used and the nature of the stressors from which individuals recovered differed (see Table 5). In the youth samples, Bai and Repetti (2018) defined affective recovery as the degree of recovery from school problems at bed time. Children whose positive or negative affect at bedtime was no longer significantly affected by the problems were considered showing resilience. In the other youth studies, recovery time was defined as the time until negative/positive affect after a stressful events on was no longer significantly different from baseline negative/positive affect. Specifically, Kuranova et al. (2021) investigated the speed of affect recovery from unpleasant events across multiple days, whereas Kuranova et al. (2020) and Lachowicz et al. (2024) estimated the recovery time after an unpleasant event or activity within a day (see Table 5).

In the adult studies, similar approaches to estimate affective recovery after the stressor were taken. In six ESM studies, the recovery of positive or negative affect after the experience of a stressor was investigated *within a day*, assessing recovery over minutes or hours. These studies either estimated the recovery speed, i.e., amount of time until the effect of stress on negative/positive affect is no longer significantly different from the baseline negative/positive affect, or the degree of recovery by investigating the remaining effect of stress on affect later in the day. Five other studies used a multilevel model was used to estimate the effect of a stressor on *next day's* negative or positive affect, reflecting recovery over one day (Ekas & Whitman, 2011; Hamilton et al., 2008; Lee et al., 2022; Leger et al., 2018; Ong et al., 2006). Finally, the remaining five studies assessed recovery across multiple days using different complex statistical models, e.g., to assess the dampening or dissipation of negative affect after experiencing stress (Bergeman & Deboeck, 2014; Congard et al., 2011; Deboeck & Bergeman, 2013; Fleuren et al., 2023; Montpetit et al., 2010).

#### *Results and associations of affective recovery in youth*

Examining how these dynamic operationalizations of daily resilience as affective recovery were related to different psychosocial variables and mental health or well-being, we again start with a discussion of findings in youth ( $< 25$  years). These associations were only investigated at the between-person level, i.e., providing insight into which child or adolescent shows more affective recovery in daily life compared to others. Bai and Repetti (2018) reported that children who recovered more from their school problems had fewer self-reported depressive symptoms and fewer internalizing problems as reported by their parents. Relatedly, adolescents who recovered more quickly from stress experienced due to a difficult activity reported less severe anxiety symptoms (Lachowicz et al., 2024). However, affective recovery time was not related to anxiety symptoms three year later, indicating the instability of resilience. Furthermore, this operationalization of affective recovery was not related to depressive symptoms. In contrast, Kuranova et al. (2020) compared the recovery time in two groups of adolescents based on the 1-year change in depressive symptoms, i.e., stable and increasing symptoms. At baseline, the group of adolescents with increasing symptom levels recovered slower from increased negative affect after a stressful event (recovery time:  $\sim 180$  min) compared to individuals with no increase in symptoms ( $\sim 90$  min, Kuranova et al., 2020). Finally, Kuranova et al. (2021) found recovery speed after an unpleasant event across multiple days to be unrelated to trait resilience ( $r = -0.16$ ), self-reported daily resilience ( $r = -0.01$ ), and psychopathological symptoms a year later ( $\beta = -0.01$ ).

Overall, based on a limited number of studies in youth and acknowledging the limitation that within-person studies were entirely lacking, daily resilience as the speed or degree of affective recovery seems to be related to the mental health of the child or adolescents, although the predictive value for later psychopathology remains inconclusive.

#### *Results and associations of affective recovery in adults*

Compared to the studies and findings in youth, more associations between daily resilience as the affective recovery after daily stressors and other measures have been reported in adults (see Table 5). Some studies investigated under which circumstances adults showed more affective recovery. At this within-person level, affective recovery within a day was positively associated with less daily negative affect and perceived stress, but not with physical activity (Almeida et al., 2020). Additionally, a small association with sleep latency was found, i.e., on days with more recovery, adults had less difficulty falling asleep (Marcusson-Clavertz et al., 2022). Similarly, affective recovery over one day was positively related to sleep quality in women with fibromyalgia. After a poor night of sleep, the yesterday's stressful event still influences today's positive affect, whereas this association disappears after a good night of sleep, indicating recovery (Hamilton et al., 2008). Finally, adults showed more negative affect recovery on days with more positive affect (Ekas & Whitman, 2011; Ong et al., 2006).

The other studies provided insight into which adults show greater affective recovery in daily life compared to others. At this between-person level, multiple studies compared affective recovery times after a stressful or unpleasant event within a day in different groups based on mental health status. Longer affect recovery times were found for early-stage psychosis patients compared to chronic psychosis patients and healthy controls (recovery time:  $\sim 270$  min vs  $\sim 90$  min, Vaessen et al., 2019), adults with depression

compared to healthy controls (~78–89 vs ~56 min, De Calheiros Velozo et al., 2023), and mental health patients and at-risk individuals compared to healthy controls (~180 vs ~90 min, Ader et al., 2022). Furthermore, older adults showed greater affective recovery 30–60 minutes after the stressor compared to younger adults, whereas no age differences in negative affect recovery during the 2.5–5 hour period were found, suggesting catching up by younger adults (Scott et al., 2017).

In addition, affective recovery over one day or multiple days was positively related to trait resilience in older adults (Bergeman & Deboeck, 2014; Montpetit et al., 2010; Ong et al., 2006). Adults showing more affective recovery reported fewer physical health problems 10 years later (Leger et al., 2018), and a higher number of daily stressors, but not higher optimism (Lee et al., 2022). Furthermore, adults with a faster affect recovery reported more positive affect, but were not different on trait anxiety (Congard et al., 2011). Faster recovery from high perceived stress was also positively related to social coping, support from family and friends, environmental mastery, self-reported health and self-esteem, and negatively to depression (Bergeman & Deboeck, 2014), and neuroticism (Deboeck & Bergeman, 2013). Finally, individuals with high burnout levels showed steeper affective recovery after a COVID-19 event compared to individuals with lower burnout levels, suggesting more resilience (Fleuren et al., 2023). However, employees with lower burnout scores were also less affected by the COVID-19 event, i.e., they needed less recovery.

Answering the research question how daily resilience is related to broader measures of well-being, the findings demonstrate that daily resilience, when operationalized as affective recovery from daily stressors, is related to various psychosocial variables in adults. The within-person results show that days on which adults show faster affective recovery, are characterized by more positive affect, better sleep and less negative affect and stress. The between-person results show that individuals with mental health problems show slower affective recovery. Faster recovery was associated with higher trait resilience, and psychosocial variables such as social coping, support from family and friends, neuroticism and self-esteem.

## Discussion

Resilience refers to the dynamic process of adaptation to or recovery from stressors and results in positive well-being and mental health outcomes (Ioannidis et al., 2020; Rutter, 2012). This adaptation or recovery does take place after major life events, as is mostly researched in the field of resilience. In addition, recovery also takes place in the flow of everyday life, after experiencing the little hassles and small stressors that every person encounters. Despite increasing technological opportunities to measure the daily lives of individuals with Experience Sampling methods, little is known about the underlying dynamic nature of daily resilience to these daily stressors, and how this is linked to various outcomes, related to well-being and mental health. We aimed to advance our understanding of daily resilience through systematically reviewing studies measuring daily resilience in real-time, using Experience Sampling and daily diary methods.

In this preregistered systematic review, we summarized the available literature on daily resilience across the lifespan. We included 36 studies of which 11 investigated daily resilience in youth, i.e., children, adolescents, or young adults, and 25 in adult samples. Daily resilience was operationalized in two ways: self-reported ability to recover from daily stressors and, indirectly, as

the recovery of positive or negative affect after experiencing daily stressors, i.e., affective recovery. The results of the included studies indicate that an individual's ability to recover from daily stressors depends on the context and their available mental resources. At the same time, there was consensus that daily resilience, in youth and adults, was related to mental health and several psychosocial variables, including positive affect, optimism, personality, support from friends/family, and coping.

### Operationalization of daily resilience

The review of the literature highlighted that there are broadly two operationalizations of daily resilience: direct self-report items of perceived ability to recover from daily stressors, as well as a more indirect measure of the degree or speed of positive or negative affect recovery after experiencing daily stressors, i.e., how quickly is positive or negative affect back to baseline after the experience of stress.

The studies examining self-reported ability to recover from daily stressors mostly used items adapted from existing trait resilience questionnaires. The low to moderate ICCs (range = .39–.80) indicate that a substantial part of the variance in daily self-reported ability to recover from daily stressors was due to within-person fluctuations, justifying the use of ESM to assess self-reported resilience multiple times per week or day. However, self-reported daily resilience measures have limitations. When translating trait questionnaires to ESM items, only adding an indication of momentary status (i.e., "Right now" or "Today") does not necessarily translate the item to momentary states (Myin-Germeys et al., 2018). For example, the item "*I tend to bounce back quickly after hard times*" from the Brief Resilience Scale can be interpreted to reflect a more general self-appraised resilience or coping process, which is not expected to vary a lot over the day or week.

Furthermore, an important aspect for ESM self-report items is assessing behavior or processes that people are not necessarily aware of. The dynamic process of daily resilience may occur without individuals, and particularly children and adolescents, being fully aware. Questions like "*Today, I could rely on myself to overcome challenges*" are often too reflective or abstract, whereas questions about the present state, e.g. "*Right now, I feel like I can deal with whatever comes*" are more direct inquiries and easier for participants to reflect upon. Although easier to answer, the self-report items for daily resilience capture the subjective experience, but cannot adequately capture the dynamic concept of resilience. Items regarding handling or dealing with stress (e.g., *Today, I felt I could handle many things at a time* (Martinez-Corts et al., 2015) reflect concepts related to resilience, including coping and emotion regulation (Trudel-Fitzgerald et al., 2023). Such reflections on coping are not related to daily resilience as a process of recovery from stress. Furthermore, statements such as "*My daily life is full of things that keep me interested*" (Harpoth et al., 2020) or "*I am cheerful*" (Schwerdtfeger & Dick, 2019) lean more towards reflecting positive emotions rather than specifically assessing the capacity to recover from stressors in everyday life. Therefore, although self-reported measures of perceived ability to recover from daily stressors can add valuable information about the subjective coping with daily stressors and resilience factors, more research into the content and psychometric qualities of ESM items of such daily measures are needed.

The other approach of measuring daily resilience as the degree or speed of affective recovery after experiencing daily stressors can capture the process of daily resilience. Requiring data from

multiple time points, these measures are computed by combining responses to ESM items on the experience of stressors and positive or negative affect multiple times per day or week. The exact definition and operationalization of daily resilience as recovery from daily stressors varied. Some studies operationalized daily resilience as the absence of negative affect the day after experiencing daily stressors, whereas other studies used survival analyses to determine the time of returning to baseline affect after stress. The first operationalization uses a crude estimation of recovery time (i.e., recovery over one night), whereas the survival analyses studies can be more precise, down to hours or minutes. Additionally, stressors were defined in different ways, including experienced stressfulness of a negative event or perceived momentary stress. For example, some studies focused on stressors with specific contexts, such as school-related challenges, while others used broader measures of overall feelings of stress without tying them to a specific event or context. This heterogeneity in stressors limits the opportunity to investigate how daily resilience differs depending on the type or source of stress. Overall, these differences in operationalizations of affective recovery should be considered when interpreting and comparing the results of the studies.

Moreover, only four studies assessed daily resilience as affective recovery in children or adolescents. From a developmental perspective, daily resilience early in life might be especially important, because of the impact of mental health problems during childhood and adolescence on later development and risk of psychopathology in adulthood (Clayborne et al., 2019; Kim-Cohen et al., 2003; Patel et al., 2007; Thapar et al., 2012). Therefore, although the approach of measuring the speed of affective recovery as a proxy for daily resilience is promising, more research is needed. Future research should investigate the within-person and contextual factors, and between-person factors that influence the ability of youth to show daily resilience. Furthermore, the optimal method for measuring daily resilience in children and adolescents needs to be determined, taking into account the burden on young people.

Whether these two broader categories of operationalizations of daily resilience tap into similar phenomena is an open question. Only one study among 68 adolescents included a direct investigation between daily self-reported ability to recover from daily stressors and speed of affect recovery across days. Interestingly, the two were not related ( $r = -.01$ , Kuranova et al., 2021). Although this finding should be replicated, this indicates that approaches to measure daily resilience capture different aspects related to daily resilience. Self-reported ability to recover from daily stressors reflects subjective experiences of coping with stress or resilience-related factors, whereas daily resilience as the speed of affective recovery after daily stressors captures the dynamic process of recovery over time.

### Associations of daily resilience

Well-known from research to resilience after major life events or adversity, different *resilience factors* help individuals to be able to recover from stressors, e.g., optimism or social support from friends of family (van Harmelen et al., 2017). The second research question in this review was to test whether these and other factors are also associated with daily resilience. Across the included studies, associations with several psychosocial factors and well-being and mental health outcomes were included. Although self-reported ability to recover from daily stressors does not adequately

capture the process of recovery after stress, the associations between daily self-reported ability to recover from daily stressors and other psychosocial measures could give insights and starting points for future research on factors influencing daily resilience. It is important to note, however, that the reliance on self-report for both daily resilience and related psychosocial or affective measures introduces a potential common method bias (Podsakoff et al., 2003, 2024). For instance, the reporter's affective or mental health status may alter perceptions, and therefore drive the associations between daily resilience and other measures.

The findings for youth and adults for self-reported ability to recover from daily stressors were similar. At a within-person level, the studies showed that higher self-appraised daily resilience was reported when individuals also reported more positive affect and/or less negative affect. Similarly, positive within-person relations with daily hope, optimism, and meaning in life were found, and negative relations with daily somatic symptoms, depletion, and heart rate variability. At the between-person level, individuals who reported higher ability to recover from daily stressors also reported less depressive symptoms, less loneliness, more trait resilience, or scored higher on conscientiousness and agreeableness compared to others.

Focusing on daily resilience as the affective recovery after daily stressors, only a few studies in adults, and none in youth, investigated differences in daily resilience at the within-person level. Faster or more affective recovery was found in moments or days with less negative affect (Almeida et al., 2020) and after nights with above-average sleep (Hamilton et al., 2008). The remaining studies in youth and adults investigated who is more likely to show faster affective recovery after daily stressors. In youth, faster affective recovery was related to better mental health, i.e., fewer depressive or anxiety symptoms. Similarly, in adults, slower affective recovery on a day to day basis was consistently reported in individuals at risk for mental health problems or disorders compared with healthy controls.

These results suggest that daily resilience as affective recovery is influenced by factors related to the availability of mental or physical resources for recovery. For example, the findings of Hamilton et al. (2008) suggest that sleep good improves affective recovery. Similarly, faster recovery has been associated with more social support from family and friends and higher self-esteem (Bergeman & Deboeck, 2014). These factors can be seen as resources that help individuals recover from daily stressors. Unfortunately, the lacking within-person studies in youth samples limit the ability to determine under which circumstances youth recover faster.

Combined with the potential predictive value of daily resilience for psychopathology symptoms in the following year (Kuranova et al., 2020), these findings suggest that speed of recovery can be used as a signal of lower mental health stability or deterioration. Assessing daily resilience has thus the potential to identify individuals at risk of mental health problems and allow for timely intervention. However, replication and further research is needed to support these preliminary findings, particularly in children and adolescents.

### Implications and future directions

Our review shows that ESM and diary studies can provide detailed information about fluctuations in daily resilience and insights into the momentary and daily influences on daily resilience, from day to day or even hour to hour. However, both self-report studies of daily

resilience and studies of affective recovery included heterogeneous items/conceptualisations, limiting the comparability of results and overall conclusions. Future research to the optimal method to measure daily resilience is needed, as well as research to the factors influencing daily resilience.

Research on daily resilience in real time using ESM is relatively new. The majority of the included studies were from the last five years, i.e., 67% were published in 2019–2024. The samples were not diverse with respect to age, ethnicity, and gender. All studies were performed in Western countries in Europe and the US, limiting the cultural diversity of the samples and the generalizability of the findings on daily resilience to other cultures. Furthermore, most of the studies included more women than men. As is known from research to resilience after major life events or adversity, resilience may differ in men compared to women (Hodes & Epperson, 2019; Kalisch et al., 2017; van Harmelen et al., 2017), indicating the need for comparisons. Similarly, resilience, but also mental health and well-being, differs across the lifespan (Infurna, 2021; Ioannidis et al., 2020; Masten & Wright, 2009; Orben et al., 2022; Southwick et al., 2011; Werner & Smith, 2001; Werner, 1995; de Vries et al., 2024; van Harmelen et al., 2017). From a developmental perspective, both resilience to major life events and smaller daily stressors is particularly important in youth because of increased stress and the impact of mental health problems during influences on further development and risks of psychopathology later in life (Clayborne et al., 2019). Although we compared the results of studies to daily resilience in youth and adults, no study directly compared the findings in different age groups across the life span. Therefore, as the field grows and more studies on daily resilience are designed, more diverse samples need to be included and results compared in different subgroups.

Measures of the degree or speed of affective recovery can capture the resilience process in daily life. However, intensive data from individuals is required to compute this degree or speed of recovery. Therefore, promising correlates of affective recovery that are easier to measure should also be investigated as well. In a recent review, Ong and Leger (2022) discussed several of such daily resilience correlates, including dampened reactivity to stress (i.e., immediate affective response at the time or closely after the occurrence of a stressor), toughness/inoculation (i.e., influence of stressor exposure on responses to subsequent stressors), and richness of stressors (i.e., variety of stressors in daily life). Although not capturing the dynamic process of recovery, these daily resilience correlates have been linked to mental health and well-being as well, although research is limited, particularly in adolescents. To fully understand the dynamics of daily resilience and the potential of these measures for early detection of deterioration in mental health, future research is needed that combines these different measures and correlates of daily resilience.

Furthermore, in addition to self-reported data on affective recovery from daily stressors, there is a need to assess recovery from daily stressors integrating the affective, cognitive, physiological, and (objective) behavioral measurements, to capture the full recovery from stress in daily life. For example, measures such as heart rate variability, cortisol reactivity, or sleep patterns can capture other aspects of daily resilience (O'Donohue et al., 2019; Weber et al., 2022). This review focuses on the affective recovery from daily stressors. However, reviews and research on the other aspects, as well as the integration of the different aspects of recovery from daily stressors are the focus of the Stress-in-Action consortium (<https://stress-in-action.nl/>). We aim to provide a more complete understanding of the daily processes around stress.

In addition, the relationship between daily resilience and resilience as a long-term outcome after major life events or chronic stress remains an important question for future research. Results from this review suggest that daily resilience is associated with fewer mental health issues and potentially also resilience to major stressors. Daily resilience could thus be a mechanism for building and maintaining long-term resilience over time. However, more research is needed to establish the direction of effects of these processes. With new opportunities to collect data in daily life, we could investigate in more detail the mechanisms and factors underlying daily resilience and resilience to major events or chronic stress. For example, studies are needed that combine ESM bursts with long-term follow-ups. This could lead to a more nuanced understanding of how resilience functions across time scales and contexts.

Ultimately, the insights from ESM studies to the influences on daily resilience and what enables individuals to recover quickly from stress can help to develop interventions to help individuals show daily resilience to stressors on a day to day basis. For example, interventions that target resilience in the moment, i.e. Ecological Momentary Interventions (EMI) or Just-in-Time Adaptive Interventions, can be developed. These intervention types are designed to deliver personalized interventions based on what the participant needs in a certain moment and context. For example, Reininghaus et al.'s (2023) intervention is an EMI where help-seeking youth learned new compassion-focused exercises and were prompted to complete these exercises when they scored high on stress or negative affect. Initial evaluations showed small increases in momentary self-reported daily resilience after this intervention. Future research should also evaluate the effects of interventions on the process of daily resilience, i.e., affective recovery. With a focus on the young, but taking a life course perspective, interventions aimed at supporting adolescents in building daily resilience to stress reflect a proactive approach to preventing mental health problems before they emerge or worsen. Such interventions are needed in today's changing world, characterized by increasing levels of experienced stress and rising numbers of young people with mental health problems.

## Conclusion

In our fast-paced world, stress has become an inevitable part of our daily lives. Insights into the dynamic process of daily resilience can help to earlier detect individuals at risk of mental health problems after stress and inform targeted interventions and societal initiatives to support individuals through strengthening their resilience to smaller daily stressors and potentially resilience to major events or adversity as well. In this systematic review, we showed that daily resilience can be effectively measured in daily life using daily diaries and ESM studies. Self-reported measures of perceived ability to recover from daily stressors can add valuable information of the subjective coping with stress and related resilience factors. Daily resilience as affective recovery from stress can capture the dynamic process of daily resilience and can be effectively measured in everyday life with novel data-collection techniques such as smartphone-based Experience Sampling. Daily resilience was related to better mental health in both youth and adults. In adult samples, moreover, the ability to recover in daily life from stressors was influenced by factors related to the availability of resources in that moment or day, such as a good night of sleep, more social support, or feelings of optimism. Even though adolescents in particular are facing high levels of pressure

and stress, no studies, thus far, have examined how youth manage to recover in daily life. Given that daily stressors may pile up to mental health problems there is an urgent need for similar studies in adolescents to assess in which circumstances, and with which resources, they can show daily resilience.

**Supplementary material.** The supplementary material for this article can be found at <http://doi.org/10.1017/S0954579425000197>.

**Acknowledgements.** The authors wish to thank Christa Niehot from the Erasmus MC Medical Library for developing and updating the search strategies.

**Author contribution.** LdV conceptualized the study. JZ and LdV performed the literature search and extracted the data. JZ and LdV wrote the first draft of the manuscript. All authors reviewed and edited the manuscript.

**Funding statement.** This research is funded by a grant from the Stichting Erasmus Trustfonds awarded to L.P. de Vries [97,030.2023.102.114/189/RB] and is part of the PROTECT ME project funded by Convergence, the alliance between Erasmus Medical Centre Rotterdam, Erasmus University Rotterdam and Delft University of Technology. In addition, the work is part of Stress in Action, which is financially supported by the Dutch Research Council and the Dutch Ministry of Education, Culture and Science (NWO gravitation grant number 024.005.010). A-L van Harmelen is funded by the Social Resilience and Security programme at Leiden University and a VIDI award from the Netherlands Council for Scientific research. L. Keijsers is funded by the European Union (ERC, PARADOx, 101043536).

**Competing interests.** ALVH receives consultancy fees from the Augeo foundation.

## References

Ader, L., Schick, A., Simons, C., Delespaul, P., Myin-Germeys, I., Vaessen, T., & Reininghaus, U. (2022). Positive affective recovery in daily life as a momentary mechanism across subclinical and clinical stages of mental disorder: Experience sampling study. *JMIR Mental Health*, 9(11), e37394. <https://doi.org/10.2196/37394>

Almeida, D. M. (2005). Resilience and vulnerability to daily stressors assessed via diary methods. *Current Directions in Psychological Science*, 14(2), 64–68. <https://doi.org/10.1111/j.0963-7214.2005.00336.x>

Almeida, D. M., Marcusson-Clavertz, D., Conroy, D. E., Kim, J., Zawadzki, M. J., Sliwinski, M. J., & Smyth, J. M. (2020). Everyday stress components and physical activity: Examining reactivity, recovery and pileup. *Journal of Behavioral Medicine*, 43(1), 108–120. <https://doi.org/10.1007/s10865-019-00662-z>

Anderson, K., & Priebe, S. (2021). Concepts of resilience in adolescent mental health research. *Journal of Adolescent Health*, 69(5), 689–695. <https://doi.org/10.1016/j.jadohealth.2021.03.035>

Bai, S., & Repetti, R. L. (2018). Negative and positive emotion responses to daily school problems: Links to internalizing and externalizing symptoms. *Journal of Abnormal Child Psychology*, 46(3), 423–435. <https://doi.org/10.1007/s10802-017-0311-8>

Bergeman, C. S., & Deboeck, P. R. (2014). Trait stress resistance and dynamic stress dissipation on health and well-being: The reservoir model. *Research in Human Development*, 11(2), 108–125. <https://doi.org/10.1080/15427609.2014.906736>

Bergh, O., & Silverman, M. J. (2018). Undergraduate student research awards effects of music therapy in the form of patient-preferred live music on mood and pain with patients on a solid organ transplant unit: A randomized pilot study. *Music Therapy Perspectives*, 36(1), 129–130. <https://doi.org/10.1093/mtp/mix027>

Block, J., & Kremen, A. M. (1996). IQ and ego-resiliency: conceptual and empirical connections and separateness. *Journal of Personality and Social Psychology*, 70(2), 349. <https://doi.org/10.1037/0022-3514.70.2.349>

Bonanno, G. A. (2012). Uses and abuses of the resilience construct: Loss, trauma, and health-related adversities. *Social Science and Medicine*, 74(5), 753–756. <https://doi.org/10.1016/j.socscimed.2011.11.022>

Brogly, C., Lizotte, D. J., Mitchell, M., Speechley, M., MacDougall, A., Huner, E., Anderson, K. K., & Bauer, M. A. (2024). An app-based ecological momentary assessment of undergraduate mental health during the COVID-19 pandemic in Canada (Smart Healthy Campus Version 2.0). Longitudinal study, *PLOS Digital Health*, 3(5), e0000239. <https://doi.org/10.1371/journal.pdig.0000239>

Cahill, S., Hager, R., & Shryane, N. (2023). Patterns of resilient functioning in early life: Identifying distinct groups and associated factors. *Development and Psychopathology*, 36(4), 1–21. <https://doi.org/10.1017/S0954579423001165>

Charles, S. T., Piazza, J. R., Mogle, J., Sliwinski, M. J., & Almeida, D. M. (2013). The wear and tear of daily stressors on mental health. *Psychological Science*, 24(5), 733–741. <https://doi.org/10.1177/0956797612462222>

Clayborne, Z. M., Varin, M., & Colman, I. (2019). Systematic review and meta-analysis: Adolescent depression and long-term psychosocial outcomes. *Journal of the American Academy of Child & Adolescent Psychiatry*, 58(1), 72–79. <https://doi.org/10.1016/j.jaac.2018.07.896>

Congard, A., Dauvier, B., Antoine, P., & Gilles, P. Y. (2011). Integrating personality, daily life events and emotion: Role of anxiety and positive affect in emotion regulation dynamics. *Journal of Research in Personality*, 45(4), 372–384. <https://doi.org/10.1016/J.JRSP.2011.04.004>

Connor, K. M., & Davidson, J. R. (2003). Development of a new resilience scale: The Connor-Davidson resilience scale (CD-RISC). *Depression and Anxiety*, 18(2), 76–82. <https://doi.org/10.1002/da.10113>

Csikszentmihalyi, M., & Larson, R. (1987). Validity and reliability of the experience-sampling method. *Journal of Nervous & Mental Disease*, 175(9), 526–537. [https://doi.org/10.1007/978-94-017-9088-3\\_3](https://doi.org/10.1007/978-94-017-9088-3_3)

Daly, M. (2022). Prevalence of depression among adolescents in the U.S. From 2009 to 2019: Analysis of trends by sex, race/Ethnicity, and income. *Journal of Adolescent Health*, 70(3), 496–499. <https://doi.org/10.1016/J.JADOHEALTH.2021.08.026>

De Calheiros Velozo, J., Lafit, G., Viechtbauer, W., van Amelsvoort, T., Schruers, K., Marcelis, M., Goossens, L., Simons, C. J. P., Delespaul, P., Claes, S., Myin-Germeys, I., & Vaessen, T. (2023). Delayed affective recovery to daily-life stressors signals a risk for depression. *Journal of Affective Disorders*, 320, 499–506. <https://doi.org/10.1016/J.JAD.2022.09.136>

de Vries, L. P., Baselmans, B. M. L., & Bartels, M. (2021). Smartphone-based ecological momentary assessment of well-being: A systematic review and recommendations for future studies. *Journal of Happiness Studies*, 22(5), 2361–2408. <https://doi.org/10.1007/S10902-020-00324-7>

de Vries, L. P., Pelt, D. H. M., & Bartels, M. (2024). *The stability and change of well-being across the lifespan: A longitudinal twin-sibling study* (pp. 1–13). Psychological Medicine, <https://doi.org/10.1017/S0033291724000692>

Deboeck, P. R., & Bergeman, C. S. (2013). The reservoir model: A differential equation model of psychological regulation. *Psychological Methods*, 18(2), 237–256. <https://doi.org/10.1037/A0031603>

Delongis, A., Coyne, J. C., Dakof, G., Folkman, S., & Lazarus, R. S. (1982). Relationship of daily hassles, uplifts, and major life events to health status. *Health Psychology*, 1(2), 119–136. <https://doi.org/psycnet.apa.org/doi/10.1037/0278-6133.1.2.119>

Dong, F., Ablah, E., Nelson, C., Shah, S., & Khan, A. (2013). Validation of a modified CD-RISC: Including previously unaccounted for resilience variables. *Kansas Journal of Medicine*, 6(1), 11–20.

Eckenrode, J. (1984). Impact of chronic and acute stressors on daily reports of mood. *Journal of Personality and Social Psychology*, 46(4), 907–918. <https://doi.org/10.1037/0022-3514.46.4.907>

Ekas, N. V., & Whitman, T. L. (2011). Adaptation to daily stress among mothers of children with an Autism Spectrum Disorder: The role of daily positive affect. *Journal of Autism and Developmental Disorders*, 41(9), 1202–1213. <https://doi.org/10.1007/s10803-010-1142-4>

Fleuren, B. P. I., Nübold, A., Uitdewilligen, S., Verduyn, P., & Huisheger, U. R. (2023). Troubles on troubled minds: An intensive longitudinal diary study on the role of burnout in the resilience process following acute stressor exposure. *European Journal of Work and Organizational Psychology*, 32(3), 373–388. <https://doi.org/10.1080/1359432X.2022.2161369>

Fritz, J., de Graaff, A. M., Caisley, H., van Harmelen, A.-L., & Wilkinson, P. O. (2018). A systematic review of amenable resilience factors that moderate and/or mediate the relationship between childhood adversity and mental health in

young people. *Frontiers in Psychiatry*, 9, 341825. <https://doi.org/10.3389/fpsyg.2018.00230>

Hamaker, E. L., & Wichers, M. (2017). No time like the present: Discovering the hidden dynamics in intensive longitudinal data. *Current Directions in Psychological Science*, 26(1), 10–15. <https://doi.org/10.1177/0963721416666518>

Hamilton, N. A., Affleck, G., Tennen, H., Karlson, C., Luxton, D., Preacher, K. J., & Templin, J. L. (2008). Fibromyalgia: The role of sleep in affect and in negative event reactivity and recovery. *Health Psychology*, 27(4), 490–497. <https://doi.org/10.1037/0278-6133.27.4.490>

Harpoth, T. S. D., Kongerslev, M. T., Trull, T. J., Hepp, J., Bateman, A. W., & Simonsen, E. (2020). Associations of positive and negative emotions with ego-resiliency and quality of life in borderline personality disorder: A daily diary study. *Personality Disorders: Theory, Research, and Treatment*, 11(1), 13–23. <https://doi.org/10.1037/PER0000350>

Harpoth, T. S. D., Yeung, E. W., Trull, T. J., Simonsen, E., & Kongerslev, M. T. (2021). Ego-resiliency in borderline personality disorder and the mediating role of positive and negative affect on its associations with symptom severity and quality of life in daily life. *Clinical Psychology & Psychotherapy*, 28(4), 939–949. <https://doi.org/10.1002/CPP.2548>

Heininga, V. E., & Oldehinkel, A. J. (2024). Predictors, daily correlates, and effects on future functioning of resilience in patients with Major Depressive Disorder (MDD). *Journal of Affective Disorders Reports*, 16, 100703. <https://doi.org/10.1016/J.JADR.2023.100703>

Hill, Y., Meijer, R. R., Van Yperen, N. W., Michelakis, G., Barisch, S., & Den Hartigh, R. J. R. (2021). Nonergodicity in protective factors of resilience in athletes. *Sport, Exercise, and Performance Psychology*, 10(2), 217–223. <https://doi.org/10.1037/SPY0000246>

Hodes, G. E., & Epperson, C. N. (2019). Sex differences in vulnerability and resilience to stress across the life span. *Biological Psychiatry*, 86(6), 421–432. <https://doi.org/10.1016/J.BIOPSYCH.2019.04.028>

Hoorelbeke, K., Van den Berg, N., Wichers, M., & Koster, E. H. W. (2019). Between vulnerability and resilience: A network analysis of fluctuations in cognitive risk and protective factors following remission from depression. *Behaviour Research and Therapy*, 116, 1–9. <https://doi.org/10.1016/J.BRAT.2019.01.007>

Infurna, F. J. (2021). Utilizing principles of life-span developmental psychology to study the complexities of resilience across the adult life span. *The Gerontologist*, 61(6), 807–818. <https://doi.org/10.1093/GERONT/GNAB086>

Ioannidis, K., Askelund, A. D., Kievit, R. A., & Van Harmelen, A. L. (2020). The complex neurobiology of resilient functioning after childhood maltreatment. *BMC Medicine*, 18(1), 1–16. <https://doi.org/10.1186/s12916-020-1490-7>

Jennings, R. E., Lanaj, K., & Kim, Y. J. (2023). Self-compassion at work: A self-regulation perspective on its beneficial effects for work performance and wellbeing. *Personnel Psychology*, 76(1), 279–309. <https://doi.org/10.1111/PEPS.12504>

Kalisch, R., Baker, D. G., Basten, U., Boks, M. P., Bonanno, G. A., Brummelman, E., Chmitorz, A., Fernández, G., Fiebach, C. J., Galatzer-Levy, I., Geuze, E., Groppe, S., Helmreich, I., Hendl, T., Hermans, E. J., Jovanovic, T., Kubiak, T., Lieb, K., Lutz, B. . . Kleim, B. (2017). The resilience framework as a strategy to combat stress-related disorders. *Nature Human Behaviour*, 1(11), 784–790. <https://doi.org/10.1038/s41562-017-0200-8>

Kalisch, R., Cramer, A. O. J., Binder, H., Fritz, J., Leertouwer, I., Lunansky, G., Meyer, B., Timmer, J., Veer, I. M., & Van Harmelen, A.-L. (2019). Deconstructing and reconstructing resilience: A dynamic network approach. *Perspectives on Psychological Science*, 14(5), 765–777. <https://doi.org/10.5281/zenodo.3228687>

Kanner, A. D., Coyne, J. C., Schaefer, C., & Lazarus, R. S. (1981). Comparison of two modes of stress measurement: Daily hassles and uplifts versus major life events. *Journal of Behavioral Medicine*, 4(1), 1–39. <https://doi.org/10.1007/BF00844845>

Kendler, K. S., Karkowski, L. M., & Prescott, C. A. (1999). Causal relationship between stressful life events and the onset of major depression. *American Journal of Psychiatry*, 156(6), 837–841. <https://doi.org/10.1176/ajp.156.6.837>

Kessler, R. C. (1997). The effects of stressful life events on depression. *Annual Review of Psychology*, 48(1), 191–214. <https://doi.org/10.1146/annurev.psych.48.1.191>

Kim-Cohen, J., Caspi, A., Moffitt, T. E., Harrington, H. L., Milne, B. J., & Poulton, R. (2003). Prior Juvenile diagnoses in adults with mental disorder: Developmental follow-back of a prospective-longitudinal cohort. *Archives of General Psychiatry*, 60(7), 709–717. <https://doi.org/10.1001/ARCHPSYC.60.7.709>

King, L., Jolicoeur-Martineau, A., Laplante, D. P., Szekely, E., Levitan, R., & Wazana, A. (2021). Measuring resilience in children: A review of recent literature and recommendations for future research. *Current Opinion in Psychiatry*, 34(1), 10–21. <https://doi.org/10.1097/YCO.0000000000000663>

Kuranova, A., Booij, S. H., Menne-Lothmann, C., Decoster, J., van Winkel, R., Delespaul, P., De Hert, M., Derom, C., Thiery, E., Rutten, B. P. F., Jacobs, N., van Os, J., Wigman, J. T. W., & Wichers, M. (2020). Measuring resilience prospectively as the speed of affect recovery in daily life: A complex systems perspective on mental health. *BMC Medicine*, 18(1), 36. <https://doi.org/10.1186/s12916-020-1500-9>

Kuranova, A., Booij, S. H., Oldehinkel, A. J., Wichers, M., Jeronimus, B., & Wigman, J. T. W. (2021). Reflections on psychological resilience: A comparison of three conceptually different operationalizations in predicting mental health. *European Journal of Psychotraumatology*, 12(1), 1956802. <https://doi.org/10.1080/20008198.2021.1956802>

Lachowicz, A., Houben, M., Vaessen, T., Lafit, G., Achterhof, R., Akcaoglu, Z., Bamps, E., Hagemann, N., Hiekkaranta, A. P., Hermans, K., Janssens, J., Lecei, A., Kirtley, O., & Myin-Germeys, I. (2025). The link between delayed affective recovery from daily stressors and anxiety symptoms in youth. *International Journal of Stress Management*. Advance online publication. <https://dx.doi.org/10.1037/str0000347>

Lass-Hennemann, J., Sopp, M. R., Ruf, N., Equit, M., Schäfer, S. K., Wirth, B. E., & Michael, T. (2023). Generation climate crisis, COVID-19, and Russia-Ukraine-War: Global crises and mental health in adolescents. *European Child and Adolescent Psychiatry*, 1, 1–14. <https://doi.org/10.1007/S00787-023-02300-X/TABLES/4>

Lee, L. O., Grodstein, F., Trudel-Fitzgerald, C., James, P., Okuzono, S. S., Koga, H. K., Schwartz, J., Spiro, A., Mroczek, D. K., & Kubzansky, L. D. (2022). Optimism, daily stressors, and emotional well-being over two decades in a cohort of aging men. *The Journals of Gerontology: Series B*, 77(8), 1373–1383. <https://doi.org/10.1093/GERONB/GBAC025>

Leger, K. A., Charles, S. T., & Almeida, D. M. (2018). Let it go: Lingering negative affect in response to daily stressors is associated with physical health years later. *Psychological Science*, 29(8), 1283–1290. <https://doi.org/10.1177/0956797618763097>

Lohner, M. S., & Aprea, C. (2021). The resilience journal: Exploring the potential of journal interventions to promote resilience in university students. *Frontiers in Psychology*, 12, 702683. <https://doi.org/10.3389/fpsyg.2021.702683>

LoSavio, S. T., Cohen, L. H., Laurenceau, J. P., Dasch, K. B., Parrish, B. P., & Park, C. L. (2011). Reports of stress-related growth from daily negative events. *Journal of Social and Clinical Psychology*, 30(7), 760–785. <https://doi.org/10.1521/JSCP.2011.30.7.760>

Luthans, F., Youssef, C. M., & Avolio, B. J. (2007). Psychological capital: Investing and developing positive organizational behavior. *Positive Organizational Behavior*, 1(2), 9–24. <https://doi.org/10.4135/9781446212752.n2>

Luthans, F., Avolio, B. J., Avey, J. B., & Norman, S. N. (2007). Positive psychological capital: Measurement and relationship with performance and satisfaction. *Personnel Psychology*, 60(3), 2541–2572. <https://doi.org/10.1111/j.1744-6570.2007.00083.x>

Marcusson-Clavertz, D., Sliwinski, M. J., Buxton, O. M., Kim, J., Almeida, D. M., & Smyth, J. M. (2022). Relationships between daily stress responses in everyday life and nightly sleep. *Journal of Behavioral Medicine*, 45(4), 518–532. <https://doi.org/10.1007/s10865-021-00281-3>

Martinez-Corts, I., Demerouti, E., Bakker, A. B., & Boz, M. (2015). Spillover of interpersonal conflicts from work into nonwork: A daily diary study. *Journal of Occupational Health Psychology*, 20(3), 326–337. <https://doi.org/10.1037/A0038661>

Masten, A. S. (2015). Pathways to integrated resilience science. *Psychological Inquiry*, 26(2), 187–196. <https://doi.org/10.1080/1047840X.2015.1012041>

Masten, A. S., Best, K. M., & Garmezy, N. (1990). Resilience and development: Contributions from the study of children who overcome adversity.

*Development and Psychopathology*, 2(4), 425–444. <https://doi.org/10.1017/S0954579400005812>

**Masten, A. S., Lucke, C. M., Nelson, K. M., & Stallworthy, I. C.** (2021). Resilience in development and psychopathology: Multisystem perspectives. *Annual Review of Clinical Psychology*, 17(1), 521–549. <https://doi.org/10.1146/ANNUREV-CLINPSY-081219-120307>

**Masten, A. S., & Wright, M. O.** (2009). Resilience over the lifespan. In J. W. Reich, A. J. Zautra, & J. S. Hall (Eds.), *Handbook of Adult Resilience* (pp. 213–237). The Guilford Press, <https://psycnet.apa.org/record/2010-10101-011>

**McHugh, M. L.** (2012). Interrater reliability: The kappa statistic. *Biochimia Medica*, 22(3), 276–282.

**McLaughlin, K. A., Colich, N. L., Rodman, A. M., & Weissman, D. G.** (2020). Mechanisms linking childhood trauma exposure and psychopathology: A transdiagnostic model of risk and resilience. *BMC Medicine*, 18(1), 1–11. <https://doi.org/10.1186/S12916-020-01561-6> 2020.

**Mertens, M., Kraiss, J., & Vaessen, T.** (2023). A higher purpose: How individuals with different belief systems exhibit daily-life resilience [Bachelor thesis]. University of Twente. <http://essay.utwente.nl/95375/>

**Mesman, E., Vreeker, A., & Hillegers, M.** (2021). Resilience and mental health in children and adolescents: An update of the recent literature and future directions. *Current Opinion in Psychiatry*, 34(6), 586–592. <https://journals.lww.com/10.1097/YCO.0000000000000741>

**Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G.** (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Journal of Clinical Epidemiology*, 151(4), 264–269. <https://doi.org/10.1016/j.jclinepi.2009.06.005>

**Monroe, S. M.** (1983). Major and minor life events as predictors of psychological distress: Further issues and findings. *Journal of Behavioral Medicine*, 6(2), 189–205. <https://doi.org/10.1007/BF00845380>

**Montpetit, M. A., Bergeman, C. S., Deboeck, P. R., Tiberio, S. S., & Boker, S. M.** (2010). Resilience-as-process: Negative affect, stress, and coupled dynamical systems. *Psychology and Aging*, 25(3), 631–640. <https://doi.org/10.1037/A0019268>

**Myin-Germeys, I., Kasanova, Z., Vaessen, T., Vachon, H., Kirtley, O., Viechtbauer, W., & Reininghaus, U.** (2018). Experience sampling methodology in mental health research: New insights and technical developments. *World Psychiatry*, 17(2), 123–132. <https://doi.org/10.1002/WPS.20513>

**Myin-Germeys, I., & Kuppens, P.** (2022). The open handbook of experience sampling methodology: A step-by-step guide to designing, conducting, and analyzing ESM studies. In I. Myin-Germeys & P. Kuppens (Eds.), (2nd ed.), Center for Research on Experience Sampling and Ambulatory Methods Leuven.

**O'Donohue, J. S., Mesagno, C., & O'Brien, B.** (2019). How can stress resilience be monitored? A systematic review of measurement in humans. *Current Psychology*, 40(6), 2853–2876. <https://doi.org/10.1007/S12144-019-00226-9> 2019.

**Ong, A. D., Bergeman, C. S., Bisconti, T. L., & Wallace, K. A.** (2006). Psychological resilience, positive emotions, and successful adaptation to stress in later life. *Journal of Personality and Social Psychology*, 91(4), 730–749. <https://doi.org/10.1037/0022-3514.91.4.730>

**Ong, A. D., & Leger, K. A.** (2022). Advancing the study of resilience to daily stressors. *Perspectives on Psychological Science*, 17(6), 1591–1603. <https://doi.org/10.1177/17456916211071092>

**Orben, A., Lucas, R. E., Fuhrmann, D., & Kievit, R. A.** (2022). Trajectories of adolescent life satisfaction. *Royal Society Open Science*, 9(8), 211808. <https://doi.org/10.1098/RSOS.211808>

**Parrish, B. P., Cohen, L. H., & Laijrenceaij, J. P.** (2011). Prospective relationship between negative affective reactivity to daily stress and depressive symptoms. *Journal of Social and Clinical Psychology*, 30(3), 270–296. <https://doi.org/10.1521/JSCP.2011.30.3.270>

**Patel, V., Flisher, A. J., Hetrick, S., & McGorry, P.** (2007). Mental health of young people: A global public-health challenge. *The Lancet*, 369(9569), 1302–1313. [https://doi.org/10.1016/S0140-6736\(07\)60368-7](https://doi.org/10.1016/S0140-6736(07)60368-7)

**Pluess, M.** (2024). Beyond developmental psychopathology: Positive child development. *Development and Psychopathology*, 36(5), 1–9. <https://doi.org/10.1017/S0954579424000294>

**Podsakoff, P. M., MacKenzie, S. B., Lee, J.-Y., & Podsakoff, N. P.** (2003). Common method biases in behavioral research: A critical review of the literature and recommended remedies. *Journal of Applied Psychology*, 88(5), 879–903. <https://doi.org/10.1037/0021-9010.88.5.879>

**Podsakoff, P. M., Podsakoff, N. P., Williams, L. J., Huang, C., & Yang, J.** (2024). Common method bias: It's bad, it's complex, it's widespread, and it's not easy to fix. *Annual Review of Organizational Psychology and Organizational Behavior*, 11, 17–61. <https://doi.org/10.1146/ANNUREV-ORGPSCY-110721-040030>

**Poletti, M., Preti, A., & Raballo, A.** (2023). From economic crisis and climate change through COVID-19 pandemic to Ukraine war: A cumulative hit-wave on adolescent future thinking and mental well-being. *European Child and Adolescent Psychiatry*, 32(9), 1815–1816. <https://doi.org/10.1007/S00787-022-01984-X>

**Racine, N., McArthur, B. A., Cooke, J. E., Eirich, R., Zhu, J., & Madigan, S.** (2021). Global prevalence of depressive and anxiety symptoms in children and adolescents during COVID-19: A meta-analysis. *JAMA Pediatrics*, 175(11), 1142–1150. <https://doi.org/10.1001/JAMAPEDIATRICS.2021.2482>

**Reininghaus, U., Paetzold, I., Rauschenberg, C., Hirjak, D., Banaschewski, T., Meyer-Lindenberg, A., Boehnke, J. R., Boecking, B., & Schick, A.** (2023). Effects of a novel, transdiagnostic ecological momentary intervention for prevention, and early intervention of severe mental disorder in youth (EMIcompass): Findings from an exploratory randomized controlled trial. *Schizophrenia Bulletin*, 49(3), 592–604. <https://doi.org/10.1093/SCHBUL/SBAC212>

**Rutter, M.** (2012). Resilience as a dynamic concept. *Development and Psychopathology*, 24(2), 335–344. <https://doi.org/10.1017/S0954579412000028>

**Schneiders, J., Nicolson, N. A., Berkhof, J., Feron, F. J., Van Os, J., & Devries, M. W.** (2006). Mood reactivity to daily negative events in early adolescence: Relationship to risk for psychopathology. *Developmental Psychology*, 42(3), 543–554. <https://doi.org/10.1037/0012-1649.42.3.543>

**Schultze-Lutter, F., Schimmelemann, B. G., & Schmidt, S. J.** (2016). Resilience, risk, mental health and well-being: Associations and conceptual differences. *European Child & Adolescent Psychiatry*, 25(5), 459–466. <https://doi.org/10.1007/s00787-016-0851-4>

**Schwerdtfeger, A. R., & Dick, K.** (2019). Episodes of momentary resilience in daily life are associated with HRV reductions to stressful operations in firefighters: An ambulatory assessment approach using bayesian multilevel modeling. *The Journal of Positive Psychology*, 14(5), 593–602. <https://doi.org/10.1080/17439760.2018.1497689>

**Scott, S. B., Ram, N., Smyth, J. M., Almeida, D. M., & Sliwinski, M. J.** (2017). Age differences in negative emotional responses to daily stressors depend on time since event. *Developmental Psychology*, 53(1), 177–190. <https://doi.org/10.1037/DEV0000257>

**Seery, M. D., & Quinton, W. J.** (2016). Understanding resilience: From negative life events to everyday stressors. *Advances in Experimental Social Psychology*, 54, 181–245. <https://doi.org/10.1016/BS.AESP.2016.02.002>

**Smith, B. W., Dalen, J., Wiggins, K., Tooley, E., Christopher, P., & Bernard, J.** (2008). The brief resilience scale: Assessing the ability to bounce back. *International Journal of Behavioral Medicine*, 15, 194–200. <https://doi.org/10.1080/10705500802222972>

**Southwick, S. M., Bonanno, G. A., Masten, A. S., Panter-Brick, C., & Yehuda, R.** (2014). Resilience definitions, theory, and challenges: Interdisciplinary perspectives. *European Journal of Psychotraumatology*, 5(1), 25338. <https://doi.org/10.3402/ejpt.v5.25338>

**Southwick, S. M., Litz, B. T., Charney, D., & Friedman, M. J.** (2011). *Resilience and Mental Health: Challenges Across the Lifespan*. Cambridge University Press.

**Sytine, A. I., Britt, T. W., Sawhney, G., Wilson, C. A., & Keith, M.** (2019). Savoring as a moderator of the daily demands and psychological capital relationship: A daily diary study. *The Journal of Positive Psychology*, 14(5), 641–648. <https://doi.org/10.1080/17439760.2018.1519590>

**Tedeschi, R. G., & Calhoun, L. G.** (1996). The posttraumatic growth inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress*, 9, 455–471. <https://doi.org/10.1007/BF02103658>

**Thapar, A., Collishaw, S., Pine, D. S., & Thapar, A. K.** (2012). Depression in adolescence. *The Lancet*, 379(9820), 1056–1067. [https://doi.org/10.1016/S0140-6736\(11\)60871-4](https://doi.org/10.1016/S0140-6736(11)60871-4)

**Trudel-Fitzgerald, C., Boucher, G., Morin, C., Mondragon, P., Guimond, A. J., Nishimi, K., Choi, K. W., & Denckla, C.** (2023). Coping and emotion

regulation: A conceptual and measurement scoping review. *Canadian Psychology / Psychologie canadienne*, 65(3), 149–162. <https://doi.org/10.1037/cap0000377>

Uhlhaas, P. J., Davey, C. G., Mehta, U. M., Shah, J., Torous, J., Allen, N. B., Avenevoli, S., Bella-Awusah, T., Chanen, A., Chen, E. Y. H., Correll, C. U., Do, K. Q., Fisher, H. L., Frangou, S., Hickie, I. B., Keshavan, M. S., Konrad, K., Lee, F. S., Liu, C. H. . . . Wood, S. J. (2023). Towards a youth mental health paradigm: A perspective and roadmap. *Molecular Psychiatry*, 28(8), 3171–3181. <https://doi.org/10.1038/s41380-023-02202-z>.

Unger, H., Kraiss, J., & Vaessen, T. (2023). Temporal dynamics of momentary mental resilience, loneliness and self-esteem in adults from the general population: An experience sampling study [Bachelor thesis]. University of Twente. <http://essay.utwente.nl/95348/>

Vaessen, T., van Nierop, M., Decoster, J., Delespaul, P., Derom, C., de Hert, M., Jacobs, N., Menne-Lothmann, C., Rutten, B., Thiery, E., van Os, J., van Winkel, R., Wichers, M., & Myin-Germeys, I. (2017). Is sensitivity to daily stress predictive of onset or persistence of psychopathology? *European Psychiatry*, 45, 167–173. <https://doi.org/10.1016/J.EURPSY.2017.07.002>

Vaessen, T., Viechtbauer, W., van der Steen, Y., Gayer-Anderson, C., Kempton, M. J., Valmaggia, L., McGuire, P., Murray, R., Garety, P., Wykes, T., Morgan, C., Lataster, T., Lataster, J., Collip, D., Hernaus, D., Kasanova, Z., Delespaul, P., Oorschot, M., Claes, S. . . . Myin-Germeys, I. (2019). Recovery from daily-life stressors in early and chronic psychosis. *Schizophrenia Research*, 213, 32–39. <https://doi.org/10.1016/J.SCHRES.2019.03.011>

van Harmelen, A.-L. (2022). *Veerkracht bestaat niet*. Oratie, Leiden University.

van Harmelen, A.-L., Kievit, R. A., Ioannidis, K., Neufeld, S., Jones, P. B., Bullmore, E., Dolan, R., Fonagy, P., & Goodyer, I. (2017). Adolescent friendships predict later resilient functioning across psychosocial domains in a healthy community cohort. *Psychological Medicine*, 47(13), 2312–2322. <https://doi.org/10.1017/S0033291717000836>

van Roekel, E., Keijsers, L., & Chung, J. M. (2019). A review of current ambulatory assessment studies in adolescent samples and practical recommendations. *Journal of Research on Adolescence*, 29(3), 560–577. <https://doi.org/10.1111/jora.12471>

Wagner, B. M., Compas, B. E., & Howell, D. C. (1988). Daily and major life events: A test of an integrative model of psychosocial stress. *American Journal of Community Psychology*, 16(2), 189–205. <https://doi.org/10.1007/BF00912522>

Wagnild, G. M., & Young, H. M. (1993). Development and psychometric evaluation of the resilience scale. *Journal of nursing measurement*, 1(2), 165–17847.

Walsh, W. A., Dawson, J., & Mattingly, M. J. (2010). How Are We Measuring Resilience Following Childhood Maltreatment? Is the Research Adequate and Consistent? What is the Impact on Research, Practice, and Policy? *Trauma, Violence, & Abuse*, 11(1), 27–41. <https://doi.org/10.1177/1524838009358892>

Weber, J., Angerer, P., & Apolinário-Hagen, J. (2022). Physiological reactions to acute stressors and subjective stress during daily life: A systematic review on ecological momentary assessment (EMA) studies. *PLOS ONE*, 17(7), e0271996. <https://doi.org/10.1371/JOURNAL.PONE.0271996>

Werner, E. E. (1995). Resilience in development. *Current Directions in Psychological Science*, 4(3), 81–84. <https://doi.org/10.1111/1467-8721.ep10772327>

Werner, E. E., & Smith, R. S. (2001). *Journeys from Childhood to Midlife: Risk, Resilience, and Recovery*. Cornell University Press.

WHO (2021). *Mental Health of Adolescents*. World Health Organization. Retrieved 10 April 2024, from <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health>

Wiederhold, B. K. (2022). The escalating crisis in adolescent mental health. *Cyberpsychology, Behavior, and Social Networking*, 25(2), 81–82. <https://doi.org/10.1089/CYBER.2022.29237.EDITORIAL>

Wright, A. G. C., Aslinger, E. N., Bellamy, B., William, E. A. E., & Woods, C. (2019). Daily stress and hassles. In *The Oxford Handbook of Stress and Mental Health* (pp. 27–44). Oxford University Press.

Yang, Y. (2020). Daily stressor, daily resilience, and daily somatization: The role of trait aggression. *Personality and Individual Differences*, 165, 110141. <https://doi.org/10.1016/J.PAID.2020.110141>

Zautra, A. J., Guarnaccia, C. A., & Dohrenwend, B. P. (1986). Measuring small life events. *American Journal of Community Psychology*, 14(6), 629. <https://doi.org/10.1007/BF00931340>