



Universiteit
Leiden

The Netherlands

Illuminating host defence against mycobacterial infection: interactions with autophagy and LC3-associated phagocytosis

Muñoz Sánchez, S.

Citation

Muñoz Sánchez, S. (2026, February 3). *Illuminating host defence against mycobacterial infection: interactions with autophagy and LC3-associated phagocytosis*. Retrieved from <https://hdl.handle.net/1887/4288590>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4288590>

Note: To cite this publication please use the final published version (if applicable).



CHAPTER 1

General Introduction & Thesis outline

1.1 Mycobacteria and immune defence: a delicate balance

Tuberculosis (TB) is the world's leading infectious disease killer, causing a total of 1.25 million deaths in 2023. TB is present in all countries and age groups, though half of the cases are distributed among Bangladesh, China, India, Indonesia, Nigeria, Pakistan, Philippines and South Africa¹. The bacterium *Mycobacterium tuberculosis* (*Mtb*) is the causative agent of TB. This pathogen has coevolved with humans over thousands of years^{2,3}, specialising in evasion, modulation and exploitation of the host immune response^{4,5}. *Mtb* is transmitted through aerosol droplets from sick individuals. Once inhaled by the new host, *Mtb* is phagocytosed by alveolar macrophages, where, by undermining cellular defence mechanisms such as lysosomal trafficking, *Mtb* is capable of surviving intracellularly^{6,7}. As such, bacteria can establish a replicative niche as well as exploit macrophages as vehicles for dissemination⁸.

There are over 200 species in the *Mycobacterium* genus⁹, and they are classified into different groups: TB-causing, leprosy-causing and nontuberculous mycobacteria (NTM). The TB-causing group includes species capable of causing disease in humans or animals, like *Mtb* and *M. bovis*, respectively. NTM can cause disease in immunocompromised individuals, like *M. marinum* (*Mm*) and *M. avium*¹⁰. These diseases manifest among others as lung infections, skin infections and disseminated infections. *Mm*, which causes skin infection in humans, is of particular interest because the type of disease that it causes in cold-blooded animals shows strong similarities to human TB. In this thesis, we use *Mm* infections in zebrafish larvae as a model to get a better understanding of host-pathogen interactions during the early stages of infection, where the pathogen establishes its replicative niche in macrophages.

Mycobacterial infection success is, above all, due to the ability of mycobacteria to survive in diverse intracellular environments. Several virulence factors mediate intracellular survival. For example, it has been shown that both *Mtb* and *Mm* avoid acidified compartments but also have acidic resistance, which enables them to withstand eventual exposure to acidification to some extent^{11,12}. The mycobacterial membrane serine protease MarP is one of the virulence factors that enables evasion of acidification, as studies demonstrated that MarP-deficient mycobacteria are killed specifically in lysosomes¹¹. Once degradation is avoided, bacteria can replicate, eliciting the macrophage's lytic cell death or necrosis. This allows the bacteria to infect other macrophages and disseminate.

To halt infection dissemination and promote mycobacterial elimination, the host creates a physical immunological barrier, the granuloma, the pathological hallmark of TB. Granulomas are organised aggregates of macrophages, multinucleated giant cells, epithelioid and foamy cells, neutrophils, and lymphocytes.⁶ Granulomas succeed in restricting and containing the infection, which requires an effective adaptive immune response. Immunocompromised individuals with HIV infection present CD4⁺T cell dysfunction, and therefore, they fail to form well-organised granulomas and develop active TB^{13,14}. Mycobacteria have developed strategies to overcome the host defences in the granuloma and manipulate them for their growth. Research performed in the zebrafish-*Mm* infection model enabled the visualisation of early granuloma formation and the identification of host and bacterial factors contributing to this process¹⁵⁻¹⁸. In infected macrophages, the mycobacterial type VII secretion system ESX-1 and its effector protein early secreted antigenic target 6 kDa (ESAT-6) induce apoptotic cell death and recruitment of uninfected macrophages¹⁹. ESX-1/ESAT-6 has been proposed to stimulate the recruitment of new macrophages by inducing the secretion of matrix metalloproteinase 9 (MMP9) in epithelial cells surrounding the nascent granuloma, resulting in cellular expansion of the granuloma²⁰. Newly recruited macrophages phagocytose dying infected cells and their bacterial content; with this, mycobacteria ensure the expansion of their replicative niche.

When infected macrophages in the granuloma die by necrosis instead of apoptosis, bacteria are released within the granuloma environment, forming an acellular necrotic core called a caseous core. In this region, bacteria are surrounded by layers of foamy macrophages that provide a niche for dormant bacteria^{14,21}. At the necrotic caseum, the passive diffusion of antibiotics is limited, producing regions of suboptimal antibiotic concentrations and monotherapy, contributing to the selection of drug-resistant strains²²⁻²⁴. Drug-resistant *Mtb* strains are a major concern for global health. In 2023, 175,923 people were diagnosed and treated for Multidrug-resistant TB (MDR-TB)¹. The appearance of drug-resistant *Mtb* variants has been blamed on inadequate implementation of control measures. Still, since MDR-TB evolve in well-functioning health systems, more factors need to be considered²⁵. Elements of the host-pathogen interaction are increasingly recognised to influence the evolution of drug resistance^{3,25}. Increased understanding of host-pathogen interaction could enable the development of better or new therapies for the prevention and treatment of TB^{26,27}.




Figure 1.1: *Mycobacterium tuberculosis* pathogenesis.

Mtb is transmitted through aerosol droplets from sick individuals. Once inhaled by the new host, *Mtb* is phagocytosed by alveolar macrophages. Infected macrophages die by apoptosis, causing the recruitment of immune cells. Newly recruited macrophages phagocytose dying infected cells and their bacterial content; with this, mycobacteria ensure the expansion of their replicative niche. To halt infection dissemination and promote mycobacterial elimination, the host creates a physical immunological barrier, the granuloma. When infected macrophages in the granuloma die by necrosis, bacteria are released within the granuloma environment. Finally, the granuloma collapses, and the infection is disseminated. Schematic created with BioRender.

1.2 One beneficiary, many obstructions: How mycobacteria halt intracellular vesicle trafficking

Macrophages are the first responders against invading microbes, including infection by mycobacterial species. Pathogen-associated molecular patterns (PAMPs) on the *Mtb* surface, such as glycolipids, lipoproteins, and carbohydrates, activate host cell surface and intracellular pathogen recognition receptors (PRRs), including Toll-like receptors (TLRs), Nod-like receptors (NLRs), C-type lectin receptors (CLRs), and cyclic GMP-AMP synthase (cGAS)-STING^{28,29}. These receptors are involved in the activation of innate immune response mechanisms against mycobacterial infection, consisting of the phagolysosomal pathway, autophagy, inflammasome activation, and apoptosis³⁰. Mycobacteria are phagocytosed in a phagosome, whose aim is to traffic the bacteria to lysosomal degradation. Effective pathogen transport relies on a highly dynamic interaction between phagosomes and intracellular vesicles, where phagosomes undergo structural and compositional changes referred to as phagosome maturation.³¹ Phosphoinositides (PtdIns) play a significant role in phagolysosome formation and maturation. Through recruitment and interaction with effector proteins, they regulate the rearrangement of the actin cytoskeleton, ion and organelle transport, signal transduction, and membrane fusion-fission events mediating phagosome maturation and fusion with lysosomes³²⁻³⁴. After phagocytosis, early phagosomes fuse with early endosomes (EE), acquiring the RAS-associated protein RAB5A (Rab5) GTPase, which regulates fusion and traffic of EE. GTP-bound Rab5 recruits effector vacuolar protein sorting 34 homolog (Vps34), a phosphatidylinositol (PI) 3-kinase (PI3K), that generates phosphatidylinositol-3-phosphate (PtdIns3P)³⁵. Rab5 and PtdIns3P recruit EEA1, which docks early phagosomes and EE together, priming them to fuse via soluble NSF attachment protein receptor (SNARE). Conversion of early to late phagosomes requires the recruitment of RAS-associated protein RAB7A (Rab7) GTPase, which catalyses the fusion with LE and lysosomes. Finally, phagosome-lysosome fusion occurs via SNAREs associated with lysosomes. As the phagolysosome is shaped, it acquires v-ATPase H⁺ and nicotinamide adenine dinucleotide phosphate (NADPH) oxidase activities, which contribute to the acidification and the production of reactive oxygen species (ROS) in the phagolysosome, respectively³¹.

Mtb has evolved virulence factors that alter phagosome dynamics and maturation, allowing it to replicate within infected macrophages. *Mtb* is capable of blocking phagosome maturation by affecting phagosome acidification, the acquisition of hydrolytic enzymes and antimicrobial peptides^{6,8,14}. To interfere with this pathway at the Rab GTPases level, *Mtb* secretes the nucleotide diphosphate

Considering the importance of PtdIns in phagolysosome formation, the enzymes catalysing PtdIns production play critical roles in the innate immune response. As previously mentioned, PtdIns3P, together with Rab5, recruits EEA1 and facilitates fusion events between early phagosomes and EE. PtdIns3P can be removed from the phagosome due to its conversion by the enzyme phosphoinositide kinase, FYVE-type zinc finger containing lipid kinase (PIKfyve), which converts PtdIns(3)P to PtdIns(3,5)P₂⁴³. PIKfyve inhibition impairs phagosome maturation, resulting in reduced phagosome-lysosome fusion and degradative capacity^{44,45}. PIKfyve has been implicated in the pathogenesis of intracellular pathogens, such as *Coxiella burnetii*^{46,47}, *Legionella pneumophila*⁴⁸, and *Salmonella Typhimurium*^{49,50}. In these infections, PIKfyve inhibition restricted the expansion of the pathogen-containing vacuoles, indicating that these pathogens exploit PIKfyve to transform the phagosome into a replicative niche. The role of PIKfyve in mycobacterial infections has remained unknown and is addressed in this thesis.

1.3 Autophagy to the rescue?: The need for timely intervention in anti-mycobacterial immunity

Inside the phagosome, *Mtb* focuses on ensuring a successful intracellular lifecycle. Like many other aspects of *Mtb* virulence, this is determined by the ESX-1 type VII secretion system⁵¹. ESX-1 mediates phagosomal membrane damage that enables *Mtb* to access nutrients and deliver effectors to the cytosol. Damaged *Mtb* phagosomes expose their intraluminal content, activating the cytosolic surveillance pathway. Galectins target host glycans inside the phagosome^{52,53}, and bacteria are labelled with ubiquitin by SMURF⁵⁴ and parkin⁵⁵ ubiquitin ligases. Galectins and ubiquitin are recognised by selective autophagy adaptors, such as Nuclear Domain 10 Protein 52 (NDP52), optineurin (OPTN), and sequestosome 1 (SQSTM1, also known as p62), and are directed to autolysosomal degradation⁵⁶. Phagosomal damage also allows *Mtb* molecules to be released into the cytosol and recognised by cytosolic PRRs. *Mtb* DNA^{57,58} and cyclic-di-adenosine monophosphate⁵⁹ (c-di-AMO) are recognised by cytosolic PRRs, cyclic GMP-AMP synthase⁶⁰⁻⁶² (cGAS) and stimulator of interferon genes protein⁶³ (STING), inducing type I IFN⁶⁴ response and autophagy.

Autophagy is a cellular pathway important for maintaining cellular homeostasis as well as for immunity. By delivering cytoplasmic contents to the lysosome for degradation, it provides adaptation to nutrient starvation, selectively degrades proteins and foreign substances, clears damaged organelles, and

counteracts ageing⁶⁵. In the immune system, autophagy plays protective roles during microbial infections as a cell-autonomous defence mechanism of innate immune cells, by controlling inflammation, and by generating peptides for antigen presentation^{66,67}. Upon mycobacterial infection, selective autophagy is part of the host macrophages' immune response. As discussed above, mycobacteria are labelled with ubiquitin^{68,69} and recognised by autophagy adapters like sequestosome 1 (SQSTM1)⁷⁰ and optineurin (OPTN)^{71,72}. These adapters recruit the microtubule-associated protein 1 light chain 3 (LC3), which coats the double membrane of nascent phagophores (cytosolic isolation membranes), causing the engulfment of the bacteria into autophagosomes. In the same way as in phagosome maturation, PtdIns play a role in autophagosome maturation. Autophagosomes form at parts of the endoplasmic reticulum (ER) enriched in phosphatidylinositol-3-phosphate (PtdIns3P). FAK family kinase-interacting protein of 200 KDa (FIP200) interacts with the ER-enriched domain⁷³ and induces the assembly of the Unc-51-like autophagy activating kinase 1 (ULK1) complex⁷⁴, comprised of FIP200, autophagy-related proteins ATG13 and ATG101, and ULK1. ULK1 phosphorylates and activates phosphatidylinositol 3-kinase class III complex 1 (PI3KC3-C1)⁷⁵, allowing the phagophore expansion. PI3KC3-C1 is composed of five subunits: VPS34, PI3-kinase regulatory subunit 4 (VPS15), Beclin 1 (BECN1), ATG14, and nuclear receptor-binding factor 2 (NRBF2). Phagophore elongation and closure to a mature autophagosome require phosphatidylethanolamine (PE) to be conjugated to LC3⁷⁶. Autophagosome closure is catalysed by the endosomal sorting complex required for transport complex (ESCRT)^{77,78}, and regulated by Ras-related protein Rab-5A (Rab5)^{79,80}. Consequently, pleckstrin homology domain-containing family M member 1 (PLEKHM1), ectopic P-granules autophagy protein 5 Homolog (EPG5) and RAB7⁸¹ tether autophagosomes with lysosomes. Ultimately, two SNARE complexes, syntaxin 17 (STX17)-synaptosome-associated protein 29 (SNAP29)-vesicle-associated membrane protein 7 (VAMP7/8), and synaptobrevin homolog YKT6 (YKT6)-SNAP29-STX7, trigger the autophagosome fusion with the lysosome to degrade its content^{66,76,82}.

While autophagy is important as a host defence mechanism against mycobacterial infection⁸³⁻⁸⁵, *Mtb* has evolved ways to evade autophagic degradation and benefit from this host defence mechanism. *Mtb* disturbs autophagy initiation and flux, as well as autophagosome fusion with lysosomes. Furthermore, *Mtb* can also escape from autophagosomes back into the cytosol⁸⁶. Since autophagy initiation depends upon PI3P, the effectors discussed above that impair PI3P production, such as ManLAM, have also been found to impair autophagy. In vitro studies showed that ManLAM reduces LC3 trafficking to phagosomal membranes⁸⁷. Additionally, *Mtb* activates the host F-actin-binding

protein Coronin 1a (CORO1A), which blocks the p38 MAPK pathway required for autophagy induction through TLR signalling⁸⁸. Regarding autophagic flux, it has been shown that *Mtb* induces the expression of the host microRNA miR-33, which in turn downregulates the expression of autophagy genes including ATG5, ATG12, LC3, and lysosome-associated membrane glycoprotein 1 (LAMP1). This leads to a reduction in autophagic flux, as evidenced by a decrease in the recruitment of p62 and LC3⁸⁹. Moreover, *Mtb* secretes enhanced intracellular survival (Eis), which negatively modulates autophagy, and this inhibition depends on ROS production^{90,91}. Lastly, autophagosome-lysosome fusion blockage by *Mtb* is linked to Rab7⁹² and requires, among others, the lipid phosphatase SapM^{93,94}. SapM interacts with Rab7, inhibiting its recruitment to autophagosomes, blocking the maturation of *Mtb*-containing autophagosomes into autolysosomes⁹⁵.

Accumulating evidence establishes a crucial protective role for autophagy in the immune response against early mycobacterial infections⁹⁶⁻⁹⁹. As a result, more research is needed to elucidate the regulatory roles of autophagy during *Mtb* infection and to consider autophagy as a potential therapeutic target for TB control^{183,100}.

1.4 Is the alternative better? LC3-associated phagocytosis in anti-mycobacterial immunity

As a substitute for the autophagy pathway or a parallel pathway, LC3-associated phagocytosis (LAP)^{101,102} is a mechanism of phagocytic cells, including macrophages, where a subset of autophagy machinery is used in the LC3 lipidation to the single membrane of the phagosome (forming a LAPosome)^{101,103,104}. In comparison to intracellular autophagic cargo, LAP cargo comes from the extracellular environment¹⁰⁵, where it binds surface receptors such as fragment crystallizable gamma receptor (FcγR¹⁰⁶), TLRs¹⁰⁷, C-type lectin domain family 7 member A (CLEC7/dectin-1¹⁰⁸), or T-cell immunoglobulin mucin receptor 4 (TIM4)¹⁰⁹. Following receptor ligation, phagosome cup development is regulated by the same mechanism that regulates phagocytosis (previously discussed³⁵), and is, in contrast to phagophore formation in autophagy, independent of the ULK1 initiation complex. LAP requires the assembly of the PI3K3 complex, but its composition differs from the autophagy complex. Formation of the LAP complex uniquely requires ultraviolet radiation resistance-associated gene protein (UVRAG) and RUN domain and cysteine-rich domain-containing, Beclin 1-interacting protein (Rubicon or Rubcn), and shares with autophagy the presence of Beclin1, VPS34 and VPS15¹¹⁰. VPS34 lipid kinase produces PI3P on the surface of the closed phagosome, on which ROS production

and LC3 lipidation depend¹¹¹. ROS production in the phagosomal lumen is carried out by NADPH oxidase-2 complex (NOX2)¹¹², which associates with PI3P on the phagosome membrane through its p40^{phox} subunit¹¹³. Furthermore, NOX2-generated ROS are required for the assembly of V-ATPase subunits, allowing the interaction with ATG16L1, which drives LC3 conjugation to the phagosomal membrane¹¹⁴. Thus, both NOX2 activity and PI3P generation are needed for the assembly of the LC3 conjugation machinery, where LC3 is transferred from ATG3-LC3 to the PE lipid by the vATPase-ATG16L axis^{115,116}. Following the lipidation of LC3 on the phagosomal surface, the now denominated LAPosome fuses with the lysosome, and the phagocytosed cargo becomes degraded.

Pathogens have been shown to induce or suppress LAP activation¹¹⁷. Research on fungi such as *Candida albicans*¹¹⁸ and *Aspergillus fumigatus*¹¹⁹⁻¹²¹ has established a melanin-mediated LAP blockage across species^{122,123}, underlining LAP's relevance in fungal infection clearance¹²⁴. Studies in bacteria have shown species-specific mechanisms involved in LAP recognition and activation, with a significant molecular overlap¹¹⁰. Among the bacteria targeted by LAP are *Legionella dumoffii*,¹²⁵ *Listeria monocytogenes*^{126,127}, *Burkholderia pseudomallei*^{128,129}, *Shigella flexneri*^{130,131}, *Salmonella Typhimurium*^{114,132,133}. Each species has evolved unique strategies to deal with LAP-mediated degradation¹³⁴. Little is known about the role of LAP in mycobacterial infection. In vitro studies have shown that *Mm* resides transiently in single membrane compartments and that LC3 recruitment depends on the ESX-1 secretion system¹³⁵. The *Mtb* virulence factor CpsA has been shown to inactivate NADPH oxidase¹³⁶, protecting bacteria from LAP elimination. Further research is needed to elucidate the underlying mechanisms which initiate LAP and lead to LC3 recruitment onto mycobacteria-containing phagosomes, as well as the relevance of LAP as a defence against mycobacterial infection.

Rubcn is of special interest in this regard, as it has been characterised as an autophagy inhibitor by impeding PI3P production and LC3 lipidation¹³⁷⁻¹³⁹, and at the same time, as a LAP activator and essential component¹⁰¹. In LAP, Rubcn stabilises NOX2 complex through direct binding with p22^{phox} subunit, thus facilitating ROS production and stimulating the subsequent interaction between v-ATPase and ATG16L1, which drives LC3 conjugation to the phagosomal membrane^{114,140}. In autophagy, Rubcn interacts with VPS34 via its RUN domain, suppressing its lipid kinase activity and negatively regulating autophagosome formation^{137,141,142}. Lastly, Rubcn interacts with UVRAG through its coiled-coil domain. UVRAG interacts with the C-VPS/HOPS complex in late endosomes, which in turn activates Rab7 by promoting GTP binding¹⁴³. Rab7 activation is crucial for the transition from early to late endosomes in endosome maturation. UVRAG-Rubcn association in early endosomes, where Rubcn is enriched, impedes UVRAG-C-VPS/HOPS binding

and, consequently, Rab7 activation. As a result, UVRAG sequestration by Rubcn negatively regulates endocytic trafficking and autophagy¹⁴⁴.

Rubicon has been shown to promote host defence against several pathogens, including *Listeria monocytogenes*^{126,127}, *Aspergillus fumigatus*^{119,120} and *Salmonella Typhimurium*¹³². In addition, it has been reported that the maturation of *Mtb*-containing phagosomes is negatively regulated by Leucine-rich repeat kinase 2 (LRRK2) through recruiting the PI3KC3 complex and Rubcn¹⁴⁵. However, the role of Rubcn in mycobacterial infection remains to be further established.




Figure 1.3: Molecular mechanisms of phagocytosis, LC3-associated phagocytosis and autophagy during *Mtb* infection.

Mtb activates host cell surface PRR and is phagocytosed in a single membrane phagosome. After phagocytosis, early phagosomes fuse with EE, acquiring Rab5. Sequential recruitment of Rab7 by fusion with LE catalyses the fusion with lysosomes. In LAP, NOX2, stabilised by Rubcn, assembles on the *Mtb*-phagosome, inducing ROS production in the phagosomal lumen. ROS allow the assembly of V-ATPase and its interaction with ATG16L1, which drives LC3 conjugation to the phagosomal single membrane, forming the LAPosome. *Mtb* virulence factor ESX-1 promotes phagosomal damage, activating the cytosolic surveillance pathway. *Mtb* are labelled with ubiquitin and recognised by autophagy adapters (SLR). These adapters recruit LC3 to the double membrane of the phagophore. This elongates and closes, forming the autophagosome, which is targeted to lysosomal degradation. EE: early endosome; LE: late endosome; PRR: pattern recognition receptor; SLR: sequestosome 1-like receptor, also known as selective autophagy receptor. Schematic created with BioRender.

1.5 When one door closes, a transparent window opens: Illuminating mycobacterial host-pathogen interactions in zebrafish

Even though TB has afflicted human populations for millennia, many unknowns persist about this highly successful pathogen. Technological progress has made it possible to design more complex experiments in order to reduce the gap between disease models and human disease. In this context, studies using animal models are crucial in that they have provided information on the disease mechanism, while at the same time, they have been used for pre-clinical testing of TB treatments. Animal models used for TB research include *Mus musculus* (mouse), *Oryctolagus cuniculus* (rabbit), *Cavia porcellus* (guinea pig), and non-human primates (NHP). Although no animal model can recapitulate TB in its entirety, their collective contributions have been significant¹⁴⁶. Among these, the mouse has been the most commonly used model for vaccine research¹⁴⁷ and drug testing against TB¹⁴⁸. However, differences in immunological responses between the mouse model and humans, such as difficulties in mimicking caseating granuloma formation, have promoted the development of new animal models. For the past two decades, *Danio rerio* (zebrafish) has become an alternative for studying mycobacterial infection^{149,150}. Human and zebrafish have similar innate and adaptive immune systems, and more than 80% of human disease-associated genes are present in zebrafish^{151,152}. The Zebrafish TB model is based on the Zebrafish-*Mm* natural host-pathogen interaction. *Mm* and *Mtb* genomes share an 85% identity¹⁵³, and *Mm* has been shown to cause infection conducive to granuloma formation in larvae¹⁵⁴ and adult individuals¹⁵⁵. The embryo/larva model is used to study the innate immune response against mycobacterial infection in isolation from adaptive immunity. The embryo and larva external development and transparency enable microinjection of fluorescently labelled *Mm* strains into transgenic zebrafish (with fluorescently labelled macrophages or neutrophils)¹⁵⁶, facilitating imaging and analysis of early host-pathogen interactions in vivo. The Zebrafish-*Mm* infection model has contributed to many aspects of the host-pathogen interaction, including the understanding of the role of granuloma formation^{19,157,158} and vascularisation^{159,160}, infection dissemination¹⁶¹⁻¹⁶³, and the intracellular mechanisms of infection^{11,164} and host defence^{71,165-167}. Significantly, the use of the model led to understanding that many of the cellular pathological effects observed in mycobacterial infection, including the initiation of granulomas, are dependent on the ESX-1 secretion system^{168,169}.

As discussed in detail in Chapter 2, the zebrafish model has proven very useful to investigate autophagy's role in infection^{170,171}. Its capacity for whole-organism

in vivo imaging provides a unique advantage for the autophagy field, enabling the observation of host-pathogen interactions in a physiological context at an intracellular level.




Figure 1.4: Zebrafish-*Mm* tail fin infection model.

Microinjection of fluorescently mCrimson-labelled *Mm* (150 CFU, pseudo colour magenta) strains into the tail fin area of 3dpf zebrafish larvae carrying the construct (*mpeg1:mCherry-F*) that labels macrophages (pseudo colour yellow). Confocal laser-scanning microscopy images of 1, 4, and 24 hpi. At 1 hpi, macrophages are already recruited at the site of infection and phagocytosis of bacteria can be observed. At 4 hpi, all the bacteria clusters in the site of infection appeared to be phagocytosed, and *Mm*-phagosomes are visible. At 24 hpi, the bacterial cluster sizes and numbers are increased. Scale bars: 20 μ m. hpi: hours post-infection.

1.6 Thesis Outline

Since *Mtb* was identified as the causative agent of TB almost 150 years ago, research has been moving unceasingly to find a solution to this never-ending pandemic. Although much progress has been made on the understanding of the molecular pathogenesis of *Mtb* and related mycobacterial pathogens, several aspects remain unknown, limiting novel therapeutic development. This thesis focuses on adding knowledge to the host immune response during early mycobacterial infection, focusing on mycobacteria's autophagy-mediated degradation. This is achieved by coupling chemical and genetic inhibition to the zebrafish model of tuberculosis.

In **Chapter 1**, we present TB molecular pathogenesis and the involvement of autophagy and LAP in the immune response against mycobacterial infection.

Additionally, we introduce the zebrafish TB model as a unique model to study host-pathogen interactions in a whole-organism physiological context.

In **Chapter 2**, we review how research using zebrafish infection models has provided in vivo evidence of the significant role of autophagy and LAP as part of the defences against intracellular bacterial pathogens. These studies have increased the understanding of autophagy-associated host-pathogen interactions, which could favour the design of autophagy-modulating strategies oriented towards the degradation of specific pathogens.

In **Chapter 3**, we report on high-resolution live imaging data from the zebrafish TB model to characterise mycobacteria-autophagy interactions during the early stages of infection. By microinjecting fluorescent *Mm* into the tail fin tissue of GFP-Lc3 transgenic zebrafish larvae, we were able to visualise phagocytosed *Mm* clusters and LC3-positive *Mm*-containing vesicles within the first hour of infection. LC3 associations with *Mm*-vesicles were transient and remarkably heterogeneous, alternating between spacious, compact and composite morphologies.

In **Chapter 4**, we focus on the involvement of the phosphoinositide kinase PIKfyve in mycobacterial infection. PIKfyve catalyses the generation of PtdIns(3,5)P₂, which is critical for the maturation of autophagosomes and the proper function of lysosomes. We studied the role of PIKfyve in the zebrafish TB model using two chemical inhibitors of the enzymatic activity of PIKfyve, YM201636 and Apilimod. We found that in the infected macrophages, PIKfyve mediates maturation of (auto) phagosomal *Mm*-containing vesicles, increasing the acidification, and ultimately enhancing the cell autonomous resistance to mycobacterial infection.

In **Chapter 5**, we address the role of the RUN domain-containing protein Rubicon (Rubcn) in mycobacterial infection. Rubcn negatively regulates autophagy and stimulates LAP. Since mycobacterial virulence factors target autophagy and LAP, the efficacy of these pathways as host defence mechanisms remains unknown. We studied the role of Rubcn in the zebrafish-TB model by affecting *rubcn* transcript levels by morpholino KD and *rubcn* mRNA overexpression. We also generated a CRISPR-Cas9 zebrafish mutant line. Our results suggest that the lack of Rubcn affects *Mm* normal vesicle trafficking, impairing host defence. Our model provides evidence of Rubcn's protective role in early *Mm* infection.

In **Chapter 6**, the results obtained in this thesis are summarised and discussed in a broader context.

1.7 References

- 1 WHO. Global tuberculosis Report 2024. (Geneva: World Health Organization, 2024).
- 2 Comas, I. *et al.* Out-of-Africa migration and Neolithic coexpansion of *Mycobacterium tuberculosis* with modern humans. *Nat Genet* **45**, 1176-1182 (2013). <https://doi.org/10.1038/ng.2744>
- 3 Gagneux, S. Ecology and evolution of *Mycobacterium tuberculosis*. *Nature Reviews Microbiology* **16**, 202-213 (2018). <https://doi.org/10.1038/nrmicro.2018.8>
- 4 Cambier, C. J., Falkow, S. & Ramakrishnan, L. Host evasion and exploitation schemes of *Mycobacterium tuberculosis*. *Cell* **159**, 1497-1509 (2014). <https://doi.org/10.1016/j.cell.2014.11.024>
- 5 Chandra, P., Grigsby, S. J. & Philips, J. A. Immune evasion and provocation by *Mycobacterium tuberculosis*. *Nat Rev Microbiol* **20**, 750-766 (2022). <https://doi.org/10.1038/s41579-022-00763-4>
- 6 Bussi, C. & Gutierrez, M. G. *Mycobacterium tuberculosis* infection of host cells in space and time. *FEMS Microbiol Rev* **43**, 341-361 (2019). <https://doi.org/10.1093/femsre/fuz006>
- 7 Upadhyay, S., Mittal, E. & Philips, J. A. Tuberculosis and the art of macrophage manipulation. *Pathog Dis* **76** (2018). <https://doi.org/10.1093/femspd/fty037>
- 8 Sia, J. K. & Rengarajan, J. Immunology of *Mycobacterium tuberculosis* Infections. *Microbiol Spectr* **7** (2019). <https://doi.org/10.1128/microbiolspec.GPP3-0022-2018>
- 9 Diab, M. K., Soliman, T. H. A., Mohamed, A. M. & Elsemman, I. E. A novel in silico molecular tool for comprehensive differentiation of *Mycobacterium* species. *Scientific Reports* **15** (2025). <https://doi.org/10.1038/s41598-025-89148-0>
- 10 Turenne, C. Y. Nontuberculous mycobacteria: Insights on taxonomy and evolution. *Infect Genet Evol* **72**, 159-168 (2019). <https://doi.org/10.1016/j.meegid.2019.01.017>
- 11 Levitte, S. *et al.* Mycobacterial Acid Tolerance Enables Phagolysosomal Survival and Establishment of Tuberculous Infection In Vivo. *Cell Host & Microbe* **20**, 250-258 (2016). <https://doi.org/10.1016/j.chom.2016.07.007>
- 12 Gouzy, A., Healy, C., Black, K. A., Rhee, K. Y. & Ehrt, S. Growth of *Mycobacterium tuberculosis* at acidic pH depends on lipid assimilation and is accompanied by reduced GAPDH activity. *Proceedings of the National Academy of Sciences* **118**, e2024571118 (2021). <https://doi.org/10.1073/pnas.2024571118>
- 13 Boom, W. H., Schaible, U. E. & Achkar, J. M. The knowns and unknowns of latent *Mycobacterium tuberculosis* infection. *J Clin Invest* **131** (2021). <https://doi.org/10.1172/JCI136222>
- 14 Kinsella, R. L. *et al.* Perspectives and Advances in the Understanding of Tuberculosis. *Annu Rev Pathol* **16**, 377-408 (2021). <https://doi.org/10.1146/annurev-pathol-042120-032916>
- 15 Ramakrishnan, L. The Zebrafish Guide to Tuberculosis Immunity and Treatment. *Cold Spring Harbor Symposia on Quantitative Biology* **78**, 179-192 (2013). <https://doi.org/10.1101/sqb.2013.78.023283>
- 16 Tobin, D. M. & Ramakrishnan, L. Comparative pathogenesis of *Mycobacterium marinum* and *Mycobacterium tuberculosis*. *Cellular Microbiology* **10**, 1027-1039 (2008). <https://doi.org/10.1111/j.1462-5822.2008.01133.x>
- 17 Cronan, M. R. & Tobin, D. M. Fit for consumption: zebrafish as a model for tuberculosis. *Disease Models & Mechanisms* **7**, 777-784 (2014). <https://doi.org/10.1242/dmm.016089>

- 18 Van Leeuwen, L. M., Van Der Sar, A. M. & Bitter, W. Animal Models of Tuberculosis: Zebrafish. *Cold Spring Harbor Perspectives in Medicine* **5**, a018580-a018580 (2015). <https://doi.org/10.1101/cshperspect.a018580>
- 19 Davis, J. M. & Ramakrishnan, L. The role of the granuloma in expansion and dissemination of early tuberculous infection. *Cell* **136**, 37-49 (2009). <https://doi.org/10.1016/j.cell.2008.11.014>
- 20 Volkman, H. E. *et al.* Tuberculous granuloma induction via interaction of a bacterial secreted protein with host epithelium. *Science* **327**, 466-469 (2010). <https://doi.org/10.1126/science.1179663>
- 21 Russell, D. G., Cardona, P. J., Kim, M. J., Allain, S. & Altare, F. Foamy macrophages and the progression of the human tuberculosis granuloma. *Nat Immunol* **10**, 943-948 (2009). <https://doi.org/10.1038/ni.1781>
- 22 Moreno-Gamez, S. *et al.* Imperfect drug penetration leads to spatial monotherapy and rapid evolution of multidrug resistance. *Proc Natl Acad Sci U S A* **112**, E2874-2883 (2015). <https://doi.org/10.1073/pnas.1424184112>
- 23 Sarathy, J. P. *et al.* Prediction of Drug Penetration in Tuberculosis Lesions. *ACS Infect Dis* **2**, 552-563 (2016). <https://doi.org/10.1021/acsinfecdis.6b00051>
- 24 Strydom, N. *et al.* Tuberculosis drugs' distribution and emergence of resistance in patient's lung lesions: A mechanistic model and tool for regimen and dose optimization. *PLoS Med* **16**, e1002773 (2019). <https://doi.org/10.1371/journal.pmed.1002773>
- 25 Gygli, S. M., Borrell, S., Trauner, A. & Gagneux, S. Antimicrobial resistance in *Mycobacterium tuberculosis*: mechanistic and evolutionary perspectives. *FEMS Microbiology Reviews* **41**, 354-373 (2017). <https://doi.org/10.1093/femsre/fux011>
- 26 Borah, K., Xu, Y. & McFadden, J. Dissecting Host-Pathogen Interactions in TB Using Systems-Based Omic Approaches. *Frontiers in Immunology* **12** (2021). <https://doi.org/10.3389/fimmu.2021.762315>
- 27 Abreu, R., Giri, P. & Quinn, F. Host-Pathogen Interaction as a Novel Target for Host-Directed Therapies in Tuberculosis. *Frontiers in Immunology* **11** (2020). <https://doi.org/10.3389/fimmu.2020.01553>
- 28 Kleinnijenhuis, J., Oosting, M., Joosten, L. A., Netea, M. G. & Van Crevel, R. Innate immune recognition of *Mycobacterium tuberculosis*. *Clin Dev Immunol* **2011**, 405310 (2011). <https://doi.org/10.1155/2011/405310>
- 29 Liu, C. H., Liu, H. & Ge, B. Innate immunity in tuberculosis: host defense vs pathogen evasion. *Cellular & Molecular Immunology* **14**, 963-975 (2017). <https://doi.org/10.1038/cmi.2017.88>
- 30 Lerner, T. R., Borel, S. & Gutierrez, M. G. The innate immune response in human tuberculosis. *Cellular Microbiology* **17**, 1277-1285 (2015). <https://doi.org/10.1111/cmi.12480>
- 31 Fountain, A., Inpanathan, S., Alves, P., Verdawala, M. B. & Botelho, R. J. Phagosome maturation in macrophages: Eat, digest, adapt, and repeat. *Adv Biol Regul* **82**, 100832 (2021). <https://doi.org/10.1016/j.jbior.2021.100832>
- 32 Balla, T. Phosphoinositides: tiny lipids with giant impact on cell regulation. *Physiol Rev* **93**, 1019-1137 (2013). <https://doi.org/10.1152/physrev.00028.2012>
- 33 Nguyen, J. A. & Yates, R. M. Better Together: Current Insights Into Phagosome-Lysosome Fusion. *Frontiers in Immunology* **12** (2021). <https://doi.org/10.3389/fimmu.2021.636078>

- 34 Choy, C. H., Han, B. K. & Botelho, R. J. Phosphoinositide Diversity, Distribution, and Effector Function: Stepping Out of the Box. *Bioessays* **39** (2017).
<https://doi.org/10.1002/bies.201700121>
- 35 Levin, R., Grinstein, S. & Canton, J. The life cycle of phagosomes: formation, maturation, and resolution. *Immunological Reviews* **273**, 156-179 (2016).
<https://doi.org/10.1111/imr.12439>
- 36 Sun, J. *et al.* Mycobacterial nucleoside diphosphate kinase blocks phagosome maturation in murine RAW 264.7 macrophages. *PLoS One* **5**, e8769 (2010).
<https://doi.org/10.1371/journal.pone.0008769>
- 37 Roberts, E. A., Chua, J., Kyei, G. B. & Deretic, V. Higher order Rab programming in phagolysosome biogenesis. *J Cell Biol* **174**, 923-929 (2006).
<https://doi.org/10.1083/jcb.200603026>
- 38 Seto, S., Tsujimura, K. & Koide, Y. Rab GTPases regulating phagosome maturation are differentially recruited to mycobacterial phagosomes. *Traffic* **12**, 407-420 (2011).
<https://doi.org/10.1111/j.1600-0854.2011.01165.x>
- 39 Fratti, R. A., Chua, J., Vergne, I. & Deretic, V. Mycobacterium tuberculosis glycosylated phosphatidylinositol causes phagosome maturation arrest. *Proc Natl Acad Sci U S A* **100**, 5437-5442 (2003). <https://doi.org/10.1073/pnas.0737613100>
- 40 Vergne, I., Chua, J. & Deretic, V. Mycobacterium tuberculosis phagosome maturation arrest: selective targeting of PI3P-dependent membrane trafficking. *Traffic* **4**, 600-606 (2003). <https://doi.org/10.1034/j.1600-0854.2003.00120.x>
- 41 Puri, R. V., Reddy, P. V. & Tyagi, A. K. Secreted Acid Phosphatase (SapM) of Mycobacterium tuberculosis Is Indispensable for Arresting Phagosomal Maturation and Growth of the Pathogen in Guinea Pig Tissues. *PLoS ONE* **8**, e70514 (2013).
<https://doi.org/10.1371/journal.pone.0070514>
- 42 Vergne, I. *et al.* Mechanism of phagolysosome biogenesis block by viable Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* **102**, 4033-4038 (2005).
<https://doi.org/10.1073/pnas.0409716102>
- 43 Hazeki, K. *et al.* Essential roles of PIKfyve and PTEN on phagosomal phosphatidylinositol 3-phosphate dynamics. *FEBS Letters* **586**, 4010-4015 (2012).
<https://doi.org/10.1016/j.febslet.2012.09.043>
- 44 Kim, G. H. E., Dayam, R. M., Prashar, A., Terebiznik, M. & Botelho, R. J. PIKfyve Inhibition Interferes with Phagosome and Endosome Maturation in Macrophages. *Traffic* **15**, 1143-1163 (2014). <https://doi.org/10.1111/tra.12199>
- 45 Rivero-Rios, P. & Weisman, L. S. Roles of PIKfyve in multiple cellular pathways. *Curr Opin Cell Biol* **76**, 102086 (2022). <https://doi.org/10.1016/j.ceb.2022.102086>
- 46 Martinez, E. *et al.* Coxiella burnetii effector CvpB modulates phosphoinositide metabolism for optimal vacuole development. *Proc Natl Acad Sci U S A* **113**, E3260-3269 (2016). <https://doi.org/10.1073/pnas.1522811113>
- 47 Zhao, M. *et al.* Coxiella burnetii effector CvpE maintains biogenesis of Coxiella-containing vacuoles by suppressing lysosome tubulation through binding PI(3)P and perturbing PIKfyve activity on lysosomes. *Virulence* **15**, 2350893 (2024).
<https://doi.org/10.1080/21505594.2024.2350893>
- 48 Buckley, C. M. *et al.* PIKfyve/Fab1 is required for efficient V-ATPase and hydrolase delivery to phagosomes, phagosomal killing, and restriction of Legionella infection. *PLoS Pathog* **15**, e1007551 (2019). <https://doi.org/10.1371/journal.ppat.1007551>

- 49 Kerr, M. C. *et al.* Inhibition of the PtdIns(5) kinase PIKfyve disrupts intracellular replication of Salmonella. *EMBO J* **29**, 1331-1347 (2010). <https://doi.org/10.1038/emboj.2010.28>
- 50 Galeev, A. *et al.* Proteoglycan-Dependent Endo-Lysosomal Fusion Affects Intracellular Survival of Salmonella Typhimurium in Epithelial Cells. *Front Immunol* **11**, 731 (2020). <https://doi.org/10.3389/fimmu.2020.00731>
- 51 Wong, K. W. The Role of ESX-1 in Mycobacterium tuberculosis Pathogenesis. *Microbiol Spectr* **5** (2017). <https://doi.org/10.1128/microbiolspec.TBTB2-0001-2015>
- 52 Thurston, T. L., Wandel, M. P., von Muhlinen, N., Foeglein, A. & Randow, F. Galectin 8 targets damaged vesicles for autophagy to defend cells against bacterial invasion. *Nature* **482**, 414-418 (2012). <https://doi.org/10.1038/nature10744>
- 53 Chauhan, S. *et al.* TRIMs and Galectins Globally Cooperate and TRIM16 and Galectin-3 Co-direct Autophagy in Endomembrane Damage Homeostasis. *Developmental Cell* **39**, 13-27 (2016). <https://doi.org/10.1016/j.devcel.2016.08.003>
- 54 Franco, L. H. *et al.* The Ubiquitin Ligase Smurf1 Functions in Selective Autophagy of Mycobacterium tuberculosis and Anti-tuberculous Host Defense. *Cell Host & Microbe* **21**, 59-72 (2017). <https://doi.org/10.1016/j.chom.2016.11.002>
- 55 Manzanillo, P. S. *et al.* The ubiquitin ligase parkin mediates resistance to intracellular pathogens. *Nature* **501**, 512-516 (2013). <https://doi.org/10.1038/nature12566>
- 56 Khaminets, A., Beh, C. & Dikic, I. Ubiquitin-Dependent And Independent Signals In Selective Autophagy. *Trends in Cell Biology* **26**, 6-16 (2016). <https://doi.org/10.1016/j.tcb.2015.08.010>
- 57 Manzanillo, P. S., Shiloh, M. U., Portnoy, D. A. & Cox, J. S. Mycobacterium Tuberculosis Activates the DNA-Dependent Cytosolic Surveillance Pathway within Macrophages. *Cell Host & Microbe* **11**, 469-480 (2012). <https://doi.org/10.1016/j.chom.2012.03.007>
- 58 Watson, R. O., Manzanillo, P. S. & Cox, J. S. Extracellular M. tuberculosis DNA targets bacteria for autophagy by activating the host DNA-sensing pathway. *Cell* **150**, 803-815 (2012). <https://doi.org/10.1016/j.cell.2012.06.040>
- 59 Dey, B. *et al.* A bacterial cyclic dinucleotide activates the cytosolic surveillance pathway and mediates innate resistance to tuberculosis. *Nature Medicine* **21**, 401-406 (2015). <https://doi.org/10.1038/nm.3813>
- 60 Collins, A. C. *et al.* Cyclic GMP-AMP Synthase Is an Innate Immune DNA Sensor for. *Cell Host & Microbe* **17**, 820-828 (2015). <https://doi.org/10.1016/j.chom.2015.05.005>
- 61 Watson, R. O. *et al.* The Cytosolic Sensor cGAS Detects Mycobacterium tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. *Cell Host Microbe* **17**, 811-819 (2015). <https://doi.org/10.1016/j.chom.2015.05.004>
- 62 Wassermann, R. *et al.* Differentially Activates cGAS- and Inflammasome-Dependent Intracellular Immune Responses through ESX-1. *Cell Host & Microbe* **17**, 799-810 (2015). <https://doi.org/10.1016/j.chom.2015.05.003>
- 63 Majlessi, L. & Brosch, R. Meets the Cytosol: The Role of cGAS in Anti-mycobacterial Immunity. *Cell Host & Microbe* **17**, 733-735 (2015). <https://doi.org/10.1016/j.chom.2015.05.017>
- 64 Wiens, K. E. & Ernst, J. D. The Mechanism for Type I Interferon Induction by Mycobacterium tuberculosis is Bacterial Strain-Dependent. *PLoS Pathog* **12**, e1005809 (2016). <https://doi.org/10.1371/journal.ppat.1005809>
- 65 Klionsky, D. J. *et al.* Autophagy in major human diseases. *Embo J* **40**, e108863 (2021). <https://doi.org/10.15252/embj.2021108863>

- 66 Levine, B. & Kroemer, G. Biological Functions of Autophagy Genes: A Disease Perspective. *Cell* **176**, 11-42 (2019). <https://doi.org/10.1016/j.cell.2018.09.048>
- 67 Lamark, T. & Johansen, T. Mechanisms of Selective Autophagy. *Annual Review of Cell and Developmental Biology* **37**, 143-169 (2021). <https://doi.org/10.1146/annurev-cellbio-120219-035530>
- 68 Khaminets, A., Behl, C. & Dikic, I. Ubiquitin-Dependent And Independent Signals In Selective Autophagy. *Trends Cell Biol* **26**, 6-16 (2016). <https://doi.org/10.1016/j.tcb.2015.08.010>
- 69 Franco, L. H. *et al.* The Ubiquitin Ligase Smurf1 Functions in Selective Autophagy of Mycobacterium tuberculosis and Anti-tuberculous Host Defense. *Cell Host Microbe* **22**, 421-423 (2017). <https://doi.org/10.1016/j.chom.2017.08.005>
- 70 Seto, S., Tsujimura, K., Horii, T. & Koide, Y. Autophagy adaptor protein p62/SQSTM1 and autophagy-related gene Atg5 mediate autophagosome formation in response to Mycobacterium tuberculosis infection in dendritic cells. *PLoS One* **8**, e86017 (2013). <https://doi.org/10.1371/journal.pone.0086017>
- 71 Zhang, R. *et al.* The selective autophagy receptors Optineurin and p62 are both required for zebrafish host resistance to mycobacterial infection. *PLoS Pathog* **15**, e1007329 (2019). <https://doi.org/10.1371/journal.ppat.1007329>
- 72 Ramachandran, G., Yeruva, C. V., Swarup, G. & Raghunand, T. R. A cytoprotective role for optineurin during mycobacterial infection of macrophages. *Biochem Biophys Rep* **38**, 101672 (2024). <https://doi.org/10.1016/j.bbrep.2024.101672>
- 73 Nishimura, T. *et al.* Autophagosome formation is initiated at phosphatidylinositol synthase-enriched ER subdomains. *The EMBO Journal* **36**, 1719-1735 (2017). <https://doi.org/10.15252/embj.201695189>
- 74 Turco, E. *et al.* FIP200 Claw Domain Binding to p62 Promotes Autophagosome Formation at Ubiquitin Condensates. *Mol Cell* **74**, 330-346 e311 (2019). <https://doi.org/10.1016/j.molcel.2019.01.035>
- 75 Nishimura, T. & Tooze, S. A. Emerging roles of ATG proteins and membrane lipids in autophagosome formation. *Cell Discovery* **6** (2020). <https://doi.org/10.1038/s41421-020-0161-3>
- 76 Kriegenburg, F., Ungermann, C. & Reggiori, F. Coordination of Autophagosome-Lysosome Fusion by Atg8 Family Members. *Current Biology* **28**, R512-R518 (2018). <https://doi.org/10.1016/j.cub.2018.02.034>
- 77 Takahashi, Y. *et al.* An autophagy assay reveals the ESCRT-III component CHMP2A as a regulator of phagophore closure. *Nature Communications* **9** (2018). <https://doi.org/10.1038/s41467-018-05254-w>
- 78 Zhen, Y. *et al.* ESCRT-mediated phagophore sealing during mitophagy. *Autophagy* **16**, 826-841 (2020). <https://doi.org/10.1080/15548627.2019.1639301>
- 79 Zhou, F. *et al.* Rab5-dependent autophagosome closure by ESCRT. *J Cell Biol* **218**, 1908-1927 (2019). <https://doi.org/10.1083/jcb.201811173>
- 80 Zhou, F. *et al.* A Rab5 GTPase module is important for autophagosome closure. *Plos Genet* **13** (2017). <https://doi.org/10.1371/journal.pgen.1007020>
- 81 Wang, Z. *et al.* The Vici Syndrome Protein EPG5 Is a Rab7 Effector that Determines the Fusion Specificity of Autophagosomes with Late Endosomes/Lysosomes. *Molecular Cell* **63**, 781-795 (2016). <https://doi.org/10.1016/j.molcel.2016.08.021>

- 82 Yamamoto, H., Zhang, S. & Mizushima, N. Autophagy genes in biology and disease. *Nat Rev Genet* **24**, 382-400 (2023). <https://doi.org/10.1038/s41576-022-00562-w>
- 83 Kimmey, J. M. & Stallings, C. L. Bacterial Pathogens versus Autophagy: Implications for Therapeutic Interventions. *Trends in Molecular Medicine* **22**, 1060-1076 (2016). <https://doi.org/10.1016/j.molmed.2016.10.008>
- 84 Gutierrez, M. G. *et al.* Autophagy Is a Defense Mechanism Inhibiting BCG and Mycobacterium tuberculosis Survival in Infected Macrophages. *Cell* **119**, 753-766 (2004). <https://doi.org/10.1016/j.cell.2004.11.038>
- 85 Castillo, E. F. *et al.* Autophagy protects against active tuberculosis by suppressing bacterial burden and inflammation. *Proc Natl Acad Sci U S A* **109**, E3168-3176 (2012). <https://doi.org/10.1073/pnas.1210500109>
- 86 Bernard, E. M. *et al.* M. tuberculosis infection of human iPSDM reveals complex membrane dynamics during xenophagy evasion. *J Cell Sci* **134**, jcs252973 (2020). <https://doi.org/10.1242/jcs.252973>
- 87 Shui, W. *et al.* Organelle membrane proteomics reveals differential influence of mycobacterial lipoglycans on macrophage phagosome maturation and autophagosome accumulation. *J Proteome Res* **10**, 339-348 (2011). <https://doi.org/10.1021/pr100688h>
- 88 Seto, S., Tsujimura, K. & Koide, Y. Coronin-1a inhibits autophagosome formation around Mycobacterium tuberculosis-containing phagosomes and assists mycobacterial survival in macrophages. *Cellular Microbiology* **14**, 710-727 (2012). <https://doi.org/10.1111/j.1462-5822.2012.01754.x>
- 89 Ouimet, M. *et al.* Mycobacterium tuberculosis induces the miR-33 locus to reprogram autophagy and host lipid metabolism. *Nat Immunol* **17**, 677-686 (2016). <https://doi.org/10.1038/ni.3434>
- 90 Shin, D.-M. *et al.* Mycobacterium tuberculosis Eis Regulates Autophagy, Inflammation, and Cell Death through Redox-dependent Signaling. *PLoS Pathogens* **6**, e1001230 (2010). <https://doi.org/10.1371/journal.ppat.1001230>
- 91 Kim, K. H. *et al.* Mycobacterium tuberculosis Eis protein initiates suppression of host immune responses by acetylation of DUSP16/MKP-7. *Proc Natl Acad Sci U S A* **109**, 7729-7734 (2012). <https://doi.org/10.1073/pnas.1120251109>
- 92 Chandra, P. *et al.* Mycobacterium tuberculosis Inhibits RAB7 Recruitment to Selectively Modulate Autophagy Flux in Macrophages. *Sci Rep* **5**, 16320 (2015). <https://doi.org/10.1038/srep16320>
- 93 Vergne, I. *et al.* Mechanism of phagolysosome biogenesis block by viable *Mycobacterium tuberculosis*. *Proceedings of the National Academy of Sciences* **102**, 4033-4038 (2005). <https://doi.org/10.1073/pnas.0409716102>
- 94 Saleh, M. T. & Belisle, J. T. Secretion of an Acid Phosphatase (SapM) by Mycobacterium tuberculosis That Is Similar to Eukaryotic Acid Phosphatases. *Journal of Bacteriology* **182**, 6850-6853 (2000). <https://doi.org/10.1128/jb.182.23.6850-6853.2000>
- 95 Hu, D. *et al.* Autophagy regulation revealed by SapM-induced block of autophagosome-lysosome fusion via binding RAB7. *Biochem Bioph Res Co* **461**, 401-407 (2015). <https://doi.org/10.1016/j.bbrc.2015.04.051>
- 96 Augenreich, J. *et al.* Dynamic interplay of autophagy and membrane repair during Mycobacterium tuberculosis Infection. *PLOS Pathogens* **21**, e1012830 (2025). <https://doi.org/10.1371/journal.ppat.1012830>

- 97 Aylan, B. *et al.* ATG7 and ATG14 restrict cytosolic and phagosomal *Mycobacterium tuberculosis* replication in human macrophages. *Nature Microbiology* **8**, 803-818 (2023). <https://doi.org/10.1038/s41564-023-01335-9>
- 98 Golovkine, G. R. *et al.* Autophagy restricts *Mycobacterium tuberculosis* during acute infection in mice. *Nature Microbiology* **8**, 819-832 (2023). <https://doi.org/10.1038/s41564-023-01354-6>
- 99 Michiel *et al.* The DNA Damage-Regulated Autophagy Modulator DRAM1 Links *Mycobacterial* Recognition via TLR-MYD88 to Autophagic Defense. *Cell Host & Microbe* **15**, 753-767 (2014). <https://doi.org/10.1016/j.chom.2014.05.005>
- 100 Kiliç, G., Saris, A., Ottenhoff, T. H. M. & Haks, M. C. Host-directed therapy to combat mycobacterial infections*. *Immunological Reviews* **301**, 62-83 (2021). <https://doi.org/10.1111/imr.12951>
- 101 Heckmann, B. L. & Green, D. R. LC3-associated phagocytosis at a glance. *J Cell Sci* **132** (2019). <https://doi.org/10.1242/jcs.222984>
- 102 Mehta, P., Henault, J., Kolbeck, R. & Sanjuan, M. A. Noncanonical autophagy: one small step for LC3, one giant leap for immunity. *Current Opinion in Immunology* **26**, 69-75 (2014). <https://doi.org/10.1016/j.coi.2013.10.012>
- 103 Nieto-Torres, J. L., Leidal, A. M., Debnath, J. & Hansen, M. Beyond Autophagy: The Expanding Roles of ATG8 Proteins. *Trends Biochem Sci* **46**, 673-686 (2021). <https://doi.org/10.1016/j.tibs.2021.01.004>
- 104 Galluzzi, L. *et al.* Molecular definitions of autophagy and related processes. *Embo J* **36**, 1811-1836 (2017). <https://doi.org/10.15252/embj.201796697>
- 105 Pena-Martinez, C., Rickman, A. D. & Heckmann, B. L. Beyond autophagy: LC3-associated phagocytosis and endocytosis. *Sci Adv* **8**, eabn1702 (2022). <https://doi.org/10.1126/sciadv.abn1702>
- 106 Huang, J. *et al.* Activation of antibacterial autophagy by NADPH oxidases. *Proceedings of the National Academy of Sciences* **106**, 6226-6231 (2009). <https://doi.org/10.1073/pnas.0811045106>
- 107 Sanjuan, M. A. *et al.* Toll-like receptor signalling in macrophages links the autophagy pathway to phagocytosis. *Nature* **450**, 1253-1257 (2007). <https://doi.org/10.1038/nature06421>
- 108 Tam, J. M. *et al.* Dectin-1-Dependent LC3 Recruitment to Phagosomes Enhances Fungicidal Activity in Macrophages. *The Journal of Infectious Diseases* **210**, 1844-1854 (2014). <https://doi.org/10.1093/infdis/jiu290>
- 109 Miyanishi, M. *et al.* Identification of Tim4 as a phosphatidylserine receptor. *Nature* **450**, 435-439 (2007). <https://doi.org/10.1038/nature06307>
- 110 Heckmann, B. L., Boada-Romero, E., Cunha, L. D., Magne, J. & Green, D. R. LC3-Associated Phagocytosis and Inflammation. *J Mol Biol* **429**, 3561-3576 (2017). <https://doi.org/10.1016/j.jmb.2017.08.012>
- 111 Vieira, O. V. *et al.* Distinct roles of class I and class III phosphatidylinositol 3-kinases in phagosome formation and maturation. *J Cell Biol* **155**, 19-25 (2001). <https://doi.org/10.1083/jcb.200107069>
- 112 Lam, G. Y., Huang, J. & Brumell, J. H. The many roles of NOX2 NADPH oxidase-derived ROS in immunity. *Semin Immunopathol* **32**, 415-430 (2010). <https://doi.org/10.1007/s00281-010-0221-0>

- 113 Bagaitkar, J. *et al.* PI(3)P-p40^{phox} binding regulates NADPH oxidase activation in mouse macrophages and magnitude of inflammatory responses in vivo. *Journal of Leukocyte Biology* **101**, 449-457 (2017). <https://doi.org/10.1189/jlb.3ab0316-139r>
- 114 Hooper, K. M. *et al.* V-ATPase is a universal regulator of LC3-associated phagocytosis and non-canonical autophagy. *Journal of Cell Biology* **221** (2022). <https://doi.org/10.1083/jcb.202105112>
- 115 Gammoh, N. The multifaceted functions of ATG16L1 in autophagy and related processes. *J Cell Sci* **133**, jcs249227 (2020). <https://doi.org/10.1242/jcs.249227>
- 116 Lei, Y. & Klionsky, D. J. The coordination of V-ATPase and ATG16L1 is part of a common mechanism of non-canonical autophagy. *Autophagy* **18**, 2267-2269 (2022). <https://doi.org/10.1080/15548627.2022.2100678>
- 117 Schille, S. *et al.* LC3-associated phagocytosis in microbial pathogenesis. *Int J Med Microbiol* **308**, 228-236 (2018). <https://doi.org/10.1016/j.ijmm.2017.10.014>
- 118 Li, D., Wang, L., Zhao, Z., Bai, C. & Li, X. Autophagy and LC3-associated phagocytosis contribute negatively to the killing capability of THP-1-derived macrophages against *Candida albicans* at the mid-stage. *Immunol Lett* **263**, 25-32 (2023). <https://doi.org/10.1016/j.imlet.2023.09.006>
- 119 Kymrzi, I. *et al.* Calcium sequestration by fungal melanin inhibits calcium-calmodulin signalling to prevent LC3-associated phagocytosis. *Nat Microbiol* **3**, 791-803 (2018). <https://doi.org/10.1038/s41564-018-0167-x>
- 120 Luan, J. *et al.* The Role of LC3-Associated Phagocytosis Inhibits the Inflammatory Response in *Aspergillus fumigatus* Keratitis. *Invest Ophthalmol Vis Sci* **65**, 4 (2024). <https://doi.org/10.1167/iovs.65.8.4>
- 121 Oikonomou, V. *et al.* Noncanonical Fungal Autophagy Inhibits Inflammation in Response to IFN- γ via DAPK1. *Cell Host & Microbe* **20**, 744-757 (2016). <https://doi.org/10.1016/j.chom.2016.10.012>
- 122 Chamilos, G. *et al.* Melanin targets LC3-associated phagocytosis (LAP): A novel pathogenetic mechanism in fungal disease. *Autophagy* **12**, 888-889 (2016). <https://doi.org/10.1080/15548627.2016.1157242>
- 123 Kymrzi, I. *et al.* Calcium sequestration by fungal melanin inhibits calcium-calmodulin signalling to prevent LC3-associated phagocytosis. *Nature Microbiology* **3**, 791-803 (2018). <https://doi.org/10.1038/s41564-018-0167-x>
- 124 Oikonomou, V. *et al.* Autophagy and LAP in the Fight against Fungal Infections: Regulation and Therapeutics. *Mediators of Inflammation* **2018**, 1-7 (2018). <https://doi.org/10.1155/2018/6195958>
- 125 Hubber, A. *et al.* Bacterial secretion system skews the fate of *Legionella*-containing vacuoles towards LC3-associated phagocytosis. *Scientific Reports* **7**, 44795 (2017). <https://doi.org/10.1038/srep44795>
- 126 Gluschko, A. *et al.* The beta(2) Integrin Mac-1 Induces Protective LC3-Associated Phagocytosis of *Listeria monocytogenes*. *Cell Host Microbe* **23**, 324-337 e325 (2018). <https://doi.org/10.1016/j.chom.2018.01.018>
- 127 Li, T. *et al.* *Listeria monocytogenes* upregulates mitochondrial calcium signalling to inhibit LC3-associated phagocytosis as a survival strategy. *Nat Microbiol* **6**, 366-379 (2021). <https://doi.org/10.1038/s41564-020-00843-2>

- 128 Gong, L. *et al.* The Burkholderia pseudomallei Type III Secretion System and BopA Are Required for Evasion of LC3-Associated Phagocytosis. *PLoS ONE* **6**, e17852 (2011). <https://doi.org/10.1371/journal.pone.0017852>
- 129 Krakauer, T. Living dangerously: Burkholderia pseudomallei modulates phagocyte cell death to survive. *Med Hypotheses* **121**, 64-69 (2018). <https://doi.org/10.1016/j.mehy.2018.09.028>
- 130 Campbell-Valois, F.-X., Sachse, M., Sansonetti, P. J. & Parsot, C. Escape of Actively Secreting Shigella flexneri from ATG8/LC3-Positive Vacuoles Formed during Cell-To-Cell Spread Is Facilitated by IcsB and VirA. *mBio* **6**, e02567-02514 (2015). <https://doi.org/10.1128/mbio.02567-14>
- 131 Krokowski, S. & Mostowy, S. Interactions between Shigella flexneri and the Autophagy Machinery. *Frontiers in Cellular and Infection Microbiology* **6** (2016). <https://doi.org/10.3389/fcimb.2016.00017>
- 132 Masud, S. *et al.* Macrophages target Salmonella by Lc3-associated phagocytosis in a systemic infection model. *Autophagy* **15**, 796-812 (2019). <https://doi.org/10.1080/15548627.2019.1569297>
- 133 Masud, S., van der Burg, L., Storm, L., Prajsnar, T. K. & Meijer, A. H. Rubicon-Dependent Lc3 Recruitment to Salmonella-Containing Phagosomes Is a Host Defense Mechanism Triggered Independently From Major Bacterial Virulence Factors. *Front Cell Infect Microbiol* **9**, 279 (2019). <https://doi.org/10.3389/fcimb.2019.00279>
- 134 Yuan, J., Zhang, Q., Chen, S., Yan, M. & Yue, L. LC3-Associated Phagocytosis in Bacterial Infection. *Pathogens* **11** (2022). <https://doi.org/10.3390/pathogens11080863>
- 135 Lerena, M. C. & Colombo, M. I. Mycobacterium marinum induces a marked LC3 recruitment to its containing phagosome that depends on a functional ESX-1 secretion system. *Cellular Microbiology* **13**, 814-835 (2011). <https://doi.org/10.1111/j.1462-5822.2011.01581.x>
- 136 Koster, S. *et al.* Mycobacterium tuberculosis is protected from NADPH oxidase and LC3-associated phagocytosis by the LCP protein CpsA. *Proc Natl Acad Sci U S A* **114**, E8711-E8720 (2017). <https://doi.org/10.1073/pnas.1707792114>
- 137 Matsunaga, K. *et al.* Two Beclin 1-binding proteins, Atg14L and Rubicon, reciprocally regulate autophagy at different stages. *Nature Cell Biology* **11**, 385-U369 (2009). <https://doi.org/10.1038/ncb1846>
- 138 Zhong, Y. *et al.* Distinct regulation of autophagic activity by Atg14L and Rubicon associated with Beclin 1-phosphatidylinositol-3-kinase complex. *Nat Cell Biol* **11**, 468-476 (2009). <https://doi.org/10.1038/ncb1854>
- 139 Kim, Y.-M. *et al.* mTORC1 Phosphorylates UVRAG to Negatively Regulate Autophagosome and Endosome Maturation. *Molecular Cell* **57**, 207-218 (2015). <https://doi.org/10.1016/j.molcel.2014.11.013>
- 140 Yang, C. S. *et al.* Autophagy protein Rubicon mediates phagocytic NADPH oxidase activation in response to microbial infection or TLR stimulation. *Cell Host Microbe* **11**, 264-276 (2012). <https://doi.org/10.1016/j.chom.2012.01.018>
- 141 Sun, Q. M. *et al.* The RUN Domain of Rubicon Is Important for hVps34 Binding, Lipid Kinase Inhibition, and Autophagy Suppression. *J Biol Chem* **286**, 185-191 (2011). <https://doi.org/10.1074/jbc.M110.126425>

- 142 Tabata, K. *et al.* Rubicon and PLEKHM1 Negatively Regulate the Endocytic/Autophagic Pathway via a Novel Rab7-binding Domain. *Molecular Biology of the Cell* **21**, 4162-4172 (2010). <https://doi.org/10.1091/mbc.e10-06-0495>
- 143 Wandinger-Ness, A. & Zerial, M. Rab proteins and the compartmentalization of the endosomal system. *Cold Spring Harb Perspect Biol* **6**, a022616 (2014). <https://doi.org/10.1101/cshperspect.a022616>
- 144 Sun, Q., Westphal, W., Wong, K. N., Tan, I. & Zhong, Q. Rubicon controls endosome maturation as a Rab7 effector. *Proc Natl Acad Sci U S A* **107**, 19338-19343 (2010). <https://doi.org/10.1073/pnas.1010554107>
- 145 Härtlova, A. *et al.* LRRK2 is a negative regulator of phagosome maturation in macrophages. *Embo J* **37** (2018). <https://doi.org/https://doi.org/10.15252/embj.201798694>
- 146 Singh, A. K. & Gupta, U. D. Animal models of tuberculosis: Lesson learnt. *Indian J Med Res* **147**, 456-463 (2018). https://doi.org/10.4103/ijmr.IJMR_554_18
- 147 Cardona, P. J. & Williams, A. Experimental animal modelling for TB vaccine development. *Int J Infect Dis* **56**, 268-273 (2017). <https://doi.org/10.1016/j.ijid.2017.01.030>
- 148 Gupta, A. *et al.* Preparation and Preclinical Evaluation of Inhalable Particles Containing Rapamycin and Anti-Tuberculosis Agents for Induction of Autophagy. *Pharm Res* **33**, 1899-1912 (2016). <https://doi.org/10.1007/s11095-016-1926-0>
- 149 Gomes, M. C. & Mostowy, S. The Case for Modeling Human Infection in Zebrafish. *Trends in Microbiology* **28**, 10-18 (2020). <https://doi.org/10.1016/j.tim.2019.08.005>
- 150 Li, H. & Li, H. Animal Models of Tuberculosis. 139-170 (2023). https://doi.org/10.1007/978-3-031-24355-4_7
- 151 Renshaw, S. A. & Trede, N. S. A model 450 million years in the making: zebrafish and vertebrate immunity. *Disease Models & Mechanisms* **5**, 38-47 (2012). <https://doi.org/10.1242/dmm.007138>
- 152 Howe, K. *et al.* The zebrafish reference genome sequence and its relationship to the human genome. *Nature* **496**, 498-503 (2013). <https://doi.org/10.1038/nature12111>
- 153 Stinear, T. P. *et al.* Insights from the complete genome sequence of *Mycobacterium marinum* on the evolution of *Mycobacterium tuberculosis*. *Genome Res* **18**, 729-741 (2008). <https://doi.org/10.1101/gr.075069.107>
- 154 Davis, J. M. *et al.* Real-Time Visualization of *Mycobacterium*-Macrophage Interactions Leading to Initiation of Granuloma Formation in Zebrafish Embryos. *Immunity* **17**, 693-702 (2002). [https://doi.org/10.1016/s1074-7613\(02\)00475-2](https://doi.org/10.1016/s1074-7613(02)00475-2)
- 155 Swaim, L. E. *et al.* *Mycobacterium marinum* Infection of Adult Zebrafish Causes Caseating Granulomatous Tuberculosis and Is Moderated by Adaptive Immunity. *Infection and Immunity* **74**, 6108-6117 (2006). <https://doi.org/10.1128/iai.00887-06>
- 156 Benard, E. L. *et al.* Infection of zebrafish embryos with intracellular bacterial pathogens. *J Vis Exp* (2012). <https://doi.org/10.3791/3781>
- 157 Mark *et al.* Macrophage Epithelial Reprogramming Underlies *Mycobacterial* Granuloma Formation and Promotes Infection. *Immunity* **45**, 861-876 (2016). <https://doi.org/10.1016/j.immuni.2016.09.014>
- 158 Ramakrishnan, L. Looking Within the Zebrafish to Understand the Tuberculous Granuloma. *Adv Exp Med Biol* **783**, 251-266 (2013). https://doi.org/10.1007/978-1-4614-6111-1_13

- 159 Torraca, V., Tulotta, C., Snaar-Jagalska, B. E. & Meijer, A. H. The chemokine receptor CXCR4 promotes granuloma formation by sustaining a mycobacteria-induced angiogenesis programme. *Scientific Reports* **7**, 45061 (2017). <https://doi.org/10.1038/srep45061>
- 160 Oehlers, S. H. *et al.* Interception of host angiogenic signalling limits mycobacterial growth. *Nature* **517**, 612-615 (2015). <https://doi.org/10.1038/nature13967>
- 161 Cambier, C. J., O'Leary, S. M., O'Sullivan, M. P., Keane, J. & Ramakrishnan, L. Phenolic Glycolipid Facilitates Mycobacterial Escape from Microbicidal Tissue-Resident Macrophages. *Immunity* **47**, 552-565.e554 (2017). <https://doi.org/10.1016/j.immuni.2017.08.003>
- 162 Sommer, F., Torraca, V., Kamel, S. M., Lombardi, A. & Meijer, A. H. Frontline Science: Antagonism between regular and atypical Cxcr3 receptors regulates macrophage migration during infection and injury in zebrafish. *Journal of Leukocyte Biology* **107**, 185-203 (2020). <https://doi.org/10.1002/jlb.2hi0119-006r>
- 163 Torraca, V., Otto, N. A., Tavakoli-Tameh, A. & Meijer, A. H. The inflammatory chemokine Cxcl18b exerts neutrophil-specific chemotaxis via the promiscuous chemokine receptor Cxcr2 in zebrafish. *Dev Comp Immunol* **67**, 57-65 (2017). <https://doi.org/10.1016/j.dci.2016.10.014>
- 164 Varela, M., Van Der Vaart, M., Groenewoud, A. & Meijer, A. H. Extracellular mycobacterial DNA drives disease progression by triggering Caspase-11-dependent pyroptosis of infected macrophages. *bioRxiv* (2019). <https://doi.org/10.1101/514125>
- 165 Hosseini, R. *et al.* Correlative light and electron microscopy imaging of autophagy in a zebrafish infection model. *Autophagy* **10**, 1844-1857 (2014). <https://doi.org/10.4161/auto.29992>
- 166 van der Vaart, M. *et al.* The DNA damage-regulated autophagy modulator DRAM1 links mycobacterial recognition via TLR-MYD88 to autophagic defense. *Cell Host Microbe* **15**, 753-767 (2014). <https://doi.org/10.1016/j.chom.2014.05.005>
- 167 Zhang, R. *et al.* Deficiency in the autophagy modulator Dram1 exacerbates pyroptotic cell death of Mycobacteria-infected macrophages. *Cell Death & Disease* **11** (2020). <https://doi.org/10.1038/s41419-020-2477-1>
- 168 Stoop, E. J. M. *et al.* Zebrafish embryo screen for mycobacterial genes involved in the initiation of granuloma formation reveals a newly identified ESX-1 component. *Disease Models & Mechanisms* **4**, 526-536 (2011). <https://doi.org/10.1242/dmm.006676>
- 169 Chirakos, A. E., Balaram, A., Conrad, W. & Champion, P. A. Modeling Tubercular ESX-1 Secretion Using Mycobacterium marinum. *Microbiology and Molecular Biology Reviews* **84** (2020). <https://doi.org/10.1128/mmb.00082-19>
- 170 Lopez, A., Fleming, A. & Rubinsztein, D. C. Seeing is believing: methods to monitor vertebrate autophagy *in vivo*. *Open Biology* **8**, 180106 (2018). <https://doi.org/10.1098/rsob.180106>
- 171 Mathai, B., Meijer, A. & Simonsen, A. Studying Autophagy in Zebrafish. *Cells* **6**, 21 (2017). <https://doi.org/10.3390/cells6030021>