



Universiteit  
Leiden  
The Netherlands

## **Balancing the dose: a meta-analysis of preoperative radiotherapy strategies for soft tissue sarcomas**

Foppele, G.F.; Fiocco, M.; Riet, B.H.G. van; Haas, R.L.M.; Wiltink, L.M.

### **Citation**

Foppele, G. F., Fiocco, M., Riet, B. H. G. van, Haas, R. L. M., & Wiltink, L. M. (2025). Balancing the dose: a meta-analysis of preoperative radiotherapy strategies for soft tissue sarcomas. *Radiotherapy & Oncology*, 210. doi:10.1016/j.radonc.2025.111034

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/4288879>

**Note:** To cite this publication please use the final published version (if applicable).



## Balancing the dose: A meta-analysis of preoperative radiotherapy strategies for soft tissue sarcomas

G.F. Foppele<sup>a,b,\*</sup>, M. Fiocco<sup>c,d,e</sup>, B.H.G. van Riet<sup>a</sup>, R.L.M. Haas<sup>a,b,1</sup>, L.M. Wiltink<sup>a,b,1</sup>

<sup>a</sup> Department of Radiotherapy, Netherlands Cancer Institute, the Netherlands

<sup>b</sup> Department of Radiotherapy, Leiden University Medical Centre, the Netherlands

<sup>c</sup> Mathematical Institute, University of Leiden, the Netherlands

<sup>d</sup> Department of Biomedical Data Sciences, Leiden University Medical Centre, the Netherlands

<sup>e</sup> Princess Máxima Centre for Paediatric Oncology, Utrecht, the Netherlands

### ARTICLE INFO

#### Keywords:

Soft Tissue Sarcoma (STS)  
Preoperative radiotherapy  
Hypofractionation  
Conventional Radiotherapy  
Wound complications  
Radiation Toxicity

### ABSTRACT

**Purpose:** Soft tissue sarcomas (STS) are a rare and diverse group of tumors. Management of STS typically involves a multidisciplinary approach, including surgery, radiotherapy (RT), and chemotherapy, with surgery remaining the cornerstone of treatment. Radiotherapy, both preoperative and postoperative, has been shown to improve local control. Hypofractionation has shown promising results in other malignancies and may offer similar benefits in STS. This study systematically reviews the efficacy and safety of different preoperative RT fractionation schedules, focusing on major wound complications (MWC) and late toxicity.

**Methods:** A systematic review and meta-analysis were conducted following PRISMA guidelines. PubMed, Cochrane Library, and Embase were searched for studies on preoperative RT in STS. Included studies were categorized into conventional RT, moderate hypofractionation, and ultra-hypofractionation. Meta-analysis was performed on MWC and late toxicity ( $\geq$  grade 3) using a random-effects model. Statistical heterogeneity was assessed using I-squared and Tau statistics.

**Results:** Thirty studies with 2288 patients were included. The pooled overall rate of MWC was 18 % (95 % CI: 10–27 %) for conventional RT, and 29 % (95 % CI: 24–34 %) for moderate hypofractionation. The pooled overall rate of late toxicity was 5 % (95 % CI: 0–12 %) and 4 % (95 % CI: 2–6 %), respectively. Based on descriptive data, the MWC rates of the ultra-hypofractionated group did not exceed 41 %, with one study reporting a rate of 41 % and the remaining studies reporting rates no higher than 32 %. For toxicity, a wide range of toxicity rates (1–16 %) was reported.

**Conclusion:** Moderate hypofractionation appears to maintain oncological outcomes while balancing toxicity and wound complications. Reported MWC rates of ultra-hypofractionation regimens, based on descriptive data from heterogeneous individual studies remained below 41 %. However, these outcomes are not derived from pooled analysis. This approach should not yet be used in clinical practice, and randomized controlled trials are needed to establish its efficacy and safety in STS patients.

### Introduction

Soft tissue sarcomas (STS) are a rare and heterogeneous group of tumors. Their treatment is influenced by factors such as tumor histology, grade, location, size, and the patient's overall performance status. The management of STS typically involves a multidisciplinary approach, including surgery, radiotherapy (RT), and chemotherapy, with surgery

remaining the cornerstone of treatment. Radiotherapy, both preoperative and postoperative, has been shown to improve local control [1]. However, while radiotherapy improves local control, its addition is associated with an increased risk of toxicity.

Preoperative radiotherapy is associated with an increased risk of short-term complications, particularly wound complications after surgery. However, these short-term complications are generally transient

\* Corresponding author at: Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands.

E-mail address: [f.foppele@nki.nl](mailto:f.foppele@nki.nl) (G.F. Foppele).

<sup>1</sup> R.L.M. Haas and L.M. Wiltink: Contributed equally as last authors.

<https://doi.org/10.1016/j.radonc.2025.111034>

Received 12 May 2025; Received in revised form 24 June 2025; Accepted 3 July 2025

Available online 4 July 2025

0167-8140/© 2025 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

and manageable with appropriate care. In contrast, postoperative radiotherapy carries a lower immediate risk of wound complications but is associated with an increased risk of long-term toxicities such as fibrosis, edema, and joint stiffness [1–3]. To balance these risks, the recommended treatment regimen is preoperative radiotherapy followed by surgery after a waiting period of 6–8 weeks. Current guidelines recommend a preoperative dose of 50 Gray (Gy), delivered in 25 fractions [4–8].

In recent years, there has been growing interest in hypofractionation, which involves delivering larger radiation doses in fewer fractions, thereby reducing the overall treatment duration. The primary aim of hypofractionated radiotherapy is to improve patient outcomes by maintaining or even enhancing treatment efficacy, while reducing treatment burden. Clinical studies in other malignancies, including breast, rectal, and prostate cancer, have already demonstrated hypofractionation, to reduce overall treatment time without compromising local control or long-term toxicity [9–11]. These findings suggest that hypofractionation could also be safe in terms of long-term toxicity in the treatment of STS.

The rationale for exploring hypofractionation in STS is based on the fact that tumors with a low  $\alpha/\beta$  ratio, commonly seen in STS, may respond better to higher radiation doses per fraction [12]. Shortening the treatment duration has the potential to reduce the overall treatment burden on patients. Due to the rarity of STS, conducting large-scale non-inferiority, phase III randomized clinical trials (RCTs) remains challenging. Nevertheless, several phase 2 studies have demonstrated that hypofractionated treatment schedules can shorten treatment time without compromising local control or increasing wound complications [13,14].

This review aims to assess the current evidence for the use of hypofractionation in STS, by critically evaluating and comparing the clinical outcomes of different fractionation strategies, comparing conventional fractionation, moderate hypofractionation, and ultra-hypofractionation. The meta-analysis is specifically focused on late toxicity (grade 3 or higher) and wound complications as key outcomes.

## Methods

This research was performed according to 2020 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [15]. The databases PubMed (NCBI), Cochrane Library (Wiley), Embase (Ovid) and Web of Science (Clarivate) were searched on January 1st, 2025. The search terms included various combinations of relevant keywords such as ‘Radiotherapy’, ‘Soft tissue sarcomas’, ‘preoperative’ and ‘dose’. Only studies published in English were included. The full search strategy is outlined in the Supplementary (S1). The first and second reviewer screened articles together, excluded duplicates, studies on hyperthermia, postoperative RT, or trials with concurrent chemotherapy and immunotherapy. Studies with pediatric (<18 years) populations, studies including only myxoid liposarcoma, studies assessing metastatic disease and reviews were also excluded. For an overview of the in- and exclusion criteria see Table 1. In case of overlap of the study population, the paper with a larger study population or more extensive data of interest was chosen. Authors of studies included in the review who had published more than two papers were also searched to find additional studies not found through other sources. Studies were categorized into three groups based on the type of radiotherapy delivered: conventional radiotherapy, moderate hypofractionation, and ultra-hypofractionation. This classification was determined based on the number of fractions and dose per fraction, following the predefined criteria outlined in the study protocol.

The first reviewer (G.F.) examined all full texts thoroughly to maintain sufficient quality standards. The risk of bias for each study was assessed using the Risk of Bias In Non-randomized Studies of Interventions tool (ROBINS-I) developed by the Cochrane Bias Methods Group. For randomized studies, the Rob Tool was used [16,17]. After the initial evaluation by the first reviewer, the second reviewer (B.v.R.)

**Table 1**

In- and exclusion criteria of the systematic review.

Category	Inclusion criteria	Exclusion criteria
Study design	Any accept reviews Presented studies	Systematic reviews Narrative reviews Non-English
Population	Age $\geq$ 18 years Sex: any Race: any Disease: primary soft tissue sarcomas located at the extremities or trunk Histological grade: any except myxoid liposarcoma Stage: localized	Metastatic stage Non-human Myxoid liposarcoma studies only
Intervention	Conventional RT (1.8–2.2 Gy/fraction) Hypofractionated RT (>2.2 Gy/fraction) Surgical resection	Hyperthermia Postoperative RT (including boost)
Concurrent therapy		Concurrent chemotherapy Concurrent immunotherapy
Outcomes	Late toxicities including wound complications	
Date range	Until January 1st, 2025	

critically reviewed the bias assessment, list of results and data. Disagreements were resolved in team discussions. Find Supplementary (S1–2) for the critical review of the risk of bias.

Meta-analysis was performed on major wound complications (MWC) and late toxicity as outcomes for each study included in the analysis. MWC were defined by the Canadian SR-2 trial [1]. The definition of MWC included surgery for wound repair, invasive procedures without anesthesia (e.g., seroma aspiration), hospital readmission for IV antibiotics, or ongoing deep packing within 120 days of surgery. If MWC were not defined by the Canadian SR-2 trial by the author and in the published paper, definition was changed to the Canadian SR-2 trial if possible. Late toxicity of interest was grade 3 or higher, scored by the CTCAE version 5.0 [18]. Toxicity of grade 3 or higher was classified as significant medical symptoms requiring medical intervention, hospitalization, or more severe consequences, including life-threatening conditions (grade 4) or fatal outcomes (grade 5). In case of missing data, additional information was requested from the authors. If no response or no longer available data, missing data was reported. When the outcomes MWC and toxicity could not be defined according to the Canadian SR2 trial definition or the CTCAE version 5.0, or when the data was not available, the study was not included in the meta-analysis.

Studies for conventional radiotherapy included in the meta-analysis were structured based on the Equivalent Dose in 2 Gy fractions (EQD2) of normal tissue taking an  $\alpha/\beta$  ratio of 3 Gy [19]. After initial selection of data items by the first reviewer, the second reviewer checked for suitability and accuracy. Wherever multiple doses and fractions were used within the same study, the most used dose and fraction was applied as outcome for the meta-analysis.

A random effects model was employed to pool study-specific proportion of wound complication and toxicity to estimate overall proportions and their associated confidence intervals. Inverse variance method which gives more weight to larger study was used to pool outcomes for different studies. The overall effect corresponding to a fixed and random effects model are reported together in the same forest plots along with their confidence intervals. The sizes of the square boxes on the forest plot are proportional to the total number of patients in the selected trials. An overall test on heterogeneity between studies was performed for each separate meta-analysis (value I-squared in figures). To estimate the between-study variance which is represented as ‘Tau’ in the forest plots, DerSimonian-Laird’s method has been employed [20]. The analysis was performed in R software environment with the library meta [21].

## Results

### Characteristics of included studies

The PRISMA flow chart (Fig. 1) illustrates the complete progress of the search and selection. In total, we included 30 studies with a total of 2288 patients. The majority of studies ( $n = 26$ ) were included through the search strategy and database screening. Additionally, three studies were identified through snowballing, and one paper has not been published yet [22]. Of those studies, thirteen conventional radiotherapy studies were included, four studies were included with a moderate hypofractionated schedule and thirteen studies with an ultra-hypofractionated schedule [1,13,22–49].

Data extraction included essential information from all eligible studies and resulted in extensive overview tables (Supplementary S4–5, Tables 2 and 3). Tables includes information such as author, publication year, type of trial, number of patients, dose per fraction, age of the patients, median follow-up time, oncological outcomes, wound complications and late toxicity grade 3 or higher. After conducting the risk of bias assessment, one article was excluded due to a critical risk of bias. This study was not included in the meta-analysis (Supplementary S3). Meta-analysis on MWC and late toxicity was not performed for the ultra-hypofractionated group, since this group included a wide variety of radiotherapy regimens and follow-up time.

In Tables S4 and S5, different rates of survival outcomes are reported. Conventional radiotherapy showed local control rates from at least 93 %. Recurrence rates reported in the moderate hypofractionated studies showed a cumulative incidence of local recurrence of 7.6 % at 2 years minimum. In ultra-hypofractionation, the  $5 \times 5$  Gy regimen demonstrated lower local control, with a 5-year LRFS rate of 81 %. In contrast, the  $5 \times 7$  Gy and  $5 \times 8$  Gy regimens achieved a 0 % local recurrence rate at 2 years.

In total, 12 studies were included in the meta-analysis of MWC for

**Table 2**

Overview of toxicity outcomes for conventional radiotherapy, classified by CTCAE v5.0 and Canadian SR-2 trial.

Author	EQD2 (Gy)	Schedule	Median follow-up (range)	Toxicity $\geq$ grade 3 (%)	Major wound complication (%)
Conventional					
O' Sullivan et al.	50.0	$25 \times 2$	3.3 yr (0.27–5.6)	Not available	35
Canter et al.	50.0	$25 \times 2$	19 mo (4–113)	Not available	28
Shah et al.	50.0	$25 \times 2$	40 mo	Not available	23
Wang et al.	48.4	$28 \times 1.8$	6.0 yr	14.7 %	25
Studer et al.	50.0	$25 \times 2$	33 mo (4–106)	Not available	7
Abdallah et al.	50.0	$25 \times 2$	7.3 mo (6.0–9.0)	Not available	10
Bonvalot et al.	50.0	$25 \times 2$	> 24 mo	Not available	9
Green et al.	50.0	$25 \times 2$	27 mo (19 to 59)	Not available	11
Huang et al.	50.0	$25 \times 2$	14.3 mo (11.9–24.7)	0 %	16
Hui et al.	48.4	$28 \times 1.8$	4.1 yr (0.6–6.9)	3 %	Not available
Seddon et al.	50.0	$25 \times 2$	35.2 mo (32.9–36.6)	Not available	Not available
Shelby et al.	50.0	$25 \times 2$	Unknown	Not available	Not available
J. González-Viguera et al.	50.0	$25 \times 2$	24 mo (0–156)	0 %	Not available

Abbreviations: EQD2, equivalent dose in 2 Gy per fraction with an  $\alpha/\beta$  ratio of 3 Gy; Gy, Gray; yr, years; mo, months.




Fig. 1. PRISMA Flowchart depicting the selection process of articles included in the meta-analysis.

**Table 3**  
Overview of outcomes for moderate- and ultra-hypofractionated radiotherapy, classified by CTCAE v5.0 and Canadian SR-2 trial.

Author	EQD2 (Gy)	Schedule	Median follow-up (range)	Toxicity ≥ grade 3 (%)	Major wound complication (%)
<b>Moderate</b>					
Bishop et al.	50.0	15 × 2.85	43 mo (37–52)	4 %	31
Ahmed et al.	50.0	15 × 2.85	17 mo (3.1–37.5)	<i>Not available</i>	23.1
Montero et al.	68.3	15 × 3.5	14 mo (1–40)	5.5 %	39
Foppele et al.	54.0	14–15 × 3	29 mo (2–50)	3 %	33
<b>Ultra-hypofractionated</b>					
Kilic et al.	36.4	8 × 3.5	37 mo (11–66)	<i>Not available</i>	<i>Not available</i>
Kosela et al. (2021)	40.0	5 × 5	57 mo (95 % CI 55.1–61.0)	1.3 %	7
Kosela et al. (2014)	40.0	5 × 5	35 mo (6–94)	2.9 %	<i>Not available</i>
Heesen et al.	40.0	5 × 5	2.2 yr	<i>Not available</i>	13
Potkrajcic et al.	40.0	5 × 5	5.1 ± 1.6 mo	<i>Not available</i>	29.4
Mayo et al.	54.0	5 × 6	24.5 mo (17.0–35.7)	4.5 %	41
Parasai et al.	54.0	5 × 6	10.7 mo (1.7–33.2)	0 %	19
Kalbasi et al.	54.0	5 × 6	29 mo	0 %	32
Bedi et al.	70.0	5 × 7	36.4 mo (3–56)	13 %	25
Novikov et al.	70.0	5 × 7	Average 21.5 mo (13–30)	14 %	14
Kubicek et al. (2021)	88.0	5 × 8	1719 d (983–2327)	6.7 %	20
Kubicek et al. (2018)	88.0	5 × 8	279 d (54–506)	0 %	29
Leite et al.	88.0	5 × 8	20.7 mo (6.9–55.7)	16 %	28

Abbreviations: EQD2, equivalent dose in 2 Gy per fraction with an  $\alpha/\beta$  ratio of 3 Gy; Gy, Gray; yr, years; mo, months; d, days.

conventional radiotherapy and moderate hypofractionated radiotherapy. Six studies were excluded due to missing data or other definitions of wound complications, one study was excluded due to a critical risk of bias.

In the meta-analysis for conventional radiotherapy, eight studies were included with a total of 413 patients (Fig. 2). The pooled overall major wound complication rate estimated with a random effect model was equal to 18 % [95 % CI 10–27]. Heterogeneity within studies estimated by the I squared was 79.5 % and the between study variance (Tau) equal to 0.0099. Cochran’s Q test for heterogeneity provided a p-value < 0.0001.

In the analysis of the moderate hypofractionated studies, 312 patients were included across four studies (Fig. 3). The pooled overall major wound complication rate estimated with a random effect model was 29 % [95 % CI 23–34]. I squared was 18.0 %, Tau equal to 0.0006 and a p-value of 0.3009 for the Cochran’s Q test.

The MWC rates of the ultra-hypofractionated group are descriptively reported in Table 3. Regardless of the specific regimen used, the MWC rate did not exceed 41 % with one study reporting a rate of 41 % and the remaining studies reporting rates not higher than 32 %.

In total, 6 studies were included in the analysis of grade 3 or higher

toxicities for conventional radiotherapy and moderate hypofractionated radiotherapy. Thirteen studies were excluded due to missing data on only grade 3 or higher toxicities, one study was excluded due to a critical risk of bias.

In the meta-analysis of conventional radiotherapy, three studies were included with a total of 167 patients (Fig. 4). The pooled overall toxicity rate estimated with a random effect model was 5 % [95 % CI 0.0–12.1]. I2 was equal to 78.3 % and Tau 0.0029. Cochran’s Q test for heterogeneity showed a p-value equal to 0.0100.

In the analysis of the moderate hypofractionated studies, 209 patients were included across three studies (Fig. 5). The pooled overall toxicity rate estimated with a random effect model was 4 % [95 % CI 1.2–6.3]. I squared was equal to 0.0 % meaning that the variability can be explained by chance alone, Tau equal to 0 and a p-value of 0.8834 for the Cochran’s Q test.

The late toxicities of the ultra-hypofractionated schedule are reported in Tables 3 and 4. As shown in these tables, a wide range of toxicity rates (1–16 %) and median follow-up periods (5–57 months) are listed.




Fig. 2. Forest plot of the conventional radiotherapy studies and MWC.




Fig. 3. Forest plot of the moderate hypofractionated studies and MWC.




Fig. 4. Forest plot of conventional radiotherapy and late toxicity grade 3 or higher.




Fig. 5. Forest plot of the moderate hypofractionated studies and late toxicity grade 3 or higher.

**Discussion**

Moderate hypofractionated radiotherapy is a safe treatment in terms of toxicity and oncological outcomes. The outcomes of ultra-hypofractionated radiotherapy are still premature and heterogeneous, making it difficult to draw conclusions about this regimen. This study included a total of thirty studies, consisting of thirteen conventional radiotherapy studies, four studies on moderate hypofractionation, and thirteen studies on ultra-hypofractionation with multiple fractionation schedules. To the best of our knowledge, this is the first meta-analysis that incorporates conventional radiotherapy, moderate hypofractionation, and ultra-hypofractionated radiotherapy, focusing on MWC and late toxicities in the context of hypofractionation for STS.

Our analysis shows that studies on moderate hypofractionation report a comparable wound complication rate to the standard treatment of 25 × 2 Gy. The guideline for conventional radiotherapy has been based on the 35 % MWC rate reported by O’Sullivan et al. in 2002 [1]. Accordingly, studies reporting a MWC rate below 35 % may be designated as clinically safe. In our meta-analysis, the moderate hypofractionation studies show a wound complication rate of 29 %, which aligns with this historical reference. When comparing MWC with the ultra-hypofractionation group, outcomes were not meta-analyzed due to the different hypofractionation schedules. Additionally, timing of

surgery after radiotherapy varied, ranging from a few days after the final RT fraction to a waiting period of 6–8 weeks, potentially influencing complication rates. While descriptive findings of the included studies suggest that the rate of MWC does not exceed 41 %, regardless of the dose, these results are derived from individual heterogeneous studies and should be interpreted with caution.

Regarding late grade ≥ 3 toxicity, moderate hypofractionation showed comparable toxicity rates to conventional radiotherapy, with pooled estimates of 4 % (95 % CI 1.2–6.3) and 5 % (95 % CI 0.0–12.1), respectively. However, only four studies specifically reported grade 3 toxicity or higher and these studies reported heterogeneous toxicity results, varying from 0 up to 15 %. For the ultra-hypofractionated group, toxicities were not meta-analyzed. The ultra-hypofractionated studies are showing a wide variety of rates, ranging from 1 % up to 16 %. However, these rates are potentially misleading due to different follow up periods. It is important to note that the EQD2 calculations in this meta-analysis reflect the dose delivered to the planning target volume (PTV). The actual dose of the normal tissue within or near by the PTV may differ from the dose received by the tumor, potentially influencing the incidence of wound complications and late toxicities.

The oncological outcomes showed that moderate hypofractionation is associated with not only acceptable MWC rates and comparable late toxicity but also similar local control rates between 90 % and 95 %. In

ultra-hypofractionation, the 5 × 5 Gy regimen showed lower local control with a 5-year local recurrence-free survival (LRFS) of 81 %, while the 5 × 7 Gy and 5 × 8 Gy regimens show a 2-year local recurrence rate of 0 %. It is important to note that most of the studies used Kaplan-Meier's methodology to estimate the cumulative incidence of local recurrence. However, this method does not account for the presence of competing risks, such as death from other causes, which may lead to an overestimation of the cumulative incidence of local recurrence. The stronger the competition between the two competing events, the greater the bias [50].

This meta-analysis has several strengths, including its comprehensive inclusion of various studies across different hypofractionation schedules, offering broad insights into MWC and late toxicities in STS. However, there are notable limitations. Most of the included studies had small patient populations, limiting interpretation. Differences in fractionation schedules, surgery timing, patient demographics, and evolving radiotherapy techniques over time further complicate comparisons. Follow-up periods varied significantly, potentially underestimating long term toxicities. Additionally, most studies were phase 2 studies, single-center trials or retrospective studies, introducing potential biases. These factors underscore the need for caution when interpreting the findings of this meta-analysis.

In conclusion, moderate hypofractionation seems to be safe and effective. In all moderate hypofractionated studies, the MWC, late toxicities and oncological outcomes are comparable. In ultra-hypofractionation, a balance must be found between major wound complications and oncological outcomes, which is difficult to interpret due to the heterogeneity across studies. The need for larger cohorts and longer follow up is necessary to evaluate the ultra-hypofractionated regimens. As previously noted by Baldini et al., it is still necessary to avoid implementing hypofractionation outside of study settings [51]. A RCT is needed to compare outcomes in moderate and ultra-hypofractionated regimens. Currently, the randomized SCOPES trial (Short Course Of Preoperative Radiotherapy in Head and Neck-, Trunk-and Extremity Soft Tissue Sarcomas; NCT04425967; comparing 14 × 3 Gy vs 25 × 2 Gy) is accruing patients to evaluate this question.

#### Funding statement

This study was supported by the Sarcoma Foundation of America (SFA Grant 1285412) and the Dutch Cancer Society (KWF Grant 13697). The funding sources was not involved in the study design, data collection, analysis, and interpretation, writing of the manuscript, or the decision to submit the article for publication.

#### Registration and protocol

This meta-analysis was not registered and no protocol was prepared.

#### CRediT authorship contribution statement

**G.F. Foppele:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **M. Fiocco:** Writing – review & editing, Visualization, Validation, Supervision, Software, Resources, Methodology, Investigation, Formal analysis, Data curation, Conceptualization. **B.H.G. van Riet:** Writing – review & editing, Validation, Methodology, Formal analysis. **R.L.M. Haas:** Writing – review & editing, Visualization, Validation, Supervision, Resources, Project administration, Methodology, Funding acquisition, Data curation, Conceptualization. **L.M. Wiltink:** Writing – review & editing, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Formal analysis, Data curation, Conceptualization.

#### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.radonc.2025.111034>.

#### References

- [1] O'Sullivan B, Davis AM, Turcotte R, et al. Preoperative versus postoperative radiotherapy in soft-tissue sarcoma of the limbs: a randomised trial. *Lancet* 2002; 359:2235–41.
- [2] Davis AM, Sennik S, Griffin AM, et al. Predictors of functional outcomes following limb salvage surgery for lower-extremity soft tissue sarcoma. *J. Surg. Oncol.* 2000; 73:206–11.
- [3] Cannon CP, Ballo MT, Zagars GK, et al. Complications of combined modality treatment of primary lower extremity soft-tissue sarcomas. *Cancer* 2006;107: 2455–61.
- [4] Haas RL, Miah AB, LePechoux C, et al. Preoperative radiotherapy for extremity soft tissue sarcoma; past, present and future perspectives on dose fractionation regimens and combined modality strategies. *Radiother. Oncol.* 2016;119:14–21.
- [5] Salerno KE, Alektiar KM, Baldini EH, et al. Radiation Therapy for Treatment of Soft Tissue Sarcoma in Adults: Executive Summary of an ASTRO Clinical Practice Guideline. *Pract. Radiat. Oncol.* 2021;11:339–51.
- [6] Oncology NCPGi: NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Soft Tissue Sarcoma Version 1.2024, 2024.
- [7] Gronchi A, Miah AB, Dei Tos AP, et al. Soft tissue and visceral sarcomas: ESMO-EURACAN-GENTURIS Clinical Practice guidelines for diagnosis, treatment and follow-up(☆). *Ann. Oncol.* 2021;32:1348–65.
- [8] Haas RL, Delaney TF, O'Sullivan B, et al. Radiotherapy for management of extremity soft tissue sarcomas: why, when, and where? *Int. J. Radiat. Oncol. Biol. Phys.* 2012;84:572–80.
- [9] Widmark A, Gunnlaugsson A, Beckman L, et al. Ultra-hypofractionated versus conventionally fractionated radiotherapy for prostate cancer: 5-year outcomes of the HYPO-RT-PC randomised, non-inferiority, phase 3 trial. *Lancet* 2019;394: 385–95.
- [10] Roeder F, Meldolesi E, Gerum S, et al. Recent advances in (chemo-)radiation therapy for rectal cancer: a comprehensive review. *Radiat. Oncol.* 2020;15:262.
- [11] Murray Brunt A, Haviland JS, Wheatley DA, et al. Hypofractionated breast radiotherapy for 1 week versus 3 weeks (FAST-Forward): 5-year efficacy and late normal tissue effects results from a multicentre, non-inferiority, randomised, phase 3 trial. *Lancet* 2020;395:1613–26.
- [12] Haas RL, Floot BGJ, Scholten AN, et al. Cellular Radiosensitivity of Soft Tissue Sarcoma. *Radiat. Res.* 2021;196:23–30.
- [13] Bishop AJ, Mitra D, Farooqi A, et al. Moderately hypofractionated, preoperative radiotherapy in patients with soft tissue sarcomas (HYPORT-STIS): Updated local control, late toxicities, and patient-reported outcomes. *Cancer* 2025;131:e35542.
- [14] Guadagnolo BA, Bassett RL, Mitra D, et al. Hypofractionated, 3-week, preoperative radiotherapy for patients with soft tissue sarcomas (HYPORT-STIS): a single-centre, open-label, single-arm, phase 2 trial. *Lancet Oncol.* 2022;23:1547–57.
- [15] Page MA-O, McKenzie JE, Bossuyt PM, et al: The PRISMA 2020 statement: an updated guideline for reporting systematic reviews.
- [16] Sterne JAC HJ: The Risk Of Bias In Non-randomized Studies – of Interventions, Version 2 (ROBINS-I V2) assessment tool.
- [17] Sterne JAC SJ, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al.: RoB 2: a revised tool for assessing risk of bias in randomised trials.
- [18] Terminology Criteria for Adverse Events (CTCAE) Version 5.0, 2017.
- [19] Joiner MC, van der Kogel A, editors. *Basic Clinical Radiobiology*. (6th ed.), 2025.
- [20] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control. Clin. Trials* 1986; 7:177–88.
- [21] R Core Team (2023). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing V, Austria., <https://www.R-project.org/>.
- [22] Safia K. Ahmed M: Hypofractionated Preoperative Radiation Therapy for Extremity and Superficial Trunk Soft Tissue Sarcomas: Results of a Prospective, Phase II Trial. Presented at the CTOS Annual Meeting 2024; San Diego. Session 7, paper 49.
- [23] Shah D, Borys D, Martinez SR, et al. Complete pathologic response to neoadjuvant radiotherapy is predictive of oncological outcome in patients with soft tissue sarcoma. *Anticancer Res.* 2012;32:3911–5.
- [24] Gabriela Studer M, HoDab, Christoph Glanzmann, MDab, Franziska Maduz, BScNc, Beata Bode, MDD and Bruno Fuchs, MDbc: Preoperative IMRT for soft-tissue sarcoma of the extremities and trunk: low rate of wound complications. *Current Orthopaedic Practice*, 2018.
- [25] Canter RJ, Martinez SR, Tamurian RM, et al. Radiographic and histologic response to neoadjuvant radiotherapy in patients with soft tissue sarcoma. *Ann. Surg. Oncol.* 2010;17:2578–84.
- [26] Abdallah ZK, El Masry AM, Azmy SI, et al. Assessment of local complications with preoperative versus postoperative radiotherapy in patients with soft tissue sarcoma in extremities: a randomized comparative clinical trial. *J. Orthop.* 2024;53:34–40.

- [27] Bonvalot S, Rutkowski PL, Thariat J, et al. Final Safety and Health-Related Quality of Life Results of the Phase 2/3 Act.In.Sarc Study with Preoperative NBTXR3 Plus Radiation Therapy Versus Radiation Therapy in locally Advanced Soft-Tissue Sarcoma. *Int. J. radiation oncology, biology Phys.* 2022;114:422–32.
- [28] González-Viguera J, Reynés-Llompert G, Lozano A. Outcomes and computed tomography radiomic features extraction in soft tissue sarcomas treated with neoadjuvant radiation therapy. *Rep Pract Oncol Radiother* 2021;26:804–13.
- [29] Green CM, Nguyen N, Wylie J, et al. The treatment of periarticular soft tissue sarcoma following neo-adjuvant radiotherapy: a cohort study. *World J. Surg. Oncol.* 2015;13:108.
- [30] Huang Z, Li N, Tang Y, et al: Neoadjuvant radiotherapy for soft tissue sarcoma in China: a preliminary result. *ANNALS OF TRANSLATIONAL MEDICINE* 10, 2022.
- [31] Hui AC, Ngan SY, Wong K, et al. Preoperative radiotherapy for soft tissue sarcoma: the Peter MacCallum Cancer Centre experience. *Eur. J. Surg. Oncol.* 2006;32:1159–64.
- [32] Seddon B, Grange FL, Simões R, et al. The IMRiS Trial: a phase 2 Study of Intensity Modulated Radiation Therapy in Extremity Soft Tissue Sarcoma. *Int. J. Radiat. Oncol. Biol. Phys.* 2024;120:978–89.
- [33] Shelby RD, Suarez-Kelly LP, Yu PY, et al. Neoadjuvant radiation improves margin-negative resection rates in extremity sarcoma but not survival. *J. Surg. Oncol.* 2020;121:1249–58.
- [34] Wang D, Harris J, Kraybill WG, et al. Pathologic complete Response and Clinical Outcomes in Patients with Localized Soft Tissue Sarcoma Treated with Neoadjuvant Chemoradiotherapy or Radiotherapy: the NRG/RT0G 9514 and 0630 Nonrandomized Clinical Trials. *JAMA Oncol.* 2023;9:646–55.
- [35] Foppele GF, Fiocco M, Ubbels JF, et al. Pandemic driven preoperative moderate hypofractionated radiotherapy for soft tissue sarcomas. *Eur. J. Cancer* 2025;217:115234.
- [36] Montero A, Chen-Zhao X, Ciervide R, et al. Moderate hypofractionated radiation therapy and pathologic response for soft tissue sarcomas (STS) of limbs and trunk: experience from a tertiary cancer center. *Clinical & translational oncology: official publication of the Federation of Spanish Oncology Societies and of the National Cancer Institute of Mexico* 2024;26:204–13.
- [37] Kosela-Paterczyk H, Szacht M, Morysinski T, et al. Preoperative hypofractionated radiotherapy in the treatment of localized soft tissue sarcomas. *European journal of surgical oncology: the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology* 2014;40:1641–7.
- [38] Kosela-Paterczyk H, Teterycz P, Spalek MJ, et al: Efficacy and Safety of Hypofractionated Preoperative Radiotherapy for Primary Locally Advanced Soft Tissue Sarcomas of Limbs or Trunk Wall. *Cancers (Basel)* 13, 2021.
- [39] Heesen P, Di Lonardo M, Ciobanu-Carusu O, et al: Ultrahypofractionated Versus Normofractionated Preoperative Radiotherapy for Soft Tissue Sarcoma: A Multicenter, Prospective Real-World-Time Phase 2 Clinical Trial. *Cancers (Basel)* 16, 2024.
- [40] Potkrajcic V, Traub F, Hermes B, et al. Hypofractionated preoperative radiotherapy for high risk soft tissue sarcomas in a geriatric patient population. *Radiol. Oncol.* 2021;55:459–66.
- [41] Kilic L, Ekenel M, Karabulut S, et al. Neoadjuvant sequential chemoradiotherapy versus radiotherapy alone for treatment of high-risk extremity soft tissue sarcoma: a single-institution experience. *Contemporary oncology (Poznan, Poland)* 2017;21:60–5.
- [42] Mayo ZS, Parsai S, Asha W, et al. Early outcomes of ultra-hypofractionated preoperative radiation therapy for soft tissue sarcoma followed by immediate surgical resection. *Radiother. Oncol.* 2023;180:109439.
- [43] Parsai S, Lawrenz J, Kilpatrick S, et al. Early Outcomes of Preoperative 5-Fraction Radiation Therapy for Soft Tissue Sarcoma Followed by Immediate Surgical Resection. *Adv. Radiat. Oncol.* 2020;5:1274–9.
- [44] Kalbasi A, Kamrava M, Chu F-I, et al. A Phase II Trial of 5-Day Neoadjuvant Radiotherapy for Patients with High-Risk Primary Soft Tissue Sarcoma. *Clin. Cancer Res. official J. Am. Association Cancer Res.* 2020;26:1829–36.
- [45] Novikov SN, Gafton GI, Ebert MA, et al. Preoperative stereotactic ablative body radiotherapy with postoperative conventional irradiation of soft tissue sarcomas: Protocol overview with a preliminary safety report. *Radiother. Oncol.* 2021;161:126–31.
- [46] Kubicek GJ, Kim TW, Gutowski CJ, et al: Preoperative Stereotactic Body Radiation Therapy for Soft-Tissue Sarcoma: Results of Phase 2 Study. *ADVANCES IN RADIATION ONCOLOGY* 7, 2022.
- [47] Kubicek GJ, LaCouture T, Kaden M, et al. Preoperative Radiosurgery for Soft Tissue Sarcoma. *Am. J. Clin. Oncol.* 2018;41:86–9.
- [48] Leite ETT, Munhoz RR, Camargo VP, et al. Neoadjuvant stereotactic ablative radiotherapy (SABR) for soft tissue sarcomas of the extremities. *Radiother. Oncol.* 2021;161:222–9.
- [49] Bedi M, Singh R, Charlson JA, et al. Is 5 the New 25? Long-Term Oncologic Outcomes From a Phase II, Prospective, 5-Fraction Preoperative Radiation Therapy Trial in Patients With Localized Soft Tissue Sarcoma. *Adv. Radiat. Oncol.* 2022;7:100850.
- [50] Keurentjes JC, Fiocco M, Schreurs BW, et al. Revision surgery is overestimated in hip replacement. *Bone Joint Res* 2012;1:258–62.
- [51] Baldini EH, Guadagnolo BA, Salerno KE, et al. Hypofractionated Preoperative Radiation Should not yet Be used as Standard of Care for Extremity and Truncal Soft Tissue Sarcoma. *J. Clin. Oncol.* 2024;42:4240–5.