



Universiteit  
Leiden  
The Netherlands

## Higher physical activity levels are related to faecal microbiota diversity and composition in young adults

Ortiz-Alvarez, L.; Xu, H.W.; Ruiz-Campos, S.; Acosta, F.M.; Migueles, J.H.; Vilchez-Vargas, R.; ... ; Martinez-Tellez, B.

### Citation

Ortiz-Alvarez, L., Xu, H. W., Ruiz-Campos, S., Acosta, F. M., Migueles, J. H., Vilchez-Vargas, R., ... Martinez-Tellez, B. (2025). Higher physical activity levels are related to faecal microbiota diversity and composition in young adults. *Biology Of Sport*, 42(1), 123-135. doi:10.5114/biolsport.2025.139850

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/4299658>

**Note:** To cite this publication please use the final published version (if applicable).

# Higher physical activity levels are related to faecal microbiota diversity and composition in young adults

**AUTHORS:** Lourdes Ortiz-Alvarez<sup>1,2</sup>, Huiwen Xu<sup>1,2</sup>, Samuel Ruiz-Campos<sup>3,4</sup>, Francisco M. Acosta<sup>1,5,6,7</sup>, Jairo H. Migueles<sup>1,8</sup>, Ramiro Vilchez-Vargas<sup>9</sup>, Alexander Link<sup>9</sup>, Julio Plaza-Díaz<sup>2,10,11</sup>, Angel Gil<sup>2,11,12,13</sup>, Idoia Labayen<sup>14</sup>, Jonatan R. Ruiz<sup>1,13\*</sup>, Borja Martinez-Tellez<sup>1,3,4,15\*</sup>

<sup>1</sup> PROFITH (PROmoting FITness and Health through Physical Activity) Research Group, Sport and Health University Research Institute (iMUDS), Department of Physical and Sports Education, Faculty of Sport Sciences, University of Granada, Granada, Spain

<sup>2</sup> Department of Biochemistry and Molecular Biology II, School of Pharmacy, University of Granada, Granada, Spain

<sup>3</sup> Department of Nursing, Physiotherapy and Medicine and SPORT Research Group (CTS-1024), CERNEP Research Center, University of Almería, Almería, Spain

<sup>4</sup> Biomedical Research Unit, Torrecárdenas University Hospital, Almería, 04009, Spain

<sup>5</sup> Turku PET Centre, University of Turku, Turku, Finland

<sup>6</sup> Turku PET Centre, Turku University Hospital, Turku, Finland

<sup>7</sup> InFLAMES Research Flagship Center, University of Turku, Turku, Finland

<sup>8</sup> Department of Biosciences and Nutrition, Karolinska Institute, Karolinska, Sweden

<sup>9</sup> Department of Gastroenterology, Hepatology and Infectious Diseases, Otto-von-Guericke-University Magdeburg, Magdeburg, Germany

<sup>10</sup> Children's Hospital of Eastern Ontario Research Institute, Ottawa, ON K1H 8L1, Canada

<sup>11</sup> Institute of Nutrition and Food Technology "José Mataix", Biomedical Research Center, Parque Tecnológico Ciencias de la Salud, University of Granada, Armilla, Granada, Spain

<sup>12</sup> CIBEROBN, Biomedical Research Networking Center for Physiopathology of Obesity and Nutrition, Carlos III Health Institute, Madrid, Spain

<sup>13</sup> Instituto de Investigación Biosanitaria, ibs.Granada, Granada, Spain

<sup>14</sup> Institute for Sustainability & Food Chain Innovation (ISFOOD), Department of Health Sciences, Public University of Navarra, Campus de Arrosadía, Pamplona, Spain

<sup>15</sup> Department of Medicine, Division of Endocrinology, and Einthoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, The Netherlands

\* Joint last authors

**ABSTRACT:** Increasing physical activity (PA) is recognised as an efficacious approach for preventing and treating cardiometabolic diseases. Recently, the composition of microorganisms living within the gut has been proposed as an important appropriate target for treating these diseases. Whether PA is related to faecal microbiota diversity and composition in humans remains to be ascertained. Thus, we examined the association of the time spent in objectively measured PA with faecal microbiota diversity and composition in young adults. A cross-sectional study enrolled 88 young adults aged  $22.0 \pm 2.3$  years (72.7% women), whose time spent in PA at different intensities was objectively measured with a wrist-worn accelerometer for 7 consecutive days. Faecal microbiota diversity and composition were analysed with hypervariable tag sequencing of the V3–V4 region of the 16S rRNA gene. The mean Euclidean Norm of the raw accelerations Minus One (mg) during waking time, considered as overall PA, and the time spent in vigorous PA were positively correlated with alpha diversity indexes (all  $\rho \geq 0.23$ ,  $P \leq 0.034$ ). Regarding faecal microbiota composition, participants with low time spent in vigorous PA had higher relative abundance of the *Gammaproteobacteria* class ( $q = 0.021$ ,  $FDR = q$ -value) compared to the participants with high time spent in vigorous PA, and lower relative abundance of the *Porphyromonadaceae* family ( $q = 0.031$ ) and the *Alistipes* genus ( $q = 0.015$ ) compared to the individuals with high and intermediate time spent in vigorous PA, respectively. Our results suggest that PA, especially of vigorous intensity, is related to faecal microbiota diversity and the *Gammaproteobacteria* class and *Porphyromonadaceae* family in young adults.

**CITATION:** Ortiz-Alvarez L, Xu H, Ruiz-Campos S et al. Higher physical activity levels are related to faecal microbiota diversity and composition in young adults. *Biol Sport*. 2025;42(1):123–135.

Received: 2023-11-27; Reviewed: 2024-02-17; Re-submitted: 2024-04-05; Accepted: 2024-05-08; Published: 2024-06-04.

Corresponding author:

Jonatan Ruiz Ruiz

Email: [ruizj@ugr.es](mailto:ruizj@ugr.es)

Borja Martinez Tellez

Email: [borjammt@ual.es](mailto:borjammt@ual.es)

## ORCID:

Lourdes Ortiz-Alvarez  
0000-0002-6827-2358

Huiwen Xu  
0000-0002-2308-7140

Samuel Ruiz-Campos  
0009-0008-5544-887X

Francisco M. Acosta  
0000-0002-4792-0969

Jairo H. Migueles  
0000-0003-0366-6935

Ramiro Vilchez-Vargas  
0000-0002-7220-258X

Alexander Link  
0000-0002-9514-4562

Julio Plaza-Díaz  
0000-0002-5171-9408

Angel Gil  
0000-0001-7663-0939

Idoia Labayen  
0000-0002-4334-3287

Jonatan R. Ruiz  
0000-0002-7548-7138

Borja Martinez-Tellez  
0000-0001-8783-1859

## Key words:

Physical activity  
Gastrointestinal microbiome  
Shorts-chain fatty acids  
Obesity  
Activity monitor

## INTRODUCTION

Lifestyle physical activity (PA) is associated with a myriad of physiological adaptations that benefit human health. PA is one of the most effective strategies to prevent and combat cardiometabolic alterations and is related to a 27% decrease in mortality risk [1]. However, the underlying mechanisms that explain how PA enhances cardiometabolic health remain to be elucidated.

Gut microbiota refers to microbial communities colonising the gastrointestinal tract [2], indispensable in regulating the host nutrition, metabolic function and immunological response [3, 4]. Dysbiosis arises from an imbalance within microbial communities [5], influenced by various factors, including dietary patterns, sedentary or unhealthy lifestyles [6–8], and medication use [9]. This imbalance, in turn, correlates with conditions such as obesity and cardiometabolic diseases [10]. Recent evidence suggests that PA is one of humans' most significant lifestyle factors influencing gut microbiota diversity and composition [11, 12]. For instance, case-controlled studies showed that faecal microbiota from athletes [13, 14] was much more diverse and had a higher proportion of several bacterial taxa than healthy sedentary individuals. Similarly, in football athletes, it was found that increased levels of PA promoted greater diversity of the faecal microbiota via the production of short-chain fatty acids by gut bacteria, enhancing overall health [15]. Another cross-sectional study observed that premenopausal women meeting the PA World Health Organization recommendations had a greater relative abundance of *Akkermansia* and *Faecalibacterium* genera than sedentary women [16]. Evidence indicates that *Akkermansia* and *Faecalibacterium* genera are associated with reduced inflammation and, therefore, may play a role in preventing the development of cardiometabolic diseases [17]. Based on that, a recent study showed that individuals with higher levels of PA showed a different Mediterranean pattern and faecal microbiota composition than individuals with obesity who reported lower levels of PA [18]. Most studies investigating the relationship between PA and faecal microbiota composition have used self-reported questionnaires to determine PA levels [13, 14, 19]. However, these instruments have the disadvantage of misclassifying PA levels and thus compromise the ability to detect valid associations between PA levels and faecal microbiota composition [20]. Based on the aforementioned studies using self-reported data, we hypothesise that increased levels of PA, at different intensities, are associated with elevated faecal microbiota diversity and a greater prevalence of beneficial bacteria. Thus, through the utilization of objective measures of PA in the present study, we aimed to explore the association between the time spent in objectively measured PA at different intensities with faecal microbiota diversity and composition in a cohort of young individuals.

## MATERIALS AND METHODS

### *Design study and participants*

A total of 92 (65 women) young healthy adults, aged 18–25 years, were included in the present cross-sectional study. This study was

carried out within the framework of the ACTIBATE study [21], an exercise-based randomized controlled trial (Clinical Trials.gov ID: NCT02365129). All assessments were performed in Granada (Spain) between October and November in 2016. Inclusion criteria were: being engaged in less than 20 min of moderate-vigorous PA on less than 3 days/week, having a stable body weight over the last 3 months (< 3 kg change), not smoking, not taking any medication (including antibiotics in the last 3 months), not presenting any acute or chronic illness and not being pregnant. The study protocol and experimental design were applied in accordance with the last revised ethical guidelines of the Declaration of Helsinki. The study was approved by the Ethics Committee on Human Research of the University of Granada (no. 924) and the Servicio Andaluz de Salud (Centro de Granada, CEI-Granada); all participants signed informed consent.

### *Physical activity assessment*

PA variables were objectively measured with one accelerometer on the non-dominant wrist (ActiGraph GT3X+, Pensacola, FL), during 7 consecutive days (24 h/day) [21]. Detailed information about how to wear the accelerometer was given to participants, including the instruction to remove it in daily water-based activities, such as washing dishes or showering.

The sampling frequency of 100 Hz was selected to store the raw accelerations of the accelerometers [22]. We exported and converted the raw accelerations to the “.csv” format using ActiLife v.6.13.3 software (ActiGraph, Pensacola, FL, US). Afterwards, the “ggir” [23] package in R software was used to process the raw “.csv” files. This processing consisted of: (i) local gravity data auto-calibration of accelerations according to the local gravitational acceleration [24], (ii) calculation of the Euclidean Norm of the raw accelerations Minus One G with negative values rounded to 0 (ENMO) calculated elsewhere [25], (iii) detection of non-wear time based on the raw acceleration of the three axes, (iv) determination of MAL detection of sustained functioning of the accelerometer by means of abnormal high accelerations incompatible with human movement (i.e., related to device malfunctioning), (v) imputation of non-wear time and abnormal high accelerations, (vi) identification of waking and sleeping time based on the automated algorithm guided by the participants' daily reports [26], and (vii) estimation of sedentary time and the time spent in light PA, moderate PA, vigorous PA, and moderate to vigorous PA using age-specific cut-points for a wrist-worn accelerometer, for Euclidean Norm Minus One (ENMO) [27]. We measured the mean ENMO (mg) during waking time, which is considered an overall indicator of the PA (overall PA). For the analyses we only included the participants who wore the accelerometers for  $\geq 16$  h/day during at least 4 days (including at least 1 weekend day).

*Faecal microbiota analyses**Stool collection and DNA extraction*

The participants collected approximately 50 g of a faecal sample in plastic sterile containers, which were transported in portable coolers to the research centre. Faecal samples were stored at  $-80^{\circ}\text{C}$  until extraction of DNA. The QIAamp DNA Stool Mini Kit (QIAGEN, Barcelona, Spain) was used for extraction of DNA, following the manufacturer's instructions. The samples were incubated at  $95^{\circ}\text{C}$  to ensure lysis of both gram-positive and gram-negative bacteria. Then, we quantified DNA with a NanoDrop ND1000 spectrophotometer (Thermo Fisher Scientific, DE, USA). Finally, DNA purity was determined by measuring the ratio of absorbance at  $\text{A}260/280$  nm and  $\text{A}260/230$  nm.

*Sequencing analysis*

DNA extracted was amplified by polymerase chain reaction (PCR) by primer pairs – forward primer (5'CCTACGGGNGGCWGCAG3') and reverse primer (5'GACTACHVGGGTATCTAATCC3') – targeting the V3 and V4 hypervariable regions of the bacterial 16S rRNA gene. All PCRs were executed in  $25\ \mu\text{L}$  reaction volumes incorporating  $12.5\ \mu\text{L}$  of 2X KAPA HiFi Hotstart ready mix (KAPA Biosystems, Woburn, MA, USA),  $5\ \mu\text{L}$  of each forward and reverse primer ( $1\ \mu\text{M}$ ) and  $2.5\ \mu\text{L}$  of extracted DNA (10 ng) under the following cycling circumstances: (a) denaturation at  $95^{\circ}\text{C}$  for 3 min, (b) cycles of denaturation at  $95^{\circ}\text{C}$  for 30 s, (c) annealing at  $55^{\circ}\text{C}$  for 30 s, (d) elongation at  $72^{\circ}\text{C}$  for 30 s, (e) a final extension at  $72^{\circ}\text{C}$  for 5 min. To purify the PCR products from free primers and primer dimers we used AMPure XP beads (Beckman Coulter, Indianapolis, IN, USA). Next, the index PCR attached dual indices and Illumina sequencing adapters using the Nextera XT Index Kit (Illumina, San Diego, CA, USA), on a thermal cycler using the requirements previously mentioned. After that, AMPure XP beads (Beckman Coulter, Indianapolis, IN, USA) were used for purification of the pooled PCR products. The resultant amplicons were sequenced at MiSeq (Illumina, USA), using a paired-end ( $2 \times 300$  nt) Illumina MiSeq sequencing system (Illumina, San Diego, CA, USA).

*Bioinformatics analysis*

We analysed the FASTQ files with the "dada2" [28] package in R software, obtaining 11,659,014 paired ends with an average of  $126,728 \pm 33,395$  reads per sample. The cut-off of 10,000 reads was surpassed for all samples. Samples were resampled to a minimum sequencing depth of 30,982 reads using the "phyloseq" [29] package in R software, returning 11,158 phylotypes.

The "Classifier" function from the Ribosomal Database Project (RDP) was used to assign the taxonomic affiliation of phylotypes, based on the naïve Bayesian classification [30] with a pseudo-bootstrap threshold of 80%. We obtained a total of 209 genera belonging to 16 different phyla. The "seqmatch" [31] function from RDP was performed to define the discriminatory power of each sequence read with the purpose of annotating species assignments; we

executed annotation according to previously published criteria [32]. Microbial communities were analysed at different taxonomic levels (phylum to genus), calculating relative abundances, expressed as percentages.

We performed the analyses with those bacteria with more than 0.5% on average in their relative abundance.

Next, alpha and beta diversities were estimated based on the identified microbial communities. Alpha diversity takes into account the number of different phylotypes and relative abundances of a single sample [33], whereas beta diversity shows differences in microbial community composition between individuals, which is the degree to which samples differ from one another [34]. Alpha diversity was assessed based on Chao richness, Shannon, inverse Simpson and evenness Camargo indexes with the "microbiome" [35] package in R software. Chao richness estimates diversity according to the number of different phylotypes in the community [36]; that is, higher Chao richness indicates higher diversity in the community. Shannon diversity increases as both the richness and the evenness of the community increase [37]; the inverse of Simpson diversity is calculated from the classical Simpson diversity and indicates richness in a community with uniform evenness [38], and evenness Camargo determines the equitability of phylotype frequencies in a community [39]. Beta diversity was measured quantitatively using permutational multivariate analysis of variance (PERMANOVA) based on Bray-Curtis dissimilarity.

*Anthropometric and body composition measurements*

Participants' weight and height were measured, without shoes and wearing the standard clothes, using a SECA scale and stadiometer (model 799, Electronic Column Scale, Hamburg, Germany). Body mass index (BMI) was calculated as weight (kg)/height ( $\text{m}^2$ ). Body composition was evaluated by dual energy X-ray absorptiometry (DEXA, HOLOGIC, Discovery Wi, Marlborough, MA). The lean mass index (LMI) and fat mass index (FMI) were calculated as lean body mass and fat body mass, respectively, in kg, divided by height in  $\text{m}^2$ . The fat mass percentage was determined as the body fat mass divided by the total body mass and multiplied by 100.

*Cardiometabolic profile*

Fasting blood samples were collected for assessment of the cardiometabolic profile. Serum glucose, total cholesterol, high density lipoprotein-cholesterol (HDL-C) and triglycerides were measured following standard methods using an AU5832 automated analyser (Beckman Coulter Inc., Brea CA, USA). Low-density lipoprotein cholesterol (LDL-C) was estimated as:  $[\text{total cholesterol} - \text{HDL-C} - (\text{triglycerides}/5)]$ , in mg/dL. Serum insulin was measured using the Access Ultrasensitive Insulin chemiluminescent immunoassay kit (Beckman Coulter Inc., Brea CA, USA). The homeostatic model assessment for insulin resistance (HOMA-IR) index was calculated as  $(\text{insulin} (\mu\text{U}/\text{mL}) \times \text{glucose} (\text{mmol}/\text{L}))/22.5$ .

*Dietary assessment*

Dietary intake was registered using three non-consecutive 24-hour recalls, 2 weekdays and a weekend day. These 24-hour recalls were performed in the laboratory via face-to-face interviews with dietitians. To improve the accuracy of food quantification, we used coloured photographs of different portion sizes of food during the interviews [40]. All 24-hour recalls were analysed for total energy (kcal), fat, proteins, carbohydrates, and fibre intake (g) by EvalFINUT software, which is based on the United States Department of Agriculture (USDA) and “Base de Datos Española de Composición de Alimentos” (BEDCA) databases.

*Statistical analysis*

This is a secondary study derived from the ACTIBATE trial [41]; therefore, there is not a sample size calculation for this study. Data normality was explored using the D’Agostino & Pearson omnibus, visual histograms and Q-Q plots (data not shown). None of the variables followed a normal distribution; therefore data were presented as median ± interquartile range and non-parametric tests were used for all analyses. Moreover, no sex interaction was detected (all  $P > 0.05$ ), so both sexes were pooled together. Spearman correlations were performed to investigate the correlation between the PA variables and faecal microbiota diversity, using the “psych” [42] and “corrplot” [43] packages in R software. Since faecal microbiota diversity can be modified by several factors including sex [44], BMI [45] and dietary intake [46], we repeated the aforementioned correlations adjusted for sex, BMI and dietary intake in separate models (data not shown). Moreover, we repeated this analysis by adjusting for accelerometer non-wear time and glucose levels in separate models (data not shown) as possible confounders of PA variables. Overall PA and the time spent in vigorous PA were computed as tertiles according to number of participants with SPSS (SPSS v. 22.0, IBM SPSS Statistics, IBM Corp. Armonk, NY), because they were the only variables with a significant correlation with faecal microbiota diversity. The tertiles’ values for overall PA were low (13.45–29.44 mg), intermediate (30.02–35.41 mg), and high (35.49–67.10 mg), whereas for the time spent in vigorous PA the values were low (0.02–0.83 min/day), intermediate (0.87–2.67 min/day), and high (2.75–14.40 min/day). Tertiles of overall PA and the time spent in vigorous PA were compared using one-way PERMANOVA with 9,999 permutations for significance testing with the Paleontological Statistics (Past3) software [47] for the calculation of beta diversity. Kruskal-Wallis tests were performed to investigate whether there were significant differences in body composition, dietary intake and cardiometabolic profile as well as faecal microbiota alpha diversity and composition outcomes across tertiles of overall PA and the time spent in vigorous PA. Analysis of covariance was used to compare the relative abundance of genera across tertiles of the time spent in vigorous PA adjusted for protein intake with the data transformed by Blom’s formula. All P values were corrected by Benjamini and Hochberg

multiple testing to control the false discovery rate (FDR, shown as q-values) [48]. The level of significance was set at  $P < 0.05$  and  $q < 0.05$ . R software (V.3.6.0; <http://www.r-project.org>) and GraphPad Prism version 8.0.0 for Windows (GraphPad Software, San Diego, California, USA, (<http://www.graphpad.com>)) were used for the statistical analysis and graphical plots.


**RESULTS**

*Characteristics of participants*

A total of 92 participants had data from analysis of faecal microbiota diversity and composition, but only 88 participants (24 men, age =  $22.0 \pm 2.0$ ; and 64 women, age =  $21.6 \pm 2.0$ ) had valid PA measurements (as they wore the accelerometer for  $< 16$  h/day during at least 4 days), who were finally included in the analyses. Table 1 shows the descriptive characteristics of the included participants (age 21.7 (19.8–23.9) years and BMI 23.6 (21.6–28.1  $\text{kg/m}^2$ )), of whom 72.7% were women. We performed tertiles of overall PA and the time spent in vigorous PA and we observed that, generally, body composition, dietary intake and cardiometabolic profile were similar across them (Table S1), with the exception of protein intake and glucose levels ( $P = 0.018$  and  $P = 0.003$ , respectively; Table S1).

*Relationship between physical activity and faecal microbiota diversity*

Overall PA and the time spent in vigorous PA were positively correlated with alpha diversity indexes, more specifically with Shannon



**FIG. 1.** Spearman correlation of physical activity variables with faecal microbiota diversity. Boxes only represent the statistically significant ( $P < 0.05$ ) correlations and the values within the boxes show the Spearman correlation coefficient. Blue boxes indicate a positive correlation whereas red squares indicate a negative correlation between physical activity variables and faecal microbiota diversity indexes. mg: milli-gravitational units; PA: physical activity.

TABLE 1. Descriptive characteristics of the participants.

	All (N = 88)	Men (N = 24)	Women (N = 64)
Age (years)	21.7 (19.8–23.9)	22.0 (19.8–24.5)	21.6 (19.8–23.7)
<i>Body composition</i>			
Body mass index (kg/m <sup>2</sup> )	23.6 (21.6–28.1)	25.8 (22.8–32.5)	23.3 (21.0–27.4)
Lean mass index (kg/m <sup>2</sup> )	13.9 (12.6–15.7)	17.4 (15.3–18.4)	13.2 (12.5–14.3)
Fat mass index (kg/m <sup>2</sup> )	8.7 (6.3–11.5)	7.9 (4.5–11.3)	9.2 (6.5–11.5)
Fat mass (%)	36.7 (31.2–42.6)	32.5 (20.8–37.1)	39.3 (32.9–42.8)
<i>Sedentary and physical activity time</i>			
Non wear time (min/week)	8.6 (0.0–22.2)	8.3 (0.0–28.9)	8.6 (0.0–22.2)
Overall PA (mg)	31.8 (27.2–37.2)	31.0 (24.8–35.3)	32.8 (28.2–37.4)
Sedentary Time (min/day)	785.2 (753.2–820.7)	798.8 (770.0–837.7)	779.2 (752.6–820.2)
Light PA (min/day)	113.6 (99.2–139.0)	109.8 (90.7–124.2)	118.8 (99.7–145.3)
Moderate PA (min/day)	90.0 (67.9–109.5)	84.1 (57.7–103.1)	93.1 (69.7–113.1)
Vigorous PA (min/day)	1.2 (0.7–3.8)	1.3 (0.5–3.3)	1.2 (0.7–4.6)
Moderate to Vigorous PA (min/day)	91.3 (70.5–111.3)	86.3 (58.1–104.6)	95.6 (72.2–116.4)
<i>Dietary intake</i>			
Energy intake (kcal/day)	1847.7 (1580.8–2184.4)	2085.2 (1783.3–2641.3)	1798.2 (1553.6–2090.2)
Fat (g/day)	84.5 (70.0–100.2)	97.8 (71.5–111.8)	82.1 (69.2–96.7)
Proteins (g/day)	72.6 (61.4–90.9)	102.2 (75.3–118.9)	67.0 (59.8–79.4)
Carbohydrates (g/day)	198.2 (159.3–229.6)	200.4 (172.5–244.9)	198.2 (149.2–226.2)
Fiber (g/day)	16.2 (11.7–19.8)	16.3 (13.4–20.5)	16.2 (11.2–19.4)
<i>Cardiometabolic profile</i>			
Glucose (mg/dL)	87.0 (84.0–92.0)	91.0 (85.0–97.0)	87.0 (84.0–92.0)
Insulin (μUI/mL)	6.7 (4.9–10.7)	6.9 (4.8–12.4)	6.7 (4.9–9.5)
HOMA index	1.4 (1.1–2.4)	1.5 (1.0–3.0)	1.4 (1.1–2.2)
Total Cholesterol (mg/dL)	161.0 (141.5–189.0)	153.0 (139.0–174.0)	166.0 (143.0–197.0)
HDL-C (mg/dL)	51.0 (46.0–58.3)	46.0 (40.0–51.0)	53.0 (48.0–63.0)
LDL-C (mg/dL)	94.0 (78.0–114.0)	90.0 (81.0–104.0)	96.0 (78.0–114.0)
Triglycerides (mg/dL)	71.0 (52.8–106.3)	77.0 (60.0–115.0)	68.0 (52.0–98.0)
<i>Faecal microbiota</i>			
<b>Alpha diversity indexes</b>			
Richness Chao	391.6 (339.8–508.1)	374.1 (333.9–448.1)	423.5 (344.1–538.7)
Shannon diversity	4.2 (4.0–4.5)	4.2 (4.0–4.5)	4.3 (4.0–4.5)
Inverse Simpson diversity	31.7 (23.1–42.7)	31.7 (22.5–49.1)	31.7 (23.2–41.5)
Evenness Camargo	0.2 (0.2–0.3)	0.2 (0.2–0.3)	0.2 (0.2–0.3)
<b>Composition (Phylum)</b>			
<i>Actinobacteria</i> (%)	1.1 (0.6–1.9)	1.1 (0.4–1.7)	1.1 (0.6–2.2)
<i>Bacteroidetes</i> (%)	41.3 (34.7–44.8)	41.8 (39.9–45.2)	40.9 (34.3–44.8)
<i>Firmicutes</i> (%)	47.2 (42.1–52.5)	48.7 (43.3–52.0)	45.6 (41.3–53.1)
<i>Proteobacteria</i> (%)	5.1 (3.8–8.2)	6.7 (4.2–8.8)	4.6 (3.5–7.3)
<i>Verrucomicrobia</i> (%)	0.1 (0.0–2.0)	0.0 (0.0–0.8)	0.2 (0.0–3.5)

Data are presented as median (interquartile range). BMI: body mass index; FMI: fat mass index; HDL-C: high-density lipoprotein cholesterol; HOMA index: homeostatic model assessment index; LDL-C: low-density lipoprotein cholesterol; LMI: Lean mass index; mg: mili-gravitational units; PA: Physical activity.

and inverse Simpson diversity indexes (all  $\rho \geq 0.23$ ,  $P \leq 0.034$ ; Fig. 1). Only the time spent in vigorous PA was positively correlated with the Chao richness index ( $\rho = 0.24$ ,  $P = 0.023$ ; Fig. 1). However, we did not observe any significant correlation between other PA variables and alpha diversity indexes (all  $P > 0.05$ ; Fig. 1). The results were similar when sex, BMI, energy and macronutrient intake, as well as accelerometer non-wear time and glucose levels, were included as confounders in separate models (data not shown). Moreover, we found that individuals with high time spent in vigorous PA had a higher Chao richness, Shannon and inverse Simpson diversity indexes than individuals with low and intermediate time spent in vigorous PA (all  $P \leq 0.038$ ; data not shown). However, there were no differences across tertiles of overall PA and the time spent in

vigorous PA in the beta diversity at any taxonomic levels (all  $P \geq 0.060$ ; Table 2).


*Relationship between physical activity variables and faecal microbiota composition*

We analysed the differences across tertiles of overall PA and the time spent in vigorous PA on faecal microbiota composition at all taxonomic levels. There were no significant differences across tertiles of overall PA on the relative abundance of all bacteria at the different taxonomic levels (all  $P > 0.05$ ; Fig. 2). Similarly, we observed no differences across tertiles of time spent in vigorous PA on the relative abundance of bacteria at the phylum taxonomic level (all  $P \geq 0.318$ ; Fig. 3A). However, we observed that


**TABLE S1.** Characteristics of participants according to tertiles of overall PA and the time spent in vigorous PA.

	Overall PA (mg)			P	Time spent in vigorous PA (min/day)			P
	Low (13.5–29.4) n = 29	Intermediate (30.0–35.4) n = 30	High (35.5–67.1) n = 29		Low (0.0–0.8) n = 29	Intermediate (0.9–2.7) n = 30	High (2.8–14.4) n = 29	
Age (years)	22.3 ± 2.5	22.0 ± 2.0	21.6 ± 2.5	0.461	22.3 ± 2.3	22.2 ± 2.3	21.4 ± 2.2	0.239
Sex, N				0.501				0.555
Men	10	8	6		8	10	6	
Women	19	22	23		21	20	23	
<i>Body composition</i>								
BMI (kg/m <sup>2</sup> )	25.6 ± 5.4	24.2 ± 4.7	25.2 ± 4.2	0.548	24.5 ± 5.0	25.7 ± 4.8	24.8 ± 4.6	0.531
LMI (kg/m <sup>2</sup> )	14.4 ± 2.5	13.9 ± 2.3	14.9 ± 2.0	0.150	14.0 ± 2.3	14.5 ± 2.2	14.6 ± 2.4	0.380
FMI (kg/m <sup>2</sup> )	9.3 ± 3.4	8.7 ± 2.7	8.9 ± 3.3	0.805	9.0 ± 3.5	9.1 ± 3.0	8.8 ± 2.9	0.946
Fat mass (%)	37.1 ± 7.7	36.7 ± 6.5	35.4 ± 8.9	0.823	36.8 ± 8.5	36.6 ± 7.5	35.7 ± 7.3	0.734
<i>Dietary intake</i>								
Energy intake (kcal/day)	1985 ± 597.1	1805 ± 382.1	1992 ± 453.2	0.294	1827 ± 602.5	2055 ± 412.1	1892 ± 411.0	0.080
Fat (g/day)	91.0 ± 36.7	81.4 ± 21.9	89.3 ± 22.8	0.344	82.6 ± 37.2	92.1 ± 23.8	86.6 ± 20.0	0.171
Proteins (g/day)	78.9 ± 27.2	75.2 ± 19.6	80.6 ± 23.0	0.597	<b>71.3 ± 24.6*</b>	<b>85.2 ± 22.4*</b>	77.9 ± 21.3	<b>0.018</b>
Carbohydrates (g/day)	208.6 ± 61.5	188.1 ± 57.5	211.7 ± 67.3	0.215	196.7 ± 67.6	216.0 ± 51.5	194.8 ± 67.0	0.129
Fiber (g/day)	16.9 ± 6.2	16.8 ± 8.0	17.0 ± 5.1	0.529	16.6 ± 7.2	17.7 ± 6.7	16.4 ± 5.6	0.654
<i>Cardiometabolic profile</i>								
Glucose (mg/dL)	89.2 ± 7.5	88.7 ± 6.1	86.8 ± 6.4	0.502	<b>89.6 ± 6.3*</b>	<b>90.3 ± 7.0†</b>	<b>84.7 ± 5.5*†</b>	<b>0.003</b>
Insulin (μU/mL)	10.8 ± 9.3	7.4 ± 3.3	7.5 ± 4.3	0.149	9.4 ± 6.2	9.3 ± 8.1	7.0 ± 3.9	0.139
HOMA index	2.5 ± 2.6	1.6 ± 0.8	1.6 ± 1.0	0.132	2.1 ± 1.6	2.2 ± 2.3	1.5 ± 0.9	0.059
Total Cholesterol (mg/dL)	171.8 ± 38.1	167.6 ± 32.6	167.0 ± 42.1	0.757	163.0 ± 32.9	179.9 ± 47.2	163.3 ± 27.4	0.370
HDL-C (mg/dL)	51.4 ± 14.1	54.4 ± 9.3	52.6 ± 12.1	0.241	51.0 ± 13.0	55.2 ± 12.6	52.3 ± 9.8	0.448
LDL-C (mg/dL)	100.8 ± 29.1	97.5 ± 27.0	98.7 ± 29.1	0.893	95.0 ± 25.6	105.4 ± 34.1	96.5 ± 23.2	0.599
Triglycerides (mg/dL)	100.6 ± 65.2	89.6 ± 63.4	80.7 ± 63.0	0.068	84.6 ± 46.9	113.8 ± 91.9	72.3 ± 27.5	0.242

Data are presented as means ± standard deviations. Symbol (\*) indicates significant differences between low and high tertiles, whereas symbol (†) shows significant differences between intermediate and high tertiles by means of Kruskal-Wallis. BMI: body mass index; FMI: fat mass index; HDL-C: high-density lipoprotein cholesterol; HOMA index: homeostatic model assessment index; LDL-C: low-density lipoprotein cholesterol; LMI: Lean mass index; mg: milli-gravitational units; PA: Physical activity.



**FIG. 2.** Faecal microbiota composition according to tertiles of overall physical activity (PA) levels (L: low, 13.45–29.44 mg; I: intermediate, 30.02–35.41 mg; H: high, 35.49–67.10 mg). Panels indicate relative abundance of the faecal microbiota at phylum (A), class (B), order (C), family (D) and genus (E) taxonomic levels according to tertiles of overall PA. Stacked bar represents percentage abundance. Kruskal-Wallis test was used to test for each pairwise comparison, correcting for multiple comparisons FDR ( $q < 0.05$ ) (GraphPad Prism 8.00).



**FIG. 3.** Faecal microbiota composition according to tertiles of the time spent in vigorous physical activity (PA) (L: low, 0.02–0.83 min/day; I: intermediate, 0.87–2.67 min/day; H: high, 2.75–14.40 min/day). Panels indicate relative abundance of the faecal microbiota at phylum (A), class (B), order (C), family (D) and genus (E) taxonomic levels according to tertiles of the time spent in vigorous PA. Stacked bar represents percentage abundance. Symbols \* and o mean statistical significance differences between low and high time spent in vigorous PA, and symbol • represents statistical significance differences between low and intermediate time spent in vigorous PA. Kruskal-Wallis test was used to test for each pairwise comparison, correcting for multiple comparisons FDR ( $q < 0.05$ ) (GraphPad Prism 8.00).

individuals with low time spent in vigorous PA had a higher relative abundance of the *Gammaproteobacteria* class (*Proteobacteria* phylum) than individuals with intermediate time spent in vigorous PA ( $q = 0.021$ ,  $FDR = q$ -value; Fig. 3B). Moreover, individuals with high time spent in vigorous PA had higher relative abundance of unclassified *Firmicutes* class (*Firmicutes* phylum) and *Porphyromonadaceae* family (*Bacteroidetes* phylum) than individuals with low time spent in vigorous PA ( $q = 0.027$ , and  $q = 0.031$ , respectively; Fig. 3B and D). Finally, we found that individuals with intermediate time spent in vigorous PA had a higher relative abundance of the *Alistipes* genus (*Bacteroidetes* phylum) than individuals with low time spent in vigorous PA ( $q = 0.015$ ; Fig. 3E). Interestingly, these same participants presented differences in protein intake ( $q = 0.011$ ; Table S1). Thus, we repeated the analyses after adjusting for the protein intake and the differences in the relative abundance of the *Alistipes* genus disappeared ( $P = 0.080$ ; Table S2).

## DISCUSSION

In the present study, overall PA and the time spent in vigorous PA were found to be positively correlated with alpha diversity indexes in young adults. Moreover, there were differences across the tertiles of time spent in vigorous PA in the relative abundance of the *Gammaproteobacteria* class (*Proteobacteria* phylum), *Porphyromonadaceae* family and *Alistipes* genus (both *Bacteroidetes* phylum). These findings indicate that PA may play a role in faecal microbiota diversity and composition in young adults, although further studies are needed to confirm these findings.

Our results showing the positive correlation between PA and alpha diversity agree with recent findings [14, 49, 50]. However, the mechanisms by which PA may promote higher faecal microbiota diversity are unknown. A possible explanation could be the changes in the gastrointestinal tract due to intrinsic adaptations of performing PA [51]. Interestingly, from an ecological perspective, microbial diversity may be a key factor in allowing an ecosystem to continue operating properly [52]. In fact, greater species diversity has been associated with a healthy phenotype's host [53]. This is due to the

potential effects that the bacteria can exert via metabolites, such as short-chain fatty acids and neurotransmitters locally and extra-intestinal tissues in the host [54].

Our data showed that the participants with low time spent in vigorous PA had higher relative abundance of the *Gammaproteobacteria* class than individuals with higher time spent in vigorous PA. The relative abundance of the *Gammaproteobacteria* class (*Proteobacteria* phylum) has been reported to be increased in obese mice [55] and individuals with obesity [56], and disease states such as metabolic diseases and intestinal inflammation [57]. In fact, many common human pathogens, known as sulphur producers [58], are found in the *Gammaproteobacteria* class, for example, *Escherichia*, *Shigella*, and *Yersinia* genera [58]. In agreement with our findings, sedentary women and participants with low cardiorespiratory fitness [59] had a higher relative abundance of the *Gammaproteobacteria* class than active women and participants with high cardiorespiratory fitness, respectively. Similarly, a very recent study performed in > 8,000 individuals using accelerometers observed that PA

**TABLE 2.** Beta diversity across tertiles of overall PA and the time spent in vigorous PA at all taxonomic levels.

Taxonomic levels	Tertiles of overall PA (mg)		Tertiles of the time spent in vigorous PA (min/day)	
	Pseudo-F	P	Pseudo-F	P
Phylum	0.502	0.789	0.967	0.443
Class	0.926	0.508	1.224	0.262
Order	0.959	0.475	1.282	0.221
Family	1.132	0.306	1.476	0.106
Genus	0.968	0.492	1.524	0.060

PERMANOVA using 9,999 permutations for significance testing ( $p$ -value < 0.05). mg: milli-gravitational units; PA: physical activity; Pseudo-F: statistic, larger number indicate greater separation [64] across tertiles of total PA and vigorous PA levels.

**TABLE S2.** Differences in the relative abundance of *Alistipes* genus across tertiles of the time spent in vigorous PA.

	Time spent in vigorous PA (min/day)			P for model 1	P for model 2
	Low (0.0–0.8) n = 29	Intermediate (0.9–2.7) n = 30	High (2.8–14.4) n = 29		
<i>Alistipes</i> genus (%)	<b>4.3 ± 2.9*</b>	<b>6.3 ± 3.3*</b>	5.2 ± 3.0	<b>0.026</b>	0.080

Data are presented as means ± standard deviations. Symbol (\*) indicates significant differences between low and intermediate tertiles. Model 1: P value from Kruskal Wallis test. Model 2: P values were obtained from one-way analyses of variance adjusted for protein intake (g) with data transformed by Blom's formula. PA: Physical activity.

levels were associated differently with faecal microbiota composition, suggesting that the higher the PA level is, the higher is the diversity [60]. Moreover, several studies have shown that exercise seems to decrease the relative abundance of the *Gammaproteobacteria* class [61, 62]. Thus, our results suggest that performing less than 1 min/day of vigorous PA could be related to having a higher relative abundance of the *Gammaproteobacteria* class, bacteria considered health-detrimental.

In contrast, we observed that individuals with high and intermediate time spent in vigorous PA had a higher relative abundance of the *Porphyromonadaceae* family and *Alistipes* genus (both *Bacteroidetes* phylum) than individuals with lower time spent in vigorous PA. Accordingly, in a cross-sectional study of professional martial arts athletes, the relative abundance of the *Porphyromonadaceae* family was higher in the higher-level athletes in comparison with the lower-level athletes [63]. Moreover, regular swimming training [64] and voluntary wheel running [65], both in mice, were able to increase the relative abundance of the *Porphyromonadaceae* family. In fact, it has recently been found that lean individuals had a significantly higher relative abundance of the *Porphyromonadaceae* and *Rikenellaceae* families than individuals with obesity [45]. Of note is the fact that the *Alistipes* genus belongs to the *Rikenellaceae* family. In resistance-trained mice, the relative abundance of the *Alistipes* genus was positively correlated with resistance performance [66]. In humans, the relative abundance of the *Alistipes* genus is increased after consuming an animal-based diet intake, rich in protein, for 5 days [64]. Certain species that belong to the *Alistipes* genus are involved in amino acid metabolism; specifically, they can hydrolyse tryptophan to indole [67]. Since tryptophan is an essential amino acid that cannot be produced by animal cells, humans rely on dietary intake, mainly proteins, for incorporating it into the organism [68]. In our study, the individuals with intermediate time spent in vigorous PA had higher protein intake than individuals with low time spent in vigorous PA. In fact, when the protein intake was included as a confounder, the differences in the relative abundance of the *Alistipes* genus between these individuals disappeared. Considering the relationship between the *Alistipes* genus and protein metabolism [67], and the results observed in the present study, it seems possible that these differences were explained by protein intake. Therefore, our data suggest that spending time on vigorous PA, in the range 3–14 min/day, could be related to having a higher relative abundance of *Porphyromonadaceae* family bacteria, whereas the protein intake seems to modulate the relative abundance of the *Alistipes* genus in individuals with intermediate time spent in vigorous PA. Even so, the possible effect of time spent in vigorous PA on the relative abundance of the *Gammaproteobacteria* class, *Porphyromonadaceae* family and *Alistipes* genus deserves further analysis.

#### Limitations and strengths

A limitation to consider in the current study is that it followed a cross-sectional design, which prevents a causal interpretation of our results.

Well-designed randomized controlled trials should be carried out to elucidate the role of PA in faecal microbiota diversity and composition. In addition, we do not know whether our findings apply to older people or individuals presenting any metabolic disease. As for strengths of this study, we sequenced the microbiota composition using the latest technology (Illumina platform) and annotations were made with RDP to the genus taxon level. Moreover, PA was objectively measured by accelerometry during 7 consecutive days (24 h/day) [21], and we used a cut-point-free approach to assess overall PA since PA intensities estimated from cut-points might be biased by poor calibration studies [69].

#### CONCLUSIONS

Our data showed that overall PA and time spent in vigorous PA were positively correlated with faecal microbiota diversity in young adults. Moreover, the individuals with low time spent in vigorous PA presented higher relative abundance of the *Gammaproteobacteria* class, whereas the individuals with high time spent in vigorous PA had higher relative abundance of the *Porphyromonadaceae* family. Altogether, these findings suggest that PA, especially of vigorous intensity, is related to faecal microbiota diversity and the *Gammaproteobacteria* class and *Porphyromonadaceae* family in young adults. Further studies are needed to confirm this relationship.

#### Acknowledgements

This study is part of a PhD thesis conducted within the Biomedicine Doctoral Studies Program of the University of Granada, Spain.

#### Financial Support

The study was supported by the Spanish Ministry of Economy and Competitiveness via Fondo de Investigación Sanitaria del Instituto de Salud Carlos III (PI13/01393) and PTA 12264-I, Retos de la Sociedad (DEP2016-79512-R), and European Regional Development Funds (ERDF), by the Spanish Ministry of Education (FPU16/05159 and FPU17/01523), the Fundación Iberoamericana de Nutrición (FINUT), the Redes Temáticas De Investigación Cooperativa RETIC (Red SAMID RD16/0022), InFLAMES Flagship Programme of the Academy of Finland (decision number: 337530), Fundación Alfonso Martín Escudero and Ramon y Cajal Fellowship (RYC2022-036473-I), AstraZeneca HealthCare Foundation, the University of Granada Plan Propio de Investigación 2016-Excellence actions: Unit of Excellence on Exercise and Health (UCEES), and by the Junta de Andalucía, Consejería de Conocimiento, Investigación y Universidades (ERDF, SOMM17/6107/UGR). AL and RVV are supported by the funds of European Commission through the “European funds for regional development” (EFRE) as well as by the regional Ministry of Economy, Science and Digitalization of Saxony-Anhalt as part of the “Autonomy in old Age” (AiA) research group for “LiLife” Project (Project ID: ZS/2018/11/95324). We would like to thank the team of Data Integration Center of University Medicine Magdeburg for local data-analysis solutions; they are supported by MIRACUM

and funded by the German Federal Ministry of Education and Research (BMBF) within the “Medical Informatics Funding Scheme” (FKZ 01ZZ1801H).

### Clinical Trial Information

ClinicalTrials.gov no. NCT02365129 (registered 18 February 2015).

### Author Contributions

L.O.-A., B.M.-T. and J.R.R. designed the research; L.O.-A. and B.M.-T. conducted the research; F.M.A., J.H.M., R.V.-V., A.L., J.P.-D., A.G. and I.L. provided essential reagents or materials; L.O.-A., H.X., F.M.A.,

J.H.M. and B.M.-T. analysed data or performed the statistical analysis; L.O.-A., B.M.-T. and S.R.-C. wrote the paper; J.R.R. and B.M.-T. had primary responsibility for final content. All authors critically reviewed and approved the final manuscript.

### Conflict of interest declaration

The authors have no conflicts of interest.

### Data Availability Statement

The datasets generated during the current study are available from the corresponding author on reasonable request.

## REFERENCES

- Myers J, Kokkinos P, Nyelin E. Physical activity, cardiorespiratory fitness, and the metabolic syndrome. *Nutrients*. 2019; 11(7):1652.
- Rosenbaum M, Knight R, Leibel RL. The gut microbiota in human energy homeostasis and obesity. *Trends Endocrinol Metab*. 2015; 26(9):493–501.
- Hooper LV, Gordon JI. Commensal host-bacterial relationships in the gut. *Science*. 2001; 292(5519):1115–1118.
- Monda V, Villano I, Messina A, Valenzano A, Esposito T, Moscatelli F, Viggiano A, Cibelli G, Chieffi S, Monda M, Messina G. Exercise modifies the gut microbiota with positive health effects. *Oxid Med Cell Longev*. 2017; 2017:1–8.
- Iebba V, Totino V, Gagliardi A, Santangelo F, Cacciotti F, Trancassini M, Mancini C, Cicerone C, Corazzari E, Pantanella F, Schippa S. Eubiosis and dysbiosis: The two sides of the microbiota. *New Microbiol*. 2016; 39(1):1–12.
- David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ. Diet rapidly and reproducibly alters the human gut microbiome. *Nature*. 2014; 505(7484):559–563.
- Moscatelli F, De Maria A, Marinaccio LA, Monda V, Messina A, Monacis D, Toto G, Limone P, Monda M, Messina G, Monda A, Polito R. Assessment of lifestyle, eating habits and the effect of nutritional education among undergraduate students in southern Italy. *Nutrients*. 2023; 15(13):1–13.
- Castellanos N, Diez GG, Antúnez-Almagro C, Bailén M, Bressa C, González Soltero R, Pérez M, Larrosa M. A critical mutualism – Competition interplay underlies the loss of microbial diversity in sedentary lifestyle. *Front Microbiol*. 2020; 10:3142.
- Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR, Fernandez KC, Dose H, Mori H, Patil KR, Bork P, Typas A. Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature*. 2018; 555(7698):623–628.
- Kelly TN, Bazzano LA, Ajami NJ, He H, Zhao J, Petrosino JF, Correa A, He J. Gut microbiome associates with lifetime cardiovascular disease risk profile among bogalusa heart study participants. *Circ Res*. 2016; 119(8):956–964.
- Ortiz-Alvarez L, Xu H, Martínez-Tellez B. Influence of exercise on the human gut microbiota of healthy adults: A systematic review. *Clin Transl Gastroenterol*. 2020; 11(2):1–9.
- Mitchell CM, Davy BM, Hulver MW, Neilson AP, Bennett BJ, Davy KP. Does exercise alter gut microbial composition? A systematic review. *Med Sci Sports Exerc*. 2019; 51(1):160–167.
- Clarke SF, Murphy EF, O’Sullivan O, Lucey AJ, Humphreys M, Hogan A, Hayes P, O’Reilly M, Jeffery IB, Wood-Martin R, Kerins DM, Quigley E, Ross RP, O’Toole PW, Molloy MG, Falvey E, Shanahan F, Cotter PD. Exercise and associated dietary extremes impact on gut microbial diversity. *Gut*. 2014; 63(12):1913–1920.
- Barton W, Penney NC, Cronin O, Garcia-Perez I, Molloy MG, Holmes E, Shanahan F, Cotter PD, O’Sullivan O. The microbiome of professional athletes differs from that of more sedentary subjects in composition and particularly at the functional metabolic level. *Gut*. 2018; 67(4):625–633.
- Petri C, Mascherini G, Izzicupo P, Rosati D, Carboneschi M, Smeazzetto S, Arrones LS. Gut microbiota and physical activity level: characterization from sedentary to soccer players. *Biol Sport*. 2024; 41(3):169–176.
- Bressa C, Bailén-Andrino M, Pérez-Santiago J, González-Soltero R, Pérez M, Montalvo-Lominchar MG, Maté-Muñoz JL, Domínguez R, Moreno D, Larrosa M. Differences in gut microbiota profile between women with active lifestyle and sedentary women. *PLoS One*. 2017; 12(2):1–20.
- Khan TJ, Ahmed YM, Zamzami MA, Siddiqui AM, Khan I, Baothman OAS, Mehanna MG, Kuerban A, Kaleemuddin M, Yasir M. Atorvastatin treatment modulates the gut microbiota of the hypercholesterolemic patients. *OMICS*. 2018; 22(2):154–163.
- Ruiz-Limón P, Muralidharan J, Gomez-Perez AM, Murri M, Vioque J, Corella D, Fitó M, Vidal J, Salas-Salvadó J, Torres-Collado L, Coltell O, Atzeni A, Castañer O, Bulló M, Bernal-López MR, Moreno-Indias I, Tinahones FJ. Physical activity shifts gut microbiota structure in aged subjects with overweight/obesity and metabolic syndrome. *Biol Sport*. 2023; 41(3):47–60.
- Hughes R, Burd N, Khan N, Pindus D, Holscher H. Associations between physical activity and gut microbiota composition in adults with overweight and obesity. *Curr Dev Nutr*. 2022; 6(Suppl 1):1011.
- Shephard RJ. Limits to the measurement of habitual physical activity by questionnaires. *Br J Sports Med*. 2003; 37(3):197–206.
- Sanchez-Delgado G, Martínez-Tellez B, Olza J, Aguilera CM, Labayen I, Ortega FB, Chillón P, Fernandez-Reguera C, Alcántara JMA, Martínez-Avila WD, Muñoz-Hernandez V, Acosta FM, Prados-Ruiz J, Amaro-Gahete FJ, Hidalgo-García L, Rodríguez L, Ruiz YAK, Ramírez-Navarro A, Muros-de Fuentes MA, García-Rivero Y, Sanchez-Sanchez R, de Dios Beas Jimenez J, de Teresa C, Navarrete S, Lozano R, Brea-Gomez E, Rubio-Lopez J, Ruiz MR, Cano-Nieto A, Llamas-Elvira JM, Jimenez Rios JA, Gil A, Ruiz JR. Activating brown adipose tissue through exercise (ACTIBATE) in young adults: Rationale, design and methodology. *Contemp Clin Trials*. 2015; 45(Pt B):416–425.
- Migueles JH, Cadenas-Sanchez C, Ekelund U, Delisle Nyström C, Mora-Gonzalez J, Löf M, Labayen I, Ruiz JR, Ortega FB. Accelerometer data collection and processing criteria to

- assess physical activity and other outcomes: A systematic review and practical considerations. *Sports Med.* 2017; 47(9):1821–1845.
23. Migueles JH, Rowlands A V., Huber F, Sabia S, Van Hees VT. GGIR: A research community-driven open source R package for generating physical activity and sleep outcomes from multi-day raw accelerometer data. *J Meas Phys Behav.* 2019; 2(3):188–196.
  24. Van Hees VT, Fang Z, Langford J, Assah F, Mohammad A, Da Silva ICM, Trenell MI, White T, Wareham NJ, Brage S. Autocalibration of accelerometer data for free-living physical activity assessment using local gravity and temperature: An evaluation on four continents. *J Appl Physiol.* 2014; 117(7):738–744.
  25. Acosta FM, Martinez-Tellez B, Sanchez-Delgado G, Migueles JH, Contreras-Gomez MA, Martinez-Avila WD, Merchan-Ramirez E, Alcantara JMA, Amaro-Gahete FJ, Llamas-Elvira JM, Ruiz JR. Association of objectively measured physical activity with brown adipose tissue volume and activity in young adults. *J Clin Endocrinol Metab.* 2019; 104(2):223–233.
  26. Van Hees VT, Sabia S, Anderson KN, Denton SJ, Oliver J, Catt M, Abell JG, Kivimäki M, Trenell MI, Singh-Manoux A. A novel, open access method to assess sleep duration using a wrist-worn accelerometer. *PLoS One.* 2015; 10(11):1–13.
  27. Hildebrand M, Hansen BH, van Hees VT, Ekelund U. Evaluation of raw acceleration sedentary thresholds in children and adults. *Scand J Med Sci Sports.* 2017; 27(12):1814–1823.
  28. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods.* 2016; 13(7):581–583.
  29. McMurdie PJ, Holmes S. Phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One.* 2013; 8(4):1–11.
  30. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol.* 2007; 73(16):5261–5267.
  31. Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, Brown CT, Porras-Alfaro A, Kuske CR, Tiedje JM. Ribosomal database project: Data and tools for high throughput rRNA analysis. *Nucleic Acids Res.* 2014; 42(Database issue):D633–642.
  32. Schulz C, Schütte K, Koch N, Vilchez-Vargas R, Wos-Oxley ML, Oxley APA, Vital M, Malfertheiner P, Pieper DH. The active bacterial assemblages of the upper GI tract in individuals with and without helicobacter infection. *Gut.* 2018; 67(2):216–225.
  33. Finotello F, Mastrorilli E, Di Camillo B. Measuring the diversity of the human microbiota with targeted next-generation sequencing. *Brief Bioinform.* 2018; 19(4):679–692.
  34. Lozupone CA, Knight R. Species divergence and the measurement of microbial diversity. *FEMS Microbiol Rev.* 2008; 32(4):557–578.
  35. Leo Lahti, Sudarshan Shetty. *Microbiome R package.* 2019.
  36. Gotelli NJ, Colwell RK. Estimating species richness. In: *Biological diversity: Frontiers in measurement and assessment.* Oxford, United Kingdom: Oxford University Press; 2011. p. 39–54.
  37. Magurran A. E. *Measuring biological diversity.* Oxford, United Kingdom: Blackwell's; 2004. p. 184–189.
  38. Simpson E. H. Measurement of diversity. *Nature.* 1949; 163:688.
  39. Camargo JA. New diversity index for assessing structural alterations in aquatic communities. *Bull Environ Contam Toxicol.* 1992; 48(3):428–434.
  40. Ruiz López MD, Artacho Martín-Lagos R. *Guía para estudios dietéticos: Álbum fotográfico de alimentos.* 1<sup>st</sup> ed. Granada, Spain: Editorial Universidad de Granada; 2010. p. 124–128.
  41. Sanchez-Delgado G, Martinez-Tellez B, Olza J, Aguilera CM, Labayen I, Ortega FB, Chillón P, Fernandez-Reguera C, Alcantara JMA, Martinez-Avila WD, Muñoz-Hernandez V, Acosta FM, Prados-Ruiz J, Amaro-Gahete FJ, Hidalgo-Garcia L, Rodriguez L, Ruiz YAK, Ramirez-Navarro A, Muros-de Fuentes MA, García-Rivero Y, Sanchez-Sanchez R, de Dios Beas Jimenez J, de Teresa C, Navarrete S, Lozano R, Brea-Gomez E, Rubio-Lopez J, Ruiz MR, Cano-Nieto A, Llamas-Elvira JM, Jimenez Rios JA, Gil A, Ruiz JR. Activating brown adipose tissue through exercise (ACTIBATE) in young adults: Rationale, design and methodology. *Contemp Clin Trials.* 2015; 45(Pt B):416–425.
  42. William Revelle. *Psych: Procedures for personality and psychological research.* 2017.
  43. Wei T, Simko V, Levy M, Xie Y, Jin Y, Zemla J, Freidank M, Cai J, Protivinsky T. R package corplot: Visualization of a correlation matrix. 2022.
  44. Kim YS, Unno T, Kim BY, Park MS. Sex differences in gut microbiota. *World J Mens Health.* 2020; 38(1):48–60.
  45. Palmas V, Pisanu S, Madau V, Casula E, Deledda A, Cusano R, Uva P, Vascellari S, Loviselli A, Manzin A, Velluzzi F. Gut microbiota markers associated with obesity and overweight in Italian adults. *Sci Rep.* 2021; 11(1):5532.
  46. Zmora N, Suez J, Elinav E. You are what you eat: diet, health and the gut microbiota. *Nat Rev Gastroenterol Hepatol.* 2019; 16(1):35–56.
  47. Hammer Ø, Harper D. *Paleontological data analysis.* Malden, United States: Blackwell Publishing; 2006. p. 351.
  48. Benjamini Y, Krieger AM, Yekutieli D. Adaptive linear step-up false discovery rate controlling procedures. *Biometrika.* 2006; 93(3):491–507.
  49. Kulecka M, Fraczek B, Mikula M, Zeber-Lubecka N, Karczmarski J, Paziewska A, Ambrozkiwicz F, Jagusztyń-Krynicka K, Cieszczyk P, Ostrowski J. The composition and richness of the gut microbiota differentiate the top Polish endurance athletes from sedentary controls. *Gut Microbes.* 2020; 11(5):1374–1384.
  50. Imdad S, So B, Jang J, Park J, Lee SJ, Kim JH, Kang C. Temporal variations in the gut microbial diversity in response to high-fat diet and exercise. *Sci Rep.* 2024; 14(1):3282.
  51. Estaki M, Pither J, Baumeister P, Little JP, Gill SK, Ghosh S, Ahmadi-Vand Z, Marsden KR, Gibson DL. Cardiorespiratory fitness as a predictor of intestinal microbial diversity and distinct metagenomic functions. *Microbiome.* 2016; 4(1):1–13.
  52. Laureto LMO, Cianciaruso MV, Samia DSM. *Functional diversity: an overview of its history and applicability.* *Natureza & Conservação.* 2015; 13(2):112–116.
  53. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and resilience of the human gut microbiota. *Nature.* 2012; 489(7415):220–230.
  54. Marttinen M, Ala-Jaakkola R, Laitila A, Lehtinen MJ. Gut microbiota, probiotics and physical performance in athletes and physically active individuals. *Nutrients.* 2020; 12(10):1–32.
  55. Etxeberria U, Hijona E, Aguirre L, Milagro FI, Bujanda L, Rimando AM, Martínez JA, Portillo MP. Pterostilbene-induced changes in gut microbiota composition in relation to obesity. *Mol Nutr Food Res.* 2017; 61(1):1–37.
  56. Bai J, Hu Y, Bruner DW. Composition of gut microbiota and its association with body mass index and lifestyle factors in a cohort of 7–18 years old children from the American Gut Project. *Pediatr Obes.* 2019; 14(4):1–10.
  57. Morgan XC, Tickle TL, Sokol H, Gevers D, Devaney KL, Ward D V., Reyes JA, Shah SA, LeLeiko N, Snapper SB, Bousvaros A, Korzenik J, Sands BE, Xavier RJ, Huttenhower C. Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. *Genome Biol.* 2012; 13(9):1–18.
  58. Shin NR, Whon TW, Bae JW. Proteobacteria: Microbial signature of dysbiosis in gut microbiota. *Trends Biotechnol.* 2015; 33(9):496–503.

59. Yang Y, Shi Y, Wiklund P, Tan X, Wu N, Zhang X, Tikkanen O, Zhang C, Munukka E, Cheng S. The association between cardiorespiratory fitness and gut microbiota composition in premenopausal women. *Nutrients*. 2017; 9(8):792.
60. Baldanzi G, Sayols-Baixeras S, Ekblom-Bak E, Ekblom Ö, Dekkers KF, Hammar U, Nguyen D, Ahmad S, Ericson U, Arvidsson D, Börjesson M, Johanson PJ, Smith JG, Bergström G, Lind L, Engström G, Ärnlov J, Kennedy B, Orho-Melander M, Fall T. Accelerometer-based physical activity is associated with the gut microbiota in 8416 individuals in SCAPIS. *EBioMedicine*. 2024; 100:104989.
61. Quiroga R, Nistal E, Estébanez B, Porras D, Juárez-Fernández M, Martínez-Flórez S, García-Mediavilla MV, de Paz JA, González-Gallego J, Sánchez-Campos S, Cuevas MJ. Exercise training modulates the gut microbiota profile and impairs inflammatory signaling pathways in obese children. *Exp Mol Med*. 2020; 52(7):1048–1061.
62. Munukka E, Ahtiainen JP, Puigbó P, Jalkanen S, Pahkala K, Keskitalo A, Kujala UM, Pietilä S, Hollmén M, Elo L, Huovinen P, D'Auria G, Pekkala S. Six-week endurance exercise alters gut metagenome that is not reflected in systemic metabolism in over-weight women. *Front Microbiol*. 2018; 9:1–16.
63. Liang R, Zhang S, Peng X, Yang W, Xu Y, Wu P, Chen J, Cai Y, Zhou J. Characteristics of the gut microbiota in professional martial arts athletes: A comparison between different competition levels. *PLoS One*. 2019; 14(12):1–13.
64. de Oliveira Neves VG, de Oliveira DT, Oliveira DC, Oliveira Perucci L, dos Santos TAP, da Costa Fernandes I, de Sousa GG, Barboza NR, Guerra-Sá R. High-sugar diet intake, physical activity, and gut microbiota crosstalk: Implications for obesity in rats. *Food Sci Nutr*. 2020; 8(10):5683–5695.
65. Liu TW, Park YM, Holscher HD, Padilla J, Scroggins RJ, Welly R, Britton SL, Koch LG, Vieira-Potter VJ, Swanson KS. Physical activity differentially affects the cecal microbiota of ovariectomized female rats selectively bred for high and low aerobic capacity. *PLoS One*. 2015; 10(8):1–17.
66. Fernández J, Fernández-Sanjurjo M, Iglesias-Gutiérrez E, Martínez-Cambor P, Villar CJ, Tomás-Zapico C, Fernández-García B, Lombó F. Resistance and endurance exercise training induce differential changes in gut microbiota composition in murine models. *Front Physiol*. 2021; 12.
67. Parker BJ, Wearsch PA, Veloo ACM, Rodriguez-Palacios A. The genus *alisticipes*: Gut bacteria with emerging implications to inflammation, cancer, and mental health. *Front Immunol*. 2020; 11:1–15.
68. Wyatt M, Greathouse KL. Targeting dietary and microbial tryptophan-indole metabolism as therapeutic approaches to colon cancer. *Nutrients*. 2021; 13(4):1–23.
69. Migueles JH, Cadenas-Sanchez C, Tudor-Locke C, Löf M, Esteban-Cornejo I, Molina-Garcia P, Mora-Gonzalez J, Rodriguez-Ayllon M, Garcia-Marmol E, Ekelund U, Ortega FB. Comparability of published cut-points for the assessment of physical activity: Implications for data harmonization. *Scand J Med Sci Sports*. 2019; 29(4):566–574.