



Universiteit
Leiden
The Netherlands

Uncovering the impact of center of rotation of angulation location on high tibial osteotomy in knee osteoarthritis: a potential pathway for improved outcomes

Janssen, Y.; Nguyen, H.C.; Custers, R.J.H.; Egmond, N. van; Kruyt, M.C.; Sakkers, R.J.B.; ... ; Bax, E.A.

Citation

Janssen, Y., Nguyen, H. C., Custers, R. J. H., Egmond, N. van, Kruyt, M. C., Sakkers, R. J. B., ... Bax, E. A. (2026). Uncovering the impact of center of rotation of angulation location on high tibial osteotomy in knee osteoarthritis: a potential pathway for improved outcomes. *Cartilage*, 1-8. doi:10.1177/19476035261420279

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/4300368>

Note: To cite this publication please use the final published version (if applicable).

Uncovering the Impact of Center of Rotation of Angulation Location on High Tibial Osteotomy in Knee Osteoarthritis: A Potential Pathway for Improved Outcomes

CARTILAGE

1–8

© The Author(s) 2026

DOI: 10.1177/19476035261420279

journals.sagepub.com/home/CAR



Yannick Janssen¹, H. Chien Nguyen², Roel J. H. Custers¹, Nienke van Egmond¹, Moyo C. Kruyt¹, Ralph J. B. Sakkers¹, Jaap Thooft², Margreet Kloppenburg³, Francisco J. Blanco⁴, Ida K. Haugen⁵, Francis Berenbaum⁶, Simon C. Mastbergen⁷, Harrie Weinans^{1,8}, and Eva A. Bax¹

Abstract

Objective. Lower limb malalignment accelerates the progression of knee osteoarthritis (KOA). Knee realignment osteotomy is a well-established treatment for unicompartmental KOA with malalignment. Traditional planning in KOA patients corrects deformities with an osteotomy at the metaphysis but overlooks Paley's approach, which targets the center of rotation angulation (CORA). Osteotomy at the metaphysis may induce secondary translational deformities, which remain unstudied in KOA patients. This study aims to identify the CORA in KOA patients with tibial malalignment. **Methods.** Thirty tibiae (10 varus, 10 neutral, 10 valgus) from the IMI-APPROACH cohort were analyzed using computed tomography (CT) scans. The CORA, defined as the intersection of the proximal and distal mechanical axes, was identified. Translational deformity was calculated by multiplying the CORA-to-osteotomy distance by the tangent of the correction angle. **Results.** Among the varus tibiae, 9 out of 10 CORAs were located in the diaphysis, while 8 out of 10 valgus tibiae had their CORA in the diaphysis. When osteotomies were performed in the proximal metaphysis instead of the CORA location, secondary translational deformities of up to 3 cm were induced. **Conclusion.** In KOA patients with tibial malalignment, the CORA is predominantly located in the diaphysis rather than in the proximal metaphysis, where osteotomies are typically performed. This discrepancy leads to iatrogenic translational deformities. Future research should investigate the clinical impact of these deformities to optimize osteotomy planning and potentially improve long-term surgical outcomes.

Keywords

CORA, lower limb malalignment, osteoarthritis, osteotomy

Introduction

Osteoarthritis (OA) affected 595 million people globally in 2019,¹ with knee osteoarthritis (KOA) being the most common type.² As obesity and life expectancy rise,^{1,3–6} KOA prevalence increases.² Lower limb malalignment is a risk factor for KOA progression,^{7–9} shifting the knee's mechanical axis and accelerating cartilage degeneration.^{8–11} Realignment osteotomies are a well-established treatment for younger patients to postpone knee arthroplasty,^{12–14} which is important as they have a higher risk of revision surgery.¹⁵

Preoperative planning for knee osteotomy is essential for optimal outcomes.¹⁶ Early methods like the Fujisawa point,

Miniaci line, and Dugdale method,¹⁷ have evolved into the current planning nomenclature proposed by Paley.¹⁸ Modern planning methods focus on calculating the desired wedge height for knee osteotomies, with corrections typically performed at the tibial and femoral metaphyseal ends.^{17,19,20} Interestingly, Paley's¹⁸ approach extends beyond this conventional method by emphasizing the correction of deformities at their origin, a concept well-known in reconstructive surgery.

Building on this concept, the center of rotation angulation (CORA) is the point where the mechanical axes of a deformed bone intersect.^{18,21} By correcting deformities at the CORA, the risk of introducing secondary translational deformities is minimized.¹⁸ Tibial osteotomies for unicompartmental KOA



Creative Commons CC BY: This article is distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by/4.0/>) which permits any use, reproduction and distribution of

the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

are preferably performed at the proximal tibial end, as this region is characterized by superior bone healing due to its high trabecular density and vascularization²²⁻²⁴ compared with the mid-diaphysis. Consequently, in these procedures, potential translation of the bone is often assumed and remains unaddressed. Therefore, our study aims to identify the CORA of patients diagnosed with KOA and associated tibial malalignment. This will provide crucial insights into the potential occurrence of iatrogenic deformities.

Methods

Patients

In the prospective Applied Public-Private Research enabling OsteoArthritis Clinical Headway (IMI-APPROACH) cohort, 297 KOA participants from 5 European centers were included.²⁵⁻²⁹ Some of these participants also exhibited malalignment of the femur or tibia. Detailed inclusion and exclusion criteria have been previously published.³⁰ The study was approved by Institutional Review Boards, in accordance with all relevant ethical and legal regulations. The study was registered under clinicaltrials.gov number: NCT03883568, and informed consent was obtained from all participants.

Imaging Assessment

All patients underwent low-dose whole-body computed tomography (CT) scans. The tibia and fibula were segmented from the CT scans using validated software (Mimics; Materialise, Leuven, Belgium). Bone geometry analyses were performed in 3-matic (Materialise). The analyses involved a semi-automated method as the functions were scripted in Python language (3-match plugin).

3D tibial coordinate system. A patient-specific 3D coordinate system was constructed per tibiae. The mechanical axis ran from the tibial eminences to the distal tibial plafond. The transversal plane had its origin at the tibial eminences, with the mechanical axis as normal vector. The sagittal plane,




Figure 1. Definition of the CORA and the mMPA. **(A)** The CORA (center of rotation angulation) is the intersection of the proximal mechanical axis (PMA, black dotted line) and the distal mechanical axis (DMA, gray line) of the tibia. **(B)** The mMPA (mechanical medial proximal tibial angle) is the angle formed between the mechanical axis of the tibia and the tangent of the proximal tibial plateau on the coronal plane.

perpendicular to the transversal plane, crossed the posterior cruciate ligament attachment and medial tuberosity border (Akagi's line).³¹ The coronal plane, perpendicular to both transversal and sagittal planes, originated at the tibial eminences.

CORA calculation. The method fitted a plane to the articulating surface of the medial and lateral tibial plateaus (proximal tibial plane) and projected a line distally originating from the center of the eminence spines at an angle of 87° (coronal view) relative to the proximal tibial plane (proximal mechanical axis (PMA)) (Fig. 1A). In addition, a plane was fitted to the articulating surface of the distal tibial plafond (distal tibial plane), and a line was projected proximally at an 89° angle (coronal view) relative to the distal

¹Department of Orthopaedic Surgery, University Medical Centre Utrecht, Utrecht, The Netherlands

²3D Lab, University Medical Centre Utrecht, Utrecht, The Netherlands

³Departments of Rheumatology and Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

⁴Grupo de Investigación de Reumatología (GIR), INIBIC-Complejo Hospitalario Universitario de A Coruña, Centro Interdisciplinar de Química y Biología, CICA-UDC, Universidad de A Coruña, A Coruña, Spain

⁵Center for Treatment of Rheumatic and Musculoskeletal Diseases (REMEDY), Diakonhjemmet Hospital, Oslo, Norway

⁶Department of Rheumatology, Sorbonne University, INSERM CRSA, AP-HP Saint-Antoine Hospital, Paris, France

⁷Department of Rheumatology & Clinical Immunology, University Medical Center Utrecht, Utrecht, The Netherlands

⁸Department Biomechanical Engineering, Faculty of Mechanical Engineering, TU Delft, Delft, The Netherlands

Investigation performed at University Medical Center Utrecht, The Netherlands

Corresponding Author:

Roel J. H. Custers, Department of Orthopaedic Surgery, University Medical Centre Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands.
Email: r.j.h.custers@umcutrecht.nl

tibial plane (distal mechanical axis (DMA)), originating from the center of the distal tibial plafond (**Fig. 1A**). The crossing of the two lines represented the CORA of the coronal tibial deformity, and the location of this CORA was calculated with respect to the tibial eminence center (knee joint center) (**Fig. 1A**).

Bone deformity. Coronal bone deformities of the tibia, including the mechanical medial proximal tibial angle (mMPTA), was assessed using Paley's¹⁸ method. The mMPTA was defined as the angle (coronal view) between the mechanical axis of the tibia and the tangent to the proximal tibial plateau (**Fig. 1B**). Neutral mMPTA ranged from 85° to 90°. A varus mMPTA was defined as <85°, while a valgus mMPTA was >90°. A total of 30 tibiae were included: 10 with a neutral mMPTA, 10 with a varus mMPTA, and 10 with a valgus mMPTA. Participants with tibial deformity were randomly selected to ensure a representative distribution of the mMPTA across the different alignment groups (neutral, varus, and valgus).

The Concept of Secondary Translational Deformities in Osteotomies

In osteotomies, translation refers to the sideward displacement of the distal segment of the tibia. This occurs when the osteotomy is not performed at the CORA, which is the optimal location for performing an osteotomy without creating a secondary deformity.²¹ The reason for this translation lies in the kinematics of the correction. A bone deformity creates an angular malalignment, and when an osteotomy is performed at a distance from the CORA, the bone must be corrected by rotating or angulating the distal segment to realign it with the proximal segment. This causes a sideward shift in the direction of the angular correction. Stated otherwise, when an osteotomy is performed outside the CORA, angular correction alone can render the PMA and DMA parallel while still malaligned, necessitating medial translation of the distal segment for complete alignment (see **Figure 2**).

Translation is influenced by 2 factors: the distance from the CORA to the osteotomy site³² and the correction angle.³² Greater distance and larger correction angles increase translational displacement. The translation can be calculated by

$$T = d \times \tan(\theta) \quad (1)$$

In this equation, T refers to the sideward displacement of the distal segment, d is the distance from the CORA to the osteotomy site, and θ is the correction angle (degrees) (**Fig. 2**). For a varus tibia, the correction aimed to realign the mMPTA to 90°, whereas for a valgus tibia, the correction targeted an mMPTA of 85°. This aligns with clinical practice, where we apply slight overcorrection in KOA patients.³³




Figure 2. The concept of iatrogenic translational deformities in osteotomies when the osteotomy is performed outside the CORA. The PMA is shown in red, the DMA in blue, and the CORA is marked by the green circle. **(A)** The tibia with deformity. **(B)** High tibial osteotomy performed away from the CORA leads to parallel PMA and DMA. **(C)** Translation is required to realign the PMA and DMA. **(D)** No translation is necessary when the osteotomy is performed at the CORA.

CORA Location

In addition to assessing the magnitude of the secondary deformity, the location of the CORA was also evaluated. The location of the CORA on the tibia was defined as a percentage of the tibial length, with 100% representing the proximal end and 0% representing the distal end. This method allowed for a descriptive analysis of CORA positions in tibiae with varus and valgus deformities. By comparing these positions, differences in CORA location between the 2 deformity groups were identified.

Statistical Analysis

All statistical analyses were conducted using Statistical Package for the Social Sciences (SPSS) Version 29.0 software. Descriptive statistics were computed, including means and standard deviations (SD) for continuous variables, and numbers and percentages for categorical data.

Results

Locations of CORA in the Tibia

A total of 30 tibiae were included in this study, consisting of 10 individuals with varus alignment of the tibiae, 10 with neutral alignment of the tibiae, and 10 with valgus alignment, based on the mMPTA, all of whom presented with early-stage KOA. Most patients were female (73%), and the




Figure 3. Examples of CORA determinations in different tibiae. **(A)** Neutral tibia, where the PMA (solid line) and DMA (dotted line) run nearly parallel, indicating no CORA. **(B)** Varus tibia with an example where the CORA is in both the metaphyseal and diaphyseal regions. **(C)** Varus tibia with another example of the CORA located in both the metaphyseal and diaphyseal regions.

mean age was 62.9 ± 8.0 years. The mean mMPTA was $87.3^\circ \pm 3.3^\circ$.

In neutral tibial alignment, the mean mMPTA was $87.5^\circ \pm 1.4^\circ$. Since there was no deformity in these bones, the PMA and DMA (**Fig. 1**) were nearly parallel, and no CORA was present (**Fig. 3A**). In varus alignment, the mean mMPTA was $83.4^\circ \pm 1.0^\circ$. The CORA in the varus group was located within the proximal metaphyseal region in 1 case and within the diaphyseal region in 9 cases (**Fig. 3B**). In the valgus alignment group, the mean mMPTA was $91.0^\circ \pm 0.7^\circ$. The CORA of the valgus group was located within the metaphyseal region in 2 cases and within the diaphyseal region in 8 cases (**Fig. 3C**).

In the varus alignment group, the mean CORA was located at $54.9 \pm 18.8\%$ of the tibial length, ranging from 11.9 to 78.5% (**Fig. 4**). In the valgus alignment group, the mean CORA was situated at $33.0 \pm 15.9\%$ of the tibial length, with a range from 4.1 to 51.6% (**Fig. 4**). In the varus group, the CORA was located more distally than in the valgus group (**Fig. 4**).

Secondary Translational Deformities in Osteotomies

For an angular correction of the varus mMPTA to 90° , the mean translation when performing the osteotomy on a varus tibia at the metaphysis, as opposed to at the CORA site, was $2.09 \text{ cm} \pm 0.85 \text{ cm}$, with a range from 0.17 cm to 3.04 cm. In contrast, for an angular correction of the mMPTA to 85° when correcting a valgus tibia, the mean translation was




Figure 4. CORA locations relative to tibial length, presented separately for varus and valgus tibiae, with 100% representing the proximal end and 0% representing the distal end.




Figure 5. Secondary translation deformity values are plotted against the distances measured between the CORA and the high tibial osteotomy plane. Distinctions were made for different correction angles, highlighting the relationship between the secondary translation and the correction magnitude.

$0.96 \text{ cm} \pm 0.56 \text{ cm}$, with a range from 0.23 cm to 1.81 cm. **Figure 5** illustrates the relationship between the distances from the CORA to the high tibial osteotomy plane and the corresponding calculated secondary translational deformity.

Discussion

This study aimed to identify the CORA in KOA patients with tibial malalignment and explore secondary translation in tibial osteotomies not performed at the CORA, all conducted in a cohort comprising patients with KOA. A key finding was that 85% of the CORAs were in the diaphysis, leading to secondary translational deformities of up to 3 cm, a novel finding not previously reported in the knee osteotomy literature. Identifying and targeting the CORA during

osteotomy planning may improve knee realignment and postoperative outcomes.

The clinical impact of these secondary translational deformities is unclear, as no studies have focused on the effect of CORA locations in KOA patients with tibial malalignment. Potential consequences may include altered gait mechanics, increased joint wear, and unfavorable force distribution within the affected compartment; however, these hypotheses require further investigation. Most existing research on osteotomies in KOA focuses on the mechanical hip-knee-ankle angle.^{9,34,35} What we do know is that osteotomy is an effective treatment for relatively young KOA patients, as it can delay the need for knee arthroplasty by over 10 years.^{12,13,36,37} Predictors for an increased likelihood of conversion to arthroplasty include radiographic OA severity, pain, female sex, age, and body mass index (BMI), with radiographic OA severity being the strongest predictor.^{36,38} While osteotomies are effective, improving long-term outcomes is crucial, especially as KOA increases in prevalence.² Optimizing osteotomy techniques could improve knee function and patient satisfaction, but further research is needed to validate these findings and assess their impact on clinical outcomes and the long-term effectiveness of osteotomies in KOA treatment.

In addition to the 30 tibiae analyzed, 10 extra tibiae exhibited translational deformities (S-shaped) without prior osteotomy (Appendix). These deformities can be conceptualized as a displacement in which the distal segment shifts relative to the proximal segment,^{18,21} which is also observed when an osteotomy is not performed at the CORA. In valgus mMPTA patients, the deformities show a varus mechanical lateral distal tibial angle. In such cases, 2 osteotomies with opposing corrections at each level are recommended.²¹ These translational deformities, common in pediatric patients,³⁹⁻⁴¹ can cause abnormal gait, joint wear, and premature OA.⁴¹ To prevent these long-term complications, corrective osteotomies are often performed in children to align the bones properly.

Historically, osteotomies are performed in the metaphysis^{17,42} due to better bone healing compared with the diaphysis.²²⁻²⁴ The diaphysis consists primarily of dense cortical bone with less robust intraosseous blood supply, whereas the metaphysis is characterized by more metabolically active and vascular trabecular bone.²² This difference likely contributes to higher rates of nonunion and hinge fractures in diaphyseal osteotomies.^{23,24} Despite the risk of translation, the benefits of fracture healing often outweigh this concern. In practice, secondary translation is often overlooked, as the mechanical axis is shifted, which is the primary goal of the procedure.

The concept of secondary translational deformity has been described in the literature,^{18,21,32,41} but no studies have focused on CORA location in KOA patients with tibial malalignment. Barksfield and Monsell³² concluded that

translational deformities can be predicted by the angular correction and distance from the CORA. However, their study did not focus on KOA patients and involved smaller simulated distances. Our study shows that these distances can reach up to 3 cm, leading to larger translational deformities that require further investigation—although this was shown in a relatively small sample size. Therefore, future research is needed to evaluate CORA locations in a larger cohort of KOA patients with lower limb malalignment who are candidates for osteotomy. Future research should also assess the long-term effects of these deformities on knee function, survival rates, and patient outcomes. Moreover, the influence of these deformities on force distribution in the affected compartment should be explored, and whether performing osteotomies away from the CORA enhances or diminishes this effect.

High tibial osteotomy is indicated primarily to correct varus or valgus malalignment in unicompartmental KOA.⁴³ Feucht *et al.*⁴⁴ demonstrated that mild varus malalignment often results from a deformity in the joint line convergence angle (JLCA), not the bones themselves. In these cases, correcting the JLCA along with tibial realignment is recommended. Osteotomy in the proximal tibia is advantageous for JLCA deformities, but whether osteotomy is the right approach for JLCA deformities alone remains unclear. In clinical practice, during a high tibial osteotomy, correction of the JLCA is performed alongside tibial realignment. For patients undergoing osteotomy without bone deformities but with a deformity of the JLCA, performing the osteotomy in the proximal tibia is advantageous, as the CORA would lie within the knee joint for these patients. However, the question remains whether osteotomy is the appropriate indication for patients with only a deformity of the JLCA.³⁸ Further research is needed to better understand the optimal approach for patients with JLCA deformities.





Several limitations of this study should be acknowledged. The study included 30 tibiae, a small sample size, with the goal of raising awareness among orthopedic surgeons that the CORA in tibial malalignment is typically not always located in the proximal metaphysis. In addition, the mean age of our cohort was higher than the typical age at which high tibial osteotomy is performed, which may affect the generalizability of our findings. Our study focused solely on patients with tibial malalignment. Future research should also examine its potential impact on femoral double-level osteotomies. Second, CORA determination was based on CT imaging, not whole-leg radiography, which remains the clinical gold standard. However, CT scans eliminate positioning factors,⁴⁵ and Roth *et al.*⁴⁶ found minimal differences between weight-bearing and non-weight-bearing conditions for bony alignment. Third, in coronal alignment osteotomies for KOA patients, the osteotomy extends to the hinge point. Consequently, operating at the CORA inevitably induces minimal secondary translational deformity, as

the center of rotation does not coincide with the CORA. Finally, the standardized correction angles used in this study may differ from individualized surgical planning, which could lead to different magnitudes of translational shifts and alignment outcomes. In clinical practice, surgeons increasingly avoid over-corrections, instead aiming for a more neutral alignment. Moreover, our analysis focused on secondary translation following a tibial osteotomy, whereas larger corrections in current practice are often managed with double-level osteotomies, which distribute the correction across two bones and therefore reduce the degree of secondary translation.

Conclusion

This study identified the CORA in patients with KOA and tibial malalignment, highlighting the secondary translational deformities that arise because of not performing the osteotomy at the CORA. We concluded that 85% of CORAs were in the diaphysis and not located in the proximal metaphysis, leading to secondary translational deformities of up to 3 cm after high tibial osteotomy. Future studies should focus on the clinical implications to possibly improve both its effectiveness and long-term sustainability of osteotomies.

ORCID iDs

H. Chien Nguyen  <https://orcid.org/0000-0001-5436-7981>
 Francisco J. Blanco  <https://orcid.org/0000-0001-9821-7635>
 Simon C. Mastbergen  <https://orcid.org/0000-0002-8825-6486>
 Eva A. Bax  <https://orcid.org/0000-0003-3892-6987>

Ethical Considerations

All procedures performed in the IMI-Approach study were conducted in compliance with the protocol, Good Clinical Practice (GCP), the Declaration of Helsinki, and the applicable ethical and legal regulatory requirements (for all countries involved), and is registered under clinicaltrials.gov identifier: NCT03883568. Informed consent was obtained from all individual participants included in the study.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Innovative Medicines Initiative Joint Undertaking under grant agreement no 115770, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in-kind contribution. See www.imi.europa.eu and www.approachproject.eu.

Declaration of Conflicting Interests

The authors declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: These authors declare that they have no conflicts of interest: YJ, CN, RC, NvE, MCK, JT, and EB. RS serves as a member of the

Scientific Advisory Board of Anna Fonds and the Medical Advisory Board of the OI Foundation Europe (2020-2025). He is Vice President of the European Paediatric Orthopaedic Society. The author is also a minority shareholder (<5%) in Replasia BV and UPlanner BV. MK reports all support for the present manuscript from IMI APPROACH. Additional funding was received from the Dutch Arthritis Society. MK receives royalties from Wolters Kluwer and Springer Verlag and consulting fees from Pfizer, UCB, CHDR, GSK, Novartis, and Peptinov. Honoraria for lectures and educational activities were received from Novartis. MK has held leadership roles as member of the OARSI board (2017-2022), chair of the EULAR Advocacy Committee, and President of the Dutch Society for Rheumatology. FJB reports participation in clinical trials funded by Abbvie, Bristol Myers Squibb, Roche, Servier, Novartis, Horizon Therapeutics Ireland DAC, ITF Research Pharma S.L.U., GSK Research, Pfizer, Sanofi-Aventis, Grünenthal, Lilly, Merck Healthcare KGaA, LG Chem Ltd., UCB, Janssen, Amgen, Regeneron, Alkem Laboratories Ltd., Sun Pharma Global FZE, and Kiniksa Pharmaceuticals GmbH. FJB has received honoraria for lectures and educational activities from Medicamenta-Ecuador, Grünenthal, and Asofarma. Support for attending meetings and travel was provided by UCB, Abbvie, and Celgen. FJB serves on a Data Safety Monitoring or Advisory Board for Grünenthal. IH reports consulting fees from Novartis, GSK, and Grünenthal, and honoraria for lectures and educational activities from Abbvie. FB reports consulting fees from Grünenthal, GSK, Eli Lilly, Novartis, Pfizer, Servier, 4P Pharma, and Peptinov; honoraria for lectures and educational activities from Viatrix, Pfizer, and Zoetis; and support for attending meetings and travel from Nordic Pharma. FB holds patents with 4Moving Biotech and has stock or stock options in 4P Pharma and 4Moving Biotech. FB serves on Data Safety Monitoring or Advisory Boards for AstraZeneca, Sun Pharma, and Nordic Bioscience. SCM reports grants from ReumaNederland. HW has received research funding from Interreg (EFRO), OA-Inject (NWO), 3DHip (Eurostars), Dartbac (NWO), Kansen voor West (Province of Utrecht), and the EU (Rescue grant). The author is a minority shareholder in Replasia, Presurgeo, Amotio, and Preimure.

References

1. GBD 2021 Osteoarthritis Collaborators. Global, regional, and national burden of osteoarthritis, 1990–2020 and projections to 2050: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet Rheumatol.* 2023;5(9):e508-e522. doi:10.1016/S2665-9913(23)00163-7.
2. Cui A, Li H, Wang D, Zhong J, Chen Y, Lu H. Global, regional prevalence, incidence and risk factors of knee osteoarthritis in population-based studies. *EclinicalMedicine.* 2020;29-30:100587. doi:10.1016/j.eclinm.2020.100587.
3. Georgiev T, Angelov AK. Modifiable risk factors in knee osteoarthritis: treatment implications. *Rheumatol Int.* 2019;39(7):1145-57. doi:10.1007/S00296-019-04290-Z.
4. Driban JB, Harkey MS, Barbe MF, Ward RJ, MacKay JW, Davis JE, *et al.* Risk factors and the natural history of accelerated knee osteoarthritis: a narrative review. *BMC Musculoskelet Disord.* 2020;21(1):1-11. doi:10.1186/S12891-020-03367-2/TABLES/2.
5. Mehrzad R. Definition and introduction to epidemiology of obesity. In: Mehrzad R, editor. *Obesity: global impact and*

- epidemiology. Amsterdam, The Netherlands: Elsevier; 2020. p. 1-6.
6. Grinin L, Grinin A, Korotayev A. Global aging and our futures. *World Futures*. 2023;79(5):536-56. doi:10.1080/02604027.2023.2204791.
 7. Tanamas S, Hanna FS, Cicuttini FM, Wluka AE, Berry P, Urquhart DM. Does knee malalignment increase the risk of development and progression of knee osteoarthritis? a systematic review. *Arthritis Care Res (Hoboken)*. 2009;61(4):459-67. doi:10.1002/art.24336.
 8. Khury F, Oltmanns M, Unseld T, Fuchs M, Reichel H, Faschingbauer M. Which knee phenotypes exhibit the strongest correlation with cartilage degeneration? *Clin Orthop Relat Res*. 2024;482(3):500-10. doi:10.1097/CORR.0000000000002831.
 9. Sharma L, Song J, Dunlop D, Felson D, Lewis CE, Segal N, et al. Varus and valgus alignment and incident and progressive knee osteoarthritis. *Ann Rheum Dis*. 2011;69(11):1940-5. doi:10.1136/ard.2010.129742.Varus.
 10. Glyn-Jones S, Palmer AJR, Agricola R, Price AJ, Vincent TL, Weinans H, et al. Osteoarthritis. *Lancet*. 2015;386(9991):376-87. doi:10.1016/S0140-6736(14)60802-3.
 11. Hunter D, Bierma-Zeinstra S. Osteoarthritis. *Lancet*. 2019;393(10182):1745-59. doi:10.1016/S0140-6736(19)30417-9.
 12. Ollivier B, Berger P, Depuydt C, Vandenneucker H. Good long-term survival and patient-reported outcomes after high tibial osteotomy for medial compartment osteoarthritis. *Knee Surg Sports Traumatol Arthrosc*. 2021;29(11):3569-84. doi:10.1007/s00167-020-06262-4.
 13. Shivji FS, Foster A, Risebury MJ, Wilson AJ, Yassen SK. Ten-year survival rate of 89% after distal femoral osteotomy surgery for lateral compartment osteoarthritis of the knee. *Knee Surg Sports Traumatol Arthrosc*. 2021;29(2):594-9. doi:10.1007/s00167-020-05988-5.
 14. Hoorntje A, Pronk Y, Brinkman JM, van Geenen RCI, van Heerwaarden RJ. High tibial osteotomy versus unicompartmental knee arthroplasty for Kellgren–Lawrence grade 3–4 knee osteoarthritis in younger patients: comparable improvements in patient-reported outcomes, adjusted for osteoarthritis grade and sex. *Knee Surg Sports Traumatol Arthrosc*. 2023;31(11):4861-70. doi:10.1007/s00167-023-07526-5.
 15. Price AJ, Alvand A, Troelsen A, Katz JN, Hooper G, Gray A, et al. Knee replacement. *Lancet*. 2018;392(10158):1672-82. doi:10.1016/S0140-6736(18)32344-4.
 16. Matsushita T, Watanabe S, Araki D, Nagai K, Hoshino Y, Kanzaki N, et al. Differences in preoperative planning for high-tibial osteotomy between the standing and supine positions. *Knee Surg Relat Res*. 2021;33(1):8. doi:10.1186/s43019-021-00090-7.
 17. Smith JO, Wilson AJ, Thomas NP. Osteotomy around the knee: evolution, principles and results. *Knee Surg Sports Traumatol Arthrosc*. 2013;21(1):3-22. doi:10.1007/s00167-012-2206-0.
 18. Paley D. Principles of deformity correction. Berlin, Germany: Springer-Verlag GmbH & Co. KG; 2002. p. 816.
 19. Lee DC, Byun SJ. High tibial osteotomy. *Knee Surg Relat Res*. 2012;24(2):61-9. doi:10.5792/ksrr.2012.24.2.61.
 20. Sabzevari S, Ebrahimpour A, Roudi MK, Kachooei AR. High tibial osteotomy: a systematic review and current concept. *Arch Bone Jt Surg*. 2016;4(3):204-12.
 21. Paley D, Herzenberg JE, Tetsworth K, McKie J, Bhav A. Deformity planning for frontal and sagittal plane corrective osteotomies. *Orthop Clin North Am*. 1994;25(3):425-65. doi:10.1016/S0030-5898(20)31927-1.
 22. Liu DS, Snyder BD, Mahan ST. Fracture nonunion and delayed union. *J Pediatr Soc North Am*. 2024;7:100058. doi:10.1016/j.jposna.2024.100058.
 23. Miltenberg B, Puzitiello RN, Ruelos VCB, Masood R, Pagani NR, Moverman MA, et al. Incidence of complications and revision surgery after high tibial osteotomy: a systematic review. *Am J Sports Med*. 2024;52(1):258-68. doi:10.1177/03635465221142868.
 24. Nicholson JA, Makaram N, Simpson A, Keating JF. Fracture nonunion in long bones: a literature review of risk factors and surgical management. *Injury*. 2021;52(suppl 2):S3-S11. doi:10.1016/j.injury.2020.11.029.
 25. Damman W, Liu R, Kroon FPB, Reijnen M, Huizinga TWJ, Rosendaal FR, et al. Do comorbidities play a role in hand osteoarthritis disease burden? data from the hand osteoarthritis in secondary care cohort. *J Rheumatol*. 2017;44(11):1659-66. doi:10.3899/JRHEUM.170208.
 26. Wesseling J, Boers M, Viergever MA, Hilberdink WK, Lafeber FP, Dekker J, et al. Cohort profile: Cohort Hip and Cohort Knee (CHECK) study. *Int J Epidemiol*. 2016;45(1):36-44. doi:10.1093/ije/dyu177.
 27. Magnusson K, Hagen KB, Østerås N, Nordsletten L, Natvig B, Haugen IK. Diabetes is associated with increased hand pain in erosive hand osteoarthritis: data from a population-based study. *Arthritis Care Res (Hoboken)*. 2015;67(2):187-95. doi:10.1002/ACR.22460.
 28. Sellam J, Maheu E, Crema MD, Touati A, Courties A, Tuffet S, et al. The DIGICOD cohort: a hospital—based observational prospective cohort of patients with hand osteoarthritis—methodology and baseline characteristics of the population. *Joint Bone Spine*. 2021;88(4):105171. doi:10.1016/J.JBSPIN.2021.105171.
 29. Oreiro-Villar N, Fernandez-Moreno M, Cortes-Pereira E, Vázquez-Mosquera ME, Relaño S, Pértega-Díaz S, et al. Metabolic syndrome and knee osteoarthritis. Impact on the prevalence, severity incidence and progression of the disease. *Osteoarthritis Cartilage*. 2017;25:S286-S287. doi:10.1016/J.JOCA.2017.02.483.
 30. van Helvoort EM, van Spil WE, Jansen MP, Welsing PMJ, Kloppenburg M, Loef M, et al. Cohort profile: the Applied Public-Private Research enabling OsteoArthritis Clinical Headway (IMI-APPROACH) study: a 2-year, European, cohort study to describe, validate and predict phenotypes of osteoarthritis using clinical, imaging and biochemical markers. *BMJ Open*. 2020;10(7):e035101. doi:10.1136/bmjopen-2019-035101.
 31. Saffarini M, Nover L, Tandogan R, Becker R, Moser LB, Hirschmann MT, et al. The original Akagi line is the most reliable: a systematic review of landmarks for rotational alignment of the tibial component in TKA. *Knee Surg Sports Traumatol Arthrosc*. 2019;27(4):1018-27. doi:10.1007/s00167-018-5131-z.

32. Barksfield RC, Monsell FP. Predicting translational deformity following opening-wedge osteotomy for lower limb realignment. *Strategies Trauma Limb Reconstr.* 2015;10(3):167-73. doi:10.1007/s11751-015-0232-4.
33. Van den Bempt M, Van Genechten W, Claes T, Claes S. How accurately does high tibial osteotomy correct the mechanical axis of an arthritic varus knee? a systematic review. *Knee.* 2016;23(6):925-35. doi:10.1016/j.knee.2016.10.001.
34. Kawashima I, Kawai R, Ishizuka S, Hiraiwa H, Tsukahara T, Imagama S. Association between knee alignment and meniscal tear in pediatric patients with anterior cruciate ligament injury. *J Bone Joint Surg Am.* 2021;103(17):1604-10. doi:10.2106/JBJS.20.01219.
35. Song JH, Bin SI, Kim JM, Lee BS, Lee CR. Influence of varus alignment on survivorship after lateral meniscal allograft transplantation. *Am J Sports Med.* 2020;48(6):1374-8. doi:10.1177/0363546520914592.
36. Primeau CA, Birmingham TB, Leitch KM, Willits KR, Litchfield RB, Fowler PJ, *et al.* Total knee replacement after high tibial osteotomy: time-to-event analysis and predictors. *CMAJ.* 2021;193(5):E158-E166. doi:10.1503/CMAJ.200934.
37. Hevesi M, Wilbur RR, Keyt LK, Abdel MP, Levy BA, Saris DBF, *et al.* Total knee arthroplasty function at 25 years following proximal tibial osteotomy: paired outcomes of bilateral TKAs following unilateral osteotomy. *J Bone Joint Surg Am.* 2022;104(6):552-8. doi:10.2106/JBJS.21.00761.
38. Dawson MJ, Ollivier M, Menetrey J, Beaufile P. Osteotomy around the painful degenerative varus knee: a 2022 ESSKA formal consensus. *Knee Surg Sports Traumatol Arthrosc.* 2023;31(8):3041-3. doi:10.1007/s00167-022-07024-0.
39. Janoyer M. Blount disease. *Orthop Traumatol Surg Res.* 2019;105(1):S111-S121. doi:10.1016/j.otsr.2018.01.009.
40. Ramella M, Depaoli A, Menozzi GC, Gallone G, Cerasoli T, Rocca G, *et al.* Recurrence and complication rates of surgical treatment for Blount's disease in children: a systematic review and meta-analysis. *J Clin Med.* 2023;12(20):6495. doi:10.3390/jcm12206495.
41. Gupta P, Gupta V, Patil B, Verma V. Angular deformities of lower limb in children: correction for whom, when and how? *J Clin Orthop Trauma.* 2020;11(2):196-201. doi:10.1016/j.Jcot.2020.01.008.
42. Brett A. Operative correction of genu recurvatum. *J Bone Joint Surg Am.* 1935;17(4):984-9.
43. Capella M, Gennari E, Dolfin M, Saccia F. Indications and results of high tibial osteotomy. *Ann Jt.* 2017;2:33. doi:10.21037/AOJ.2017.06.06.
44. Feucht MJ, Winkler PW, Mehl J, Bode G, Forkel P, Imhoff AB, *et al.* Isolated high tibial osteotomy is appropriate in less than two-thirds of varus knees if excessive overcorrection of the medial proximal tibial angle should be avoided. *Knee Surg Sports Traumatol Arthrosc.* 2021;29(10):3299-309. doi:10.1007/s00167-020-06166-3.
45. Nguyen HC, Gielis WP, van Egmond N, Weinans H, Slump CH, Sakkera RJB, *et al.* The need for a standardized whole leg radiograph guideline: the effects of knee flexion, leg rotation, and X-ray beam height. *J Cartil Jt Preserv.* 2021;1(3):100022. doi:10.1016/j.jcjp.2021.100022.
46. Roth T, Carrillo F, Wieczorek M, Ceschi G, Esfandiari H, Sutter R, *et al.* Three-dimensional preoperative planning in the weight-bearing state: validation and clinical evaluation. *Insights Imaging.* 2021;12(1):44. doi:10.1186/s13244-021-00994-8.

Appendix




Figure A1. Examples of the translational deformities in IMI-APPROACH, where the PMA and DMA run nearly parallel. The mean mMPTA and mLDTA were 91.0 ± 0.4 and 93.9 ± 1.6 , respectively.