



Universiteit
Leiden
The Netherlands

Hidden star formation in the early Universe

Leeuwen, I. F. van

Citation

Leeuwen, I. F. van. (2026, April 9). *Hidden star formation in the early Universe*. Retrieved from <https://hdl.handle.net/1887/4301034>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4301034>

Note: To cite this publication please use the final published version (if applicable).

Hidden star formation in the early Universe

Proefschrift

ter verkrijging van
de graad van doctor aan de Universiteit Leiden,
op gezag van rector magnificus prof. dr. S. de Rijcke,
volgens besluit van het college voor promoties
te verdedigen op donderdag 9 april 2026
klokke 11:30 uur

door

Ivana Francesca van Leeuwen

geboren te Woerden, Nederland
in 1997

Promotor: Prof. dr. P.P. van der Werf

Co-promotores: Dr. R.J. Bouwens
Dr. J.A. Hodge

Promotiecommissie: Prof. dr. I.A.G. Snellen
Prof. dr. H.J.A. Rottgering
Prof. dr. M.T. Kriek
Prof. dr. J.F. Hennawi
Prof. dr. K.I. Caputi (Rijksuniversiteit Groningen)
Dr. I. Shivaie (Centro de Astrobiologia, Madrid, Spain)

ISBN: 978-94-6534-282-5

Cover design and illustration: Olivia Seymour

Printed by: ProefschriftMaken

“I now walk into the wild.”
– *Christopher McCandless (1992)*

“But there is nothing quite so tangible beyond the sky,
only the suggestion of infinite space that scares him a little,
and thrills him even more, and makes his teeth itch at the dizzying
vastness of the universe and indeed the possibility of an infinite number of
universes beyond this one, if indeed infinity could even be numbered.”
– *The Perfect Golden Circle, Ben Myers (2022)*

CONTENTS

1	Introduction	1
1.1	A brief history of the Universe	1
1.2	Galaxy evolution: star formation through cosmic time	3
1.2.1	Spectral energy distribution	3
1.2.2	Star formation rate tracers	4
1.2.2.1	Rest-ultraviolet emission	4
1.2.2.2	Infrared emission	6
1.2.2.3	Far-IR fine structure lines	6
1.2.3	Star formation rate density	7
1.3	Galaxy samples	8
1.3.1	REBELS	8
1.3.2	Companion galaxies	10
1.4	Instruments	12
1.4.1	Atacama Large Millimeter/Submillimeter Array	12
1.4.2	Hubble Space Telescope	13
1.4.3	James Webb Space Telescope	13
1.5	This thesis	13
2	Characterising the contribution of dust-obscured star formation at $z \geq 5$ using 18 serendipitously identified [CII] emitters	17
2.1	Introduction	18
2.2	Methods	20
2.2.1	Observational data	20
2.2.2	Line-search algorithm	22
2.2.3	Moment-0 and continuum maps	26
2.2.4	Rest-UV photometry	26
2.2.5	Star formation rates	29
2.3	Results	31
2.3.1	Newly identified candidate companion galaxies	31
2.3.2	[CII]-selected sample	33
2.3.3	Obscured star formation	37
2.3.4	The Intrinsic UV Luminosity Function	39
2.3.5	The SFRD at $z \gtrsim 5$	45
2.4	Discussion	47
2.4.1	Comparisons with earlier selections of [CII]-emitting Companion Galaxies	47
2.4.2	Obscured star formation derived from [CII]-Selected Samples	50
2.4.3	The use of [CII] as a SFR tracer	51

2.4.4	Characterising the obscured fraction of SFR from [CII] emitters found in overdense regions	52
2.4.5	Evolution of the SFRD	54
2.4.5.1	Comparison of total SFRD	54
2.4.5.2	Dust-obscured SFRD	56
2.5	Conclusions	57
2.A	Appendices	62
2.A.1	Low-redshift contamination	62
2.A.2	Neighbour candidates	63
2.A.3	SExtractor	63
2.A.4	ALPINE companion galaxies	68
2.A.5	Archival ALMA Programs	68
3	The ALMA REBELS survey: [OIII]_{88μm} line scans of UV-bright $z \geq 7.6$ galaxies	73
3.1	Introduction	74
3.2	Observations and data reduction	76
3.2.1	The ALMA REBELS survey	76
3.2.2	Data reduction	76
3.2.3	Coverage of redshift likelihood distribution	77
3.2.4	[OIII] _{88μm} line emission	79
3.2.5	Dust continuum measurements	82
3.3	Results	83
3.3.1	Upper limits on $L_{[\text{OIII}]_{88\mu\text{m}}}$ of REBELS sources	83
3.3.2	[OIII] _{88μm} deficit	86
3.3.3	ALMA Band 7 continuum detections	89
3.3.4	Constraints on T_{dust} using Band 6 and 7	92
3.4	Discussion	94
3.4.1	[OIII] _{88μm} non-detections	94
3.4.1.1	REBELS-04	94
3.4.1.2	The remaining REBELS sources	96
3.4.2	Dust continuum emission from $z \gtrsim 7.6$ galaxies	98
3.5	Conclusions	98
3.A	Appendices	101
3.A.1	The photometric redshift of REBELS-04	101
3.A.2	Probable noise peak close to the rest-UV position of REBELS-13	102
3.A.3	Dust SED fit of REBELS-37	105
4	A JWST/NIRCam View of [CII]_{158μm}-Selected Companion Galaxies: A Key Population for Probing Dust-Obscured Star Formation at $z > 4$	109
4.1	Introduction	110
4.2	Sample and observations	112

4.3	Methods	114
4.3.1	Aperture photometry	114
4.3.2	NIRCam grism extractions	115
4.3.3	Photometric redshift and SED fitting	119
4.4	Results	120
4.4.1	Photometry	120
4.4.2	Updated rest-UV SFRs	124
4.4.3	[CII] or low- z CO emission	124
4.4.4	[CII] as a SFR tracer	127
4.4.5	SEDs of [CII] companions	127
4.4.6	Stellar masses	129
4.4.7	Obscured star formation	131
4.5	Discussion	133
4.5.1	Confirmation that most [CII]-selected galaxies are at $z > 4$	133
4.5.2	Extremely obscured normal star-forming galaxies . . .	136
4.5.3	Robustness of the stellar mass estimates	137
4.5.4	Overdense environments	139
4.6	Conclusions	140
4.A	Appendices	143
4.A.1	QSO companion galaxies	143
4.A.2	Low redshift SED fits	144
	Nederlandse samenvatting	157
	Publications	161
	Curriculum Vitae	163
	Acknowledgements	165