



Universiteit  
Leiden

The Netherlands

## Combining surgery and systemic therapy in metastatic melanoma

Blankenstein, S.A.M.

### Citation

Blankenstein, S. A. M. (2026, April 30). *Combining surgery and systemic therapy in metastatic melanoma*. Retrieved from <https://hdl.handle.net/1887/4303011>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4303011>

**Note:** To cite this publication please use the final published version (if applicable).





# PART I

**Neoadjuvant targeted therapy in  
unresectable locally advanced melanoma**



# Chapter 2

## Neoadjuvant cytoreductive treatment with BRAF/MEK inhibition of prior unresectable regionally advanced melanoma to allow complete surgical resection, REDUCTOR: a prospective, single arm, open-label phase II trial

Stéphanie A. Blankenstein, Maartje W. Rohaan, W. Martin C. Klop, Bernies van der Hiel, Bart A. van de Wiel, Max J. Lahaye, Sandra Adriaansz, Karolina Sikorska, Harm van Tinteren, Aysegül Sari, Lindsay G. Grijpink-Ongering, Winan J. van Houdt, Michel W.J.M. Wouters, Christian U. Blank, Sofie Wilgenhof, Johannes V. van Thienen, Alexander C.J. van Akkooi, John B.A.G. Haanen

## ABSTRACT

**Objective:** To evaluate the potency of short-term neoadjuvant cytoreductive therapy with dabrafenib plus trametinib (BRAF and MEK inhibitor) to allow for radical surgical resection in patients with unresectable locally advanced melanoma.

**Summary background data:** Approximately 5% of stage III melanoma patients presents with unresectable locally advanced disease, making standard of care with resection followed by adjuvant systemic therapy impossible. Although neoadjuvant targeted therapy has shown promising results in resectable stage III melanoma, its potency to enable surgical resection in patients with primarily unresectable locally advanced stage III melanoma is still unclear.

**Methods:** In this prospective, single arm, phase II trial, patients with unresectable BRAF-mutated locally advanced stage IIIC or oligometastatic stage IV melanoma were included. After 8 weeks of treatment with dabrafenib and trametinib, evaluation by PET/CT and physical examination were used to assess sufficient downsizing of the tumor to enable resection. The primary objective was the percentage of patients who achieved a radical (R0) resection.

**Results:** Between August 2014 and March 2019, 21 patients were included. Planned inclusion of 25 patients was not reached due to slow accrual and changing treatment landscape. Despite this, the predefined endpoint was successfully met. In 18/21 (86%) patients a resection was performed, of which 17 were R0 resections. At a median follow-up of 50 months (IQR 37.7-57.1 months), median recurrence free survival was 9.9 months (95% CI 7.52-not reached) in patients undergoing surgery.

**Conclusions:** This prospective, single arm, open-label phase II trial, shows neoadjuvant dabrafenib plus trametinib as a potent cytoreductive treatment, allowing radical resection of metastases in 17/21 (81%) patients with prior unresectable locally advanced melanoma.

## INTRODUCTION

Historically, the standard of care for macroscopic regional metastatic melanoma has been complete surgical resection of lymph node metastases. However, approximately five percent of patients presents with unresectable locally advanced disease with very bulky lymph node metastases or in-transit metastases, making a complete (R0) resection unfeasible. These patients are currently given the same treatment as patients with stage IV disease and subsequently have a similar prognosis.<sup>1</sup>

The systemic treatment of patients with unresectable stage III and IV melanoma has evolved drastically over the past decade with the development of both immune checkpoint inhibitors (ICIs) and targeted therapy. In metastatic melanoma, targeting the mitogen-activated protein kinase (MAPK)-pathway with BRAF plus MEK inhibitors (BRAF/MEKi) has proven to be successful in patients with BRAF-mutated melanoma.<sup>2-5</sup> Compared to ICI, treatment with BRAF/MEKi shows higher response rates (up to 68% versus up to 59% with the combination ICI of nivolumab and ipilimumab), but most patients acquire resistance in due course (5-year progression free survival (PFS) of 19% versus 36%).<sup>4-14</sup>

Indications for these systemic therapies are broadening, since treatment in the adjuvant setting has recently become the standard of care for patients with resectable stage III melanoma, who have a substantial risk of recurrence after resection.<sup>1, 15, 16</sup> Both adjuvant nivolumab or pembrolizumab, and adjuvant BRAF/MEKi were shown to improve the recurrence free survival (RFS) in this patient population.<sup>17-19</sup> This treatment regimen is not feasible in patients with unresectable locally advanced melanoma, however, systemic therapy could possibly enable a complete resection when given in a neoadjuvant, cytoreductive setting. ICI and BRAF/MEKi have recently been demonstrated to downsize resectable stage III melanoma substantially, with very impressive major pathological responses, including complete remissions in only a very short period of time (6-12 weeks).<sup>20, 21</sup> Yet no data are available for patients with primarily unresectable stage IIIC disease. For such patients with BRAF V600 mutated melanoma, the BRAF/MEKi combination may be suitable to enable enough downsizing to allow R0 resection, given the high response rate and short time to response.

Similar to unresectable stage IIIC melanoma, systemic therapy is the first choice of treatment in most stage IV melanoma patients. However, in patients with oligometastatic stage IV melanoma ( $\leq 3$  metastatic lesions), surgical resection could be a potential curative approach for a minority of patients.<sup>22, 23</sup> Nowadays, these patients are also eligible for adjuvant systemic therapy with nivolumab.<sup>17</sup> Neoadjuvant treatment in these patients may also reduce the development of new metastatic lesions from undetected micrometastatic disease.<sup>1</sup>

The aim of this prospective, single arm, open-label phase II study is to evaluate the potency of short-term neoadjuvant cytoreductive therapy with dabrafenib plus trametinib (BRAF and MEK inhibitor respectively) to allow for radical surgical resection in patients with unresectable and BRAF-mutated, locally advanced stage III or oligometastatic stage IV melanoma.

## PATIENTS AND METHODS

### Study design and population

This study was designed as a prospective, single arm, phase II trial including 25 patients when at least four of the first 14 included patients would respond to the neoadjuvant treatment. The study was approved by the ethics committee of the Netherlands Cancer Institute (NKI, EudraCT number 2013-002616-28) and all participants provided written informed consent. The trial was conducted according to the principles of the Declaration of Helsinki and the Medical Research Involving Human Subjects Act (WMO). Patients were enrolled and treated at the Departments of Medical and Surgical Oncology at NKI (Amsterdam, The Netherlands).

Patients with unresectable BRAF-mutated stage IIIc melanoma or stage IV melanoma with  $\leq 3$  metastases, in which surgery alone was deemed not to be a feasible treatment option, were eligible for this trial. As official criteria for unresectability have not yet been defined, it is therefore considered as the result of a multisurgeon decision during multidisciplinary melanoma meetings. Staging was performed according to the 7<sup>th</sup> edition of the melanoma American Joint Committee on Cancer staging manual (AJCC 7<sup>th</sup>).<sup>24</sup> Pathologic confirmation of cutaneous melanoma harboring a BRAF V600E or -K mutation was required. Also, patients had to be treatment naïve for this disease stage and the intended operation should be considered to offer a chance of curation or substantial palliation. Other main inclusion criteria were: patients of  $\geq 18$  years of age; WHO performance status of 0 or 1; evaluable lesions on CT, MRI or PET/CT; and adequate organ functions. Main exclusion criteria were presence of central nervous system metastases; major surgery, radiotherapy or systemic therapy in the four weeks prior to inclusion; pregnancy or lactation; known Human Immunodeficiency Virus (HIV), Hepatitis B or C infection; and cardiac abnormalities.

### Treatment and procedures

All patients underwent 18F-Fluorodeoxyglucose (18F-FDG) PET/CT and MRI of the brain as baseline measurements. Neoadjuvant treatment consisted of dabrafenib 150 mg twice daily (BID) and trametinib 2 mg once daily (QD) for a period of eight weeks. Evaluation was performed with physical examination by the surgeon after 4 and 8 weeks and with PET/CT after 2 and 8 weeks of treatment. If the tumor had been downsized sufficiently (as defined by a multidisciplinary board) and no new lesions had occurred, the operation was performed within the next 2-3 weeks. In this period the dabrafenib and trametinib were continued. If the tumor was still deemed unresectable, treatment with BRAF/MEKi was continued until disease progression or intolerable toxicity.

## Outcomes

The primary endpoint of this study was the percentage of patients for whom a radical (R0) resection was achieved, defined as tumor free margins of the resection specimen, confirmed by an expert pathologist. Pathologic responses were assessed using the consensus guidelines of the International Neoadjuvant Melanoma Consortium (INMC), using the percentage of tumor bed occupied by viable tumor cells.<sup>25</sup> Secondary endpoints were RFS, PFS and overall survival (OS). RFS was calculated in patients undergoing a resection, defined as the time between surgery and time of first disease progression (PD). In all patients, PFS was determined as the interval between the date of first administration of BRAF/MEKi and the time of PD or time of death due to any cause. The time of PD is defined as the date of radiological PD scored on imaging data according to Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1.<sup>26</sup> OS is defined as the time between the date of first administration of BRAF/MEKi and date of death due to any cause. Patients not experiencing an event will be censored at the day of last contact. Metabolic response rates were evaluated on 18F-FDG PET/CT according to response criteria of the European Organization for Research and Treatment of Cancer (EORTC), and PET Response Criteria in Solid Tumors (PERCIST).<sup>27,28</sup> Radiologic response rates were evaluated on (low dose) CT following RECIST 1.1.

Additionally, adverse events (AE) were evaluated by the investigators during the study according to the National Cancer Institute's Terminology Criteria for Adverse Events version 4.03 (NCI-CTCAE v.4.03). New AEs were registered from the moment of signing informed consent until 30 days after the last study intervention and all existing AEs were followed-up until recovery or baseline levels. Only clinically significant laboratory abnormalities were reported (grade  $\geq 3$ ).

## Statistical analyses

The hypothesis is that if downsizing by dabrafenib and trametinib treatment results in an R0 resection in 45% or more patients, the treatment modality is considered effective (alternative hypothesis). An R0 resection in 20% or less patients is considered futile (null hypothesis). The sample size was calculated using the Simon optimal two-stage method using an alpha of 0.1 and beta of 10% (power 90%).<sup>29</sup> The first stage would include 14 patients and the trial continued to the second stage, with 25 patients enrolled in total, if an R0 resection was achieved in at least four patients. Statistical analyses were performed using R version 3.5.1. Descriptive statistics were used to describe baseline patient and tumor characteristics; adverse events; responses; and recurrences. Survival endpoints (RFS, PFS, and OS) were analyzed using Kaplan-Meier estimates and medians were presented with 95% confidence intervals (CI).

## RESULTS

### Patient and treatment characteristics

Between August 2014 and March 2019 a total of 21 patients were included in this study. After inclusion of the first 14 patients, sufficient downsizing of the tumor was seen in 10 patients and the trial proceeded. However, inclusion was ceased before reaching the predefined total of 25 patients due to slow accrual rate, changing treatment landscape in the running time of the trial and results that could not fail to meet the predefined endpoints anymore, as further elucidated in the discussion section. Baseline characteristics of the 21 included patients are summarized in table 1. All but one patient (95%) had unresectable stage IIIc disease at the time of inclusion. A large proportion (43%) of the included patients had a melanoma of unknown primary (MUP).

All patients completed treatment with neoadjuvant BRAF/MEKi for eight weeks, although in six patients a short interruption was necessary due to toxicity (median four days (range 1 –18 days)).

**Table 1. Baseline characteristics**

Characteristic	n	%
Age, years		
Median	53	
Range	25-76	
Sex		
Female	10	48.0
Male	11	52.0
WHO performance status		
0	20	95.2
1	1	4.8
2	0	0.0
3	0	0.0
Disease stage (AJCC 7 <sup>th</sup> edition)		
IIIc	20	95.2
IV	1	4.8
Location primary melanoma		
Extremity	7	33.3
Trunk	5	23.8
Head and neck	0	0.0
Unknown primary	9	42.9
Type		
Superficial spreading	5	23.8
Nodular	3	14.3
Acral lentiginous	2	9.5
Lentigo maligna	0	0.0
Desmoplastic	0	0.0
Unknown primary	9	42.9

Characteristic	n	%
Unknown	2	9.5
Breslow thickness		
≤1.0 mm	0	0.0
1.01-2.0 mm	6	28.6
2.01-4.0 mm	2	9.5
>4.0 mm	2	9.5
Unknown primary	9	42.9
Unknown	2	9.5
Ulceration		
Yes	2	9.5
No	10	47.6
Unknown primary	9	42.9
Site locoregional metastases		
Axillary LN	10	47.6
Axillary and cervical LN	5	23.8
Inguinal LN	0	0.0
Iliac LN	3	14.3
Inguinal and iliac LN	3	14.3
BRAF-mutation		
V600E	20	95.2
V600K	1	4.8
LDH		
≤ULN	11	52.4
>ULN	9	42.9
Unknown	1	4.8

LDH, lactatedehydrogenase; LN, lymph node, ULN, upper limit of normal.

### Clinical responses

After eight weeks of treatment, two patients had experienced PD as detected on PET/CT, both of whom developed distant metastases during the neoadjuvant treatment period. In one of these patients, an axillary lymph node dissection (LND) was performed for palliative reasons and to obtain local control, as the initial lymph node metastases had responded and became resectable, despite the development of new distant lesions. Sufficient downsizing of the tumor was seen in the remaining 19 patients and all of these proceeded to surgery. Despite significant downsizing, in one patient the tumor was still deemed unresectable during surgery due to encasement of the iliac artery and vein. Therefore, in 18/21 patients (86%) a complete macroscopic resection could be performed.

Table 2 shows radiologic and metabolic responses after eight weeks. According to RECIST, most patients (76.2%) had a partial response (PR) on CT and one patient had a radiologic complete response (CR). Metabolic responses according to EORTC and PERCIST were comparable in all patients but one, predominantly revealing a partial metabolic response (PMR, 61.9% PERCIST). This one patient with a different EORTC and PERCIST response showed

stable disease (SD) according to PERCIST, but PMR when using EORTC. Four patients (19.0%) had a complete metabolic response (CMR).

**Table 2. Response and surgery**

<b>Response</b>	<b>N</b>	<b>%</b>
Radiologic response		
Complete response	1	4.8%
Partial response	16	76.2%
Stable disease	1	4.8%
Progressive disease	2	9.5%
Missing*	1	4.8%
Metabolic response EORTC criteria		
Complete metabolic response	4	19.0%
Partial metabolic response	14	66.7%
Stable metabolic disease	0	0.0%
Progressive metabolic disease	2	9.5%
Missing*	1	4.8%
Metabolic response PERCIST		
Complete metabolic response	4	19.0%
Partial metabolic response	13	61.9%
Stable metabolic disease	1	4.8%
Progressive metabolic disease	2	9.5%
Missing*	1	4.8%
Pathologic response		
Pathologic complete response	6	28.6%
Pathologic near complete response	3	14.3%
Pathologic partial response	4	19.0%
Pathologic non response	5	23.8%
N.A.	3	14.3%
Radical excision		
R0	17	81.0%
R1	1	4.8%
R2	0	0.0%
N.A.	3	14.3%

\* In one patient lesions did not meet RECIST 1.1 criteria, additionally PERCIST and EORTC could not be assessed in this patient due to an incompatible FDG-PET at baseline. N.A., not applicable; R0, radical resection; R1, tumor positive microscopic resection margins; R2, macroscopic residual tumor

### Surgical and pathological outcomes

Since the vast majority of patients in this trial had locally advanced stage III melanoma, surgery primarily consisted of a LND. In 13 (72%) patients an axillary LND was performed, in four patients this procedure was combined with a resection of either supraclavicular or cervical lymph nodes. Five (28%) patients underwent an extensive superficial (inguinal-femoral) and deep (iliac-obturator) groin LND. In the patient with stage IV disease at inclusion, an iliac LND was combined with resection of a spermatic cord metastasis and in a second procedure resection of a lung metastasis was performed.

Of the 18 patients undergoing a resection, the expert pathologist classified 17 as an R0 resection. In one patient the resection margin was not tumor-free and a re-excision was performed. In this second specimen no vital melanoma cells could be found. Pathologic responses were evaluated in the 18 patients undergoing a per protocol resection: six patients had a pathologic CR (pCR); three a near pCR; four had a pathologic PR (pPR) and pathologic non response (pNR) was seen in five patients. In the patient that underwent a palliative axillary LND despite development of distant metastases, a pPR was still seen in the resection specimen and tumor margins were free. However, due to the development of distant metastases during neoadjuvant treatment, this procedure is not classified as a per protocol resection nor an R0 resection. Neither radiologic response on CT nor metabolic response on PET could accurately predict pathologic response, which is shown in table 3.

**Table 3. Radiologic and metabolic versus pathologic response**

Pathology	pCR	pnCR	pPR	pNR	N.A.
Imaging					
Radiologic response					
CR	0	0	0	1	0
PR	6	3	4	2	1
SD	0	0	0	1	0
PD	0	0	0	0	2
Metabolic response EORTC criteria					
CMR	1	0	0	3	0
PMR	5	3	4	1	1
SMD	0	0	0	0	0
PMD	0	0	0	0	2
Metabolic response PERCIST					
CMR	1	0	0	3	0
PMR	5	3	3	1	1
SMD	0	0	1	0	0
PMD	0	0	0	0	2

CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; CMR, complete metabolic response; PMR, partial metabolic response; SMD, stable metabolic disease; PMD, progressive metabolic disease; pCR, pathologic complete response; pPR, pathologic partial response; nPCR, pathologic near complete response; pNR, pathologic non response. \* In one patient lesions did not meet RECIST 1.1 criteria, additionally PERCIST and EORTC could not be assessed in this patient due to an incompatible FDG-PET at baseline.

Per protocol, no adjuvant therapy was given, except for adjuvant radiotherapy in two patients. One patient traveled abroad to receive adjuvant systemic therapy (one year of PD-1 blockade), since this was not part of the study treatment nor standard and reimbursed care at that time in the Netherlands.

## Adverse events


Most patients experienced some form of toxicity during neoadjuvant systemic treatment with BRAF/MEKi: only two (10%) patients experienced no treatment-related AEs. In the majority of patients the worst toxicity was grade 1 (11 patients, 52%), in four (19%) patients grade 2 and four (19%) patients experienced grade 3 toxicity. The most commonly reported AEs related to dabrafenib plus trametinib were fever (48%), chills (19%), fatigue (19%), nausea (19%) and myalgia (14%).

The majority of patients that underwent surgery (16/20, 80%) experienced a surgical complication, consisting mainly of the development of seroma (55%) and wound infections (30%). Three patients had grade 1 seroma, five patients grade 2 and three patients experienced grade 3 seroma for which elective radiologic or surgical intervention was indicated. Grade 2 wound infection was seen in two patients and four patients endured a grade 3 wound infection requiring intravenous antibiotics and hospital admittance.


## Survival outcomes

At a median follow-up of 50 months (IQR 37.7-57.1 months), the median RFS in the 18 patients undergoing surgery was 9.9 months (95% CI 7.52-not reached). Median PFS in all 21 patients was 12.4 months (95% CI 8.67 – not reached). The patient who had received adjuvant systemic therapy abroad, outside of the study protocol, was censored for PFS and RFS at time of commencement of adjuvant systemic treatment. Median OS was not reached. The 1-year OS was 100% and 2-year OS 85% (95% CI 70.0-100.0). RFS, PFS and OS are shown in figure 1a, b, and c respectively.


**Figure 1a. Recurrence free survival**



**Figure 1b. Progression free survival**



**Figure 1c. Overall survival**



## Recurrences

Recurrences were seen in half of the patients that underwent surgery (9/18). Most patients developed distant recurrences (6/9), versus three patients with locoregional recurrences as a first presentation of relapse. The three patients with locoregional recurrences were treated with surgery, but all developed distant metastases and were treated with systemic therapy at later time points. Of patients with a pCR, five (83%) did not relapse, but one (17%) did develop a recurrence. Four out of five patients with a pNR relapsed. All three patients that could not undergo a resection, due to either PD during study treatment or insufficient response, were treated with ICI, resulting in a CR in these patients.

## DISCUSSION

In this prospective, single arm, open-label phase II trial, neoadjuvant treatment with dabrafenib and trametinib has shown to be a potent cytoreductive treatment, allowing radical resection of metastases in 17/21 (81%) patients with prior unresectable locally advanced melanoma. Encouraging RFS and OS are also seen in these patients, even compared to patients with resectable stage IIID melanoma.<sup>30</sup>

Several trials have reported neoadjuvant treatment with either targeted therapy or ICI in stage III melanoma. Two studies reported treatment with neoadjuvant BRAF/ MEKi in resectable BRAF-mutated stage III melanoma. Both Amaria et al.<sup>31</sup> and Long et al.<sup>20</sup> treated patients with neoadjuvant (8 and 12 weeks respectively) and adjuvant (44 and 40 weeks respectively) BRAF/ MEKi for one year in total. Patients presented with a MUP in 14% (standard of care arm) and 29% (neoadjuvant arm) of cases in the trial by Amaria et al. The site of the primary melanoma was not described by Long et al. In our current trial, a relatively large proportion of patients (43%) presented with a MUP. This could be due to the fact that they were unaware of their melanoma and that patients were not seen in a standard follow-up program, resulting in later clinical presentation. In our current trial, the neoadjuvant treatment was given for a relatively short time compared to the trials by Amaria et al and Long et al. The cytoreductive effect of BRAF/MEK inhibitors usually occurs quickly after initiation of treatment, thus it was hypothesized that 8 weeks neoadjuvant treatment should suffice to achieve enough tumor downsizing to enable resection. A later response after this time was deemed unlikely. In the trial by Amaria et al., the RFS in patients that received this treatment was 19.7 months (95% CI 16.2-not reached), versus 2.9 months (95% CI 1.7-not reached, HR 0.016) in the standard of care arm of this trial. In the study by Long et al., the RFS was 23.3 months (95% CI 17.7-not reached). In both trials high response rates could be observed, with 85% of patients achieving a radiological response scored per RECIST 1.1 and 58% of patients achieving a pCR in de trial by Amaria et al. In the trial by Long et al., 86% of patients achieved a radiological response prior to surgery and in all patients a pathological response was seen in the resection specimen (49% pCR, 51% non-complete pathological response). These striking differences in

(complete) response rates (RR) compared to the current trial can potentially be explained by the different patient populations included in these trials. These other two trials only included patients with resectable stage IIIB-C disease, whereas our current trial only included patients with unresectable disease, presently considered as stage IIID disease following AJCC 8<sup>th</sup> edition. The differences in RFS may in turn be explained by the absence of adjuvant treatment after resection in our current trial, this in contrast to the trials by Amaria et al and Long et al.

Of the patients undergoing surgical resection, 17% presented with locoregional disease as site of first recurrence in our present trial. This is comparable to earlier reported locoregional recurrence rates in patients with resectable IIIC melanoma undergoing resection without (neo) adjuvant systemic therapy, as represented by placebo arms of recent adjuvant randomized trials.<sup>32-34</sup> However, the patients in our trial would not been able to undergo surgery, without cytoreductive treatment.

In our study, both radiologic and metabolic response prior to surgery were not able to predict pathologic response. This was also shown by Eroglu et al. in a retrospective study in a comparable patient population.<sup>32</sup> Of nine patients with a (near) pCR, only one patient had shown a corresponding metabolic CR and no patients had a radiologic CR; the remainder of these pCR patients had shown a radiologic and metabolic PR. The poor performance of radiologic response as a predictor of pCR might be explained by fibrotic tissue visible as a remaining lesion on CT, although no viable tumor cells are present. Tan et al. described that FDG-PET may be superior to CT in showing a CR in metastatic melanoma patients treated with ICI.<sup>33</sup> However, in our cohort this distinction was not as clear. Both EORTC and PERCIST criteria were designed to measure response to chemotherapy, and perhaps this cannot be directly translated to treatment with targeted therapy.

Besides neoadjuvant targeted therapy, different schemes of neoadjuvant anti-PD-1 or the combination ICI with nivolumab and ipilimumab have been reported in patients with resectable palpable stage III melanoma as well, as described by Blank et al.<sup>34</sup>, Rozeman et al.<sup>21</sup>, Amaria et al.<sup>35</sup>, and Huang et al.<sup>36</sup> In combination ICI studies, very high pathologic RR were seen of 73-80% and to date, pathologic response seemed predictive of RFS, as none of the patients who achieved a pathologic response relapsed in the OpACIN-neo study.<sup>21, 34, 35</sup>

As yet, no direct comparison between (neo)adjuvant targeted therapy and ICI has been performed in resectable stage III melanoma. Although response rates are high in the neoadjuvant setting for ICI, it is uncertain whether this can be translated to the population of patients with unresectable stage III melanoma. Both BRAF/ MEKi and ipilimumab/nivolumab have shown rapid responses and high objective RR in stage IV melanoma.<sup>4, 11</sup> In our current trial we have shown that 18/21 patients had a radiological and metabolic response after only 8 weeks of dabrafenib/trametinib treatment and radical resection could be achieved in 17/21 patients. The majority of patients did not develop distant metastases during this treatment period, despite being at very high risk for metastatic disease and with a median follow-up of 50 months 43%

was still disease free. Whether similar or better results could be obtained for unresectable stage III disease with short term neoadjuvant ICI is currently unknown.

2 Due to the rapid developments in the systemic treatment of stage IV melanoma and the oftentimes quickly progressive nature of stage IV disease, only one patient with stage IV disease was included in this study. Currently, the standard of care in these patients is first line systemic treatment with PD-1 blockade or combined ipilimumab/nivolumab and surgery is only considered in select cases of residual disease or oligoprogression.<sup>1</sup>

This single arm, open-label phase II trial is, to our knowledge, the first prospective trial treating unresectable locally advanced melanoma patients with neoadjuvant cytoreductive targeted therapy, of which the results look very promising. A limitation of this trial however, is the small patient cohort. This is partly due to the specific patient population, as most patients have either limited locoregional disease only or bulky locoregional metastases combined with multiple distant metastases. Also, during the course of this trial, BRAF/MEKi became available for unresectable stage III melanoma patients outside of clinical trials, reducing the number of referrals from other sites. Due to the slow accrual rate, the decision was made to cease the trial prematurely to retain its clinical relevance. An unplanned interim analysis of the, up until then, 21 included patients showed that predefined endpoints had already been exceeded. Moreover, since the neoadjuvant treatment with BRAF/MEKi was effective in most patients and adjuvant treatment had become standard of care in resected stage III melanoma during the course of this study, many patients inquired about the possibility of receiving adjuvant treatment after resection. Due to the combination of the slow accrual rate, results of the unplanned interim analysis and the changing treatment landscape with adjuvant systemic therapy as the new standard of care, the decision was made to cease inclusion and report the outcomes of this study.

Dabrafenib plus trametinib has shown to be a potent neoadjuvant cytoreductive treatment in this select population of BRAF-mutated unresectable locally advanced melanoma patients. Patients with no recurrence remained disease-free for a prolonged period of time. However, when recurrences were seen, this usually occurred quickly after surgery. This may present a window of opportunity for adjuvant therapy with ICI, possibly in combination with BRAF/MEKi, in order to achieve improved and more durable RFS. Further clinical trials are needed to explore the additional benefit of such adjuvant therapy after previous neoadjuvant BRAF/MEKi.

## CONCLUSION

In this prospective single arm, open-label phase II trial, neoadjuvant dabrafenib and trametinib has shown to be a potent cytoreductive treatment, allowing radical resection of metastases in 17/21 (81%) patients with prior unresectable locally advanced melanoma. If relapses occurred, this was usually quickly (within months) after surgery. This could present an opportunity for tailored adjuvant therapy.

## REFERENCES

1. Michielin O, van Akkooi ACJ, Ascierto PA, et al. Cutaneous melanoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2019; 30(12):1884-1901.
2. Ascierto PA, McArthur GA, Dreno B, et al. Cobimetinib combined with vemurafenib in advanced BRAF(V600)-mutant melanoma (coBRIM): updated efficacy results from a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2016; 17(9):1248-60.
3. Long GV, Flaherty KT, Stroyakovskiy D, et al. Dabrafenib plus trametinib versus dabrafenib monotherapy in patients with metastatic BRAF V600E/K-mutant melanoma: long-term survival and safety analysis of a phase 3 study. *Ann Oncol* 2017; 28(7):1631-1639.
4. Robert C, Grob JJ, Stroyakovskiy D, et al. Five-Year Outcomes with Dabrafenib plus Trametinib in Metastatic Melanoma. *N Engl J Med* 2019; 381(7):626-636.
5. Dummer R, Ascierto PA, Gogas HJ, et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2018; 19(5):603-615.
6. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010; 363(8):711-23.
7. Ascierto PA, Del Vecchio M, Robert C, et al. Ipilimumab 10 mg/kg versus ipilimumab 3 mg/kg in patients with unresectable or metastatic melanoma: a randomised, double-blind, multicentre, phase 3 trial. *Lancet Oncol* 2017; 18(5):611-622.
8. Ascierto PA, Long GV, Robert C, et al. Survival Outcomes in Patients With Previously Untreated BRAF Wild-Type Advanced Melanoma Treated With Nivolumab Therapy: Three-Year Follow-up of a Randomized Phase 3 Trial. *JAMA Oncol* 2019; 5(2):187-194.
9. Larkin J, Minor D, D'Angelo S, et al. Overall Survival in Patients With Advanced Melanoma Who Received Nivolumab Versus Investigator's Choice Chemotherapy in CheckMate 037: A Randomized, Controlled, Open-Label Phase III Trial. *J Clin Oncol* 2018; 36(4):383-390.
10. Weber JS, Gibney G, Sullivan RJ, et al. Sequential administration of nivolumab and ipilimumab with a planned switch in patients with advanced melanoma (CheckMate 064): an open-label, randomised, phase 2 trial. *Lancet Oncol* 2016; 17(7):943-955.
11. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Five-Year Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N Engl J Med* 2019; 381(16):1535-1546.
12. Hodi FS, Chesney J, Pavlick AC, et al. Combined nivolumab and ipilimumab versus ipilimumab alone in patients with advanced melanoma: 2-year overall survival outcomes in a multicentre, randomised, controlled, phase 2 trial. *Lancet Oncol* 2016; 17(11):1558-1568.
13. Robert C, Ribas A, Schachter J, et al. Pembrolizumab versus ipilimumab in advanced melanoma (KEYNOTE-006): post-hoc 5-year results from an open-label, multicentre, randomised, controlled, phase 3 study. *Lancet Oncol* 2019; 20(9):1239-1251.
14. Hamid O, Puzanov I, Dummer R, et al. Final analysis of a randomised trial comparing pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory advanced melanoma. *Eur J Cancer* 2017; 86:37-45.
15. Schadendorf D, van Akkooi ACJ, Berking C, et al. Melanoma. *Lancet* 2018; 392(10151):971-984.
16. Blankenstein SA, van Akkooi ACJ. Adjuvant systemic therapy in high-risk melanoma. *Melanoma Res* 2019; 29(4):358-364.
17. Weber J, Mandala M, Del Vecchio M, et al. Adjuvant Nivolumab versus Ipilimumab in Resected Stage III or IV Melanoma. *N Engl J Med* 2017; 377(19):1824-1835.
18. Eggermont AMM, Blank CU, Mandala M, et al. Adjuvant Pembrolizumab versus Placebo in Resected Stage III Melanoma. *N Engl J Med* 2018; 378(19):1789-1801.
19. Hauschild A, Dummer R, Schadendorf D, et al. Longer Follow-Up Confirms Relapse-Free Survival Benefit With Adjuvant Dabrafenib Plus Trametinib in Patients With Resected BRAF V600-Mutant Stage III Melanoma. *J Clin Oncol* 2018; Jco1801219.

20. Long GV, Saw RPM, Lo S, et al. Neoadjuvant dabrafenib combined with trametinib for resectable, stage IIIB-C, BRAF(V600) mutation-positive melanoma (NeoCombi): a single-arm, open-label, single-centre, phase 2 trial. *Lancet Oncol* 2019; 20(7):961-971.
21. Rozeman EA, Mazières AM, van Akkooi ACJ, et al. Identification of the optimal combination dosing schedule of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma (OpACIN-neo): a multicentre, phase 2, randomised, controlled trial. *Lancet Oncol* 2019; 20(7):948-960.
22. Sosman JA, Moon J, Tuthill RJ, et al. A phase 2 trial of complete resection for stage IV melanoma: results of Southwest Oncology Group Clinical Trial S9430. *Cancer* 2011; 117(20):4740-06.
23. Howard JH, Thompson JF, Mozzillo N, et al. Metastectomy for distant metastatic melanoma: analysis of data from the first Multicenter Selective Lymphadenectomy Trial (MSLT-I). *Ann Surg Oncol* 2012; 19(8):2547-55.
24. Balch CM, Gershenwald JE, Soong SJ, et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol* 2009; 27(36):6199-206.
25. Tetzlaff MT, Messina JL, Stein JE, et al. Pathological assessment of resection specimens after neoadjuvant therapy for metastatic melanoma. *Ann Oncol* 2018; 29(8):1861-1868.
26. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009; 45(2):228-47.
27. Young H, Baum R, Cremerius U, et al. Measurement of clinical and subclinical tumour response using [18F]-fluorodeoxyglucose and positron emission tomography: review and 1999 EORTC recommendations. European Organization for Research and Treatment of Cancer (EORTC) PET Study Group. *Eur J Cancer* 1999; 35(13):1773-82.
28. Wahl RL, Jacene H, Kasamon Y, et al. From RECIST to PERCIST: Evolving Considerations for PET response criteria in solid tumors. *J Nucl Med* 2009; 50 Suppl 1:122s-50s.
29. Simon R. Optimal two-stage designs for phase II clinical trials. *Control Clin Trials* 1989; 10(1):1-10.
30. Gershenwald JE, Scolyer RA, Hess KR, et al. Melanoma staging: Evidence-based changes in the American Joint Committee on Cancer eighth edition cancer staging manual. *CA Cancer J Clin* 2017; 67(6):472-492.
31. Amaria RN, Prieto PA, Tetzlaff MT, et al. Neoadjuvant plus adjuvant dabrafenib and trametinib versus standard of care in patients with high-risk, surgically resectable melanoma: a single-centre, open-label, randomised, phase 2 trial. *Lancet Oncol* 2018; 19(2):181-193.
32. Eroglu Z, Eatrides J, Naqvi SMH, et al. Neoadjuvant BRAF-targeted therapy in regionally advanced and oligo-metastatic melanoma. *Pigment Cell Melanoma Res* 2020; 33(1):86-95.
33. Tan AC, Emmett L, Lo S, et al. Utility of 1-year FDG-PET (PET) to determine outcomes from anti-PD-1 (PD1) based therapy in patients (pts) with metastatic melanoma (MM). *Journal of Clinical Oncology* 2018; 36(15\_suppl):9517-9517.
34. Blank CU, Rozeman EA, Fanchi LF, et al. Neoadjuvant versus adjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma. *Nat Med* 2018; 24(11):1655-1661.
35. Amaria RN, Reddy SM, Tawbi HA, et al. Neoadjuvant immune checkpoint blockade in high-risk resectable melanoma. *Nat Med* 2018; 24(11):1649-1654.
36. Huang AC, Orlovski RJ, Xu X, et al. A single dose of neoadjuvant PD-1 blockade predicts clinical outcomes in resectable melanoma. *Nat Med* 2019; 25(3):454-461.

