



Universiteit
Leiden

The Netherlands

Single-cell dissection of immune system aging and acute myeloid leukemia

Karakaslar, E.O.

Citation

Karakaslar, E. O. (2026, June 16). *Single-cell dissection of immune system aging and acute myeloid leukemia*. Retrieved from <https://hdl.handle.net/1887/4306748>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4306748>

Note: To cite this publication please use the final published version (if applicable).



Chapter 5

Mapping AML Heterogeneity: Multi-cohort transcriptomic analysis identifies novel clusters and divergent ex-vivo drug responses

Jeppe F. Severens^{1,2,3}, E. Onur Karakaslar^{1,2,3}, Bert A. van der Reijden⁴, Elena Sánchez-López^{3,5}, Redmar R. van den Berg⁴, Constantijn J.M. Halkes⁶, Peter van Balen⁶, Hendrik Veelken^{3,6}, Marcel J.T. Reinders^{1,2,3}, Marieke Griffioen⁶, Erik B. van den Akker^{1,2,3,*}

¹ Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

² Pattern Recognition & Bioinformatics, Delft University of Technology, Delft, The Netherlands

³ Leiden Center for Computational Oncology, Leiden University Medical Center, Leiden, The Netherlands

⁴ Department of Laboratory Medicine, Radboud University Medical Center, Nijmegen, The Netherlands

⁵ Center for Proteomics and Metabolomics, Leiden University Medical Center, Leiden, The Netherlands

⁶ Department of Hematology, Leiden University Medical Center, Leiden, The Netherlands

5.1 Abstract

Subtyping of acute myeloid leukaemia (AML) is predominantly based on recurrent genetic abnormalities, but recent literature indicates that transcriptomic phenotyping holds immense potential to further refine AML classification. Here we integrated five AML transcriptomic datasets with corresponding genetic information to provide an overview (n=1224) of the transcriptomic AML landscape. Consensus clustering identified 17 robust patient clusters which improved identification of *CEBPA*-mutated patients with favourable outcomes, and uncovered transcriptomic subtypes for *KMT2A* rearrangements (2), *NPM1* mutations (5), and AML with myelodysplasia-related changes (AML-MRC) (5). Transcriptomic subtypes of *KMT2A*, *NPM1* and AML-MRC showed distinct mutational profiles, cell type differentiation arrests and immune properties, suggesting differences in underlying disease biology. Moreover, our transcriptomic clusters show differences in ex-vivo drug responses, even when corrected for differentiation arrest and superiorly capture differences in drug response compared to genetic classification. In conclusion, our findings underscore the importance of transcriptomics in AML subtyping and offer a basis for future research and personalised treatment strategies. Our transcriptomic compendium is publicly available and we supply an R package to project clusters to new transcriptomic studies.

5.2 Introduction

In acute myeloid leukaemia (AML), recurrent genetic abnormalities (RGA) have been identified through systematic genomic studies^{1–5}. Based on these RGAs, the World Health Organization (WHO 2022) and International Consensus Classification (ICC 2022) define several AML subtypes, as well as a heterogeneous subtype of AML with myelodysplasia-related changes (AML-MRC)^{6,7}. RGAs are essential for risk-stratification and are increasingly targeted with drugs^{8,9}.

AML subclassification is genetics-based, but transcriptomics holds immense potential to refine AML classification further^{1–3,10–12}. Transcriptomic studies have led to the discovery of *CEBPA*-mutated AML^{13,14}, and *NPM1*-mutated AML subtypes with different cell differentiation arrests and ex-vivo drug responses^{15,16}. Similar stratification would be beneficial for AML-MRC, given its heterogeneity^{17,18}. Still, a comprehensive examination of AML subtypes defined by gene expression has yet to be performed. Furthermore, the differentiation arrest state is known to modify drug response in AML¹⁹, and failing to account for this effect when comparing drug responses could skew conclusions.

Therefore, we integrated five mRNAseq datasets with corresponding genetic aberration data and annotated cases according to WHO and ICC 2022 standards. We outline AML's transcriptomic landscape and define transcriptional subtypes with distinct gene expressions, genetic aberrations, and cell type arrests. We relate the clusters to ex-vivo drug responses independently of differentiation arrest and show how they superiorly capture differences in response compared to genetic classification. We provide all harmonised data and a transcriptional cluster predictor for future research. Our study underscores the importance of incorporating transcriptomic data in AML classification.

5.3 Results

Multi-cohort AML gene expression compendium

We collected 1224 RNAseq samples from adult (BEAT, TARGET, TCGA, Leucegene, LUMC) and paediatric (TARGET) cohorts with corresponding genetic and clinical data (**Figure 5.1A**). We quantified gene expression with the same pipeline and corrected counts for cohort, sex and source tissue (**Supplemental Figure 1**). Sample classification by their genetic data according to the WHO (**Figure 5.1C**) and ICC was successful for 97% of the samples. In line with previous reports, frequencies of the AML subtypes were similar for the adult cohorts but different between paediatric and adult cohorts (**Supplemental Table 1**), confirming that our dataset is representative of the AML landscape^{40,5}.

Transcriptomics define 17 AML clusters

Next, we assigned AML cases to 17 transcriptional clusters using consensus clustering (**Figure Supplemental Figures 2 & 3**). We named the clusters based on genetic diagnoses (**Figure 5.1B**, **Supplemental Figure 4 & 5**). As expected, the distribution over the clusters was different for paediatric and adult cohorts, exemplified by the large percentage of paediatric samples in the *KMT2A*^T clusters (26%), and adult samples in the *NPM1*^T (93%) and AML-MRC^T (94%) clusters (**Supplemental Table 1**, **Supplemental Figure 6**). However, samples of identical AML genetic subtypes from adult and paediatric cohorts clustered together, indicating that the 17 clusters capture differences in AML biology.




Figure 5.1: **Transcriptomic analysis further stratifies AML.** **A** Flowchart of the used data and methods. **B** Sankey plot showing the assignment of WHO 2022 diagnoses over the identified clusters. **C** tSNE visualisation of the gene expression of patient samples. Each dot represents a patient. The samples are coloured according to the WHO 2022 subtyping of AML. **D** The same tSNE visualisation as in **C**, but samples are coloured according to the 17 clusters. **E** Dot plots that show enriched aberrations in the 17 clusters. The dots are coloured according to the adjusted p-value. The dots are sized according to the sample fraction with the aberration in the cluster. The x-axis shows the aberrations, and the y-axis shows the clusters. We only visualised enriched aberrations that occurred in at least 10% of the patients in a cluster.

We examined clustering robustness using the stability score (**Supplemental Figure 7**). Median clustering stability was high (0.97-1.00), with AML-MRC(3)^T showing the lowest stability. A correlation test revealed a significant but weak correlation ($\rho = 0.18$, $p\text{-value} < .001$) between blast percentage and clustering robustness, but blast percentage varied greatly in clusters. tSNEs generated using different MVGs (**Supplemental Figure 8**) were stable from 500 to 2500 features. These results show that clustering was robust and only weakly influenced by blast percentage.

We developed a transcriptional cluster predictor using uncorrected counts as input (accuracy = 0.90), demonstrating the persistence of expression patterns. The quality of the predictor was further improved (accuracy = 0.95) by including a reject option (10% rejected) (**Supplemental Figure 9**).

Next, we tested for enrichment of mutations, fusions and cytogenetic aberrations ($n=102$) (**Figure 5.1E**, p -values and frequencies in **Supplemental Table 2**). Four transcriptomic clusters corresponded to singular genetic AML subtypes: RUNX1::RUNX1T1^T (RUNX1::RUNX1T1: 94%, FDR<.001), CBFB::MYH11^T (CBFB::MYH11: 95%, FDR<.001), PML::RARA^T (PML::RARA: 100%, FDR<.001), and NUP98^T (NUP98::NSD1: 45%, FDR<.001). Risk-stratification for survival based on transcriptional subtypes performed similarly to genetics (**Supplemental Figure 10 & 11**). We identified no enrichment for BCR::ABL1 and DEK::NUP214, possibly due to their limited occurrence. For KMT2A rearrangements, CEBPA mutations, NPM1 mutations, and AML-MRC, we found evidence that transcriptomics can refine subtyping, as described below.

Transcriptome analysis identifies two KMT2A-related clusters

The WHO classification defines a single KMT2A-rearranged subtype (KMT2A-r), while the ICC recognises KMT2A::MLLT3 and other KMT2A fusions as distinct^{6,7}. We identified two KMT2A fusion clusters. KMT2A(1)^T was significantly enriched for KMT2A::MLLT3 (31%, FDR<.001), KMT2A::MLLT10 (19%, FDR<.001) and any KMT2A fusion (67%, FDR<.001), while KMT2A(2)^T was enriched for KMT2A::MLLT4 (67%, FDR<.001) and high MECOM expression (80%, FDR<.001) (**Figure 5.2A & B**). Interestingly, we found cases with NPM1 mutations and trisomy 8/8q localised in KMT2A(1)^T, indicating that these lead to KMT2A fusion-like gene expression.

The genes *LAMP5*, and *ADCY9* showed high expression in KMT2A(1)^T and low expression in KMT2A(2)^T (**Figure 5.2B**), and all have been shown to contribute to AML pathogenesis^{41,42}. Additionally, the transcription factor (TF) *ETV2* was highly expressed in KMT2A(1)^T, while the TF *ERG* displayed high expression in KMT2A(2)^T. The cell type scores revealed KMT2A(1)^T to have a significantly higher promonocyte-like score (FDR<.001), while KMT2A(2)^T was more hematopoietic stem cell (HSC)-like (FDR<.001) (**Figure 5.2B, Supplemental Figure 12**). FAB annotations showed similar results for KMT2A(1)^T, which had a high M5 (monocytic leukaemia) fraction (90%), while KMT2A(2)^T was more mixed (**Figure 5.2C**). Overall, we found that gene expression-based separation of KMT2A-r did not align with the ICC 2022 classification.

The CEBPA^T cluster indicates a favourable prognosis

As acknowledged in the ELN2022, patients with a CEBPA bZIP inframe mutation have a favourable prognosis^{8,43}. We identified a transcriptional CEBPA^T cluster significantly enriched for mutated CEBPA cases (72%, FDR<.001), with 42% of the samples having a CEBPA bZIP indel, either as single mutation or combined with an N-terminal frameshift mutation (**Figure 5.3A & B**). The remaining samples contained other mutations in the bZIP area or N-terminal region or had no detectable CEBPA mutation. Of note, a single CEBPA bZIP indel case resided outside the CEBPA^T cluster. This patient had an IDH-R132 mutation with a VAF=0.47, while the CEBPA bZIP




Figure 5.2: Transcriptome analysis identifies two *KMT2A*-related clusters. **A** tSNE visualisation of patient samples, coloured according to *KMT2A*-fusion or *NPM1* mutation and trisomy 8. **B** Waterfall plot of aberrations in the *KMT2A* clusters, including the percentage of samples with the aberration. The plot is combined with heatmaps showing the expression of marker genes and cell type scores. The columns are samples, which are split according to transcriptional clusters. **C** Fraction of FAB annotations per cluster. HSC = hematopoietic stem cells, Prog. = progenitor, GMP = granulocyte-monocyte progenitor, Prom. = promonocytes, Mono. = monocytes, cDC = conventional dendritic cells

in-frame mutation had a VAF=0.21. This finding suggests that the *IDH*-R132 mutation dominates the expression pattern, placing this case in cluster AML-MRC(1)^T. Conversely, all CEBPA^T cluster patients showed similar favourable outcomes (log-rank test: p-value=.80), irrespective of whether the CEBPA bZIP inframe mutation was detected (Figure 5.3C). The CEBPA^T cluster thus marks patients with a favourable outcome regardless of CEBPA mutation detection, which the CEBPA^T expression profile can detect.




Figure 5.3: The CEBPA^T cluster indicates a favourable prognosis. **A** Waterfall plot and gene expression heatmap of all samples in the CEBPA^T cluster and samples with a CEBPA mutation located outside the CEBPA^T cluster. **B** tSNE visualisation of patient samples, coloured according to the type of CEBPA mutation. For samples with multiple CEBPA mutations, we used the ordering as in **A** to decide which mutation to display. **C** Kaplan-Meier curve of the survival of BEAT and TCGA patients in and outside the CEBPA^T cluster.

Gene expression profiling identifies five transcriptional NPM1-related clusters

The 2022 WHO and ICC classifications include one subtype of *NPM1*-mutated AML^{6,7}. However, we identified five clusters enriched for mutated *NPM1* (Figure 5.4). We observed elevated expression of *HOXA3*, *HOXB5*, and *MEIS1* (Figure 5.4A), which has been earlier associated with *NPM1* mutations⁴⁴. Interestingly, *NPM1* mutation-lacking samples generally also exhibited high expression of these genes, suggesting that there are alternative mishaps that disrupt these genes leading to *NPM1* mutated-like AML.

NPM1(1)^T exhibited the highest percentage (95%, FDR<.001) of *NPM1* mutated samples and was significantly enriched for *IDH1*-R132 (25%, FDR<.001), *IDH2*-R140 (37%, FDR<.001), and *TET2* (33%, FDR<.001) co-mutations (Figure 5.4A). NPM1(2)^T samples were enriched for *FLT3*-ITD mu-

Results

tations (84%, FDR<.001), but *FLT3*-ITD was also enriched in NPM1(1)^T, NPM1(3)^T and NPM1(4)^T (42-43%, all FDR<.001). Additionally, NPM1(4)^T and NPM1(5)^T had a significantly lower variant allele frequency for mutated *NPM1* (Figure 5.4B). We found two *NPM1::MLF1* cases in our compendium, which both clustered in NPM1(3)^T. *NPM1::MLF1* has been shown to localise in the cytoplasm⁴⁵, like mutated *NPM1*, possibly leading to a similar expression profile as *NPM1*-mutated cases.

Each of the *NPM1*-related clusters exhibited unique marker genes (Figure 5.4A). For instance, *FTO* expression was high in NPM1(1)^T. Additionally, *LYRM1*, *ADAM8*, and *DNAJC13* were elevated in NPM1(2)^T, NPM1(4)^T, and NPM1(5)^T, respectively. NPM1(3)^T had a less distinct expression pattern, suggesting a more heterogeneous cluster. Also, we observed differential expression of TFs (*RUNX1*, *PRDM16*, *SPI1*)^{46,47} – even in samples lacking the *NPM1* mutation – and TF expression aligned with cell differentiation stages.


NPM1(1)^T and NPM1(2)^T displayed a HSC-like expression pattern, NPM1(3)^T was mixed, whereas NPM1(4)^T and NPM1(5)^T were more differentiated (Figure 5.4A, Supplemental Figure 12). FAB annotations showed additional differences, with NPM1(5)^T containing fewer M4 (myelomonocytic leukaemia) but more M5 (monocytic leukaemia) cases than NPM1(4)^T (Figure 5.4C & D). Using scores for HLA I and HLA II antigen-presenting cells³⁹ we found NPM1(1)^T to have significantly lower HLA I (FDR<.001) and HLA II (FDR<.001) scores than the other clusters (Figure 5.4E & F). NPM1(1)^T and NPM1(5)^T patients were significantly older (FDR<.05), while NPM1(3)^T patients were younger (FDR<.01) (Supplemental Figure 6F). Our findings emphasise the existence of distinct *NPM1*-related subsets, highlighting the limitations of relying solely on genetic classification.

Gene expression profiling identifies five transcriptional AML-MRC-related clusters

The ICC 2022 divides AML-MRC into three groups based on *TP53* mutations, myelodysplasia-related gene mutations, and cytogenetic abnormalities⁷. Our study identified five gene expression-based AML-MRC related clusters (Figure 5.5), with varying fractions of *TP53* mutations, MRC gene mutations, and cytogenetic abnormalities.

Despite sharing these mishaps, each cluster had unique characteristics (Figure 5.5A). AML-MRC(1)^T was characterised by *IDH1*-R132 (49%, FDR<.001) and *IDH2*-R170 (27%, FDR<.001) mutations. A *DNMT3A* and *IDH1/2* mutated subtype has been reported⁵, but 41% of the AML-MRC(1)^T cases lacked *DNMT3A* mutations. AML-MRC(2)^T, AML-MRC(3)^T, and AML-MRC(4)^T were all enriched (FDR<.001) for *TP53* mutations, cytogenetic abnormalities and high *MECOM* expression, and AML-MRC(3)^T also contained a large fraction of mutated MRC genes (65%, FDR<0.001). AML-MRC(5)^T stood out with the highest fraction of mutated MRC genes cases (81%, FDR<.001) and the lowest fraction of *TP53* mutations (6%) and cytogenetic abnormalities (34%).

Mapping AML Heterogeneity: Multi-cohort transcriptomic analysis identifies novel clusters and divergent ex-vivo drug responses



Results

Figure 5.4: **Gene expression profiling identifies five transcriptional *NPM1*-related clusters.** **A** Waterfall plot of aberrations in the *NPM1*-related clusters, including the percentage of samples with the aberration. The plot is combined with heatmaps showing the expression of marker genes and cell type scores. The columns are samples, which are split according to transcriptional clusters. **B** Boxplot showing the scaled variant allele frequency (VAF) of mutated *NPM1* from the BEAT, Leucegene, and LUMC cohorts. The VAF was scaled per gene and study to allow for a combined analysis. We used a two-sided Wilcoxon test to test for statistical differences and Benjamini-Hochberg to adjust p-values for multiple testing. **C** tSNE visualisation of patient samples, coloured according to the FAB annotation. Only *NPM1*-related clusters are coloured; the rest are in grey. **D** Fraction of FAB annotations per cluster. **E** and **F** show boxplots of HLA I and HLA II antigen presenting cell scores, respectively. Tests were performed as in **B**, but were only done between the two KMT2A^T, the five *NPM1^T and the five AML-MRC^T clusters. FDR values: * < 0.05, ** < 0.01, *** < 0.005, **** < 0.001. HSC = hematopoietic stem cells, Prog. = progenitor, GMP = granulocyte-monocyte progenitor, Prom. = promonocytes, Mono. = monocytes, cDC = conventional dendritic cells, VAF = variant allele frequency*

We found marker genes for all clusters (Figure 5.5A). For instance, high *SRSF12* marked AML-MRC(1)^T, and *LINC00865* marked AML-MRC(5)^T. AML-MRC(2)^T presented high glycophorin genes and *UROD* expression, suggesting an association with acute erythroid leukaemia^{48–50}. Distinct cell differentiation scores further highlighted differences (Figure 5.5A–C, Supplemental Figure 12). For example, AML-MRC(1)^T showed high progenitor-like scores, with 65% M1 (minimal maturation) cases, and AML-MRC(3)^T showed a more differentiated pattern, with 54% M2 (significant maturation) cases. AML-MRC(2)^T was the only cluster with M6 (erythroid leukaemia) – in line with high expression of erythrocyte cell markers – and M7 (megakaryocytic leukaemia) cases. Additionally, high cytolytic cell infiltration has been reported for AML-MRC cases³⁹. Using the same score (Figure 5.5D) we found that the cytolytic infiltration was significantly (FDR<.05) lower for AML-MRC(1)^T and higher for AML-MRC(2)^T compared to the other AML-MRC clusters. Our results demonstrate that different AML-MRC transcriptomic clusters can be identified, showing genetic enrichments that do not necessarily align with the ICC 2022 classification.

AML clusters exhibit cell type-independent differences in ex-vivo drug responses


Finally, we assessed the drug sensitivity of the transcriptional AML subtypes. Using ex-vivo drug response data, we discovered 101 drug-cluster combinations with significantly lower resistance (FDR<.05), of which 21 combinations remained statistically significant (FDR<.05) when adjusting for cell differentiation status (Figure 5.6, Supplemental Figure 13, Supplemental Table 3).

The ex-vivo drug responses between *NPM1*-related clusters were often divergent, exemplified by venetoclax and selumetinib (Figure 5.6A,B). *NPM1*(1)^T, *NPM1*(2)^T and *NPM1*(3)^T mostly responded positively to tyrosine kinase inhibitors and CDK kinase inhibitors. *NPM1*(4)^T and *NPM1*(5)^T samples were more sensitive to PI3K and MAPK kinase inhibitors. We also found drugs where only one cluster was responsive, exemplified by axitinib for *NPM1*(4)^T where this effect remained significant (FDR<.05) when controlled for cell type scores (Figure 6C).

Several drugs demonstrated favourable ex-vivo responses in KMT2A(1)^T compared to the other clusters, exemplified by idelalisib. For KMT2A(2)^T we found no significant responsive drugs, but testing was limited due to small cluster size. For the AML-MRC clusters, most drugs showed strong resistance. Still, specific drugs were more effective for AML-MRC(1)^T, AML-MRC(4)^T, and AML-MRC(5)^T (Figure 5.6A, B), suggesting potential for targeted treatments in this diverse, high-risk patient group.

Next, we examined if transcriptional clusters provide insights beyond genetic classifications. Comparing the AUCs of each drug between groups, we found 71 drugs with significantly different (FDR<.05) median AUCs between the clusters, while only 21 drugs were significantly different between ICC 2022 classes (Figure 5.6D). Additionally, 57 of the 101 cluster-drug combinations remained significant (FDR<.05) when cluster membership and ICC 2022 diagnosis were included in a LM, suggesting that the transcriptional clusters offer information beyond genetic classification.

Mapping AML Heterogeneity: Multi-cohort transcriptomic analysis identifies novel clusters and divergent ex-vivo drug responses



Results




Figure 5.6: AML clusters exhibit cell type-independent differences in ex-vivo drug responses **A** Heatmap coloured according to the median scaled area under the curve (AUC) of the ex-vivo drug response per drug and cluster. On the left is the drug name, and on the right is the drug family. A green colour indicates a lower median AUC for the drug for the samples in the cluster compared to the other clusters, indicating a strong drug response. Red indicates a higher median AUC, meaning a weak drug response. **B** Boxplots showing ex-vivo drug responses for a selection of drugs. We performed significance testing using a two-sided Wilcoxon test. FDR values: * < 0.05, ** < 0.01, *** < 0.05, **** < 0.001. **C** Plots of the multivariate linear models with cluster membership and the six cell type scores as independent variables and AUC as the dependent variable. On the x-axis, the coefficient of the variables in the models is shown, and the y-axis shows the $-\log_{10}$ of the p-value for each variable. The shown p-values are not corrected and are for visualisation to indicate variable importance in the multivariate model. The corrected p-values of the cluster membership variable are shown in **Supplemental Table 3**. The red line indicates a p-value of 0.05. **D** Barplots of the top 40 drugs with highest corrected p-values for Kruskal-Wallis tests between ex-vivo drug response and clusters or ICC 2022 diagnosis to test if there were significant differences in the median AUCs. All p-values were corrected with Benjamini-Hochberg.

Overall, our findings offer novel opportunities for targeted therapy in AML. We observed effective drug responses even after adjusting for differentiation status, possibly allowing gene expression-based subtypes to guide treatment strategies.

5.4 Discussion

This study presents an overview of transcriptomics in AML and provides a framework for transcriptional subtyping. We integrate multiple cohorts to identify 17 robust transcriptional subtypes that subclassify ~75% of our datasets' patients. We make the harmonised data and a cluster predictor publicly available, facilitating future research.

For the CEBPA^T cluster, we show that patients without a detected *CEBPA* bZIP indel mutation still have similar favourable survival. Patients without the canonical *CEBPA* mutation in the CEBPA^T cluster could be explained by undetected mutations, given the complexities of CEBPA sequencing. Also, *CEBPA* hypermethylation has been described to lead to a similar expression profile⁵¹. The use of the CEBPA^T gene signature for risk stratification could be a relevant alternative to detect these favourable-outcome patients.

KMT2A(1)^T mainly featured *KMT2A* with the fusion partners *MLLT3*, *MLLT10* and *MLLT1* – all TFs in the super elongation complex whose perturbation leads to disrupted hematopoietic lineage commitment⁵². In contrast, KMT2A(2)^T featured *KMT2A::MLLT4*, which is thought to cause leukaemia by promoting self-association⁵³. Interestingly, *MLLT3*, *MLLT10* and *MLLT1* all fuse a specific region of *KMT2A*, but *MLLT4* shows less specificity⁵⁴. Collectively, these results suggest that two types of oncogenic mechanisms involving *KMT2A* fusions exist that may be marked with unique gene expression patterns.

We identified five *NPM1*-related clusters, further underpinning findings of transcriptional heterogeneity among *NPM1*-mutated patients^{15,16,55}, but also providing additional insight into co-mutations and detailed subtypes. We observed several samples from *NPM1*-related clusters that lacked the *NPM1* mutation. Several rare *NPM1* fusions, like *NPM1::MLF1*, have been described to lead to cytoplasmic localisation of *NPM1*, comparable to the canonical *NPM1* frameshift^{45,56}. These non-canonical mishaps could lead to a *NPM1*-mutated-like presentation and similarities in survival and drug response should be explored. Additionally, NPM1(1)^T was mutually exclusive enriched for *IDH1/2* and *TET2* co-mutations. *IDH1/2* mutations lead to an aberrant alpha-ketoglutarate metabolism and are functionally complementary to *TET2* loss-of-function mutations⁵⁷. While NPM1(1)^T and NPM1(5)^T show significant enrichments for *TET2*, only NPM1(1)^T shows this mutual exclusivity with *IDH1/2*. This suggests that only NPM1(1)^T is driven by aberrant alpha-ketoglutarate metabolism, which should be further studied using metabolomics.

Similarly, cytogenetic abnormalities, AML-MRC mutations and high MECOM expression were found in all AML-MRC clusters, but lead to different gene expression. A possible explanation could be clonal architecture and the differentiation state of the cell acquiring the leukemic aberration, both known to influence the biology of the resultant leukaemia^{58,59}. To our knowledge, we are the first to show different gene expression-based subgroups in AML-MRC, with divergent drug responses. Accurate identification of these clusters requires gene expression analysis, showing the relevance of our work.

We found no additional survival differences between other clusters. However, data availability limited the survival analysis, and different treatment protocols across studies could have led to

Methods

confoundment. Survival differences between transcriptional subtypes should thus be further explored in one large cohort. However, we did find marked differences in drug responses between the clusters. Ideally, new studies should test in-patient efficacy of drugs with good ex-vivo responses in transcriptional subtypes. Furthermore, transcriptional AML subtyping could aid specialists in the complex field of clinical care and lead to multidisciplinary tailored-based treatment advice⁶⁰.

In conclusion, the transcriptional subtypes reveal heterogeneity in AML not captured by genetic classification. Integration of transcriptomics into AML research and diagnostics could improve disease understanding and lead to more treatment options.

5.5 Methods

Transcriptomic data

We acquired transcriptomics data of primary AML patients from blood or bone marrow from BEAT^{3,20} (n = 425), TARGET² (n = 145), TCGA¹ (n = 150), and Leucegene^{11,21–23} (n = 399), and our in-house LUMC²⁴ dataset (n = 95). Data statements and methods for transcriptome sequencing are available in the referenced studies.

We acquired quantified gene expression for BEAT, TARGET and TCGA from www.portal.gdc.cancer.gov (release 36) and implemented the same pipeline for Leucegene and LUMC to harmonise quantification. In short, FASTQ files were aligned and quantified with STAR²⁵ to the GRCh38 reference genome²⁶, using Gencode v36²⁷ as the gene annotation index which included 60 600 genes.

Gene expression count data were corrected with Combat-Seq²⁸ for the variables “cohort”, “sex”, and “tissue”. We split Leucegene for the batch correction into Leucegene.stranded and Leucegene.unstranded, since different sequencing libraries were used. We removed 8057 genes that were not detected in all cohorts, leaving 52 603 genes. Finally, we removed genes detected in less than 200 samples or with less than 300 counts leaving 41 862 genes for our final dataset. We normalised the corrected count data using the geometric mean and variance stabilising transformation (VST)²⁹ and quantified the remaining cohort-specific variation using kBET³⁰.

Genetic and patient data

We acquired genetic data for the samples from the referenced studies in the form of mutation and fusion calling, and cytogenetics data and clinical data on sex, age, blast percentage, and survival. Blacklisted fusions as reported by Arriba³¹ were removed from the fusion calling data.

We harmonised the data by standardising the annotation of gene names, fusion genes, and karyotyping. Using genetic data, we subclassified samples according to the WHO 2022⁶ and ICC 2022⁷. Samples for which we found no RGA and all genetic data available were annotated as “No RGA found”. We classified samples with missing data and “No RGA found” as “Inconclusive”.

Clustering

We employed consensus clustering^{32,33} on the batch-adjusted gene expression. First, we created a weighted nearest neighbour graph³⁴ using the 2000 most variable genes (MVGs). MVGs were selected via the median absolute deviation from samples with a blast percentage over 70% to minimise tumour microenvironment effects. Using the Leiden algorithm³⁵ – with seed and resolution varied per iteration – we generated 300 cluster assignments from the graph for each n.clusters

ranging from 10 to 20, totaling 3300 assignments.

From these 3300 assignments we created a consensus matrix with values ranging between 0 and 1 based on pairwise co-clustering. We then converted this matrix into a distance matrix (1 - consensus matrix) and conducted Ward.D2 hierarchical clustering. The final cluster count was determined based on the individual separation of WHO classes and clusters displaying differential traits.

Cluster stability

To evaluate per sample clustering stability, we devised a stability score. We constructed a consensus matrix for each n_clusters (300 assignments) and subtracted each co-clustering value from 1 if it was below 0.5. Then, we took the mean of all values per sample as the stability score, which ranged from 0.5 to 1, with higher scores indicating less clustering ambiguity. To investigate correlation between cluster stability and blast percentage we performed a Spearman correlation test. Additionally, we generated tSNEs using 100 to 2500 MVGs to visually assess cluster stability.

Cluster prediction

To predict cluster assignments we trained a one-vs-rest SVM per cluster. As input we used the uncorrected gene expression of the 2000 MVGs used for clustering. To select hyperparameters and evaluate performance we utilised 5x5 nested cross-fold validation.

To improve predictions we included a reject option using a minimum distance to the decision boundary. We determined this distance by looping over possible minimum values for the predictions of the inner fold. We selected the minimum value with the highest Kappa for the accepted inner fold samples and an accuracy < 0.5 for the rejected inner folds samples.

The final model was trained on the whole dataset, using 5-fold cross-validation to select hyperparameters and the minimum decision boundary distance.

Differential gene expression analysis

Differential gene expression analysis between the clusters was performed using DESeq2²⁹ using the corrected gene counts. We performed one-versus-rest analyses to identify differentially expressed genes in one cluster compared to all others. We annotated genes as transcription factors or coding for cell surface proteins using public databases^{36,37}.

Aberration enrichment analysis

To test if aberrations occurred more in a cluster than in others we first removed aberrations found in only one cohort or which occurred in less than 1% of the samples. We also included high *MECOM* expression in the analysis (VST expression > 6 , based on the tail of a *MECOM* expression density plot). We tested for enrichment per aberration by performing a one-sided Fisher exact test for one cluster versus all others and adjusted p-values using the Benjamini-Hochberg (BH) procedure. We considered aberrations with a false discovery ratio (FDR) < 0.05 enriched.

Survival analysis

We performed survival analysis using right censored overall survival data by generating Kaplan Meier (KM) curves on BEAT and TCGA survival data, comparing different groups of patients with the log rank test. We also performed Cox-regression using BEAT, TARGET and TCGA survival data for different patient groups and included cohort, sex and age as co-variables to analyse hazard ratios.

Acknowledgements

Expression based score

We created cell type scores to assess the differentiation arrest of AML samples, using the mean expression of 30 marker genes for six haematological cells³⁸. Additionally, we created immune phenotype scores for cytolytic infiltration and HLA I and HLA II antigen presenting cells using the mean expression of marker genes³⁹.

Drug response analysis

To analyse drug response differences, we used ex-vivo drug response data of 331 BEAT³ samples, quantified as area under the curve (AUC). We excluded drugs with less than 200 samples or missing data for any cluster, leaving 103 of the 123 drugs. We used a Kruskal-Wallis test for each drug with the AUC as response and clusters as groups to compare the average drug response per cluster. Drugs with a significant difference ($FDR < .05$) were analysed with one-sided Wilcoxon tests to identify clusters with low AUCs. Additionally, we performed a Kruskal-Wallis test for each drug with the ICC 2022 diagnosis as groups, to compare with clusters as groups.

Multivariate linear models (LM) were evaluated per drug to test if clusters were sensitive to a drug when adjusted for cell type, with AUC as response and cluster membership (one-versus-rest) and the six cell type scores as explaining variables. Similarly, we fitted LMs but with cluster membership and ICC 2022 diagnoses as variables. We considered clusters sensitive to a drug if the cluster membership's FDR was below 0.05 and the LM coefficient was negative. All p-values were corrected using BH.

5.6 Acknowledgements

This project was funded by a strategic investment of the Leiden University Medical Center, embedded within the Leiden Oncology Center, and executed within the Leiden Center for Computational Oncology. E.B.A. was funded by a personal grant from the Dutch Research Council (NWO; VENI: 09150161810095). The funding bodies had no role in the study design, the collection, analysis, and interpretation of data, the writing of the manuscript, and the decision to submit the manuscript for publication.

5.7 Author Contributions

M.J.R., M.G., and E.B.A. conceived and designed the project; E.B.A. acquired funding; E.B.A. performed project administration; M.G., E.B.A., H.V., R.R.B., C.J.M.H., P.B. performed oversight and management of resources (data generation, collection, transfer, infrastructure, data processing); J.F.S. performed computational and statistical analyses; J.F.S., E.B.A., M.G., E.O.K., E.S.-L. performed analyses and interpretation; J.F.S. performed and structured data visualisation; M.J.R., M.G. and E.B.A. provided supervision and scientific direction; J.F.S. wrote the manuscript; and all authors critically reviewed the manuscript and figures.

DATA AVAILABILITY

The datasets generated and/or analysed during the current study are available at <http://www.osf.io/wq7gx>.

CODE AVAILABILITY

All code used to generate results is available on reasonable request. The predictor is available at www.github.com/jeppe-severens/AMLmapR as an R package.

5.8 References

1. The Cancer Genome Atlas Research Network. Genomic and Epigenomic Landscapes of Adult De Novo Acute Myeloid Leukemia. *N. Engl. J. Med.* **368**, 2059–2074 (2013).
2. Farrar, J. E. *et al.* Genomic Profiling of Pediatric Acute Myeloid Leukemia Reveals a Changing Mutational Landscape from Disease Diagnosis to Relapse. *Cancer Res.* **76**, 2197–2205 (2016).
3. Tyner, J. W. *et al.* Functional genomic landscape of acute myeloid leukaemia. *Nat.* **2018** 5627728 **562**, 526–531 (2018).
4. Papaemmanuil, E. *et al.* Genomic Classification and Prognosis in Acute Myeloid Leukemia. *N. Engl. J. Med.* **374**, 2209–2209 (2016).
5. Tazi, Y. *et al.* Unified classification and risk-stratification in Acute Myeloid Leukemia. *Nat. Commun.* **13**, 4622 (2022).
6. Khoury, J. D. *et al.* The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia* **36**, 1703–1719 (2022).
7. Arber, D. A. *et al.* International Consensus Classification of Myeloid Neoplasms and Acute Leukemias: integrating morphologic, clinical, and genomic data. *Blood* **140**, 1200–1228 (2022).
8. Döhner, H. *et al.* Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood* **140**, 1345–1377 (2022).
9. Burd, A. *et al.* Precision medicine treatment in acute myeloid leukemia using prospective genomic profiling: feasibility and preliminary efficacy of the Beat AML Master Trial. *Nat. Med.* **26**, 1852–1858 (2020).
10. Valk, P. J. M. *et al.* Prognostically Useful Gene-Expression Profiles in Acute Myeloid Leukemia. *N. Engl. J. Med.* **350**, 1617–1628 (2004).
11. Lavallée, V.-P. *et al.* The transcriptomic landscape and directed chemical interrogation of MLL-rearranged acute myeloid leukemias. *Nat. Genet.* **47**, 1030–1037 (2015).
12. Mou, T. *et al.* The transcriptome-wide landscape of molecular subtype-specific mRNA expression profiles in acute myeloid leukemia. *Am. J. Hematol.* **96**, 580–588 (2021).
13. Wouters, B. J. *et al.* Double CEBPA mutations, but not single CEBPA mutations, define a subgroup of acute myeloid leukemia with a distinctive gene expression profile that is uniquely associated with a favorable outcome. *Blood* **113**, 3088–3091 (2009).
14. Taskesen, E. *et al.* Prognostic impact, concurrent genetic mutations, and gene expression features of AML with CEBPA mutations in a cohort of 1182 cytogenetically normal AML patients: further evidence for CEBPA double mutant AML as a distinctive disease entity. *Blood* **117**, 2469–2475 (2011).
15. Mer, A. S. *et al.* Biological and therapeutic implications of a unique subtype of NPM1 mutated AML. *Nat. Commun.* **12**, 1054 (2021).
16. Cheng, W.-Y. *et al.* Transcriptome-based molecular subtypes and differentiation hierarchies improve the classification framework of acute myeloid leukemia. *Proc. Natl. Acad. Sci.* **119**, e2211429119 (2022).
17. de Leeuw, D. C., Ossenkoppele, G. J. & Janssen, J. J. W. M. Older Patients with Acute Myeloid Leukemia Deserve Individualized Treatment. *Curr. Oncol. Rep.* **24**, 1387–1400 (2022).
18. Improved relative survival in older patients with acute myeloid leukemia over a 30-year period in the Netherlands: a long haul is needed to change nothing into something | Leukemia. <https://www.nature.com/articles/s41375-021-01503-y>.
19. Zeng, A. G. X. *et al.* A cellular hierarchy framework for understanding heterogeneity and predicting drug response in acute myeloid leukemia. *Nat. Med.* **28**, 1212–1223 (2022).
20. Bottomly, D. *et al.* Integrative analysis of drug response and clinical outcome in acute myeloid leukemia. *Cancer Cell* **40**, 850–864.e9 (2022).
21. Macrae, T. *et al.* RNA-Seq reveals spliceosome and proteasome genes as most consistent transcripts in human cancer cells. *PLoS One* **8**, e72884 (2013).
22. Lavallée, V.-P. *et al.* RNA-sequencing analysis of core binding factor AML identifies recurrent ZBTB7A mutations and defines RUNX1-CBFA2T3 fusion signature. *Blood* **127**, 2498–2501 (2016).
23. Pabst, C. *et al.* GPR56 identifies primary human acute myeloid leukemia cells with high repopulating potential in vivo. *Blood* **127**, 2018–2027 (2016).

References

24. Arindrarto, W. *et al.* Comprehensive diagnostics of acute myeloid leukemia by whole transcriptome RNA sequencing. *Leuk.* 2020 351 35, 47–61 (2020).
25. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinforma. Oxf. Engl.* 29, 15–21 (2013).
26. Genome Reference Consortium. <https://www.ncbi.nlm.nih.gov/grc>.
27. Frankish, A. *et al.* GENCODE 2021. *Nucleic Acids Res.* 49, D916–D923 (2021).
28. Zhang, Y., Parmigiani, G. & Johnson, W. E. ComBat-seq: batch effect adjustment for RNA-seq count data. *NAR Genomics Bioinforma.* 2, lqaa078 (2020).
29. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550 (2014).
30. Büttner, M., Miao, Z., Wolf, F. A., Teichmann, S. A. & Theis, F. J. A test metric for assessing single-cell RNA-seq batch correction. *Nat. Methods* 16, 43–49 (2019).
31. Uhrig, S. *et al.* Accurate and efficient detection of gene fusions from RNA sequencing data. *Genome Res.* 31, 448–460 (2021).
32. Monti, S., Tamayo, P., Mesirov, J. & Golub, T. Consensus Clustering: A Resampling-Based Method for Class Discovery and Visualization of Gene Expression Microarray Data. *Mach. Learn.* 52, 91–118 (2003).
33. Jeub, L. G. S., Sporns, O. & Fortunato, S. Multiresolution Consensus Clustering in Networks. *Sci. Rep.* 8, 3259 (2018).
34. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. Preprint at <https://doi.org/10.48550/arXiv.1802.03426> (2020).
35. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected communities. *Sci. Rep.* 9, 5233 (2019).
36. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* 172, 650–665 (2018).
37. Bausch-Fluck, D. *et al.* A Mass Spectrometric-Derived Cell Surface Protein Atlas. *PLOS ONE* 10, e0121314 (2015).
38. van Galen, P. *et al.* Single-Cell RNA-Seq Reveals AML Hierarchies Relevant to Disease Progression and Immunity. *Cell* 176, 1265–1281.e24 (2019).
39. Dufva, O. *et al.* Immunogenomic Landscape of Hematological Malignancies. *Cancer Cell* 38, 380–399.e13 (2020).
40. Patel, J. P. *et al.* Prognostic Relevance of Integrated Genetic Profiling in Acute Myeloid Leukemia. *N. Engl. J. Med.* 366, 1079–1089 (2012).
41. Gracia-Maldonado, G., Clark, J., Mulloy, J. C. & Kumar, A. R. LAMP5 - a Novel Target of MLL-Fusion Proteins Is Required for the Propagation of Leukemia. *Blood* 128, 1512 (2016).
42. Milan, T. *et al.* Epigenetic changes in human model KMT2A leukemias highlight early events during leukemogenesis. *Haematologica* 107, 86–99 (2022).
43. Wakita, S. *et al.* Prognostic impact of CEBPA bZIP domain mutation in acute myeloid leukemia. *Blood Adv.* 6, 238–247 (2022).
44. Brunetti, L. *et al.* Mutant NPM1 maintains the leukemic state through HOX expression. *Cancer Cell* 34, 499–512.e9 (2018).
45. Falini, B. *et al.* Aberrant subcellular expression of nucleophosmin and NPM-MLF1 fusion protein in acute myeloid leukaemia carrying t(3;5): A comparison with NPMc+ AML. *Leukemia* 20, 368–371 (2006).
46. Aguilo, F. *et al.* Prdm16 is a physiologic regulator of hematopoietic stem cells. *Blood* 117, 5057 (2011).
47. Imperato, M. R., Cauchy, P., Obier, N. & Bonifer, C. The RUNX1–PU.1 axis in the control of hematopoiesis. *Int. J. Hematol.* 101, 319–329 (2015).
48. Hollox, E. J. & Louzada, S. Genetic variation of glycoporphins and infectious disease. *Immunogenetics* (2022) doi:10.1007/s00251-022-01280-7.
49. Greaves, M. F., Sieff, C. & Edwards, P. A. W. Monoclonal Antiglycophorin as a Probe for Erythroleukemias. *Blood* 61, 645–651 (1983).
50. Andersson, L. C., Jokinen, M. & Gahmberg, C. G. Induction of erythroid differentiation in the human leukaemia cell line K562. *Nature* 278, 364–365 (1979).
51. Hollink, I. H. I. M. *et al.* Characterization of CEBPA mutations and promoter hypermethylation in pediatric acute myeloid leukemia. *Haematologica* 96, 384–392 (2011).
52. Mohan, M., Lin, C., Guest, E. & Shilatifard, A. Licensed to elongate: a molecular mechanism for MLL-based leukaemogenesis. *Nat. Rev. Cancer* 10, 721–728 (2010).
53. Liedtke, M., Ayton, P. M., Somervaille, T. C. P., Smith, K. S. & Cleary, M. L. Self-association mediated by the Ras association 1

Mapping AML Heterogeneity: Multi-cohort transcriptomic analysis identifies novel clusters and divergent ex-vivo drug responses

domain of AF6 activates the oncogenic potential of MLL-AF6. *Blood* **116**, 63–70 (2010).

54. Meyer, C. *et al.* The KMT2A recombinoome of acute leukemias in 2023. *Leukemia* **37**, 988–1005 (2023).

55. Mason, E. F., Kuo, F. C., Hasserjian, R. P., Seegmiller, A. C. & Pozdnyakova, O. A distinct immunophenotype identifies a subset of NPM1-mutated AML with TET2 or IDH1/2 mutations and improved outcome. *Am. J. Hematol.* **93**, 504–510 (2018).

56. Martelli, M. P. *et al.* Novel NPM1 exon 5 mutations and gene fusions leading to aberrant cytoplasmic nucleophosmin in AML. *Blood* **138**, 2696–2701 (2021).

57. Figueroa, M. E. *et al.* Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell* **18**, 553–567 (2010).

58. Morita, K. *et al.* Clonal evolution of acute myeloid leukemia revealed by high-throughput single-cell genomics. *Nat. Commun.* **11**, 5327 (2020).

59. Krivtsov, A. V. *et al.* Cell of origin determines clinically relevant subtypes of MLL-rearranged AML. *Leukemia* **27**, 852–860 (2013).

60. Willemsen, A. E. C. A. B. *et al.* Molecular tumour boards and molecular diagnostics for patients with cancer in the Netherlands: experiences, challenges, and aspirations. *Br. J. Cancer* **121**, 34–36 (2019).

5.9 Appendix

The appendix was shortened due to environmental considerations. To minimize unnecessary printing and reduce material use, extensive data including 13 supplementary figures and 4 supplementary tables are provided online at: <https://doi.org/10.1038/s41375-024-02137-6>.