



Universiteit
Leiden

The Netherlands

Optimizing care in lumbar radiculopathy and neurogenic claudication: from injection to inference, and from clinician to algorithm

Verheijen, E.J.A.

Citation

Verheijen, E. J. A. (2026, June 30). *Optimizing care in lumbar radiculopathy and neurogenic claudication: from injection to inference, and from clinician to algorithm*. Retrieved from <https://hdl.handle.net/1887/4307337>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4307337>

Note: To cite this publication please use the final published version (if applicable).

Part I

**Optimizing care for
lumbar radiculopathy**

1

General introduction and outline

Lumbosacral radiculopathy, often used interchangeably with sciatica in the literature despite referring more broadly to a condition of the lumbar or sacral nerve roots, is a neurological disorder with a reported lifetime prevalence of up to 43% [1]. Generally, it is characterized by unilateral radicular pain extending along one of the exiting nerve roots and is caused by either lumbosacral disc herniation (LDH) or degenerative spinal stenosis (LSS) due to age-related disc degeneration, hypertrophy of the ligamentum flavum and zygapophyseal joint, and osteophyte formation [2]. Radicular pain is frequently accompanied by functional impairments, back pain and motor-sensory deficits [3]. Originally, nerve root compression was assumed to be the primary cause of symptoms. However, research over the past decades has demonstrated that a significant proportion of patients exhibit severe symptoms despite the absence of nerve root compression on imaging. Conversely, some patients present with clear radiological evidence of nerve root impingement but experience little to no symptoms [4-6]. As a result, the general idea has changed and it is now accepted that the pathophysiology of lumbosacral radiculopathy is a multifaceted interplay involving inflammatory processes and immunological responses in conjunction with mechanical compression [7-10]. Exposure of the nerve root to nucleus pulposus material has been shown to elicit an inflammatory response, including cytokine release and recruitment of pro-inflammatory cells, which exacerbates pain and neural dysfunction. Stenosis can result in nerve root oedema and inflammation caused by local ischemia as a result from congestion of venous blood around the nerve root inducing cytokine release. Additionally, vertebral end plate devascularization may strengthen this response [10-13].

Disease progression, however, usually differs as patients with LDH are younger on average and have a more acute onset of symptoms which may aggravate when coughing, sneezing or leg-straightening, while patients with LSS often are older and experience gradually increasing symptoms that may exacerbate during standing or walking. However, for both patient groups, the impact on daily functioning can be severe, necessitating effective management strategies [14]. Since lumbosacral radiculopathy due to LDH may follow a favorable, self-limiting course, and patient outcomes after one year of follow-up have been shown to be similar between surgical and non-surgical groups, initial treatment is conservative for several weeks to months before considering imaging and non-conservative interventions [15-18]. There is less consensus on the optimal treatment approach for LSS, as its symptoms usually follow a more chronic course, though some patients may still experience spontaneous resolution of their symptoms [19-21]. In both patient groups, oral pain medication often provides inadequate relief, and the substantial impact of the condition leads to significant medical and socio-economic costs [22].

One of the primary interventional treatments for lumbosacral radiculopathy is an epidural steroid injection (ESI), which aims to mitigate inflammation and provide symptomatic relief. The origins of this technique date back to 1895 when Bier anesthetized the lower body of one of his residents by injecting a cocaine solution into the intrathecal space [23]. Building on this, Cathelin and Sicard in the early 1900s adapted the approach by using the caudal route to remain within the epidural space [24,25], and it was not until the 1950s that Robecchi and Capra introduced the first periradicular steroid injection with hydrocortisone, laying the foundation for modern ESI [26]. In addition to the caudal approach, interlaminar and transforaminal techniques have been developed. Particularly the transforaminal epidural steroid injection (TEI) approach is now widely employed as it is considered the most selective one of techniques to deliver corticosteroids near the affected nerve root.

Given its potential to reduce inflammatory mediators and modulate immune responses [11,27-29], TEI is commonly administered as a minimally invasive alternative for patients with lumbar radiculopathy who do not respond adequately to conservative treatment [30]. However, its efficacy remains a topic of debate due to inconsistent findings in the literature. While some studies report significant pain relief and functional improvement, others suggest only limited benefits compared to placebo treatments. A 2012 systematic review and meta-analysis by Pinto et al. found a slight favour for corticosteroids but questioned the clinical utility [31]. Since then, several additional randomized controlled trials have been conducted, specifically for the transforaminal approach, highlighting the need for a renewed evaluation of the effectiveness of TEI.

The effectiveness of TEI in patients with lumbosacral radiculopathy may depend on the underlying etiology; however, this remains uncertain. The majority of studies have focused on patients with MRI-confirmed LDH, while relatively few have evaluated TEI outcomes in those with radiculopathy due to other etiologies or in the absence of a compressive lesion on imaging [32,33]. The studies that have compared the effect of TEI between patients with LDH and those with other etiologies suggest comparable outcomes, though findings vary based on the outcome measure and duration of follow-up [34-36]. If TEI provides symptom relief independent of the underlying pathology, routine pre-treatment MRI scans may not be necessary. One study specifically indicated that MRI prior to TEI may be redundant, as it did not significantly influence treatment effectiveness or clinical decision-making [36]. However, the overall body of evidence remains constrained by the small number of studies and the lack of prospective data.

Beyond the underlying etiology of symptoms, no consensus exists regarding other factors associated with TEI effectiveness. Some studies suggest that patients with a shorter duration of symptoms may experience greater benefit from TEI compared to those with a more chronic history [32], or that early pain relief following TEI could serve as a prognostic indicator of long-term outcomes [37,38]. Various demographic, clinical, and radiological variables have been explored; however, no strong conclusions have been drawn, largely due to the considerable heterogeneity in patient populations, epidural steroid techniques, and outcome measures. This lack of consistency has precluded the development of a predictive model for TEI treatment success and, consequently, the implementation of more patient-tailored therapeutic strategies for patients suffering from lumbosacral radiculopathy.

AIMS AND OUTLINE OF THIS THESIS:

1. The only systematic review and meta-analysis comparing the effectiveness of a transforaminal epidural steroid injection (TEI) to placebo – either a transforaminal injection with saline or a sham-injection in the para-spinal muscle – dates back to 2012 [31]. Since then, multiple randomized controlled trials have been conducted, primarily evaluating transforaminal injections. Given these new studies, a reassessment is warranted to determine whether the additional data alter the existing conclusions. Therefore, this first aim is to systematically review and update the current evidence on TEI efficacy compared to placebo in patients with lumbosacral radiculopathy (Chapter 2).
2. TEI has become an increasingly popular treatment for patients with lumbosacral radiculopathy. Since this condition can be caused by a variety of spinal pathology, it is essential to determine whether TEI effectiveness depends on the underlying etiology. Hence, the second aim is to assess the effectiveness of TEI in patients with MRI-confirmed lumbar disc herniation (LDH) compared to those with alternative causes or no compressive lesion on imaging, using data from a large retrospective cohort (Chapter 3).
3. To optimize treatment strategies, it is essential to identify factors that positively or negatively affect patient outcome after treatment with TEI. Therefore, the third aim is to comprehensively review all studies that have examined demographic, clinical, or radiological parameters and their association with the outcome of a single transforaminal injection in patients with lumbar radiculopathy secondary to disc herniation or spinal stenosis (Chapter 4).
4. Current literature suggests that patients with a shorter duration of symptoms may benefit more from TEI. Moreover, there is some evidence that

routine MRI before TEI may be redundant. At present, patients are treated conservatively for several weeks to months before referral to a neurologist and subsequent MRI examination. It is usually only after these steps that they become eligible for TEI or surgical intervention. During this period, symptoms often remain inadequately managed, impairing daily function and contributing to absenteeism. Given these findings, our fourth aim is to investigate whether patients may benefit from early TEI administration. To address this, we present the protocol for a randomized controlled trial comparing the effectiveness of TEI to usual care in patients experiencing acute lumbar radiculopathy symptoms (Chapter 5).

5. The identification of prognostic factors for TEI is significantly limited by the lack of large cohorts and prospective data. Furthermore, substantial heterogeneity in study designs and treatment protocols—combining outcomes from caudal, interlaminar, and transforaminal injections using various corticosteroids—complicates personalizing treatment strategies. To achieve this, the fifth aim is to develop a prediction model for TEI success. We present the protocol for a large, multi-center, prospective cohort study designed to assess the effects of TEI in patients with MRI-confirmed disc herniation or stenosis, congruent with clinical findings, and to identify demographic, clinical and radiological parameters associated with treatment outcomes (Chapter 6).
6. Vertebral endplate changes, known as Modic changes (MC), have been proposed as markers of inflammatory processes and are associated with disc herniation and less favorable outcomes after surgery [39-46]. It has been suggested that the absence of MC correlates with reduced radicular pain and improved clinical outcomes, whereas their presence may aggravate symptoms [46]. Consequently, in patients with MC at the level of disc herniation, inflammation may play a more significant role, and TEI may be more effective. The sixth aim is to assess this association using prospective data from our large cohort study (Chapter 7).

REFERENCES

1. Konstantinou K, Dunn KM (2008) Sciatica: review of epidemiological studies and prevalence estimates. *Spine (Phila Pa 1976)* 33 (22):2464-2472. doi:10.1097/BRS.0b013e318183a4a2
2. Waldman SD (2019) 84 - Spinal Stenosis. In: Waldman SD (ed) *Atlas of Common Pain Syndromes (Fourth Edition)*. Elsevier, Philadelphia, pp 324-327. doi:https://doi.org/10.1016/B978-0-323-54731-4.00084-0
3. Koes BW, van Tulder MW, Peul WC (2007) Diagnosis and treatment of sciatica. *Bmj* 334 (7607):1313-1317. doi:10.1136/bmj.39223.428495.BE
4. el Barzouhi A, Vleggeert-Lankamp CL, Lycklama à Nijeholt GJ, Van der Kallen BF, van den Hout WB, Jacobs WC, Koes BW, Peul WC (2013) Magnetic resonance imaging in follow-up assessment of sciatica. *N Engl J Med* 368 (11):999-1007. doi:10.1056/NEJMOa1209250
5. van Rijn JC, Klemetso N, Reitsma JB, Majoie CB, Hulsmans FJ, Peul WC, Bossuyt PM, Heeten GJ, Stam J (2006) Symptomatic and asymptomatic abnormalities in patients with lumbosacral radicular syndrome: Clinical examination compared with MRI. *Clin Neurol Neurosurg* 108 (6):553-557. doi:10.1016/j.clineuro.2005.10.003
6. Karppinen J, Malmivaara A, Tervonen O, Pääkkö E, Kurunlahti M, Syrjälä P, Vasari P, Vanharanta H (2001) Severity of symptoms and signs in relation to magnetic resonance imaging findings among sciatic patients. *Spine (Phila Pa 1976)* 26 (7):E149-154. doi:10.1097/00007632-200104010-00015
7. Stafford MA, Peng P, Hill DA (2007) Sciatica: a review of history, epidemiology, pathogenesis, and the role of epidural steroid injection in management. *Br J Anaesth* 99 (4):461-473. doi:10.1093/bja/aem238
8. Wang K, Bao JP, Yang S, Hong X, Liu L, Xie XH, Wu XT (2016) A cohort study comparing the serum levels of pro- or anti-inflammatory cytokines in patients with lumbar radicular pain and healthy subjects. *Eur Spine J* 25 (5):1428-1434. doi:10.1007/s00586-015-4349-4
9. Valat JP, Genevay S, Marty M, Rozenberg S, Koes B (2010) Sciatica. *Best Pract Res Clin Rheumatol* 24 (2):241-252. doi:10.1016/j.berh.2009.11.005
10. Dower A, Davies MA, Ghahreman A (2019) Pathologic Basis of Lumbar Radicular Pain. *World Neurosurg* 128:114-121. doi:10.1016/j.wneu.2019.04.147
11. McLain RF, Kapural L, Mekhail NA (2005) Epidural steroid therapy for back and leg pain: mechanisms of action and efficacy. *Spine J* 5 (2):191-201. doi:10.1016/j.spinee.2004.10.046
12. Mulleman D, Mammou S, Griffoul I, Watier H, Goupille P (2006) Pathophysiology of disk-related sciatica. I.--Evidence supporting a chemical component. *Joint Bone Spine* 73 (2):151-158. doi:10.1016/j.jbspin.2005.03.003
13. Lee BH, Moon SH, Suk KS, Kim HS, Yang JH, Lee HM (2020) Lumbar Spinal Stenosis: Pathophysiology and Treatment Principle: A Narrative Review. *Asian Spine J* 14 (5):682-693. doi:10.31616/asj.2020.0472
14. Deyo RA, Mirza SK (2016) CLINICAL PRACTICE. Herniated Lumbar Intervertebral Disk. *N Engl J Med* 374 (18):1763-1772. doi:10.1056/NEJMcp1512658
15. Chiu CC, Chuang TY, Chang KH, Wu CH, Lin PW, Hsu WY (2015) The probability of spontaneous regression of lumbar herniated disc: a systematic review. *Clin Rehabil* 29 (2):184-195. doi:10.1177/0269215514540919

16. Peul WC, van Houwelingen HC, van den Hout WB, Brand R, Eekhof JA, Tans JT, Thomeer RT, Koes BW (2007) Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med* 356 (22):2245-2256. doi:10.1056/NEJMoa064039
17. Kreiner DS, Hwang SW, Easa JE, Resnick DK, Baisden JL, Bess S, Cho CH, DePalma MJ, Dougherty P, 2nd, Fernand R, Ghiselli G, Hanna AS, Lamer T, Lisi AJ, Mazanec DJ, Meagher RJ, Nucci RC, Patel RD, Sembrano JN, Sharma AK, Summers JT, Taleghani CK, Tontz WL, Jr., Toton JF (2014) An evidence-based clinical guideline for the diagnosis and treatment of lumbar disc herniation with radiculopathy. *Spine J* 14 (1):180-191. doi:10.1016/j.spinee.2013.08.003
18. Lorio M, Kim C, Araghi A, Inzana J, Yue JJ (2020) International Society for the Advancement of Spine Surgery Policy 2019-Surgical Treatment of Lumbar Disc Herniation with Radiculopathy. *Int J Spine Surg* 14 (1):1-17. doi:10.14444/7001
19. Anderson DB, Luca K, Jensen RK, Eyles JP, Van Gelder JM, Friedly JL, Maher CG, Ferreira ML (2021) A critical appraisal of clinical practice guidelines for the treatment of lumbar spinal stenosis. *Spine J* 21 (3):455-464. doi:10.1016/j.spinee.2020.10.022
20. Katz JN, Zimmerman ZE, Mass H, Makhni MC (2022) Diagnosis and Management of Lumbar Spinal Stenosis: A Review. *Jama* 327 (17):1688-1699. doi:10.1001/jama.2022.5921
21. Kreiner DS, Shaffer WO, Baisden JL, Gilbert TJ, Summers JT, Toton JF, Hwang SW, Mendel RC, Reitman CA (2013) An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spinal stenosis (update). *Spine J* 13 (7):734-743. doi:10.1016/j.spinee.2012.11.059
22. Lambeek LC, van Tulder MW, Swinkels IC, Koppes LL, Anema JR, van Mechelen W (2011) The trend in total cost of back pain in The Netherlands in the period 2002 to 2007. *Spine (Phila Pa 1976)* 36 (13):1050-1058. doi:10.1097/BRS.0b013e3181e70488
23. Bier A (1899) Versuche über Cocainisirung des Rückenmarkes. *Deutsche Zeitschrift für Chirurgie* 51 (3):361-369. doi:10.1007/BF02792160
24. Sicard A (1901) Les injections medicamenteuses extra-durales par voie sacrococcygienne. *Compt Rend Soc De Biol* 53:396-398
25. Cathelin F (1901) Une nouvelle voie d'injection rachidienne: methode des injections epidurales par le procede du canal sacre-applications a l'homme. *Compt Rend Soc De Biol* 53:452-453
26. Robecchi A, Capra R (1952) [Hydrocortisone (compound F); first clinical experiments in the field of rheumatology]. *Minerva Med* 43 (98):1259-1263
27. Kantrowitz F, Robinson DR, McGuire MB, Levine L (1975) Corticosteroids inhibit prostaglandin production by rheumatoid synovia. *Nature* 258 (5537):737-739. doi:10.1038/258737a0
28. Johansson A, Hao J, Sjölund B (1990) Local corticosteroid application blocks transmission in normal nociceptive C-fibres. *Acta Anaesthesiol Scand* 34 (5):335-338. doi:10.1111/j.1399-6576.1990.tb03097.x
29. Collighan N, Gupta S (2010) Epidural steroids. *Continuing Education in Anaesthesia, Critical Care and Pain* 10 (1):1-5. doi:10.1093/bjaceaccp/mkp043
30. Manchikanti L, Falco FJ, Singh V, Pampati V, Parr AT, Benyamin RM, Fellows B, Hirsch JA (2012) Utilization of interventional techniques in managing chronic pain in the Medicare population: analysis of growth patterns from 2000 to 2011. *Pain Physician* 15 (6):E969-982

31. Pinto RZ, Maher CG, Ferreira ML, Hancock M, Oliveira VC, McLachlan AJ, Koes B, Ferreira PH (2012) Epidural corticosteroid injections in the management of sciatica: a systematic review and meta-analysis. *Ann Intern Med* 157 (12):865-877. doi:10.7326/0003-4819-157-12-201212180-00564
32. MacVicar J, King W, Landers MH, Bogduk N (2013) The effectiveness of lumbar transforaminal injection of steroids: a comprehensive review with systematic analysis of the published data. *Pain Med* 14 (1):14-28. doi:10.1111/j.1526-4637.2012.01508.x
33. Smith CC, McCormick ZL, Mattie R, MacVicar J, Duszynski B, Stojanovic MP (2020) The Effectiveness of Lumbar Transforaminal Injection of Steroid for the Treatment of Radicular Pain: A Comprehensive Review of the Published Data. *Pain Med* 21 (3):472-487. doi:10.1093/pm/pnz160
34. Maus TP, El-Yahchouchi CA, Geske JR, Carter RE, Kaufmann TJ, Wald JT, Diehn FE (2016) Imaging Determinants of Clinical Effectiveness of Lumbar Transforaminal Epidural Steroid Injections. *Pain Med* 17 (12):2176-2184. doi:10.1093/pm/pnw043
35. Ng LC, Sell P (2004) Outcomes of a prospective cohort study on peri-radicular infiltration for radicular pain in patients with lumbar disc herniation and spinal stenosis. *Eur Spine J* 13 (4):325-329. doi:10.1007/s00586-003-0649-1
36. Cohen SP, Gupta A, Strassels SA, Christo PJ, Erdek MA, Griffith SR, Kurihara C, Buckenmaier CC, 3rd, Cornblath D, Vu TN (2012) Effect of MRI on treatment results or decision making in patients with lumbosacral radiculopathy referred for epidural steroid injections: a multicenter, randomized controlled trial. *Arch Intern Med* 172 (2):134-142. doi:10.1001/archinternmed.2011.593
37. El-Yahchouchi C, Wald J, Brault J, Geske J, Hagen C, Murthy N, Kaufmann T, Thielen K, Morris J, Diehn F, Amrami K, Carter R, Shelerud R, Maus T (2014) Lumbar transforaminal epidural steroid injections: does immediate post-procedure pain response predict longer term effectiveness? *Pain Med* 15 (6):921-928. doi:10.1111/pme.12347
38. Joswig H, Neff A, Ruppert C, Hildebrandt G, Stienen MN (2016) The Value of Short-Term Pain Relief in Predicting the One-Month Outcome of Lumbar Transforaminal Epidural Steroid Injections. *World Neurosurg* 96:323-333. doi:10.1016/j.wneu.2016.09.016
39. Dudli S, Miller S, Demir-Deviren S, Lotz JC (2018) Inflammatory response of disc cells against *Propionibacterium acnes* depends on the presence of lumbar Modic changes. *Eur Spine J* 27 (5):1013-1020. doi:10.1007/s00586-017-5291-4
40. Viswanathan VK, Shetty AP, Rajasekaran S (2020) Modic changes - An evidence-based, narrative review on its patho-physiology, clinical significance and role in chronic low back pain. *J Clin Orthop Trauma* 11 (5):761-769. doi:10.1016/j.jcot.2020.06.025
41. Heggli I, Laux CJ, Mengis T, Karol A, Cornaz F, Herger N, Aradi-Vegh B, Widmer J, Burkhard MD, Farshad-Amacker NA, Pfammatter S, Wolski WE, Brunner F, Distler O, Farshad M, Dudli S (2023) Modic type 2 changes are fibroinflammatory changes with complement system involvement adjacent to degenerated vertebral endplates. *JOR Spine* 6 (1):e1237. doi:10.1002/jsp2.1237
42. Dudli S, Sing DC, Hu SS, Berven SH, Burch S, Deviren V, Cheng I, Tay BKB, Alamin TF, Ith MAM, Pietras EM, Lotz JC (2017) ISSLS PRIZE IN BASIC SCIENCE 2017: Intervertebral disc/bone marrow cross-talk with Modic changes. *Eur Spine J* 26 (5):1362-1373. doi:10.1007/s00586-017-4955-4

43. Modic MT, Steinberg PM, Ross JS, Masaryk TJ, Carter JR (1988) Degenerative disk disease: assessment of changes in vertebral body marrow with MR imaging. *Radiology* 166 (1 Pt 1):193-199. doi:10.1148/radiology.166.1.3336678
44. Dudli S, Karol A, Giudici L, Heggli I, Laux CJ, Spirig JM, Wanivenhaus F, Betz M, Germann C, Farshad-Amacker N, Brunner F, Distler O, Farshad M (2022) CD90-positive stromal cells associate with inflammatory and fibrotic changes in modic changes. *Osteoarthritis Cartilage* 4 (3):100287. doi:10.1016/j.ocarto.2022.100287
45. Albert HB, Kjaer P, Jensen TS, Sorensen JS, Bendix T, Manniche C (2008) Modic changes, possible causes and relation to low back pain. *Medical Hypotheses* 70 (2):361-368. doi:10.1016/j.mehy.2007.05.014
46. Djuric N, Yang X, Ostelo R, van Duinen SG, Lycklama À Nijeholt GJ, van der Kallen BFW, Peul WC, Vleggeert-Lankamp CLA (2019) Disc inflammation and Modic changes show an interaction effect on recovery after surgery for lumbar disc herniation. *Eur Spine J* 28 (11):2579-2587. doi:10.1007/s00586-019-06108-9