



Universiteit
Leiden
The Netherlands

Safety of synthetic and biological DMARDs: a systematic literature review informing the 2025 update of the EULAR recommendations for the management of rheumatoid arthritis

Laskou, F.; Konzett, V.; Smolen, J.S.; Edwards, C.J.; Aletaha, D.; Heijde, D. van der; ... ; Kerschbaumer, A.

Citation

Laskou, F., Konzett, V., Smolen, J. S., Edwards, C. J., Aletaha, D., Heijde, D. van der, ... Kerschbaumer, A. (2026). Safety of synthetic and biological DMARDs: a systematic literature review informing the 2025 update of the EULAR recommendations for the management of rheumatoid arthritis. *Annals Of The Rheumatic Diseases*, 85(6), 1055-1072. doi:10.1016/j.ard.2026.02.024

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/4308296>

Note: To cite this publication please use the final published version (if applicable).



Rheumatoid arthritis

Safety of synthetic and biological DMARDs: a systematic literature review informing the 2025 update of the EULAR recommendations for the management of rheumatoid arthritis

Faidra Laskou^{1,*}, Victoria Konzett², Josef S Smolen², Christopher J Edwards³, Daniel Aletaha², Désirée van der Heijde⁴, Kevin L Winthrop⁵, Tsutomu Takeuchi⁶, Roberto Caporali^{7,8}, Patrick Verschueren⁹, Janet E Pope¹⁰, Kimme Hyrich¹¹, Savia de Souza¹², Tanja A Stamm^{13,14}, Jette Primdahl^{15,16}, Jan W Schoones¹⁷, Robert BM Landewé^{18,19}, Andreas Kerschbaumer^{2,20}

¹ MRC Lifecourse Epidemiology Centre Southampton, University Hospital Southampton NHS Foundation Trust, Southampton, UK

² Department of Medicine III, Division of Rheumatology, Medical University of Vienna, Vienna, Austria

³ NIHR Southampton Clinical Research Facility, University Hospital Southampton, Southampton, UK

⁴ Department of Rheumatology, Leiden University Medical Center, Leiden, The Netherlands

⁵ School of Public Health, Oregon Health & Science University, Portland, OR, USA

⁶ Department of Rheumatology and Applied Immunology/Clinical Immunology, Saitama Medical University, Moroyama-machi, Iruma, Saitama, Japan

⁷ Department of Clinical Science and Community Health, University of Milan, Milan, Italy

⁸ Department of Rheumatology and Medical Sciences, ASST Pini-CTO, Milan, Italy

⁹ Department of Rheumatology, University Hospitals Leuven and Skeletal Biology and Engineering Research Centre, KU Leuven, Leuven, Belgium

¹⁰ Schulich School of Medicine, University of Western Ontario, London, ON, Canada

¹¹ Centre for Musculoskeletal Research, The University of Manchester and NIHR Manchester Biomedical Research Centre, Manchester University NHS Trust, Manchester, UK

¹² Clinical Research Facility, King's College Hospital, London, UK

¹³ Institute of Outcomes Research, Centre for Medical Data Science, Medical University of Vienna, Vienna, Austria

¹⁴ Ludwig Boltzmann Institute for Arthritis and Rehabilitation, Vienna, Austria

¹⁵ Danish Hospital for Rheumatic Diseases, University Hospital of Southern Denmark, Sønderborg, Denmark

¹⁶ Department of Regional Health Research, University of Southern Denmark, Odense, Denmark

¹⁷ Directorate of Research & Valorisation, Leiden University Medical Center, Leiden, The Netherlands

¹⁸ Amsterdam Rheumatology Center, AMC, Amsterdam, The Netherlands

¹⁹ Rheumatology, Zuyderland Medical Center, Heerlen, The Netherlands

²⁰ Division of Immunology and Rheumatology, Department of Medicine, Stanford University, School of Medicine, Stanford, CA, USA

*Correspondence to Dr Faidra Laskou, MRC Lifecourse Epidemiology Centre Southampton, University Hospital Southampton NHS, Foundation Trust, Southampton, UK.

E-mail address: f.laskou@soton.ac.uk (F. Laskou).

Handling editor Iain McInnes

<https://doi.org/10.1016/j.ard.2026.02.024>

ARTICLE INFO

Article history:

Received 8 December 2025

Received in revised form 24 February 2026

Accepted 26 February 2026

ABSTRACT

Objectives: This study aims to perform a systematic literature review (SLR) concerning the safety of synthetic and biological disease-modifying antirheumatic drugs (DMARDs) for the 2025 update of European Alliance of Associations for Rheumatology (EULAR) recommendations for the management of rheumatoid arthritis (RA).

Methods: Medline, Embase, Cochrane CENTRAL, and Web of Science were searched for observational and randomised controlled trials with a primary endpoint of DMARD safety on the conventional synthetic (cs-), biological (b-), and targeted synthetic-DMARDs, as well as glucocorticoids, published between January 14, 2022, and January 22, 2025. Separate searches on DMARD monitoring were conducted from database inception to January 22, 2025. A comparator group was required for inclusion. All safety outcomes were included.

Results: A total of 3837 articles were identified, with 321 selected for full-text review; 71 articles were included. Across the evidence base, infections were the most frequently assessed outcome: 13 studies examined serious or hospitalised infections, usually as composite endpoints of bacterial, opportunistic, and herpes zoster infection, while 1 addressed nonserious infections. Serious infections were more common with bDMARDs than csDMARDs. Janus kinase inhibitors (JAKis) showed a higher herpes zoster risk than bDMARDs. Tuberculosis risk was not increased with JAKis compared with bDMARDs, but was higher with infliximab and adalimumab compared with etanercept. Fifteen studies evaluated malignancy, split evenly between analyses of any malignancy and those excluding nonmelanoma skin cancer (NMSC); 2 focused on melanoma and 2 on NMSC. Increased NMSC was noted in patients with RA using DMARDs compared with the general population, with no link to a specific DMARD. Cardiovascular and thromboembolic events were reported in 20 studies. No consistent evidence of increased major adverse cardiovascular events risk with JAKis compared with bDMARDs was identified. Venous thromboembolism risk appeared elevated with JAKis compared with bDMARDs, driven mainly by pulmonary embolism. Fourteen studies reported retention and adverse event–related withdrawals, and 8 assessed other specific adverse events. Gastrointestinal perforation and demyelinating disease were each reported in 3 studies. No eligible articles were identified in searches on DMARD monitoring.

Conclusions: There has been a notable increase in studies evaluating safety outcomes, with the majority of these being observational studies focusing primarily on malignancy, thromboembolic, and cardiovascular events, with most studies pertaining to JAKi safety. A substantial proportion of studies in this SLR relied on claims databases to evaluate safety outcomes, a practice that carries important methodological limitations for safety research. Surprisingly, not many studies looked into glucocorticoid safety outcomes over the past 3 years. This SLR, along with the SLR on efficacy of DMARDs, informed the 2025 update of the EULAR recommendations for management of RA with synthetic and biological DMARDs.

INTRODUCTION

In comparison with the general population, patients with rheumatoid arthritis (RA) are at an increased risk of morbidity and mortality, particularly those with more severe disease activity [1,2]. Disease-modifying antirheumatic drugs (DMARDs) provide effective treatment for RA that can alleviate pain, joint damage, and disability [3]. DMARDs currently approved for RA include conventional synthetic DMARDs (csDMARDs), biological DMARDs (bDMARDs), and targeted synthetic DMARDs (tsDMARDs) [4]. DMARDs is an umbrella term including drugs with different modes of action and molecular targets such as tumour necrosis factor (TNF) alpha, interleukin (IL)-6 receptor, T-cell costimulation (CD80/86), and B-cells (CD20). So far, Janus kinase inhibitors (JAKis) are the only tsDMARDs approved by the U.S. Food and Drug Administration and European Medicines Agency (EMA) for RA. Exploring the associated safety aspects of DMARDs remains clinically important as they influence clinical decision-making. This is particularly important following the recent regulatory changes made by EMA in its recommendations of JAKi use in immune-mediated diseases due to concerns over the increased risk of cardiovascular (CV) and thromboembolic events, and malignancies [5]. Previous

systematic literature reviews (SLRs) have shed light on the safety and efficacy aspects of drugs used to treat RA, with the latest being published in 2022 [6,7]. It is important to update the evidence on safety that has become available from randomised controlled trials (RCTs) and real-world data analyses since then.

Safety concerns associated with many DMARDs can be mitigated by the early identification of toxicity through routine laboratory monitoring [8]. Significant variation in blood monitoring frequency when using DMARDs is described [9]. Current recommendations adopt a cautious stance regarding DMARD blood monitoring frequency, primarily based on clinical trial safety data [10,11]. Considering these gaps, an SLR on DMARD monitoring was incorporated, prompted by the steering committee's decision to broaden the scope of inquiry to encompass previously underexplored aspects of clinical practice.

In order to inform the task force responsible for the 2025 update of the European Alliance of Associations for Rheumatology (EULAR) RA management recommendations, we performed an SLR to update evidence on the safety of csDMARDs, bDMARDs, and tsDMARDs in patients with RA, building upon the previous SLR performed for the corresponding 2022 update [6]. The safety SLR, as well as the SLR on DMARD monitoring, is

WHAT IS ALREADY KNOWN ON THIS TOPIC

- Since the 2022 European Alliance of Associations for Rheumatology (EULAR) recommendations for the management of rheumatoid arthritis (RA), new evidence has emerged on the safety of synthetic and biological DMARDs (bDMARDs) in RA.

WHAT THIS STUDY ADDS

- Serious infections continue to be shown to be more common with bDMARDs when compared to conventional synthetic DMARDs (csDMARDs), while Janus kinase inhibitors (JAKis) show a higher herpes zoster risk than bDMARDs.
- The risk of serious infections was found to be the same for bDMARDs and JAKis; higher rates of serious infections were reported with JAKis compared with interleukin-6 inhibitors in 1 study.
- The risk of malignancy, excluding nonmelanoma skin cancer, was similar between csDMARDs and bDMARDs and across different bDMARDs, and it was not increased with JAKi compared with tumour necrosis factor inhibitors.
- Increased nonmelanoma skin cancer was noted in patients with RA using DMARDs compared with the general population, with no link to a specific DMARD.
- In observational studies, no consistent new evidence of increased major adverse cardiovascular events risk was demonstrated for JAKi when compared with bDMARDs. No new randomised controlled trials investigating safety outcomes as a primary endpoint on JAKi have been published since the last update.
- Venous thromboembolic, mainly pulmonary embolism, risk appeared to be increased in patients treated with JAKi when compared with bDMARDs.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- This review informed the 2025 EULAR recommendations for the management of RA, highlighting new evidence on the safety of synthetic and bDMARDs.

included in this manuscript, while an SLR on efficacy, RA-associated interstitial lung disease, and pre-RA is published separately [12]. Both SLRs provided the task force with the current state of evidence.

METHODS

EULAR standard operating procedures for the development of recommendations and points to consider, as well as the Cochrane Handbook for SLRs and Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines, were adhered to in this project [13–16].

Literature search

The scope of the review was defined during an initial steering committee meeting held on November 16, 2024. Research questions were defined in Population, Intervention, Comparator, Outcome (PICO) format, and a study protocol was distributed among all committee members. Based on these, a literature search strategy was developed by an experienced librarian (JWS), and Medline, Embase, The Cochrane Central Register of Controlled Trials (CENTRAL), and Web of Science, as well as conference abstracts from the 2023 and 2024 EULAR and American College of Rheumatology annual meetings, were screened for all relevant articles without language restrictions.

PICO questions and the detailed search strategies used are provided in the Supplementary Material (Section S1). Seven research questions were formulated for the safety SLR. Observational studies, namely cohort studies/registries with >50 participants, were the main study type in the previous and current SLRs. Studies were only eligible if they included an adequate comparator group (either another DMARD or a combination DMARD therapy strategy); the sole exception was a study that used the general population as a comparator due to a lack of other available studies on pregnancy-related outcomes. Additionally, RCTs with a primary safety outcome were included. Safety data of RCTs (and related long-term extension studies), selected in the accompanying SLR addressing efficacy [12], were also included to assess the safety of drugs with or without available real-world data. Unlike the previous update, no separate SLR on glucocorticoids (GCs) was conducted, but respective trials on GCs were included in the efficacy and safety SLRs.

For the separate SLR focused on DMARD monitoring in patients with RA, both randomised controlled and observational, namely cohort, studies with at least 30 participants were included. Studies were only eligible if they included an adequate comparator group (either another DMARD or a combination DMARD therapy strategy). Details on the additional search are provided in the Supplementary Material (Section S1.2). Ten research questions were formulated for DMARD monitoring (Supplementary Material, Section S1.6).

For all other searches, study duration of at least 6 months and a sample size of at least 50 participants were considered, but smaller trials or studies were also included if the former were not available for specific research questions. Published manuscripts and conference abstracts were considered without language restriction. Publications were included from January 14, 2022, to January 22, 2025, for the update of the SLR for safety of DMARDs; for the DMARD monitoring SLR, we included all search results from database inception until January 22, 2025, as monitoring had not been examined in previous SLRs. Details on complete search strategies are provided in the Supplementary Material.

Data extraction

Two researchers (FL and VK) screened 10% of all articles in duplicates. If an agreement rate of >90% was achieved, the remaining articles were screened by 1 researcher (FL), and uncertainties were discussed with the methodologists (AK and RBML). Selected articles were fully assessed for eligibility, study and baseline characteristics, as well as safety/monitoring outcomes from the finally included articles extracted using a previously prepared data extraction form.

Assessment of risk of bias

The risk of bias (RoB) of each included study was assessed using the ‘Hayden tool’ for observational studies and The Cochrane Collaboration’s tool for RCTs (RoB 2) [17–19].

The RoB assessment in this SLR was conducted using the Hayden tool, which provides a structured framework for evaluating observational studies across key domains, including study participation, attrition, measurement of prognostic factors and outcomes, confounding, and statistical analysis. Applying this tool to real-world evidence presented several challenges, particularly given the variability in exposure and outcome definitions and the frequent presence of residual confounding that could not be fully assessed. Studies rated as high RoB commonly

demonstrated limited representativeness of the underlying population and insufficient reporting on the handling of missing data. A further recurring limitation was the lack of clarity in outcome measurement, which hindered the ability to reliably compare findings across studies.

No meta-analyses were conducted due to the high heterogeneity of studies and patient populations included. Decisions on study selection, extraction, and RoB assessment were discussed with a methodologist (either AK or RBML) whenever necessary.

RESULTS

Safety SLR

For the safety SLR, a total of 3837 abstracts were obtained from the database search, of which 3502 were screened following deduplication. A total of 321 articles were selected for full-text review, and 71 articles were finally included. No RCTs with a primary safety outcome were identified; in total, 13 RCTs from the efficacy SLR were included. Thirteen RCTs investigating DMARDs with efficacy as a primary endpoint were included; these studies were selected from the efficacy SLR that was also submitted separately [12]. We investigated the reported safety events from these RCTs, but no RCTs with a primary endpoint of a safety outcome were identified.

For DMARD monitoring, 3835 abstracts were obtained, with 3698 screened following deduplication. Thirty articles were sought for retrieval and assessment of eligibility, of which none were deemed eligible for inclusion. Flowcharts of these 2 literature searches are provided in [Figures 1 and 2](#). Studies were heterogeneous, precluding meta-analysis; therefore, results are presented descriptively.

A summary of all outcomes examined and the total number of studies included for each outcome in the safety SLR is shown in [Table 1](#). A summary of the number of included publications in the efficacy SLR for the different classes of drugs is shown in [Table 2](#). Detailed overviews of all studies included, as well as RoB analyses, baseline characteristics, and safety outcomes, are provided in the [Supplementary Material](#).

Observational studies

Infections

Twenty-one studies evaluated the risk of infections:

Thirteen studies evaluated the risk of serious infections (5 low [20–24], 2 moderate [25,26], and 6 high RoB [27–32]), and 2 studies examined the risk of opportunistic infections (high and moderate RoB, respectively [26,33]).

Six studies evaluated the risk of herpes zoster infection (1 low [24], 1 moderate [26], and 4 high RoB [28,32,34,35]), and 4 evaluated the risk of tuberculosis (TB) (1 low RoB [20], 3 high RoB [36–38]). One study assessed the risk of nonserious infections [34] (high RoB), and 2 assessed the risk of COVID-19 infection [39,40] (both high RoB). No studies examined the risk of pneumocystis pneumonia since the last SLR.

Serious infections

The risk of serious infection was increased with bDMARDs compared with csDMARDs in 2 studies (adjusted hazard ratio [aHR]: 1.74 [1.07–2.81] and 2.3 [1.6–3.3]) [21,23], with the risk remaining increased when compared with non-TNF inhibitor (TNFi) (2.3 [1.2–4.1]) and TNFi (1.9 [1.3–3.0]) separately [21] ([Table 3](#)) [20–32]. These studies investigated the risk of *Staphylococcus aureus* bacteraemia [21] and osteoarticular infection post *Staphylococcus* bacteraemia [23]. These represent an extended follow-up of the same cohort. No increased risk was observed in a study comparing leflunomide and tacrolimus with TNFi (aHR 1.03 [0.87–1.22] and 0.91 [0.77–1.08], respectively) (2 of them at low RoB) [20,22,28].

Five studies that compared different bDMARDs showed overall similar risk of serious infection across different bDMARDs [20,24,27,30,31]. In 2 low RoB studies, the risk of serious infection was similar across different bDMARDs, including TNFi, IL-6 inhibitor (IL-6i), and rituximab, while a lower risk was reported with certolizumab and sarilumab [20,24].

One study, at high RoB, showed different levels of risk across different bDMARDs compared with IL-6i; with TNFis (etanercept, adalimumab, and golimumab), the risk was increased compared with IL-6i (tocilizumab) (aHR 2.40 [1.24–4.61], 1.90 [0.75–4.83], and 1.21 [0.66–2.23], respectively), while with




Figure 1. PRISMA chart illustrating the systematic literature review (SLR) and study selection process for the DMARD safety SLR. DMARD, disease-modifying antirheumatic drug; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; RCT, randomised controlled trial.




Figure 2. PRISMA chart illustrating the systematic literature review (SLR) and study selection process for the DMARD Monitoring SLR. No articles were identified via other methods (manual search). DMARD, disease-modifying antirheumatic drug; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

abatacept, the risk was reduced (aHR 0.89 [0.49-1.62]) [27]. Lower risk with abatacept compared with TNFi was also seen in a high RoB study (aHR 0.610 [0.247-1.508]), while the risk was increased with tocilizumab compared with TNFi (aHR 2.493 [1.275-4.876]) [31].

Five studies compared the risk of serious infections between bDMARDs and JAKi [22,25,26,28,32] (1 study at low RoB [22]); the risk was similar to bDMARDs with JAKi [22,25,26,28], apart from 1 study where the risk was increased when JAKis were compared with IL-6i (aHR 3.45 [1.48-9.04]) (high RoB [32]). In 1 study, the risk of a serious infection was increased in older individuals aged >76 years with tofacitinib compared with bDMARDs [22].

Four studies on serious infections performed subgroup analysis and looked for age-specific associations; 1 study reports the hazard of a first serious infection due to tofacitinib compared with bDMARDs to be increased with increasing age by a factor of 1.05 (95% CI 1.00-1.11) per year, although the relationship was not statistically significant [22]. The increased risk for a serious infection on tofacitinib becomes convincing and potentially clinically relevant after the age of 69 (1.99 [95% CI 1.02-3.90] and 2.05 [95% CI 1.04-4.05], left and right date imputation), while at the age of 76, the risk is higher (2.85 [95% CI 1.27-6.38] and 2.87 [95% CI 1.27-6.52], left and right date imputation); results below age 65 are too uncertain to interpret [22]. Three studies report no significant differences in the incidence or risk of serious infections between JAKis and TNFis for patients aged >65 years with at least 1 atherosclerotic risk factor [25], a similar risk for serious bacterial infections and opportunistic infections among those aged ≥60 [26], and, for those aged ≥75 treated with different bDMARDs, the risk remained similar except for etanercept compared with tocilizumab (adjusted risk ratio (aRR) 0.30 [0.11-0.85] for <65 years, aRR 0.99 [0.50-1.98]

for 65-74 years, and aRR 2.40 [1.24-4.61] for >75 years) [27].

Herpes zoster

The risk for herpes zoster (HZ) was found to be more than 3 times higher among patients with RA on b/tsDMARDs compared with the general population (aHR 3.65 [3.13-4.25]) [24] (Table 4) [24,26,28,32,34,35]. The risk of HZ was assessed in 6 studies in total [24,26,28,32,34,35] (1 low RoB [24]). Two studies examined the risk of HZ across different bDMARDs and JAKis compared with etanercept [24,35]; HZ risk was increased in all studies comparing JAKis to TNFis [26,28,32,34], abatacept [35], or IL-6i [32]. The risk of serious HZ was examined only in 1 high RoB study (aHR 7.43 [3.91-14.11]) [26] comparing JAKis to TNFis. On individual JAKi, the HZ risk was found to be increased with tofacitinib (aHR 2.06 [1.38-3.08] [35] and 4.00 [1.59-10.06]) [24] and baricitinib (aHR 3.82 [2.05-7.09]) [24], while no published data on any other JAKis were identified.

When looking into recurrent HZ infection (defined as active HZ infection on DMARDs with a history of HZ infection pre-DMARD initiation), the relationship remained the same, with an increased risk observed with tofacitinib vs etanercept (aHR 3.01 [1.49-6.11]) or abatacept (aHR 3.69 [1.77-7.69]) [35]. The risk for HZ was the same with baricitinib compared with tofacitinib [28,34] in 2 studies, whereas in 1 study the risk was found to be increased with tofacitinib compared with baricitinib (aHR 1.92 [0.43-8.51]) [32].

Infliximab (aHR 1.36 [1.06-1.74]) and adalimumab (aHR 1.29 [1.02-1.64]) were associated with a significantly increased HZ risk compared with abatacept [34].

Tuberculosis

The risk of TB, although a rare event, was found to be more than 3 times higher among patients with RA on b/tsDMARDs

Table 1
Number of all observational studies and outcomes included in the current safety literature review

2022-2025 SLR Observational studies	
Main outcome	No. of studies n = 71
Infections	
Serious infections	12 (+1)
Any opportunistic infection	1 (+1)
Herpes zoster	2 (+4)
Tuberculosis	3 (+1)
<i>Pneumocystis jirovecii</i>	0
Hospitalisation due to infection	(+1)
Reactivation of hepatitis B	0
Nonserious infections	1
COVID-19	2
Malignancies	
Any malignancy	9
Any excluding NMSC	5
Any lymphoma	(+1)
Any haematological	(+3)
Leukaemia	0
Any skin	1
Melanoma	(+2)
NMSC	(+3)
Mortality	
Cardiovascular events	9
Cardiovascular events	5
MACE combined (MI, stroke, CV death)	11
Heart failure	1 (+1)
MI	1 (+5)
Stroke	(+6)
Arrhythmia	1 (+2)
Venous thromboembolism	1 (+6)
Gastrointestinal perforation	2
Neuroinflammatory events	1
Pregnancy outcomes	1 ^a
Withdrawals due to AEs	14
Glucocorticoid-associated AEs	3 ^b
Any AE	8 ^c

AE, adverse event; CV, cardiovascular; MACE, major adverse cardiovascular events; MI, myocardial infarction; NMSC, nonmelanoma skin cancer; SLR, systematic literature review.

(+in brackets) to indicate studies that examined more than one outcome.

^a Pre-eclampsia.

^b Infection, cardiovascular events, diabetes, fracture.

^c Dyslipidaemia, peripheral oedema, nonsteatosis liver disease, depression, suicide, autoimmune thyroid disease, diabetes, bone mineral density loss and fractures.

compared with the general population (aHR 3.14 [1.60-6.14]) [23]. TB was assessed in 4 studies (1 low RoB) [20]. The risk of TB was not increased with JAKis compared with TNFis or bDMARDs [37,41]. Infliximab and adalimumab increased the risk of TB compared with etanercept (aHR 3.4 [1.6-7.3] and aHR 3.3 [1.4-7.8], respectively) [20] or IL-6i (aHR 3.07 [1.33-7.1]) [36].

Opportunistic infections

Nontuberculous mycobacterial infection was assessed in 1 high RoB study; the risk was not found to be increased with TNFis compared with csDMARDs (aHR 0.517 [0.205-1.301]) [33]. Similarly, the risk of opportunistic infection (including TB, nontuberculous mycobacterial infection, and systemic fungal infections; the majority being TB) was lower with JAKis compared with TNFis (aHR 0.25 [0.09-0.73]) [26].

No *Pneumocystis jirovecii* pneumonia infection was reported in any studies.

COVID-19 infection was assessed in 2 studies; the risk of COVID-19 infection was not increased with JAKis or IL-6i compared with TNFis [39], but hospitalisation, intensive care unit

admission, and severe/fatal events due to COVID-19 infection were increased with rituximab (aHR 2.14 [1.51-3.04], 5.22 [1.77-15.41], and 1.66 [0.89-3.08], respectively) compared with csDMARDs [40].

Malignancy

Fifteen studies evaluated the risk of malignancy, of which 14 evaluated the risk of any type of malignancy (Table 5). Of those 14, 5 studies included any type of malignancy except nonmelanoma skin cancer (NMSC) (1 low [42], 1 moderate [43], and 3 high RoB [44–46]); 1 included any malignancy except haematological malignancy [28] (high RoB), and 1 included any malignancy except skin malignancy [47] (moderate RoB). Two studies examined the risk of melanoma [44,48] (high RoB). Three studies examined the risk of haematological malignancy alone [43,44,49] (high and moderate RoB). Two studies examined the risk of NMSC only [43,45], and 1 study examined the risk between methotrexate and hydroxychloroquine (HCQ) users for any new skin malignancy [48] (high and moderate RoB).

Any type of malignancy

Six studies assessed the risk of any malignancy. The risk of malignancy was similar with csDMARDs and bDMARDs [50,51], including 1 low RoB study [50]. The risk was increased with rituximab, abatacept, and JAKis compared with TNFis in 1 high RoB study [52], whereas the same risk was not found in another study examining different bDMARDs and JAKis vs TNFis [53]; in the latter, risk was increased with IL-6i and cytotoxic T-lymphocyte-associated protein 4 immunoglobulin fusion proteins (CTL4-IgG) compared with TNFis [53]. Similar risk was found between JAKis and TNFis in 3 studies [25,38,49].

Any type of malignancy except NMSC

Seven studies assessed the risk of malignancy excluding NMSC. The risk was similar between csDMARDs and bDMARDs [44,47] and across different bDMARDs [47], while an increased incidence rate was reported with JAKis compared with IL-6i (incidence rate ratio (IRR) 2.13 [0.67-7.42]) (high RoB study and not statistically significant) [46]. The risk was not increased with JAKis compared with TNFis (aHR range from 0.385 [0.095-1.552] to 1.41 [0.76-2.37]) [28,42,43,45,47], while numerically increased risk estimates were detected in 1 study with low RoB [42].

NMSC and melanoma

NMSC incidences in the JAKi, TNFi, and non-TNFi groups were all higher than in the general population [43]; the risk of NMSC has been shown to be increased with JAKis compared with TNFis in 2 studies of moderate (aHR 1.39 [1.01-1.91]) and high RoB (aHR 1.38 [0.39-4.88]); however, no differences were observed among different csDMARDs [43,45,48].

Similarly, no difference was found for the risk of melanoma with bDMARDs compared with csDMARDs, or among different csDMARDs [44,48].

Haematological malignancy

The risk of all haematological malignancies was increased with JAKis compared with TNFis in 2 studies (both high RoB) (aHR 1.90 [0.70-5.16] [42] and aHR 2.86 [0.41-20.00]) [49], while the risk was similar with bDMARDs (aHR 1.35 [0.72-2.53]), TNFi-exposed (aHR 0.77 [0.42-1.43]), and csDMARDs or in combination (aHR 1.18 [0.55-2.52]) [44].

Table 2

Number of included publications in the efficacy SLR for the different classes of drugs

Drug class	No. of publications	No. of RCTs	No. of LTEs	No. of PBO-controlled	No. of biosimilar studies
csDMARDs					
MTX	1	1			
bDMARDs					
PD-1 agonist	1	1		1	
TNFi	3				3
CD40L	1	1		1	
GM-CSF	2	2	2	2	
IL-6Ri	4	4	1		3
CD80/86	1	1		1	
tsDMARDs					
JAKi 1-3	1	1		1	
BTK	1	1		1	
JAK1	1	1	1	1	
Total	16	13	4	8	6

bDMARD, biological disease-modifying antirheumatic drug; BTK, Bruton's tyrosine kinase inhibitor; CD80/86, cluster of differentiation 80/86; csDMARD, conventional disease-modifying antirheumatic drug; CD40L, CD40 ligand; GM-CSF, granulocyte-macrophage colony-stimulating factor; IL-6Ri, interleukin-6 receptor inhibitor; JAK1, Janus kinase 1; JAKi, Janus kinase inhibitor; LTE, long-term extension; MTX, methotrexate; PBO, placebo-controlled; PD-1, programmed death-1; RCT, randomised controlled trial; SLR, systematic literature review; TNFi, tumour necrosis factor inhibitor; tsDMARD, targeted synthetic disease-modifying antirheumatic drug.

Table 3

Serious infections, comparison between different DMARDs (observational studies)

Study ID	Registry	Intervention	Control	aHR/OR	RoB
Shin et al [29], 2022	Claims database	LEF	TNFi	1.03 (0.87-1.22)	High
		TAC		0.91 (0.77-1.08)	
Harigai et al [27], 2024	MDV ^c	ETA	TCZ	2.40 (1.24-4.61)	High
		ADA		1.90 (0.75-4.83)	
		GOL		1.21 (0.66-2.23)	
		ABA		0.89 (0.49-1.62)	
Dieperink et al [21], 2022	DANBIO	bDMARDs	Non-bDMARD users	2.3 (1.6-3.3)	Low
		TNFi		1.9 (1.3-3.0)	
		non-TNFi		2.3 (1.2-4.1)	
Lauper et al [20], 2024	BSRBR-RA	ADA	ETA	1.1 (1.0-1.3)	Low
		IFX		1.3 (1.1-1.5)	
		CERT		0.8 (0.5-1.1)	
		RTX		1.0 (0.8-1.1)	
		TCZ		1.2 (1.0-1.4)	
		ABA		1.2 (0.8-1.7)	
Frisell et al [24], 2023	ARTIS	ADA	ETA	1.12 (0.96-1.31)	Low
		IFX		1.32 (1.09-1.60)	
		CERT		1.10 (0.91-1.33)	
		GOL		1.07 (0.86-1.32)	
		ABA		1.08 (0.91-1.29)	
		RTX		1.31 (1.15-1.50)	
		TCZ		0.98 (0.80-1.19)	
		SAR		0.61 (0.24-1.53)	
		BARI		0.98 (0.75-1.29)	
		TOFA		0.89 (0.47-1.69)	
Ota et al [30], 2024	Claims database	TNFi	IL-6i	0.66 (0.36-1.20)	High
Bellan et al [31], 2022	Claims database	ABA	TNFi	0.61 (0.247-1.508)	High
		TCZ		2.49 (1.275-4.876)	
Dieperink et al [23], 2024	DANBIO	bDMARDs	Non-bDMARD users	1.74 (1.07-2.81)	Low
		Current user		2.27(1.29-3.98)	
Riek et al [22], 2023	SCQM	bDMARDs	TOFA	1.05 (1.00-1.11)	Low
Mok et al [25], 2024	HK registry	JAKi	TNFi	1.08 (0.84-1.39)	Moderate
Uchida et al [28], 2023	Rheumatology departments ^a	JAKi	TNFi	0.79 (0.417-1.502)	High
Choi et al [26], 2023	Claim database	JAKi	TNFi	1.04 (0.71-1.52)	Moderate
Yoshida et al [32], 2024	FMU Rheumatology ^b	JAKi	IL-6i	3.45 (1.48-9.04)	High

ABA, abatacept; ADA, adalimumab; aHR, adjusted hazard ratio; ARTIS, Anti-Rheumatic Therapy in Sweden, BARI, baricitinib; bDMARD, biological disease-modifying antirheumatic drug; BSRBR-RA, British Society of Rheumatology Biologics Register for Rheumatoid Arthritis; c, control; CERT, certolizumab pegol; DANBIO, Danish nationwide quality registry; DMARD, disease-modifying antirheumatic drug; ETA, etanercept; FMU, Fukushima Medical University; GOL, golimumab; HK, Hong Kong biologic registry; i, intervention; IFX, infliximab; IL-6i, interleukin-6 inhibitor; JAKi, Janus kinase inhibitor; LEF, leflunomide; OR, odds ratio; RoB, risk of bias; RTX, rituximab; SAR, sarilumab; SCQM, Swiss Clinical Quality Management in Rheumatic diseases; TAC, tacrolimus; TCZ, tocilizumab; TNFi, tumour necrosis factor inhibitor; TOFA, tofacitinib.

^a Nagasaki University Hospital, Sasebo Chuo Hospital, or Ureshino Medical Center.

^b Department of Rheumatology of Fukushima Medical University Hospital, Japanese Red Cross Fukushima Hospital, and Ohta Nishinouchi General Hospital Foundation.

^c Medical Data Vision Co., Ltd (Tokyo, Japan)

Table 4
Herpes zoster, comparison between different DMARDs (observational studies)

Study ID	Registry	Intervention	Control	aHR	RoB
Frisell et al [24], 2023	ARTIS	ADA	ETA	1.48 (0.90-2.43)	Low
		IFX		1.64 (0.90-2.99)	
		CERT		1.32 (0.73-2.38)	
		GOL		1.20 (0.63-2.28)	
		ABA		0.89 (0.49-1.64)	
		RTX		1.21 (0.77-1.91)	
		TCZ		1.06 (0.52-2.17)	
		SAR		N/A	
		BARI		3.82 (2.05-7.09)	
		TOFA		4.00 (1.59-10.06)	
Jeong et al [35], 2022	Claims database	INF	ETA	1.14 (0.95-1.36)	High
		ADA		1.09 (0.94-1.25)	
		GOL		0.89 (0.69-1.14)	
		TOC		0.74 (0.58-0.94)	
		RTX		1.01 (0.52-1.97)	
		TOFA		2.06 (1.38-3.08)	
		ABA		0.84 (0.66-1.06)	
Uchida et al [28], 2023	Rheumatology departments ^a	JAKi	TNFi	0.20 (0.08-0.52)	High
Choi et al [26], 2023	Claims database	JAKi	TNFi	2.37 (2.00-2.80)	Moderate
Opdam et al [34], 2024	Claims database	JAKi	bDMARDs	2.65 (1.94-3.60)	High
Yoshida et al [32], 2024	Rheumatology departments	JAKi	IL-6i	2.83 (0.87-10.96)	High

ABA, abatacept; ADA, adalimumab; aHR, adjusted hazard ratio; ARTIS, Anti-Rheumatic Therapy in Sweden; BARI, baricitinib; bDMARD, biological disease-modifying antirheumatic drug; c, control; CERT, certolizumab pegol; DMARD, disease-modifying antirheumatic drug; ETA, etanercept; GOL, golimumab; i, intervention; IFX, infliximab; IL-6i, interleukin-6 inhibitor; JAKi, Janus kinase inhibitor; RoB, risk of bias; RTX, rituximab; SAR, sarilumab; TCZ, tocilizumab; TNFi, tumour necrosis factor inhibitor; TOFA, tofacitinib.

^a Nagasaki University Hospital, Sasebo Chuo Hospital, or Ureshino Medical Center.

Nine studies assessed the risk of malignancy while also performing subgroup analysis and looked for age-specific associations; no statistically significant trends in younger and older groups were observed [25,38,44,45,47,49–51]. Analyses stratified by age displayed numerically increased yet statistically not significant hazard ratios (HRs) between age groups (aHR 1.47 [0.79-2.74] in 50+, 1.35 [0.61-2.97] in 65+ years of age, and 1.50 [0.83-2.72] in all ages) in a study examining JAKis compared with TNFis [42].

Cardiovascular events

Twenty studies investigated the risk of CV events (major adverse cardiovascular events [MACE], heart failure, myocardial infarction, stroke, arrhythmias) (4 low RoB [24,54–56] and 16 moderate or high RoB [25,38,43,46,57–64]). Two studies described the risk of heart failure [63,65] (1 low and 1 high RoB, respectively), 6 studies assessed the risk of myocardial infarction (2 low [56,65] and 4 high RoB [45,58,63,65,66]) and stroke (2 low [56,65] and 4 high RoB [41,45,58,63]), while arrhythmias were assessed in 3 studies (1 low [65] and 2 high RoB [63,67]).

MACE

Twelve studies examined the risk of MACE, and 4 studies assessed the risk of CV events (myocardial infarction and/or heart failure, and stroke, but no CV death); the definition of a CV event varied among them [38,58,61,64] (Table 6). The risk of MACE is similar between different csDMARDs (methotrexate [MTX] vs HCQ: aHR 1.07 [0.97-1.18] [59,63]; csDMARDs with MTX vs MTX only: aHR 0.859 [0.375-1.97]; csDMARDs vs MTX: aHR 1.552 [0.766-3.144]) or between csDMARDs and bDMARDs [54–56,59]. No difference was observed with TNFis compared with csDMARDs alone (aHR 0.91 [0.72-1.17] and aHR 0.65 [0.40-1.05]) [54,55].

Similar risk of MACE was found across different bDMARDs with a reported higher rate for certolizumab (aHR 1.31 [0.99-1.73]) and rituximab (aHR 1.27 [1.04-1.55]) compared with etanercept in 1 study [24,56].

The risk of MACE was increased with JAKis compared with TNFis and IL-6i in 2 studies, 1 of moderate and 1 of high RoB (aHR 2.09 [1.10-3.95] [25] and 3.03 [0.77-15.21] [46], respectively), whereas 7 studies reported no increased risk with JAKis compared with TNFis or any bDMARDs (including 3 low RoB studies) [24,45,55–57,60,62].

No significant differences were found across different DMARDs regarding the isolated risk of heart failure, stroke, myocardial infarction, or arrhythmias.

Eight studies on MACE performed subgroup analysis and looked for age-specific associations; age-specific subgroup analysis of patients aged ≥65 years did not find an increased risk of MACE for JAKis compared with TNFis [45], JAKis compared with adalimumab [60], tofacitinib compared with TNFis [57], or JAKis, other bDMARDs and csDMARDs compared with TNFis [55], whereas the risk was found to be similar in patients >60 years of age and in those aged 50+ with a presence of at least 1 CV risk factor when JAKis were compared with TNFis [25,68]. In a high RoB study, the risk of CV disease, not specifically MACE, tended to increase with JAKis compared with that with bDMARDs in patients aged ≥65 years (aHR 1.83 [0.36-9.31] vs 1.13 [0.39-3.30] in patients <65); it is important to report that approximately 90% of participants on JAKis were aged <65, and wide CIs were seen [62]. Finally, for coronary heart disease, the incidence rates between JAKis and TNFis were insignificant in those aged ≥50 (HR 0.68 [0.26-1.81]) and in those <50 years of age (HR 1.03 [0.45-2.36]; high RoB) [38].

Venous thromboembolism

Seven studies examined the risk of thromboembolic events among patients on JAKis and bDMARDs (2 low RoB [65,69] and 5 moderate/high RoB [25,41,45,58,60]) (Table 6).

Table 5
Malignancies, comparison between different DMARDs (observational studies)

Study ID	Registry	Intervention	Control	aHR	RoB
Any malignancy					
Simon et al [50], 2024	ARTIS, BC, FORWARD, RABBIT	csDMARDs bDMARDs	ABA	1.1 (0.8-1.5) 1.0 (0.8-1.3)	Low
Choi et al [51], 2022	Claims database	TNFi	nbDMARDs	0.37 (0.255-0.563)	High
Song et al [49], 2022	Claims database	JAKi	TNFi	0.83 (0.55-1.27)	High
Fang et al [38], 2022	TNHIRD/TDR	JAKi	TNFi	1.10 (0.44-2.78)	High
Molina-Collada et al [53], 2024	BIOBADASER	JAKi	TNFi	0.7 (0.2-2.7)	
		Anti-CD20		0.3 (0.1-1.3)	
		Anti-IL-6		1.7 (0.6-4.3)	
		CTLA4-Ig		1.9 (0.8-4.7)	
		Anti-IL-17		N/A	
Sendaydiego et al [52], 2024	Claims database	RTX	TNFi	1.91 (1.17-3.14)	High
		IL-6i		1.04 (0.57-1.92)	
		ABA		1.47 (1.03-2.11)	
		JAKi		1.36 (0.94-1.96)	
Mok et al [25], 2024	HK Biologic registry	JAKi	TNFi	0.87 (0.39-1.95)	Moderate
Any malignancy except NMSC					
Seror et al [44], 2022	Claims database	bDMARDs bDMARD + csDMARD	csDMARDs	0.99 (0.86-1.14) 0.98 (0.83-1.16)	High
		TNFi exposed		1.03 (0.88-1.21)	
Westermann et al [42], 2024	DANBIO	JAKi	bDMARDs	1.41 (0.76-2.37)	Low
		TNFi	bDMARDs naïve	1.0 (0.9-1.0)	
		RTX		1.0 (0.9-1.1)	
		ABA		1.15 (0.98-1.34)	
		JAKi		1.0 (0.8-1.2)	
Uchida et al [28], 2023	Rheumatology departments ^a	JAKi	TNFi	0.38 (0.095-1.552)	High
Min et al [45], 2023	Claims database	JAKi naïve JAKi treated	TNFi	1.53 (0.81-2.87) 1.02 (0.81-1.29)	High
Huss et al [47], 2022	SRQ	TOFA BARI JAKi Non-TNFi	TNFi	1.08 (0.52-2.24) 0.92 (0.61-1.38) 0.94 (0.65-1.38) 1.12 (0.88-1.43)	Moderate
Yoshida et al [46], 2023	Rheumatology departments ^b	JAKi	IL-6i	2.13 (0.67-7.42)	High
Melanoma					
Lee et al [48], 2023	Claims database	MTX	HCQ	1.39 (0.87-2.21)	High
Seror et al [44], 2022	Claims database	bDMARDs TNFi exposed	csDMARDs	0.91 (0.43-1.79) 0.73 (0.33-1.62)	High
NMSC					
Lee et al [48], 2023	Claims database	MTX	HCQ	1.01 (0.90-1.12)	High
Huss et al [47], 2022	SRQ	JAKi TOFA BARI Non-TNFi	TNFi	1.39 (1.01-1.91) 1.56 (0.83-2.92) 1.37 (0.97-1.92) 1.00 (0.78-1.28)	Moderate
Min et al [45], 2023	Claims database	JAKi	TNFi	1.38 (0.39-4.88)	High
Haematological malignancy					
Seror et al [44], 2022	Claims database	bDMARDs bDMARDs & csDMARDs TNFi exposed	csDMARDs	1.35 (0.72-2.53) 1.18 (0.55-2.52) 0.77 (0.42-1.43)	High
Huss et al [47], 2022	SRQ	JAKi TOFA BARI Non-TNFi	TNFi	1.90 (0.70-5.16) N/A 1.96 (0.66-5.79) 1.04 (0.48-2.25)	Moderate
Song et al [49], 2022	Claims database	JAKi	TNFi	2.86 (0.41-20.00)	High

ABA, abatacept; aHR, adjusted hazard ratio; ARTIS, Anti-Rheumatic Therapy in Sweden, BC, British Columbia Canadian RA Cohort; bDMARD, biological disease-modifying antirheumatic drug; BIOBADASER, Safety Register of the Spanish Society of Rheumatology; c, control; csDMARD, conventional disease-modifying antirheumatic drug; DANBIO, Danish nationwide quality registry; DMARD, disease-modifying antirheumatic drug; FORWARD, National Databank for Rheumatic Diseases longitudinal prospective observational study; HCQ, hydroxychloroquine; HK, Hong Kong biologic registry; i, intervention; IL-17, interleukin-17; IL-6i, interleukin-6 inhibitor; JAKi, Janus kinase inhibitor; MTX, methotrexate; N/A, Not applicable; nbDMARD, nonbiologic disease-modifying antirheumatic drug; NMSC, nonmelanoma skin cancer; RABBIT, Rheumatoid Arthritis Observation of Biologic Therapy registry; RoB, risk of bias; RTX, rituximab; SRQ, Swedish Rheumatology Quality Register; TNFi, tumour necrosis factor inhibitor; TNHIRD/TDR, Taiwan National Health Insurance Research Database/Taiwan Death Registry; TOFA, tofacitinib.

^a Nagasaki University Hospital, Sasebo Chuo Hospital, or Ureshino Medical Center.

^b Department of Rheumatology of Fukushima Medical University Hospital, Japanese Red Cross Fukushima Hospital, and Ohta Nishinouchi General Hospital Foundation.

The risk of thromboembolic events was increased with JAKis and IL-6i compared with TNFis or other bDMARDs in 5 studies [25,38,45,58,65,69], while in 1 high RoB study, the risk was similar between JAKi and adalimumab [60]. The risk of overall

venous thromboembolism (VTE) was increased primarily secondary to pulmonary embolism events (aHR 3.21 [2.11-4.88]) and not deep vein thrombosis (aHR 0.83 [0.47-1.45]) in 1 low RoB study [69]. When studying specific JAKis, the risk was

increased with upadacitinib and baricitinib compared with tofacitinib (low RoB) [65].

Gastrointestinal perforation and demyelinating disease

Intestinal perforations and neuroinflammatory events were assessed in 3 studies, including patients on JAKis and bDMARDs [70–72].

In both studies evaluating the risk of gastrointestinal perforation (GIP) with JAKis compared with other bDMARDs, the risk was found to be increased with any JAKi (Reporting odds ratio (ROR) 1.98 [1.69–2.31]) and with each JAKi: baricitinib (ROR 5.38 [3.46–8.37] and aHR 2 [1.2–3.4]), upadacitinib (ROR 2.73 [2.17–3.44] and aHR 2.8 [1.1–7.2]) and tofacitinib (ROR 1.52 [1.25–1.85] and aHR 1.7 [0.9–3.1]) [70,71]. Both were high RoB studies.

No risk was observed for neuroinflammatory diseases with other TNFis compared with etanercept [72].

Others

Mortality was assessed as a primary endpoint in 3 studies, while in 6 studies it was assessed as a secondary endpoint; 7 studies examined all-cause mortality [24,38,45,53,57,62,67] (2 low and 5 high RoB), while in 2 studies, CV-related mortality was assessed [56,64] (1 low and 1 high RoB, respectively).

Fourteen studies assessed retention rates and withdrawals due to adverse events (11 low RoB [24,73–82] and 3 high RoB [25,83,84]).

Eight studies examined any other adverse events, including dyslipidaemia (high RoB), peripheral oedema [65] (low RoB), nonsteatosis liver disease, depression, suicide [24] (low RoB), autoimmune thyroid disease (high RoB), diabetes [85] (high RoB), and osteoporosis [86–88] (high RoB).

Only 1 study reported the risk of preeclampsia in patients on csDMARDs, bDMARDs, or GCs [89] (high RoB, control was the general population), and 3 studies evaluated GC-associated risks (1 low [20] and 2 high RoB [90,91]). Additional information and details on respective studies are available in the Supplementary Material (Sections S3.1.7–3.111, pages 102–128).

RCTs

Thirteen low RoB RCTs described in the efficacy SLR are included in the safety SLR. One trial evaluated csDMARDs, more specifically, methotrexate [92]. Two phase 2 [93,94] and 4 phase 3 [95–97] trials examined bDMARDs, while 1 phase 2 [98] and 1 phase 3 [99] trials included patients on tsDMARDs. Six RCTs on biosimilars are also included in this SLR [100–104]. Most RCTs were not designed and, therefore, not powered to evaluate safety outcomes. The incidence of major adverse events was low and mostly comparable between active treatment, placebo, or active comparator. No RCTs with a primary safety outcome were found.

Infections

In the sole RCT examining the risk of serious infections comparing oral methotrexate to subcutaneous methotrexate, the risk was equal in both formulations (15.4% vs 18%, respectively). No serious infections were reported in phase 2 bDMARD trials [93,94].

In phase 3 bDMARD trials, and more specifically in CONTRAST 1, comparing otilimab with tofacitinib, the risk of serious

infections was similar across different doses of otilimab and equally common to tofacitinib (range <1%–1% in week 0–12 and 3%–4% in week 12–52) or placebo (1% at week 0–12) [95]. Similar results were seen in CONTRAST 2 (range <1%–1% in week 0–12 and 2%–4% in week 12–52) [95]. When comparing otilimab with sarilumab in CONTRAST 3, more common serious infections were noted in otilimab low dose in weeks 0–24 (3%) compared with sarilumab (1%), while no serious infections were reported in high-dose otilimab. The frequency of serious infections in weeks 0–12 was reported to be the same between otilimab and sarilumab (<1%), while there were no serious infection reports in the placebo group. No reports of serious infections were reported in the AVERT 2 trial when comparing placebo with abatacept [105].

One episode of HZ was reported in the placebo group when compared with peresolimab (death protein 1 [PD-1] inhibitor) [93]. HZ was similarly common when comparing placebo with ivarmacitinib (range 0.6%–2.6%); higher dose of ivarmacitinib resulted in a slightly higher incidence of HZ (1.6% at weeks 0–24, 1.8% in weeks 24–52 for 4 mg ivarmacitinib vs 2.6% and 2.3%, respectively) [99]. From the biosimilar studies, only 1 HZ case (0.4%) was reported in the infliximab generic group [106].

Three opportunistic infections were reported in CONTRAST 1 in the high-dose otilimab group only (1 in weeks 0–12 and 2 in weeks 12–52), while 4 were reported in the otilimab group (3 in the low dose and 1 in the high dose) in CONTRAST 2, compared with 5 cases reported in the tofacitinib group by week 52.

Incidences of TB were similar between otilimab, tofacitinib, or sarilumab, but more common compared with placebo (2%–4% vs 0%, respectively) [95,96].

Malignancy

No malignancies were reported when comparing oral with subcutaneous methotrexate (0% and 0%, respectively) [92], or dazodalibep with placebo (0% and 0%, respectively) [94]. One episode of malignancy was reported in the placebo group when compared with peresolimab (4% vs 0%, respectively) [93].

No malignancies were reported in the abatacept group compared with placebo, where the rate of malignancy was numerically higher (0 [0%] vs 3 [0.7%]) [105]. One episode of malignancy was reported in the placebo group when compared with BMS-986142 (Bruton's tyrosine kinase inhibitor) (1 [<1%] vs 0 [0%]) [98].

Compared with otilimab, the incidence of malignancy was similar to that with sarilumab and tofacitinib in 2 RCTs by week 12 (<1% in all groups), but was numerically slightly higher in patients on tofacitinib by week 52 only (<1% vs 2%, respectively) and compared with placebo (<1%) [95,96]. Malignancy was also numerically higher in ivarmacitinib (0.5% by week 24 and 1.1% by weeks 24–52) compared with placebo (0% by week 24 and weeks 24–52) in 1 RCT [99].

The number of malignancies reported in biosimilar studies was also similar for infliximab, tocilizumab, and adalimumab (range 0.4%–<1%) [100,101,103,104,107,108].

VTE

No thromboembolic events were reported in csDMARDs [92], phase 2 bDMARDs studies [93,94] or tsDMARDs [98,99]. In phase 3 bDMARDs studies, 3 thromboembolic events were reported by week 12 with otilimab 150 mg QW (<1%) compared with no events with tofacitinib (0%), sarilumab (0%), or placebo (0%) [95,96]. One thromboembolic event was reported with

otilimab 90 mg QW (<1%) compared with tofacitinib (0 [0%]) by week 52, while no thromboembolic events were reported by week 24 in otilimab vs sarilumab groups (0% for both groups) [96].

MACE

CV events were reported in the sole trial on csDMARDs in both methotrexate arms (5.8% and 4% for oral and subcutaneous MTX, respectively) [92].

CV events were reported in both the placebo (<1%-1%), and otilimab groups (range 0-<1); by week 12 in csDMARD inadequate response (IR)/bDMARD IR population, 3 events (<1%) were reported with otilimab 90 mg and 150 mg, while no events were reported with placebo (0%), tofacitinib (0%), or sarilumab (0%) [95,96]. By week 52, in the MTX IR population, 4 events (<1%) were reported in otilimab 90 mg compared with 1 event (<1%) in placebo and no events in the tofacitinib group (0%) [95]. Additionally, by week 52, in csDMARD IR/bDMARD IR populations, 5 events were reported (<1%) in the otilimab 90 mg group and 3 events (<1%) in the otilimab 150 mg group compared with no events in the tofacitinib and sarilumab groups (0 [0%]) [95]. No CV events were reported in the otilimab or sarilumab groups (0%) compared with 1 event in the placebo group (1 [1%]) by week 24 [96]. One CV event (1.3%) was reported with low-dose BMS-986142 compared with placebo (0%) [98]. One event was also reported with high-dose ivarmacitinib 8 mg (0.5%) compared with placebo (0%) by week 24, and 1 event in the switch group from placebo to ivarmacitinib 4 mg from week 24 to 52 (0.6) compared with no events in the groups that continued on ivarmacitinib at 4 and 8 mg (0%) [99].

No CV events were reported with peresolimab (0%) and dazodalibep (0%) vs placebo (0%) [93,94]. Low rates of CV events were noted with biosimilar tocilizumab studies only (0.3%-1%) [100,101,108].

Mortality

One death (1 [5.9%]) was reported in the MIDORA trial, in the dazodalibep group only [94]. In CONTRAST trials 1 and 2, 34 deaths were reported in total in all groups, including 4 in the placebo group; none of these deaths was considered treatment-related, while 8 of them were COVID-19-related [95]. Deaths were equally common in CONTRAST trial 3 and the AVERT trial between intervention groups (otilimab and sarilumab [<1%], abatacept [<1%]) [96,97]. No deaths were reported in csDMARDs and tsDMARDs studies [92,98,99].

Gastrointestinal perforations

GIPs were only reported in bDMARD studies comparing otilimab with tofacitinib (1 [<1%] by week 12 with tofacitinib, and 1 [<1%] by week 52 with otilimab) [95]. GIPs were more common with higher dose ivarmacitinib compared with placebo (6 [3.2%] vs 3 [1.6%], respectively) and compared with ivarmacitinib lower dose (6 [3.2%] vs 2 [1.1%]) by week 24, while they increased numerically by week 52 (range 1.8%-48%) [99]. Two GIP events were reported in 2 tocilizumab biosimilar studies [101,107].

DMARD monitoring SLR

No RCTs or observational studies addressed the clinical utility of long-term routine laboratory toxicity surveillance.

DISCUSSION

Since the publication of the last SLR on the safety of DMARDs in 2022 [6], particularly following concerns raised by the ORAL surveillance study [109] regarding the safety of tsDMARDs, there has been a notable increase in studies evaluating safety outcomes, with the majority of these observational studies focusing primarily on malignancy, thromboembolic and CV events. However, no new RCTs evaluating the comparative safety of JAKis vs other bDMARDs with regard to MACE, VTE, or malignancy risks have been published since the last update.

Consistent with previous findings [6], the risk of serious infections remains elevated with bDMARDs compared with csDMARDs, although the risk appeared comparable across different classes of bDMARDs and tsDMARDs. Notably, higher rates of serious infections were reported with JAKi compared with IL-6i in 1 study only [32]. In line with the previous SLR, HZ was more frequently observed with JAKis than with bDMARDs [6]. In comparison, in a high-risk-of-bias study, the risk of serious infections and HZ infection was found to be lower with JAKis compared with TNFis (HR 0.79 [0.417-1.502] and HR 0.20 [0.08-0.52], respectively). These results were incongruous with the broader literature [28]. Future efforts to estimate and compare HZ risk across different JAKis are likely to be constrained by several methodological challenges. Chief among these are the widespread uptake of the recombinant zoster vaccine, synchronous and overlapping prescribing trajectories of JAKis, and vaccine availability in different health care settings and systems [110]. Together, these factors may complicate attempts to disentangle drug-specific risk profiles among JAKis and may limit the precision of comparative safety assessments. Additionally, we did not find new data published on HZ risk with filgotinib, but rather with JAKi overall or with baricitinib and tofacitinib specifically. This could reflect current clinical practice, as the latter 2 JAKis were the first JAKis approved and studied extensively [111]. The 2016 SLR suggested an increased risk of TB associated with TNFis [112]; however, while the most recent 2022 SLR found no supporting evidence for this association, recent data from a low-risk-of-bias study indicated that infliximab and adalimumab are still associated with an increased TB risk [20], while no significant differences were observed between JAKis and bDMARDs. It is important to note that most cases in the study occurred before comprehensive TB screening, including interferon γ (IFN- γ) –based immunological testing, was routinely recommended before initiating b/tsDMARDs and information on prior treatment for latent TB was not available [20]. With these in mind, it remains important to monitor for TB reactivation risk, but the overall risk remains low if mandatory screening is in place before b/tsDMARD initiation.

Another important finding that arose from the ORAL surveillance study was the higher incidence of malignancy in patients receiving tofacitinib [109]; in the open-label study of ORAL surveillance, patients with a history of CV disease and CV risk factors were at increased risk of malignancies excluding NMSC, and at increased risk of NMSC alone with tofacitinib vs TNFis [113]. Similar analysis has since been undertaken in cohort studies to assess malignancy risk in patients on different DMARDs. The risk of malignancy has been found to be lower with csDMARDs than with bDMARDs, although the risk appears consistent across different bDMARDs. Conflicting evidence regarding abatacept was noted in the previous SLR [6]; however, these findings were not corroborated in the current review. In recent years, the association between DMARD use and skin malignancies has garnered increased attention [114]. An

elevated risk of both melanoma and NMSCs has been reported among patients with RA treated with DMARDs, and specifically TNFi [112]. Nevertheless, no significant differences in NMSC have been observed between various groups of bDMARDs, while the risk of NMSC alone was found to be increased with JAKis compared with TNFis [43,45]. However, it must be noted that both studies were assessed as having at least a moderate RoB, and wide CIs associated with a moderately increased aHR. Substantial differences in the characteristics between patients initiating a JAKi and those initiating a TNFi were found, as well as a short follow-up period, which is why these results have to be interpreted with caution [43,45]. Assessing malignancy risk with DMARDs, particularly JAKi, requires long-term follow-up; thus, a well-controlled prospective study excluding confounders like smoking, cancer risk factors, and geographic variability would be valuable in understanding this risk. Another challenge lies in appropriately accounting for varying time exposure and risk within observational studies, as both treatment patterns and outcome hazard evolve over time, and observational studies may not capture these changes [115].

The previous systematic review placed significant emphasis on CV and thromboembolic events, particularly in response to safety concerns raised by the ORAL surveillance study [6,109]. Since the ORAL surveillance study was published, several observational studies have sought to address these concerns and generate evidence based on clinical practice using observational studies and real-world clinical data, included in this SLR; however, no additional RCTs have been conducted to date addressing safety outcomes as a primary endpoint.

Evidence from studies with low RoB indicates no increased risk of MACE with JAKis compared with bDMARDs. In contrast, 2 high RoB studies reported an increased risk of MACE with JAKis relative to TNFis and IL-6i [25,46]. Inflammation has been recognised as a central process in the development of CV disease, and the excess inflammation present in RA has been hypothesised to be a determinant of the observed higher risk in RA [116–119]; patients receiving JAKis might have longer disease duration at treatment start, accompanied with higher disease burden, which potentially increases overall CV risk in this patient population. These results may offer cautious reassurance to clinicians and patients engaged in shared decision-making regarding the initiation and continuation of JAKi therapy. JAKi use in patients with risk factors remains a shared decision between physician and informed patient, given the modest risk increase observed in individuals with a CV history, while further investigations outside robust RCTs may offer limited additional insights, as current guidance appears to be reasonably well established. Additionally, no CV benefit was observed with TNFis compared with csDMARDs, corroborating earlier safety findings [6,54,55].

One important safety signal identified in the 2022 SLR was the increased risk of VTE associated with high doses of tofacitinib, primarily based on data from the ORAL surveillance study [109]; these data led the regulatory authorities to recommend caution when JAKis are prescribed, especially in patients with risk factors for VTE [120]. Subsequently, real-world studies have further explored that risk since the last SLR. The risk of VTE was increased in all but 1 high RoB study with JAKis compared to bDMARDs (where the risk was found to be comparable), specifically for pulmonary embolism as reported in 1 low RoB study [69]. Interestingly, the risk was found to be increased with upadacitinib and baricitinib compared with tofacitinib [65]. The cause of this observed risk remains unclear, and future research should aim to understand the underlying mechanism/

aetiology of the above-mentioned association, though further exploration beyond well-designed RCTs may yield limited incremental insight given the current strength of available evidence.

A number of observational studies, included in this SLR, investigated age-specific associations, particularly in relation to serious infections, MACE, and malignancy, guided by the ORAL surveillance study results in relation to increased risk of the above-mentioned outcomes in patients over the age of 75 years. Across studies evaluating age-stratified risks, JAKis generally did not show increased rates of MACE, malignancy, or serious infection compared with TNFis or other bDMARDs, aside from results derived from a low RoB study notifying of a higher serious infection risk with tofacitinib at ≥ 70 years, while this risk is shown to be increased with increasing age [22]. It is, though, important to continue investigating age-specific associations to guide biologic selection in clinical practice.

A critical methodological consideration in synthesising safety data is the consistent definition of clinical outcomes across studies; for example, in the current SLR, many of the included studies failed to clearly define CV events, thereby limiting the interpretability and comparability of findings related to specific safety outcomes. This lack of standardisation underscores the need for harmonised outcome definitions in future research to enable more robust meta-analyses and evidence synthesis. Additionally, many studies in this SLR based their safety analyses on claims databases, which poses notable methodological risks given the limited clinical granularity, potential coding inaccuracies and incomplete ascertainment of adverse events inherent to such data [121].

A related issue concerns the distinction between RoB classification and the precision of study findings. It has to be highlighted that, while assessing RoB and study quality in this SLR was based on the Hayden tool, the correct interpretation of measures of uncertainty (eg, CIs) in context of the respective effect sizes is a crucial component for interpreting evidence of observational studies correctly. Large real-world data studies, in particular, may yield highly precise estimates, often reflected in narrow CIs around effect sizes, while still being vulnerable to substantial systematic bias. Such estimates (eg, odds ratios close to 1 with very tight CIs) can be misinterpreted as clinically meaningful solely on the basis of statistical significance. However, high precision does not compensate for underlying bias, and small effect sizes should be interpreted cautiously unless baseline risks are sufficiently high to confer meaningful impact.

Beyond effect sizes and their precision, the metric of association should also be considered when estimating effect magnitudes and evaluating the relevance of any study's finding in regard to its clinical relevance. In contrast to RCTs, predefined assumptions of effect sizes (or their differences) are critical, which is largely absent in observational studies. The primary aim of our safety systematic review is to identify potential safety signals rather than to adjudicate their magnitude. The identification of a signal, does not itself, determine its clinical significance. This distinction underscores the necessary step from detecting a potential association to evaluating its practical relevance for patient care.

On the other hand, the additional SLR on DMARD monitoring revealed a notable evidence gap, as no RCTs or observational studies addressed the clinical utility of long-term routine laboratory toxicity surveillance. This paucity of data has contributed to the widespread adoption of frequent monitoring practices in clinical settings derived primarily from safety data from phase 3 RCTs that then translate into label texts. Current recommendations advocate for intensive

Table 6
MACE and VTE, comparison between different DMARDs (observational studies)

Study ID	Registry	Intervention	Control	aHR/OR	RoB
MACE					
D'Andrea et al [63], 2022	Claims database	HCQ	MTX	1.07 (0.97-1.18)	High
Yoshida et al [54], 2023	CorEvitas RA ^a	TNFi & MTX NDE	MTX	0.77 (0.60-1.00)	Low
		TNFi & MTX CDE		0.91 (0.72-1.17)	
Huang et al [59], 2024	Claims database	TNFi + MTX	MTX	1.30 (0.578-2.952)	High
		Non-TNFi + MTX		0.84 (0.157-4.547)	
		csDMARDs + MTX		0.85 (0.375-1.97)	
		TNFi		1.37 (0.627-3.032)	
		Non-TNFi		0.92 (0.299-2.859)	
		csDMARDs		1.55 (0.766-3.144)	
Frisell et al [24], 2023	ARTIS	ADA	ETA	1.12 (0.89-1.41)	Low
		IFX		1.00 (0.72-1.39)	
		CERT		1.31 (0.99-1.73)	
		GOL		1.14 (0.82-1.58)	
		ABA		1.14 (0.87-1.48)	
		RTX		1.27 (1.04-1.55)	
		TCZ		0.97 (0.70-1.33)	
		SAR		N/A	
		BARI		0.83 (0.49-1.42)	
		TOFA		0.78 (0.31-1.99)	
Bower et al [56], 2023	SRQ	JAKi	TNFi	0.71 (0.51-0.99)	Low
		TOFA		0.78 (0.39-1.55)	
		BARI		0.70 (0.49-1.01)	
		UPA		N/A	
		Non-TNFi		0.98 (0.78-1.23)	
Mok et al [25], 2024	HK Biologic registry	JAKi	TNFi	2.09 (1.10-3.95)	Moderate
Min et al [45], 2023	Claims database	JAKi naïve	TNFi	0.69 (0.42-1.14)	High
		JAKi treated		0.84 (0.70-1.00)	
Yoshida et al [46], 2023	Rheumatology departments ^b	JAKi	IL-6i	3.03 (0.77-15.21)	High
Ma et al [57], 2024	Claims database	TOFA	ADA	0.99 et al [0.760-1.303]	High
Hoisnard et al [60], 2023	SNDS	JAKi	ADA	1.0 (0.7-1.5)	High
Song et al [62], 2023	Claims database	JAKi	bDMARDs	1.28 (0.53-3.11)	High
Meissner et al [55], 2023	RABBIT	JAKi	TNFi	0.89 (0.52-1.52)	Low
		Other bDMARDs		0.76 (0.45-1.27)	
		csDMARDs		1.36 (0.85-2.19)	
VTE					
Molander et al [69], 2023	SRQ	RTX	TNFi	0.94 (0.74-1.20)	Low
		IL-6i		1.25 (0.94-1.67)	
		ABA		0.89 (0.66-1.21)	
		JAKi		1.73 (1.24-2.42)	
Goldman et al [65], 2024	Claim database	JAKi	bDMARDs	2.11 (1.97-2.25)	Low
Mok et al [25], 2024	HK Biologic registry	JAKi	TNFi	3.90 (0.20-78.1)	Moderate
Min et al [45], 2023	Claim database	JAKi naïve	TNFi	1.46 (0.70-3.03)	High
		JAKi treated		1.35 (0.92-1.97)	
Fang et al [38], 2022	TNHIRD/TDR	JAKi	TNFi	0.65 (0.25-1.70)	High
Sakai et al [58], 2024	Claim database	JAKi	TNFi	2.08 (1.47-2.93)	High
Hoisnard et al [60], 2023	SNDS	JAKi	ADA	1.1 (0.7-1.6)	High

ABA, abatacept; ADA, adalimumab; aHR, adjusted hazard ratio; ARTIS, Anti-Rheumatic Therapy in Sweden; bDMARD, biological disease-modifying antirheumatic drug; BARI, baricitinib; c, control; CDE, Controlled direct effect analyses; CERT, certolizumab; csDMARD, conventional disease-modifying antirheumatic drug; GOL, Golimumab; HCQ, hydroxychloroquine; HK, Hong Kong biologic registry; i, intervention; IL-6i, interleukin-6 inhibitor; IFX, infliximab; JAKi, Janus kinase inhibitor; MACE, major adverse cardiovascular events; MTX, methotrexate; NDE, Natural direct effect analyses; OR, odds ratio; RABBIT, Rheumatoid Arthritis Observation of Biologic Therapy registry; RoB, risk of bias; RTX, rituximab; SAR, sarilumab; SNDS, Système National des Données de Santé, France; SRQ, Swedish Rheumatology Quality Register; TCZ, tocilizumab; TNFi, tumour necrosis factor inhibitor; TOFA, tofacitinib; UPA, upadacitinib; VTE, venous thromboembolism.

Nagasaki University Hospital, Sasebo Chuo Hospital, or Ureshino Medical Center.

^a Formerly known as CORRONA.

^b Department of Rheumatology of Fukushima Medical University Hospital, Japanese Red Cross Fukushima Hospital, and Ohta Nishinouchi General Hospital Foundation.

laboratory monitoring during the initial 6 months of DMARD therapy, a period associated with elevated risk of adverse reactions, followed by a transition to less frequent assessments throughout the remainder of treatment, which may extend for decades or even a lifetime [10,11,122]. To date, no studies have systematically compared alternative blood or laboratory monitoring strategies in terms of their influence on clinically meaningful outcomes. Emerging data studies suggest that intensive long-term blood monitoring offers limited

benefit, indicating that this practice may warrant reconsideration in future guideline updates [8]. Future research should investigate both patient and physician perspectives on monitoring practices, including the possibility that routine testing, despite limited medical necessity, may foster medication adherence by providing reassurance regarding drug safety. Additionally, further inquiry is warranted to identify potential legal implications and practical challenges that could impede the implementation of revised clinical guidelines.

Randomisation is the cornerstone of RCTs, rendering them the gold standard for evaluating differences in outcomes across treatment groups [123]. By randomly allocating participants, RCTs achieve a balanced distribution of both measured and unmeasured confounding variables, thereby minimising bias and enhancing internal validity [123]. Typically, the primary endpoint in RCTs pertains to efficacy, while formal comparisons of safety outcomes are often constrained by limited statistical power [123]. Moreover, RCTs frequently exclude individuals at elevated risk for adverse events and are generally of insufficient duration to detect outcomes with extended latency periods, such as malignancies [123].

Consequently, observational studies, particularly those involving unselected patient populations monitored over prolonged periods, serve as the principal source of evidence in this SLR for assessing drug safety. While these studies offer valuable insights into real-world safety profiles, they are inherently limited by the absence of randomisation [124]. This introduces the potential for confounding by indication, wherein baseline differences in risk factors between treatment groups may influence observed outcomes [125]. Safety findings from the ORAL Surveillance RCT trial demonstrated a clear dose-dependent signal [109]. However, observational studies may be limited in their ability to investigate dose-response associations with JAKis, given the use of fixed dosing regimens in certain agents (eg, upadacitinib) and restricted dose variability in some regions globally. This constraint hampers the validation of dose-response relationships in real-world settings and could represent a limitation for comparative safety evaluations. Another limitation is the geographic heterogeneity due to differences in healthcare systems, screening and prescribing practices, and vaccination schedules observed when comparing studies, which could limit generalisability [126].

Despite these limitations, observational studies remain indispensable for evaluating safety, especially for outcomes that are rare or manifest over long time horizons [127]. Ideally, a hybrid approach that integrates the methodological rigour of experimental designs with the external validity of observational research would provide the most comprehensive assessment of treatment safety. In the current systematic review, no RCTs were found to have safety events as primary endpoints.

Conclusion

Overall, the conclusions drawn from this SLR are consistent with those reported in the 2022 review, which is reassuring and reinforces the evolving consensus regarding the safety profiles of cs-, b-, and tsDMARDs in the treatment of RA. The accumulating body of evidence supports the notion that these agents can be employed with safety in clinical practice. Importantly, this SLR, when considered alongside the corresponding efficacy review, provides a comprehensive evidence base that strengthens the EULAR recommendations for the management of RA with DMARDs. Moreover, it offers a substantive reaffirmation of the theoretical framework and guiding principles upon which these clinical guidelines are founded.

Competing interests

FL received speaker fees from Alfasigma. VK received speaker's or consultancy fees from Eli Lilly, AbbVie, and AstraZeneca. JSS received honoraria for consultancies and/or speaking engagements from Abbott, AbbVie, Anaptyx, Ananda,

AstraZeneca, Celltrion, Chugai, Immunovant, Lilly, R-Pharma, Samsung, Sandoz, Shattuck labs, and UCB. CJE received honoraria, advisory boards, speaker's bureau, and research support from AbbVie, Alphasigma, AstraZeneca, BMS, Biogen, Boehringer Ingelheim, Celgene, Celltrion, Fresenius, Gilead, GSK, Janssen, Lilly, Pfizer, MSD, Novartis, Roche, Samsung, Sanofi, and UCB. DA received honoraria from Advanz, Janssen/Johnsen & Johnsen, Sanofi, Mitsubishi Tanabe, UCB, and AstraZeneca, and grants to his institution from Lilly and Johnson & Johnson. DvdH received consulting fees from AbbVie, Alfasigma, ArgenX, BMS, Eli Lilly, Grey-Wolf Therapeutics, Janssen, Novartis, Pfizer, Takeda, and UCB Pharma, and is associate editor of *Annals Rheumatic Diseases*, editorial board member of *Journal of Rheumatology*, and director of Imaging Rheumatology. KLV received honoraria for consulting from AbbVie, AstraZeneca, BMS, Eli Lilly, Galapagos, Gilead, GlaxoSmithKline (GSK), Moderna, Novartis, Roche, Sanofi, Union Chimique Belge (UCB), Zurabio, Takeda, Ouros, and Otsuka. TT received speaker's or consultancy fees from AbbVie GK, Chugai, Eli Lilly Japan, Eisai, Gilead Sciences Inc, Pfizer Japan Inc, Mitsubishi Tanabe, and Taisho Pharma. RC received consultancy and speaker fees from AbbVie, Alfasigma, AstraZeneca, GSK, Eli Lilly, Galapagos, Janssen, Novartis, Pfizer, and UCB. PV received research grants from Alfasigma and Pfizer; speaker or consultancy fees from Alfasigma, AbbVie, Galapagos, Eli Lilly, Cytrill, and Boehringer Ingelheim; and support for attending meetings from Fresenius Kabi and AbbVie. JaP declares no conflicts. KH received speaker's fees from AbbVie, grants from Pfizer and BMS, and serves as associate Editor for *Annals of the Rheumatic Diseases*. SdS declares no conflicts. JeP received speaker fees from Pfizer and Eli Lilly. JWS declares no conflicts. RBML received honoraria for lectures and payments for advisory boards from AbbVie, Jansen Pharma, Galapagos, Novartis, and UCB, and is director of Joint Imaging BV. AK received consulting fees from AbbVie, Lilly, Gilead, Janssen, and UCB; lecture fees from Lilly, Galapagos, Janssen, MSD, Novartis, and Pfizer; and congress support from Boehringer Ingelheim, Lilly, and UCB.

CRediT authorship contribution statement

Faidra Laskou: Writing – review & editing, Writing – original draft, Methodology, Formal analysis, Data curation, Conceptualization. **Victoria Konzett:** Writing – review & editing, Conceptualization. **Josef S Smolen:** Writing – review & editing, Supervision, Conceptualization. **Christopher J Edwards:** Writing – review & editing, Supervision, Conceptualization. **Daniel Aletaha:** Writing – review & editing, Conceptualization. **Désirée van der Heijde:** Writing – review & editing, Conceptualization. **Kevin L Winthrop:** Writing – review & editing, Conceptualization. **Tsutomu Takeuchi:** Writing – review & editing, Conceptualization. **Roberto Caporali:** Writing – review & editing, Conceptualization. **Patrick Verschuere:** Writing – review & editing, Conceptualization. **Janet E Pope:** Writing – review & editing, Conceptualization. **Kimme Hyrich:** Writing – review & editing, Conceptualization. **Savia de Souza:** Writing – review & editing, Conceptualization. **Tanja A Stamm:** Writing – review & editing, Conceptualization. **Jette Primdahl:** Writing – review & editing, Conceptualization. **Jan W Schoones:** Data curation. **Robert BM Landewé:** Writing – review & editing, Supervision, Methodology, Conceptualization. **Andreas Kerschbaumer:** Writing – review & editing, Supervision, Methodology, Conceptualization.

Contributors

All authors contributed and finally approved the current manuscript. FL is the guarantor of the study. The guarantor accepts full responsibility for the finished work and the conduct of the study, had access to the data, and controlled the decision to publish the manuscript together with the methodologists (AK and RBML) and the convenors of the taskforce (JSS and CJE).

Funding

European Alliance of Associations for Rheumatology (EULAR).

Patient consent for publication

Because of the nature of the analyses, this is not applicable for the current work. All patients gave informed consent to participation in the studies included and publication of study results.

Ethics approval

Because of the nature of the analyses performed here, this is not applicable to the current project. All studies included in this work were approved by the respective ethics committees.

Provenance and peer review

Not commissioned; externally peer reviewed.

Data availability statement

Not applicable, as no datasets were generated for this study. All data relevant to the study are available in public and are included in the article or uploaded as online supplemental information.

Patient and public involvement

One patient research partner (SdS) participated in the steering group and task force, and 2 additional patients contributed to the task force that developed the guidelines. Patients and/or the public were otherwise not involved in the design, or conduct, or reporting, or dissemination plans of this research. Details are provided in the Methods section.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.ard.2026.02.024](https://doi.org/10.1016/j.ard.2026.02.024).

Orcid

Faidra Laskou: <http://orcid.org/0000-0002-8481-6343>

REFERENCES

- [1] Mehta B, Pedro S, Ozen G, Kalil A, Wolfe F, Mikuls T, et al. Serious infection risk in rheumatoid arthritis compared with non-inflammatory rheumatic and musculoskeletal diseases: a US national cohort study. *RMD Open* 2019;5(1):e000935.
- [2] Finckh A, Gilbert B, Hodkinson B, Bae SC, Thomas R, Deane KD, et al. Global epidemiology of rheumatoid arthritis. *Nat Rev Rheumatol* 2022 Oct;18(10):591–602.
- [3] Aletaha D, Smolen JS. Diagnosis and management of rheumatoid arthritis: a review. *JAMA* 2018 Oct 2;320(13):1360–72.
- [4] Smolen JS, van der Heijde D, Machold KP, Aletaha D, Landewé R. Proposal for a new nomenclature of disease-modifying antirheumatic drugs. *Ann Rheum Dis* 2014 Jan;73(1):3–5.
- [5] Agency. EM. EMA confirms measures to minimise risk of serious side effects with Janus kinase inhibitors for chronic inflammatory disorders [cited 2026 Jan 14]. Available from: <https://www.ema.europa.eu/en/medicines/human/referrals/janus-kinase-inhibitors-jaki>
- [6] Sepriano A, Kerschbaumer A, Bergstra SA, Smolen JS, van der Heijde D, Caporali R, et al. Safety of synthetic and biological DMARDs: a systematic literature review informing the 2022 update of the EULAR recommendations for the management of rheumatoid arthritis. *Ann Rheum Dis* 2023;82(1):107–18.
- [7] Kerschbaumer A, Sepriano A, Bergstra SA, Smolen JS, Van Der Heijde D, Caporali R, et al. Efficacy of synthetic and biological DMARDs: a systematic literature review informing the 2022 update of the EULAR recommendations for the management of rheumatoid arthritis. *Ann Rheum Dis* 2022;82:95–106.
- [8] Ulijn E, Den Broeder AA, Boers N, Gotthardt M, Bouman CAM, Landewé R, et al. Extra-articular findings with FDG-PET/CT in rheumatoid arthritis patients: more harm than benefit. *Rheumatol Adv Pract* 2022;6(1):rkac014.
- [9] Fraser SD, Lin SX, Stammers M, Culliford D, Ibrahim K, Barrett R, et al. Persistently normal blood tests in patients taking methotrexate for RA or azathioprine for IBD: a retrospective cohort study. *Br J Gen Pract* 2022;72(720):e528–37.
- [10] Holroyd CR, Seth R, Bukhari M, Malaviya A, Holmes C, Curtis E, et al. The British Society for Rheumatology biologic DMARD safety guidelines in inflammatory arthritis. *Rheumatology (Oxford)* 2019 Feb 1;58(2):e3–42.
- [11] Fraenkel L, Bathon JM, England BR, St Clair EW, Arayssi T, Carandang K, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Rheumatol* 2021 Jul;73(7):1108–23.
- [12] Kerschbaumer A, Konzett VL, Edwards CJ, Aletaha D, van der Heijde D, Winthrop KL, et al. Efficacy of synthetic and biological DMARDs and glucocorticoids: a systematic literature review informing the 2025 update of the EULAR recommendations for the management of rheumatoid arthritis. in press - accepted for publication ARDS 18/03/2026.
- [13] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Rev* 2021 Mar 29;10(1):89.
- [14] EULAR. Standard operating procedures for task forces. [cited 2025 Nov 24]. Available from: <https://www.eular.org/sops-for-eular-post-tf-implementation-projects>
- [15] Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, editors. *Cochrane handbook for systematic reviews of interventions*. 2nd ed. Chichester, UK: John Wiley & Sons; 2019.
- [16] Sackett DL, Richardson WS, Rosenberg W, Haynes RB. *Evidence-based medicine: how to practice and teach EBM*. Churchill Livingstone; BMJ; 2005.
- [17] Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019 Aug 28;366:l4898.
- [18] Hayden JA, Côté P, Bombardier C. Evaluation of the quality of prognosis studies in systematic reviews. *Ann Intern Med* 2006;144:427–37.
- [19] Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011 Oct 18;343:d5928.
- [20] Lauper K, Kearsley-Fleet L, Galloway JB, Watson KD, BSRBR-RA Contributors Group, Hyrich KL, et al. Evaluation of serious infections, including Mycobacterium tuberculosis, during treatment with biologic disease-modifying anti-rheumatic drugs: does line of therapy matter? *Rheumatology (Oxford)* 2024;63(7):1957–64.
- [21] Dieperink SS, Mehnert F, Nørgaard M, Oestergaard LB, Benfield T, Petersen A, et al. Antirheumatic treatment, disease activity and risk of Staphylococcus aureus bacteraemia in rheumatoid arthritis: a nationwide nested case-control study. *RMD Open* 2022 Dec;8(2):e002636.
- [22] Riek M, Scherer A, Möller B, Ciurea A, von Mühlhagen I, Gabay C, et al. Serious infection risk of tofacitinib compared to biologics in patients with rheumatoid arthritis treated in routine clinical care. *Sci Rep* 2023 Oct 18;13(1):17776.
- [23] Dieperink SS, Nørgaard M, Mehnert F, Oestergaard LB, Benfield T, Torp-Pedersen C, et al. Rheumatoid arthritis and risk of osteoarticular infection and death following Staphylococcus aureus bacteraemia: a nationwide cohort study. *Rheumatology (Oxford)* 2024;63(11):2989–96.
- [24] Frisell T, Bower H, Morin M, Baecklund E, Di Giuseppe D, Delcoigne B, et al. Safety of biological and targeted synthetic disease-modifying antirheumatic drugs for rheumatoid arthritis as used in clinical practice: results from the ARTIS programme. *Ann Rheum Dis* 2023;82(5):601–10.

- [25] Mok CC, So H, Yim CW, To CH, Lao WN, Wong SPY, et al. Safety of the JAK and TNF inhibitors in rheumatoid arthritis: real world data from the Hong Kong Biologics Registry. *Rheumatology (Oxford)* 2024;63(2):358–65.
- [26] Choi SR, Shin A, Ha YJ, Lee YJ, Lee EB, Kang EH. Comparative risk of infections between JAK inhibitors versus TNF inhibitors among patients with rheumatoid arthritis: a cohort study. *Arthritis Res Ther* 2023 Jul 26;25(1):129.
- [27] Harigai M, Fujii T, Sakai R, Igarashi A, Shoji A, Yamaguchi H, et al. Risk of hospitalized infections in older elderly patients with rheumatoid arthritis treated with tocilizumab or other biological/targeted synthetic disease-modifying antirheumatic drugs: evaluation of data from a Japanese claims database. *Mod Rheumatol* 2024;34(2):287–96.
- [28] Uchida T, Iwamoto N, Fukui S, Morimoto S, Aramaki T, Shomura F, et al. Comparison of risks of cancer, infection, and MACEs associated with JAK inhibitor and TNF inhibitor treatment: a multicentre cohort study. *Rheumatology (Oxford)* 2023;62(10):3358–65.
- [29] Shin A, Lee JH, Ha YJ, Lee YJ, Lee EB, Kang EH. Infectious risk of add-on leflunomide or tacrolimus versus TNF inhibitors among patients with rheumatoid arthritis receiving background methotrexate: a population-based cohort study. *Semin Arthritis Rheum* 2022;55:152019.
- [30] Ota R, Hirata A, Hata T, Nishihara M, Neo M, Katsumata T. Incidence of serious infections in the working-age Japanese adult population with rheumatoid arthritis treated with tumor necrosis factor- α inhibitors and interleukin-6 inhibitors: a nationwide retrospective cohort study. *Pharmacotherapy* 2024;44(7):570–80.
- [31] Bellan M, Scotti L, Ferrante D, Calzaducce E, Manfredi GF, Sainaghi PP, et al. Risk of severe infection among rheumatoid arthritis patients on biological DMARDs: a population-based cohort study. *J Clin Med* 2022 May 24;11:2955.
- [32] Yoshida S, Miyata M, Suzuki E, Kanno T, Sumichika Y, Saito K, et al. Incidence rates of infections in rheumatoid arthritis patients treated with Janus kinase or interleukin-6 inhibitors: results of a retrospective, multicenter cohort study. *J Clin Med* 2024 May 1;13:3000.
- [33] Park HJ, Choi B, Song YK, Oh YJ, Lee EB, Kim IW, et al. Association of tumor necrosis factor inhibitors with the risk of nontuberculous mycobacterial infection in patients with rheumatoid arthritis: a nationwide cohort study. *J Clin Med* 2023 Nov 9;12:6998.
- [34] Opdam MAA, Broeder ND, van den Bemt BJF, Mulder K, van de Wiel KM, van Ballegooijen H, et al. Higher infection risk for JAK inhibitors tofacitinib and baricitinib compared to subcutaneous biological DMARDs. *Clin Rheumatol* 2024;43(6):2133–8.
- [35] Jeong S, Choi S, Park SM, Kim J, Ghang B, Lee EY. Incident and recurrent herpes zoster for first-line bDMARD and tsDMARD users in seropositive rheumatoid arthritis patients: a nationwide cohort study. *Arthritis Res Ther* 2022 Jul 28;24(1):180.
- [36] So MW, Kim AR, Lee SG. Drug persistence and incidence of active tuberculosis of tumor necrosis factor alpha inhibitors versus tocilizumab as the first-line biological treatment in patients with rheumatoid arthritis: a nationwide population-based retrospective cohort analysis. *Rheumatol Ther* 2024 Aug;11(4):881–95.
- [37] Song YK, Lee J, Jo J, Kwon JW. Comparison of active tuberculosis occurrence associated with Janus kinase inhibitors and biological DMARDs in rheumatoid arthritis. *RMD Open* 2024 Apr 12;10(2):e003946.
- [38] Fang YF, Liu JR, Chang SH, Kuo CF, See LC. Comparative safety of Janus kinase inhibitors and tumor necrosis factor inhibitors in patients undergoing treatment for rheumatoid arthritis. *Int J Rheum Dis* 2022;25:1254–62.
- [39] Tsai JJ, Liu LT, Chen CH, Chen LJ, Wang SI, Wei JC. COVID-19 outcomes in patients with rheumatoid arthritis with biologic or targeted synthetic DMARDs. *RMD Open* 2023 Jul;9(3):e003038.
- [40] Singh N, Madhira V, Hu C, Olex AL, Bergquist T, Fitzgerald KC, et al. Rituximab is associated with worse COVID-19 outcomes in patients with rheumatoid arthritis: a retrospective, nationally sampled cohort study from the U.S. National COVID Cohort Collaborative (N3C). *Semin Arthritis Rheum* 2023;58:152149.
- [41] Fang YF, Chang SH, Kuo CF, See LC. Safety outcomes of tocilizumab and tofacitinib treatment for rheumatoid arthritis: target trial emulation. *Int J Rheum Dis* 2024;27(11):e15406.
- [42] Westermann R, Cordtz RL, Duch K, Mellemkjaer L, Hetland ML, Burden AM, et al. Cancer risk in patients with rheumatoid arthritis treated with Janus kinase inhibitors: a nationwide Danish register-based cohort study. *Rheumatology (Oxford)* 2024;63(1):93–102.
- [43] Huss V, Bower H, Hellgren K, Frisell T, Askling J. Behalf of the ARTIS group. Cancer risks with JAKi and biological disease-modifying antirheumatic drugs in patients with rheumatoid arthritis or psoriatic arthritis: a national real-world cohort study. *Ann Rheum Dis* 2023;82:911–9.
- [44] Seror R, Lafourcade A, De Rycke Y, Pinto S, Castaneda J, Fautrel B, et al. Risk of malignancy in rheumatoid arthritis patients initiating biologics: an historical propensity score matched cohort study within the French nationwide healthcare database. *RMD Open* 2022 Jun;8(2):e002139.
- [45] Min HK, Kim H, Jeong HJ, Kim SH, Kim HR, Lee SH, et al. Risk of cancer, cardiovascular disease, thromboembolism, and mortality in patients with rheumatoid arthritis receiving Janus kinase inhibitors: a real-world retrospective observational study using Korean health insurance data. *Epidemiol Health* 2023;45:e2023045.
- [46] Yoshida S, Miyata M, Suzuki E, Kanno T, Sumichika Y, Saito K, et al. Safety of JAK and IL-6 inhibitors in patients with rheumatoid arthritis: a multicenter cohort study. *Front Immunol* 2023;14:1267749.
- [47] Huss V, Bower H, Wadström H, Frisell T, Askling J, group ARTIS. Short- and longer-term cancer risks with biologic and targeted synthetic disease-modifying antirheumatic drugs as used against rheumatoid arthritis in clinical practice. *Rheumatology (Oxford)* 2022;61(5):1810–8.
- [48] Lee H, Chen SK, Gautam N, Vine SM, He M, Desai RJ, et al. Risk of malignant melanoma and non-melanoma skin cancer in rheumatoid arthritis patients initiating methotrexate versus hydroxychloroquine: a cohort study. *Clin Exp Rheumatol* 2023;41:110–7.
- [49] Song YJ, Cho SK, You SH, Kim JY, Kim H, Jung SY, et al. Association between malignancy risk and Janus kinase inhibitors versus tumour necrosis factor inhibitors in Korean patients with rheumatoid arthritis: a nationwide population-based study. *RMD Open* 2022 Dec;8(2):e002614.
- [50] Simon TA, Suissa S, Boers M, Hochberg MC, Skovron ML, Askling J, et al. Malignancy outcomes in patients with rheumatoid arthritis treated with abatacept and other disease-modifying antirheumatic drugs: results from a 10-year international post-marketing study. *Semin Arthritis Rheum* 2024;64:152240.
- [51] Choi B, Park HJ, Song YK, Oh YJ, Kim IW, Oh JM. The risk of newly diagnosed cancer in patients with rheumatoid arthritis by TNF inhibitor use: a nationwide cohort study. *Arthritis Res Ther* 2022 Aug 9;24(1):191.
- [52] Sendaydiego X, Gold LS, Dubreuil M, Andrews JS, Reid P, Liew DFL, et al. Use of biologic or targeted synthetic disease-modifying antirheumatic drugs and cancer risk. *JAMA Netw Open* 2024;7(11):e2446336.
- [53] Molina-Collada J, Alonso F, Otero L, Bohórquez C, Díaz Torné C, Pérez García C, et al. Cancer risk with biologic and targeted synthetic DMARDs in patients with rheumatic diseases and previous malignancies: results from the BIOBADASER register. *Semin Arthritis Rheum* 2024;64:152341.
- [54] Yoshida K, Harrold LR, Middaugh N, Guan H, Stryker S, Karis E, et al. Examining the potential direct cardiovascular benefit of tumor-necrosis factor inhibitor in rheumatoid arthritis: natural and controlled direct effect analyses. *Pharmacoepidemiol Drug Saf* 2023;32(4):407–15.
- [55] Meissner Y, Schäfer M, Albrecht K, Kekow J, Zinke S, Tony HP, et al. Risk of major adverse cardiovascular events in patients with rheumatoid arthritis treated with conventional synthetic, biologic and targeted synthetic disease-modifying antirheumatic drugs: observational data from the German RABBIT register. *RMD Open* 2023 Oct;9(4):e003489.
- [56] Bower H, Frisell T, di Giuseppe D, Delcoigne B, Lindström U, Turesson C, et al. Are JAKis more effective among elderly patients with RA, smokers and those with higher cardiovascular risk? A comparative effectiveness study of b/tsDMARDs in Sweden. *RMD Open* 2023 Dec 26;9(4):e003648.
- [57] Ma XN, Shi MF, Wang SI, Feng W, Chen SL, Zhong XQ, et al. Risk of dyslipidemia and major adverse cardiac events with tofacitinib versus adalimumab in rheumatoid arthritis: a real-world cohort study from 7580 patients. *Front Pharmacol* 2024;15:1370661.
- [58] Sakai R, Tanaka E, Inoue E, Harigai M. Increased risk of cardiovascular events under the treatments with Janus kinase inhibitors versus biological disease-modifying antirheumatic drugs in patients with rheumatoid arthritis: a retrospective longitudinal population-based study using the Japanese health insurance database. *RMD Open* 2024 Jun 17;10(2):e003885.
- [59] Huang Y, Agarwal SK, Chatterjee S, Chen H, Johnson ML, Aparasu RR. Risk of incident cardiovascular events with disease-modifying anti-rheumatic drugs among adults with rheumatoid arthritis: a nested case-control study. *Clin Rheumatol* 2024;43(1):103–16.
- [60] Hoisnard L, Pina Vegas L, Dray-Spira R, Weill A, Zureik M, Sbidian E. Risk of major adverse cardiovascular and venous thromboembolism events in patients with rheumatoid arthritis exposed to JAK inhibitors versus adalimumab: a nationwide cohort study. *Ann Rheum Dis* 2023;82(2):182–8.
- [61] Popa CD, Opdam MAA, den Broeder N, van Ballegooijen H, Mulder K, van de Wiel KM, et al. Therapy with JAK inhibitors or bDMARDs and the risk of cardiovascular events in the Dutch rheumatoid arthritis population. *Rheumatology (Oxford)* 2024;63:2142–6.
- [62] Song YK, Lee G, Hwang J, Kim JW, Kwon JW. Cardiovascular risk of Janus kinase inhibitors compared with biologic disease-modifying antirheumatic drugs in patients with rheumatoid arthritis without underlying cardiovascular diseases: a nationwide cohort study. *Front Pharmacol* 2023;14:1165711.

- [63] D'Andrea E, Desai RJ, He M, Glynn RJ, Lee H, Weinblatt ME, et al. Cardiovascular risks of hydroxychloroquine vs methotrexate in patients with rheumatoid arthritis. *J Am Coll Cardiol* 2022;80:36–46.
- [64] Tong X, Shen CY, Jeon HL, Li Y, Shin JY, Chan SC, et al. Cardiovascular risk in rheumatoid arthritis patients treated with targeted synthetic and biological disease-modifying antirheumatic drugs: a multi-centre cohort study. *J Intern Med* 2023 Sep;294(3):314–25.
- [65] Goldman A, Galper BL, Druyan A, Grossman C, Sharif K, Shechtman L, et al. Adverse cardiovascular events in rheumatoid arthritis patients treated with JAK inhibitors: an analysis of postmarketing spontaneous safety reports. *Semin Arthritis Rheum* 2024;67:152461.
- [66] Delcoigne B, Ljung L, Provan SA, Grintborg B, Hetland ML, Grøn KL, et al. Short-term, intermediate-term and long-term risks of acute coronary syndrome in cohorts of patients with RA starting biologic DMARDs: results from four Nordic countries. *Ann Rheum Dis* 2022;81(6):789–97.
- [67] Quiñones ME, Joseph JK, Dowell S, Moore HJ, Karasik PE, Fonarow GC, et al. Hydroxychloroquine and risk of long QT syndrome in rheumatoid arthritis: a veterans cohort study with nineteen-year follow-up. *Arthritis Care Res (Hoboken)* 2023;75(7):1571–9.
- [68] Bower H, Frisell T, di Giuseppe D, Delcoigne B, Askling J. Comparative cardiovascular safety with Janus kinase inhibitors and biological disease-modifying antirheumatic drugs as used in clinical practice: an observational cohort study from Sweden in patients with rheumatoid arthritis. *RMD Open* 2023 Nov 23;9(4):e003630.
- [69] Molander V, Bower H, Frisell T, Delcoigne B, Di Giuseppe D, Askling J, et al. Venous thromboembolism with JAK inhibitors and other immune-modulatory drugs: a Swedish comparative safety study among patients with rheumatoid arthritis. *Ann Rheum Dis* 2023;82(2):189–97.
- [70] Goldman A, Raschi E, Druyan A, Sharif K, Lahat A, Ben-Zvi I, et al. Gastrointestinal perforations associated with JAK inhibitors: a disproportionality analysis of the FDA adverse event reporting system. *United European Gastroenterol J* 2025;13:566–75.
- [71] Hoisnard L, Meyer A, Dray-Spira R, Weill A, Zureik M, Sbidian E. Risk of gastrointestinal perforation in patients with rheumatic diseases exposed to Janus kinase inhibitors versus adalimumab: a nationwide cohort study. *Arthritis Rheumatol* 2024;76(9):1364–76.
- [72] Delcoigne B, Kopp TI, Arkema EV, Hellgren K, Provan SA, Relas H, et al. Exposure to specific tumour necrosis factor inhibitors and risk of demyelinating and inflammatory neuropathy in cohorts of patients with inflammatory arthritis: a collaborative observational study across five Nordic rheumatology registers. *RMD Open* 2023 Feb;9(1):e002924.
- [73] Mielnik P, Sexton J, Fagerli KM, Bakland G, Hu Y, Kristianslund EK, et al. Discontinuation rate of sulfasalazine, leflunomide and methotrexate due to adverse events in a real-life setting (NOR-DMARD). *Rheumatol Adv Pract* 2023;7:rkad053.
- [74] Bredemeier M, Duarte AL, Pinheiro MM, Kahlow BS, Macieira JC, Ranza R, et al. The effect of antimalarials on the safety and persistence of treatment with biologic agents or Janus kinase inhibitors in rheumatoid arthritis. *Rheumatology (Oxford)* 2024;63(2):456–65.
- [75] Jinno S, Onishi A, Hattori S, Dubreuil M, Ueda Y, Nishimura K, et al. Comparison of retention of biologics in Japanese patients with elderly-onset rheumatoid arthritis-the ANSWER cohort study. *Rheumatology (Oxford)* 2025;64:509–16.
- [76] Lauper K, Iudici M, Mongin D, Bergstra SA, Choquette D, Codreanu C, et al. Effectiveness of TNF-inhibitors, abatacept, IL6-inhibitors and JAK-inhibitors in 31 846 patients with rheumatoid arthritis in 19 registers from the 'JAK-pot' collaboration. *Ann Rheum Dis* 2022;81(10):1358–66.
- [77] Ebina K, Etani Y, Maeda Y, Okita Y, Hirao M, Yamamoto W, et al. Drug retention of biologics and Janus kinase inhibitors in patients with rheumatoid arthritis: the ANSWER cohort study. *RMD Open* 2023 Aug;9(3):e003160.
- [78] Gilbert BTP, Mongin D, Aymon R, Lauper K, Laedermann C, Perrier C, et al. Comparative effectiveness of baricitinib and alternative biological DMARDs in a Swiss cohort study of patients with RA. *BMJ Open* 2024 Mar 12;14(3):e072300.
- [79] Nakayama Y, Onishi A, Yamamoto W, Yoshikawa A, Shiba H, Yoshida N, et al. Safety of Janus kinase inhibitors compared to biological DMARDs in patients with rheumatoid arthritis and renal impairment: the ANSWER cohort study. *Clin Exp Med* 2024;24(1):97.
- [80] Nakayama Y, Watanabe R, Yamamoto W, Ebina K, Hirano T, Kotani T, et al. IL-6 inhibitors and JAK inhibitors as favourable treatment options for patients with anaemia and rheumatoid arthritis: ANSWER cohort study. *Rheumatology (Oxford)* 2024;63(2):349–57.
- [81] Onishi A, Yamada H, Yamamoto W, Watanabe R, Hara R, Katayama M, et al. Comparative effectiveness of biological disease-modifying antirheumatic drugs and Janus kinase inhibitor monotherapy in rheumatoid arthritis. *Rheumatology* 2024;63:3065–73.
- [82] Movahedi M, Choquette D, Coupal L, Cesta A, Li X, Keystone EC, et al. Discontinuation of tofacitinib and TNF inhibitors in patients with rheumatoid arthritis: analysis of pooled data from two registries in Canada. *BMJ Open* 2023 Mar 6;13(3):e063198.
- [83] Terabe K, Takahashi N, Asai S, Hirano Y, Kanayama Y, Yabe Y, et al. Effectiveness of tacrolimus concomitant with biological disease-modifying antirheumatic drugs in patients with rheumatoid arthritis. *Mod Rheumatol* 2023;33(2):292–301.
- [84] Velthuis K, Jessurun NT, Nguyen TDM, Scholl J, Jansen JRG, van Lint JA, et al. First-time adverse drug reactions, survival analysis, and the share of adverse drug reactions in treatment discontinuation in real-world rheumatoid arthritis patients: a comparison of first-time treatment with adalimumab and etanercept. *Expert Opin Drug Saf* 2023;22(6):485–92.
- [85] Su YJ, Chen HM, Chan TM, Cheng TT, Yu SF, Chen JF, et al. Disease-modifying anti-rheumatic drugs associated with different diabetes risks in patients with rheumatoid arthritis. *RMD Open* 2023 Jul;9(3):e003045.
- [86] Chen YW, Chen HH, Huang WN, Chen JP, Chen YH, Chen YM. Potential alleviation of bone mineral density loss with Janus kinase inhibitors in rheumatoid arthritis. *Clin Rheumatol* 2024;43(1):117–28.
- [87] Abtahi S, Cordtz R, Dreyer L, Driessen JHM, Boonen A, Burden AM. Biological disease-modifying antirheumatic drugs and osteoporotic fracture risk in patients with rheumatoid arthritis: a Danish cohort study. *Am J Med* 2022;135(7):879–888.e3.
- [88] Su YJ, Lin CY, Hsu CY. Analyzing the risk of osteoporosis and fracture in rheumatoid arthritis patients who have been treated with various biologics. *Int J Rheum Dis* 2024;27(2):e15055.
- [89] Secher AEP, Granath F, Grintborg B, Rom A, Hetland ML, Hellgren K. Risk of pre-eclampsia and impact of disease activity and antirheumatic treatment in women with rheumatoid arthritis, axial spondylarthritis and psoriatic arthritis: a collaborative matched cohort study from Sweden and Denmark. *RMD Open* 2022 Nov;8(2):e002445.
- [90] Delteil A, Lambert C, Pereira B, Couderc M, Malochet-Guinamand S, Pickering ME, et al. Glucocorticoid trajectories over 2 years in patients with rheumatoid arthritis in a real-life setting. *RMD Open* 2023 Oct;9(4):e003366.
- [91] Coburn BW, Baker JF, Hsu JY, Wu Q, Xie F, Curtis JR, et al. Association of cardiovascular outcomes with low-dose glucocorticoid prescription in patients with rheumatoid arthritis. *Arthritis Rheumatol* 2024;76(11):1585–93.
- [92] Tanaka Y, Okuda K, Takeuchi Y, Katayama K, Haji Y, Yamanishi Y, et al. Efficacy and tolerability of subcutaneously administered methotrexate including dose escalation in long-term treatment of rheumatoid arthritis in a Japanese population. *Mod Rheumatol* 2023;33(4):680–9.
- [93] Tuttle J, Drescher E, Simón-Campos JA, Emery P, Greenwald M, Kivitz A, et al. A phase 2 trial of peresolimab for adults with rheumatoid arthritis. *N Engl J Med* 2023;388(20):1853–62.
- [94] Kivitz A, Wang L, Alevizos I, Gunsior M, Falloon J, Illei G, et al. The MID-ORA trial: a phase II, randomised, double-blind, placebo-controlled, mechanistic insight and dosage optimisation study of the efficacy and safety of nazodalibep in patients with rheumatoid arthritis. *RMD Open* 2023;9(3):e003317.
- [95] Fleischmann RM, van der Heijde D, Strand V, Atsumi T, McInnes IB, Takeuchi T, et al. Anti-GM-CSF otilimab versus tofacitinib or placebo in patients with active rheumatoid arthritis and an inadequate response to conventional or biologic DMARDs: two phase 3 randomised trials (contrAST 1 and contrAST 2). *Ann Rheum Dis* 2023;82(12):1516–26.
- [96] Taylor PC, Weinblatt ME, McInnes IB, Atsumi T, Strand V, Takeuchi T, et al. Anti-GM-CSF otilimab versus sarilumab or placebo in patients with rheumatoid arthritis and inadequate response to targeted therapies: a phase III randomised trial (contrAST 3). *Ann Rheum Dis* 2023;82(12):1527–37.
- [97] Emery P, Tanaka Y, Bykerk VP, Bingham CO, Huizinga TWJ, Citera G, et al. The trajectory of clinical responses in patients with early rheumatoid arthritis who achieve sustained remission in response to abatacept: subanalysis of AVERT-2, a randomized phase IIb study. *Arthritis Res Ther* 2023;25(1):67.
- [98] Conaghan PG, Nowak M, Du S, Luo Y, Landis J, Pachai C, et al. Evaluation of BMS-986142, a reversible Bruton's tyrosine kinase inhibitor, for the treatment of rheumatoid arthritis: a phase 2, randomised, double-blind, dose-ranging, placebo-controlled, adaptive design study. *Lancet Rheumatol* 2023;5(5):e263–73.
- [99] Liu J, Jiang Y, Zhang S, Liu S, Su J, Lin C, et al. Ivarmacitinib, a selective Janus kinase 1 inhibitor, in patients with moderate-to-severe active rheumatoid arthritis and inadequate response to conventional synthetic DMARDs: results from a phase III randomised clinical trial. *Ann Rheum Dis* 2025;84:188–200.
- [100] Zubrzycka-Sienkiewicz A, Klama K, Ullmann M, Petit-Frere C, Baker P, Monnet J, et al. Comparison of the efficacy and safety of a proposed biosimilar MSB11456 with tocilizumab reference product in subjects with

- moderate-to-severe rheumatoid arthritis: results of a randomised double-blind study. *RMD Open* 2024;10(1):e003596.
- [101] Leng X, Leszczyński P, Jeka S, Liu SY, Liu H, Miakisz M, et al. Comparing tocilizumab biosimilar BAT1806/BIIB800 with reference tocilizumab in patients with moderate-to-severe rheumatoid arthritis with an inadequate response to methotrexate: a phase 3, randomised, multicentre, double-blind, active-controlled clinical trial. *Lancet Rheumatol* 2024;6(1):e40–50.
- [102] Ye H, Liu S, Xu J, Chai K, He D, Fang Y, et al. Efficacy and safety of CMAB008 compared with innovator infliximab in patients with moderate-to-severe rheumatoid arthritis receiving concomitant methotrexate: a randomized, double-blind, multi-center, phase III non-inferiority study. *Rheumatol Ther* 2023;10(3):757–73.
- [103] Furst DE, Jaworski J, Wojciechowski R, Wiland P, Dudek A, Krogulec M, et al. Efficacy and safety of switching from reference adalimumab to CT-P17 (100 mg/ml): 52-week randomized, double-blind study in rheumatoid arthritis. *Rheumatology (Oxford)* 2022;61(4):1385–95.
- [104] Liu Y, Liu S, Liu L, Gong X, Liu J, Sun L, et al. Fine comparison of the efficacy and safety between GB242 and infliximab in patients with rheumatoid arthritis: a phase III study. *Rheumatol Ther* 2022;9(1):175–89.
- [105] Emery P, Tanaka Y, Bykerk VP, Huizinga TWJ, Citera G, Bingham CO, et al. Sustained remission and outcomes with abatacept plus methotrexate following stepwise dose de-escalation in patients with early rheumatoid arthritis. *Rheumatol Ther* 2023;10(3):707–27.
- [106] Liu XL, Li L, Wang Q, Jiang FC, Zhang P, Guo F, et al. A novel humanized anti-interleukin-6 antibody HZ0408b with anti-rheumatoid arthritis therapeutic potential. *Front Immunol* 2022;12(10):816646.
- [107] Smolen JS, Trefler J, Racewicz A, Jaworski J, Zielińska A, Krogulec M, et al. Efficacy and safety of CT-P47 versus reference tocilizumab: 32-week results of a randomised, active-controlled, double-blind, phase III study in patients with rheumatoid arthritis, including 8 weeks of switching data from reference tocilizumab to CT-P47. *RMD Open* 2024;10(4):e004514.
- [108] Leng X, Leszczyński P, Jeka S, Liu S, Liu H, Miakisz M, et al. A phase 3, randomized, double-blind, active-controlled clinical trial to compare BAT1806/BIIB800, a tocilizumab biosimilar, with tocilizumab reference product in participants with moderate-to-severe rheumatoid arthritis with inadequate response to methotrexate: treatment period 2 analysis (week 24 to week 48). *Arthritis Res Ther* 2024;26(1):157.
- [109] Ytterberg SR, Bhatt DL, Mikuls TR, Koch GG, Fleischmann R, Rivas JL, et al. Cardiovascular and cancer risk with tofacitinib in rheumatoid arthritis. *N Engl J Med* 2022;386(4):316–26.
- [110] Galloway J, Raine T, Rivett L, Roberts J, Dews SA, Choy EH. Herpes zoster and Janus kinase inhibition in rheumatology and gastroenterology patients: managing risk and vaccination. *Clin Exp Rheumatol* 2022;40:1432–41.
- [111] Szekanecz Z, Buch MH, Charles-Schoeman C, Galloway J, Karpouzas GA, Kristensen LE, et al. Efficacy and safety of JAK inhibitors in rheumatoid arthritis: update for the practising clinician. *Nat Rev Rheumatol* 2024;20(2):101–15.
- [112] Ramiro S, Sepriano A, Chatzidionysiou K, Nam JL, Smolen JS, van der Heijde D, et al. Safety of synthetic and biological DMARDs: a systematic literature review informing the 2016 update of the EULAR recommendations for management of rheumatoid arthritis. *Ann Rheum Dis* 2017 Jun;76(6):1101–36.
- [113] Curtis JR, Yamaoka K, Chen YH, Bhatt DL, Gunay LM, Sugiyama N, et al. Malignancy risk with tofacitinib versus TNF inhibitors in rheumatoid arthritis: results from the open-label, randomised controlled ORAL Surveillance trial. *Ann Rheum Dis* 2023;82(3):331–43.
- [114] Shasti-Nazem B, Chandra S, Strouse J, Singh N. Treatment of persons with rheumatoid arthritis with a history of cancer. *Curr Rheumatol Rep* 2025 Aug 11;27(1):34.
- [115] Rop M, Maposa I, Young T, Machezano R. Causal estimation of time-varying treatments in observational studies: a scoping review of methods, applications, and missing data practices. *BMC Med Res Methodol* 2025 Aug 27;25(1):202.
- [116] Ruparel N, Chai JT, Fisher EA, Choudhury RP. Inflammatory processes in cardiovascular disease: a route to targeted therapies. *Nat Rev Cardiol* 2017 Mar;14(3):133–44.
- [117] Liao KP, Solomon DH. Traditional cardiovascular risk factors, inflammation and cardiovascular risk in rheumatoid arthritis. *Rheumatology (Oxford)* 2013 Jan;52(1):45–52.
- [118] Chodara AM, Wattiaux A, Bartels CM. Managing cardiovascular disease risk in rheumatoid arthritis: clinical updates and three strategic approaches. *Curr Rheumatol Rep* 2017 Apr;19(4):16.
- [119] Drosos GC, Vedder D, Houben E, Boekel L, Atzeni F, Badreh S, et al. EULAR recommendations for cardiovascular risk management in rheumatic and musculoskeletal diseases, including systemic lupus erythematosus and antiphospholipid syndrome. *Ann Rheum Dis* 2022 Jun;81(6):768–79.
- [120] European Medicines Agency. EMA confirms measures to minimise risk of serious side effects with Janus kinase inhibitors for chronic inflammatory disorders. [cited 2022 Nov 11]. Available from: <https://go.nature.com/48tdDis>
- [121] Tyree PT, Lind BK, Lafferty WE. Challenges of using medical insurance claims data for utilization analysis. *Am J Med Qual* 2006;21(4):269–75.
- [122] Mercer L, Abhishek A, Kavirayani A, Ahmed A, Davidson A, Foulkes A, et al. The prescription and monitoring of conventional synthetic disease-modifying anti-rheumatic drugs: British Society for Rheumatology guideline scope. *Rheumatol Adv Pract* 2024;8(3):rkae077.
- [123] Braga LH, Farrokhyar F, Dönmez Mİ, Nelson CP, Haid B, Herbst K, et al. Randomized controlled trials - the what, when, how and why. *J Pediatr Urol* 2025 Apr;21(2):397–404.
- [124] Landewé R, van der Heijde D. Primer: challenges in randomized and observational studies. *Nat Clin Pract Rheumatol* 2007;3:661–6.
- [125] Sender R, Stürmer T. Core concepts in pharmacoepidemiology: confounding by indication and the role of active comparators. *Pharmacoepidemiol Drug Saf* 2022 Mar;31(3):261–9.
- [126] Yusuf S, Wittes J. Interpreting geographic variations in results of randomized, controlled trials. *N Engl J Med* 2016 Dec 8;375(23):2263–71.
- [127] Hannan EL. Randomized clinical trials and observational studies: guidelines for assessing respective strengths and limitations. *JACC Cardiovasc Interv* 2008 Jun;1(3):211–7.