



Universiteit
Leiden
The Netherlands

Development and implementation of VR-simulated experiences to train communication skills

Hoek, K.

Citation

Hoek, K. (2026, September 4). *Development and implementation of VR-simulated experiences to train communication skills*. Retrieved from <https://hdl.handle.net/1887/4309192>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4309192>

Note: To cite this publication please use the final published version (if applicable).

4C

**DEVELOPMENT OF A PATIENT-
EMBODIED EXPERIENCE,
HOW AND WHY?**

ARTICLE INFO

Authors:


Krista Hoek
Monique van Velzen
Elise Sarton

Keywords:

VR
Therapeutic language
and communication,
Ethnographic phenomenology,
Healthcare education,
Patient-embodied experience,
Proteus-effect

DOI: 10.5220/0012755800003693

Serious Games for Education



Volume 1

ISSN 2184-5026

ISBN 978-989-758-697-2

May 2024

Pages 758-764

ABSTRACT

Introduction

The rise of immersive technologies, particularly virtual reality (VR), has significantly impacted medical education. In patient-embodied VR, through VR headsets, learners can embody a patient's perspective, offering a secure and immersive learning encounter. This communication outlines a framework for crafting patient-embodied experiences, drawing from our VR endeavours aimed at enhancing therapeutic communication skills in medical education.

Methods

Our framework includes a development process with consideration of user experience, technical implementation, content creation and validation. Central to content creation is the collaborative construction of a patient journey, involving the involved parties via storyboards and scripts distinguishing direct and indirect actions.

Results

For our patient-embodied experience, the cooperative development of the patient journey, script and storyboard included an initial version created by the main researcher after study of landmark articles on therapeutic communication and fieldwork. Validation was achieved through two group sessions with healthcare providers who consented to participate.

Conclusions and practice implications

The findings and insights presented can contribute to the growing knowledge in the field of educational VR development. They demonstrate the feasibility and potential of leveraging immersive technologies to create engaging and impactful virtual experiences. Hitherto, further validation may evaluate how they influence beliefs and attitudes of healthcare providers towards therapeutic communication.

INTRODUCTION

Medical education has been incorporating elements of experiential learning for several decades, and the specific use of immersive technologies has gained more prominence in recent years. In particular, learning with virtual reality (VR), or immersive virtual environments (IVEs) aligns with the Constructivist Theory, which suggests that learners actively construct their knowledge and understanding of the world by building upon their existing mental frameworks. In an immersive learning environment, learners are given the opportunity to explore, discover, and make meaning from their experiences (1)

In virtual reality (VR), immersive learning incorporates various sensory modalities to enhance *multisensory engagement* enabling embodiment. Embodiment is the possibility in VR to visually substitute a person's real body by a life-sized virtual one, seen from the person's own first-person perspective. In other words, when we place a VR headset on, our virtual bodies momentarily substitute our real bodies, a phenomenon known as the *Proteus effect* (2). Multisensory and motor systems interconnect during cognitive processing. During VR, embodiment can increase users' engagement and provide emotional fidelity invoking a sense of *presence* creating realistic emotional and neurocognitive responses (3, 4).

When using VR as an educational tool, learners have the ability to experience a full 360-degree view, providing them with unrestricted access and freedom to explore their surroundings. This contrasts to traditional video experiences, where the audience's perspective is limited to a fixed viewpoint as depicted in **Figure 1**. The distinctive aspect of VR lies in its participatory nature. In VR, writers, directors, and producers, referred to as builders, do not have the ability to dictate how learners engage with the story. Instead, they can only invite participation, allowing learners to choose where to direct their attention and which aspects of the story to focus on (5). This stands in contrast to conventional storytelling, where a storyteller transmits the narrative to listeners. Traditional videos typically provide viewers a fixed perspective. The screen acts as a window into the world of the video, and viewers can only observe the events and scenes from the specific viewpoint chosen by the director as shown in **Figure 1**.




Figure 1. Red outlines show the extent of what a camera can film in 90 vs 360 degrees of freedom.

Left: 90 degrees of freedom of traditional film

Right: 360 degrees of freedom of VR

In healthcare, a growing emphasis on patient-centered care, and consumerism in medicine has exemplified the importance of effective physician-patient communication (6) and the use of IVEs as an educational tool in this context. We will refer to patient-embodied VR as the specific VR application where a learner who puts on the VR headset virtually becomes the patient. Given that this immersive transformation has the capacity to affect self-perception, attitudes, convictions, and conduct in both implicit and explicit ways (2, 4, 7), our hypothesis posits that it holds substantial promise as an exceedingly effective and innovative pedagogical approach in education on communication skills.

An example of the use of patient-embodied VR can be found in our work published recently (8, 9). We developed two patient-embodied experiences to create a possibility for healthcare providers to feel what it is like to become a patient. In these VR experiences, the learner experiences the sequential stages of a patient undergoing elective general anesthesia and surgery, with nuanced shifts in language and interactions. We will describe how we have developed these experiences, and what important lessons we have learned that can be used by researchers aiming to develop-embodied VR experiences.

METHODS

This prospective exploratory research was carried out at the Leiden University Medical Center (LUMC, Leiden, the Netherlands), a large tertiary academic teaching hospital. The protocol was approved by the Institutional Science Committee and obtained a waiver from the Institutional Review Board (NWMO-LUMC).

Informed consent was obtained prior to inclusion, participation was voluntary and privacy rights were in alignment with the Declaration of Helsinki and GDPR guidelines. Data was collected and recorded between February 2019 to December 2020. Participants received no financial compensation.

Prototype iterations were systematically tested in simulated settings, followed by expert debriefings and FMEA. Each cycle informed design enhancements, ensuring usability and training effectiveness. Adjustments continued iteratively until the prototype was deemed optimized. User evaluations revealed that medical interns and clinical educators believed in the potential and usefulness of the VR simulation as a learning tool for CRM.

Encountered challenges included onboarding in the virtual environment requiring participants to familiarize themselves with effective navigation within the virtual space.

Another challenge involved translating clinical needs into programming possibilities outlined by the developers. This continuous need for translation persisted between user requirements and technical capabilities. Integration of simulation with UCD and FMEA facilitated design and redesign of the prototype.

The anaesthesiology department provided full support, and a collaborative partnership was made with another university medical center to facilitate further validation and broader implementation for other institutions enhancing scalability.

Participants and procedures

Healthcare providers working in the OR were invited to participate in the development of the patient-embodied VR experiences. They were recruited between January 2019 and May 2019. The procedure encompassed content creation, content validation, and filming and editing of the VR experiences as shown in **Figure 2**.




Figure 2. structure of the study participation

Content creation

To create the storyboard and script of the patient journey, a theoretical framework needed to consider the study of landmark articles in the concerned domain. In our case, Landmark patient-centered studies were used to include important aspects of lived patient experiences in preoperative settings (10-18).

Secondly, fieldwork of the researcher (KH) included informal interviews with healthcare providers and patients, along with observations to gather data was used to create an initial script for the patient journeys as shown in **Figure 2**.

Content validation

Group sessions are an adapted tool to validate the storyboard and script (19). In our study, a first group session was held to discuss and adapt the scripts.

Group input such as photographs, focused and selective observation notes, reflective notes and commentaries were used to write a second version of the patient journeys. A second group session implied a final validation where the scripts were discussed and adapted until consensus was reached; that is until the final script was consented by all group members (20) as shown in **Figure 2**.

The specific linguistic features were further validated by an independent hypnotherapist based on validated and relevant research (16, 18, 21-25).

Filming and editing

Asking healthcare providers to play their own role has several advantages as its economical, and the actors may identify themselves very easily with their own professional role. We asked the developers to play their own professional role, e.g. a surgeon would play the surgeon. An anaesthetic nurse volunteered to play the patient, as she had been a surgical patient several times before.

RESULTS

Case study description

Nine healthcare providers participated in the development of the patient-embodied VR experiences. Six were OR nurses, one was a surgeon and two were anaesthesiologist-hypnotherapists. One hypnotherapist who was not affiliated with our hospital did not participate in the filming of the experiences. The majority was female (82%). Demographics are shown in Table 1.

Table 1. Demographic characteristics of the VR development team and study participants

Clinical role	
Anaesthesiologist-hypnotherapist	2
CRNA (Nurse anaesthetist)	2
Preoperative holding area nurse	3
Operating room nurse	1
Surgeon	1
Years of practice	
<1year	1
1-5years	3
5-10years	1
>10years	4
Sex	
Male/Female	2/7

Development of the patient-embodied VR experiences

Between February 2019 and December 2019 two scripts and storyboards were created: the positive and negative patient-embodied VR experience.

Guidelines of several digital platforms were used (5, 26). Filming was performed using a *GoPro*® camera, editing was performed using *Movavi*® enabling the experiences to be available on most common VR headsets. Also, an online Youtube version was uploaded making it possible to view the experience online.

The final durations of the experiences were 12 minutes and 19 seconds for the negative experience, and 10 minutes and 36 seconds for the positive experience.

The story-world needed to be as consistent with reality as possible, using objects, personas, and actions to stimulate the sense of immersive *presence*.

Storyboard and script

We developed a storyboard with the main scenes as shown in **Figure 3** including all direct interactions with the patient.




Figure 3. Storyboard patient-embodied VR experience

We developed a script with ‘shadow actions’ visible if the learner would look around. An overview of the main differences in setting, direct interactions and shadow actions can be found in **Table 2**.

Table 2. Overview of main difference in setting, interactions and shadow actions

	Negative experience	Positive experience
Setting		
Preoperative holding area	No (distraction) activities for other patients No preventive measures for delirium prevention	Other patients use a tablet or read a magazine Curtains opened, visible clock, noises of monitoring minimized
Operation room	Radio on Alarms and sounds of continuous monitoring on. Surgical equipment partially displayed	Noises minimized No visible surgical equipment

Table 2. Overview of main difference in setting, interactions and shadow actions (*continued*)

	Negative experience	Positive experience
Direct interactions		
Contact with preoperative holding area	-Patient is ignored when she requests to go to the toilet -Nurse does not ask any personal questions while she places the IV cannula	-Nurse sits next to the patient -Nurse uses hypnotic distraction methods during the placement of the IV cannula -Nurse responds to the concerns of the patient.
Contact with anaesthesio-logist	Contact with anaesthesiologist with little rapport.	Anaesthesiologist asks personal questions before the patient is taken into the OR
Shadow interactions		
Preoperative holding area	OR nurse passes as the patient is brought to the preoperative holding area, he tells his colleague there is a bleeding in the OR, and he needs to hurry	OR nurse passes as the patient is brought to the preoperative holding area. He holds something in his arms, but is discrete.

Setting

In the positive experience, another patient present is able to read a magazine as shown in **Figure 4**. This is a distraction method that may increase comfort while the patients await their surgery (27).



Figure 4. Positive experience: another patient reads a magazine

Direct interactions

There are changes in attitude of the healthcare providers with or without a direct therapeutic relationship. One example is the communication style of the anaesthesiologist just before the start of the inductive phase of anaesthesia (patient is induced into a state of

unconsciousness and analgesia (pain relief) before the start of the surgical procedure) as shown in **Figure 5**.

In the negative video, the anaesthesiologist administers the medication, however, the communication shows limited effort to establish rapport (8, 17, 28). Furthermore, the potential side effects of the drugs are explicitly mentioned, thereby increasing the likelihood of the patient experiencing these adverse effects (18, 25).

In the positive experience, the anaesthesiologist uses hypnotic linguistic techniques to induce relaxation (24), enhance focus and promote a smooth start of general anaesthesia. The anaesthesiologist sits beside the patient (16).



Figure 5. Anaesthesiologist sits next to the patient, uses suggestive language to induce relaxation

Shadow actions

An example of a suggestion used in the final script was having an OR-nurse pass by to bring blood for the transfusion of another patient to enhance authenticity of the dynamics of the operating theatre. Subtle changes between the experiences are shown in **Figure 6**.

One portrays an OR-nurse that politely smiles to the patients without showing any concerns. The other portrays an OR-nurse that expresses a sense of urgency. He seems bothered, and tells the preoperative nurse he has to deliver blood. The preoperative nurse seems worried.



Figure 6. Left: the operating room (OR) nurse remains silent and smiles politely without showing his concerns. Right: the OR nurse conveys a sense of urgency to bring blood to another patient, and the preoperative nurse appears annoyed and worried

DISCUSSION AND CONCLUSION

Discussion

This study describes the development of two patient-embodied VR experiences. The specific linguistic features were validated by an independent hypnotherapist and were based on validated and relevant research (16, 18, 21-25).

Internal validity of the VR experiences was confirmed given that healthcare providers participated in the development of these experiences themselves. Between the two experiences, small changes in setting, interactions and shadow actions would implicitly suggest a more or less welcoming experience. Indeed, the development included the addition of these several elements enhancing authenticity of the VR experiences.

This type of immersive storytelling with VR offers learners a possibility to create their own story in 360°; as they were able to look around freely in VR. Additionally, healthcare providers played their own roles during the filming of the scenes.

Limitations

One of the limitations of the VR experiences lies in the limited interaction within the VR-environment. The learner had the ability to observe their surroundings; however, due to the recorded nature of the experiences, learners were not able to freely move around as they would have in a virtual environment utilizing computer-generated 3D video imagery.

We selected this setting based on the belief that a patient typically finds themselves confined to a hospital bed with limited mobility or free choices. We contend that utilizing a genuine operating theatre and real actors enhances fidelity, thereby intensifying the sense of immersion.

Innovation

The findings and insights presented in this study can contribute to the growing knowledge in the field of educational VR development. They demonstrate the feasibility and potential of leveraging immersive technologies to create engaging, authentic and impactful virtual experiences (8)

CONCLUSION

By using immersive VR technology, one can create an interactive and engaging virtual environment that allows learners to experience a simulated reality, that is the experience of being a patient.

This development process involves careful consideration of user experience, technical implementation, content creation and validation. In our opinion, key elements of the content creation should be based on the cooperative development of a patient journey, with participation of the involved parties using a storyboard and script that distinguishes between direct actions and indirect (shadow) actions.

Future directions and study validation

The findings of this study can contribute to further research and healthcare education programs aimed to use experiential learning with patient-embodied VR. We conclude with a call for further research to fully unlock the potential and drawbacks of patient-embodied VR as an educational tool and its usefulness in medical training. Also, further research may examine how patient-embodied VR might affect patient reported outcomes like preoperative anxiety.

Initial validation of our patient-embodied experience consisted of a qualitative study analyzing the lived experience of anaesthesiologists (8). Further validation may include evaluation of the effects of the implementation of a training in therapeutic communication with an integrated patient-embodied VR experience.

ACKNOWLEDGEMENTS

We thank all healthcare providers participating in the development of the VR experiences.

COMPETING INTERESTS

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

REFERENCES

1. Whitman N. A review of constructivism: understanding and using a relatively new theory. *Fam Med*. 1993;25(8):517-21.
2. Fox J, Bailenson J, Tricase L. The embodiment of sexualized virtual selves: The Proteus effect and experiences of self-objectification via avatars. *Computers in Human Behavior*. 2013;29:930-8.
3. Bem D. Self-perception theory. New York: Academic Press; 1972.
4. Navarro J, Peña J, Cebolla A, Baños R. Can Avatar Appearance Influence Physical Activity? User-Avatar Similarity and Proteus Effects on Cardiac Frequency and Step Counts. *Health Commun*. 2022;37(2):222-9.
5. O'Sullivan B, Alam F, Matava C. Creating Low-Cost 360-Degree Virtual Reality Videos for Hospitals: A Technical Paper on the Dos and Don'ts. *J Med Internet Res*. 2018;20(7):e239.
6. Santana MJ, Manalili K, Jolley RJ, Zelinsky S, Quan H, Lu M. How to practice person-centred care: A conceptual framework. *Health Expect*. 2018;21(2):429-40.
7. Bian Y, Zhou C, Tian Y, Wang P, Gao F, editors. *The Proteus Effect: Influence of Avatar Appearance on Social Interaction in Virtual Environments*. HCI; 2015.
8. Hoek K, van Velzen M, Sarton E. Patient-embodied virtual reality as a learning tool for therapeutic communication skills among anaesthesiologists: A phenomenological study. *Patient Education and Counseling*. 2023;114:107789.
9. Hoek K, van Velzen M, Sarton E. Response to correspondance on patient-embodied virtual reality as a learning tool for therapeutic communication skills among anaesthesiologists: A phenomenological study. *Patient Educ Couns*. 2023;117:107980.
10. Smith AF, Mishra K. Interaction between anaesthetists, their patients, and the anaesthesia team. *Br J Anaesth*. 2010;105(1):60-8.
11. Maranets I, Kain ZN. Preoperative anxiety and intraoperative anesthetic requirements. *Anesth Analg*. 1999;89(6):1346-51.
12. Ben-Amitay G, Kosov I, Reiss A, Toren P, Yoran-Hegesh R, Kotler M, et al. Is elective surgery traumatic for children and their parents? *J Paediatr Child Health*. 2006;42(10):618-24.
13. Kain ZN, Caldwell-Andrews AA, Maranets I, McClain B, Gaal D, Mayes LC, et al. Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. *Anesth Analg*. 2004;99(6):1648-54.
14. Kain ZN, Mayes LC, Caldwell-Andrews AA, Karas DE, McClain BC. Preoperative anxiety, post-operative pain, and behavioral recovery in young children undergoing surgery. *Pediatrics*. 2006;118(2):651-8.
15. Derksen F, Bensing J, Lagro-Janssen A. Effectiveness of empathy in general practice: a systematic review. *Br J Gen Pract*. 2013;63(606):e76-84.
16. Swayden KJ, Anderson KK, Connelly LM, Moran JS, McMahon JK, Arnold PM. Effect of sitting vs. standing on perception of provider time at bedside: a pilot study. *Patient Educ Couns*. 2012;86(2):166-71.
17. Butt MF. Approaches to building rapport with patients. *Clinical Medicine*. 2021;21(6):e662-e3.
18. Lang EV, Hatsiopoulou O, Koch T, Berbaum K, Lutgendorf S, Kettenmann E, et al. Can words hurt? Patient-provider interactions during invasive procedures. *Pain*. 2005;114(1-2):303-9.
19. Stewart D, Shamdasani P, Rook D. *Focus Groups*. Thousand Oaks, California 2007. Available from: <https://methods.sagepub.com/book/focus-groups>.
20. Briggs R, Kolfshoten G, de Vreede G-J. *Toward a Theoretical Model of Consensus Building* 2005. 12 p.

21. Lang EV, Berbaum KS. Educating interventional radiology personnel in nonpharmacologic analgesia: effect on patients' pain perception. *Acad Radiol*. 1997;4(11):753-7.
22. Lang E. Comfort Talk®: From the Waiting Room to the Treatment Suite. *Dtsch Z Zahnarztl Hypn*. 2019;25(1):22-4.
23. Watzlawick P. The language of change: Elements of therapeutic communication. New York, NY, US: W W Norton & Co; 1978. xi, 172-xi, p.
24. Boselli E, Musellec H, Bernard F, Guillou N, Hugot P, Augris-Mathieu C, et al. Effects of conversational hypnosis on relative parasympathic tone and patient comfort during axillary brachial plexus blocks for ambulatory upper limb surgery: A Quasiexperimental Pilot Study. *Int J Clin Exp Hypn*. 2018;66(2):134-46.
25. Zech N, Seemann M, Hansen E. [Nocebo effects and negative suggestion in anesthesia]. *Anaesthesist*. 2014;63(11):816-24.
26. Newton K. The Storyteller's Guide to the Virtual Reality Audience: Stanford d.school; 2016 [Available from: <https://medium.com/stanford-d-school/the-storyteller-s-guide-to-the-virtual-reality-audience-19e92da57497>].
27. Pati D, Nanda U. Influence of positive distractions on children in two clinic waiting areas. *Herd*. 2011;4(3):124-40.
28. Hall JA, Harrigan JA, Rosenthal R. Nonverbal behavior in clinician—patient interaction. *Applied and Preventive Psychology*. 1995;4(1):21-37.