



Universiteit
Leiden

The Netherlands

Exploring the clinical applicability of quantitative near-infrared fluorescence imaging with indocyanine green: towards objective and standardized assessment of tissue perfusion

Tange, F.P.

Citation

Tange, F. P. (2026, September 15). *Exploring the clinical applicability of quantitative near-infrared fluorescence imaging with indocyanine green: towards objective and standardized assessment of tissue perfusion*. Retrieved from <https://hdl.handle.net/1887/4309589>

Version: Publisher's Version


License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4309589>

Note: To cite this publication please use the final published version (if applicable).

Chapter 3





Evaluation of the Angiosome Concept Using Near-Infrared Fluorescence Imaging with Indocyanine Green

Floris P. Tange *, Bien R. Ferrari *, Pim van den Hoven Jan van Schaik, Abbey Schepers,
Catharina S.P. van Rijswijk, Rutger W. van der Meer, Hein Putter,
Alexander L. Vahrmeijer, Jaap F. Hamming, Joost R. van der Vorst

** Authors contributed equally and share first authorship*

Published in Annals of Vascular Surgery, July 2023.

Abstract

Background

The angiosome concept is defined as the anatomical territory of a source artery within all tissue layers. When applying this theory in vascular surgery, direct revascularization (DR) is preferred to achieve increased blood flow toward the targeted angiosome of the foot in patients with lower extremity arterial disease (LEAD). This study evaluates the applicability of the angiosome concept using quantified near-infrared (NIR) fluorescence imaging with indocyanine green (ICG).

Methods

This study included patients undergoing an endovascular- or surgical revascularization of the leg between January 2019 and December 2021. Preinterventional and postinterventional ICG NIR fluorescence imaging was performed. Three angiosomes on the dorsum of the foot were determined: the posterior tibial artery (hallux), the anterior tibial artery (dorsum of the foot) and the combined angiosome (second to fifth digit). The angiosomes were classified from the electronic patient records and the degree of collateralization was classified based on preprocedural computed tomography angiography and/or X-ray angiography. Fluorescence intensity was quantified in all angiosomes. A subgroup analysis based on endovascular or surgical revascularized angiosomes, and within critical limb threatening ischemia (CLTI) patients was performed.

Results

ICG NIR fluorescence measurements were obtained in 52 patients (54 limbs) including a total of 157 angiosomes (121 DR and 36 indirect revascularizations [IR]). A significant improvement of all perfusion parameters in both the directly and indirectly revascularized angiosomes was found (P-values between <0.001-0.007). Within the indirectly revascularized angiosomes, 90.6% of the scored collaterals were classified as significant. When comparing the percentual change in perfusion parameters between the directly and indirectly revascularized angiosomes, no significant difference was seen in all perfusion parameters (P-values between 0.253 and 0.881). Similar results were shown in the CLTI patients subgroup analysis, displaying a significant improvement of perfusion parameters in both the direct and indirect angiosome groups (P-values between <0.001 and 0.007), and no significant difference when comparing the percentual parameter improvement between both angiosome groups (P-values between 0.134 and 0.359). Furthermore, no significant differences were observed when comparing percentual changes of perfusion parameters in directly and indirectly revascularized angiosomes for both endovascular and surgical interventions (P-values between 0.053 and 0.899).

Conclusions

This study proves that both DR and IR of an angiosome leads to an improvement of perfusion. This suggests that interventional strategies should not only focus on creating in-line flow to the supplying angiosome. One can argue that the angiosome concept is not applicable in patients with LEAD.

Introduction

The angiosome concept was first described within the field of reconstructive surgery.¹ The authors defined an angiosome as the anatomical territory of a source artery within all tissue layers. Six angiosomes were identified in the foot, each originating from 3 source arteries and their branches: the anterior tibial artery (ATA), the posterior tibial artery (ATP), and the peroneal artery (AP). In lower extremity arterial disease (LEAD) patients with ischemic wounds, it seems rational to target the revascularization to the angiosome that supplies the site of the ulcer, known as direct revascularization (DR). Several studies, including meta-analyses, showed that angiosome-targeted revascularization increased wound healing.²⁻⁷ In contrast, a study by Rother et al. demonstrated no significant difference comparing the microvascular perfusion improvement after DR or indirect revascularization (IR) using near-infrared (NIR) fluorescence.⁸ Furthermore, literature suggests to separate surgical and endovascular interventions, as the angiosome theory might be of greater relevance in the latter.⁹ Although frequently investigated, critics consider the clinical applicability of the angiosome theory as debatable as the growth of collateral vessels in the lower extremities will interfere with the borders of the angiosomes. Both Spillerová et al. and Jongsma observed improved wound healing after IR when substantial collateralization was present. For the evaluation of the angiosome concept in vivo, reliable quantification of local tissue perfusion is essential. Different methods have been used to objectify microcirculation, including single-photon emission computed tomography (CT), light guided spectrometry and laser doppler flowmetry.^{10,11} NIR fluorescence imaging with indocyanine green (ICG) for the evaluation of tissue perfusion in patients with LEAD seems promising.¹² This technique has already been used for intraoperative guidance in multiple medical fields, including reconstructive and oncologic surgery.¹³⁻¹⁶ Furthermore, it has shown promising results in LEAD patients as quality control following revascularization.^{15,17-20} After intravenous administration of ICG, an intravascular fluorophore, the fluorescence intensity can be quantified depicting an in and outflow perfusion pattern of the imaged tissue. Combining ICG NIR fluorescence imaging and the angiosome concept can provide further insight in perfusion patterns following revascularization and guide treatment for patients with LEAD. This study compares preinterventional and postinterventional ICG NIR fluorescence measurements between DR and IR for both endovascular and surgical interventions. Additionally, the effect of collateralization on postinterventional perfusion after IR is evaluated.

Methods

This prospective cohort study was approved by the Medical Research and Ethics Committee of the Leiden University Medical Center and was performed in a single Dutch Academic Medical Center. Patients classified with Fontaine 2a to Fontaine 4 disease stage who underwent a revascularization procedure between July 2019 and December 2021 were included.


DR was defined as an achievement of inline flow to the corresponding angiosome and IR when in-line flow was not achieved. Collateral grading was performed by 2 independent interventional radiologists. The classification of the collaterals was based on the most recent preprocedural CT angiography and/or catheter angiography. The collateral grade was classified as 'substantial' when collaterals were visible for 25% of the calf length and/or estimated to be 50% of the corresponding crural artery diameter and/or when objectified as an extensive amount. Preinterventional ankle-brachial index and toe pressure measurements were performed if feasible.

ICG NIR Fluorescence Imaging

Preinterventional and postinterventional ICG NIR fluorescence images of the dorsum of the foot were obtained in accordance with a previously described protocol.²⁰ ICG NIR fluorescence imaging was performed using the Quest Spectrum Platform (Quest Medical Imaging, Middenmeer, the Netherlands). Quantification of the measured ICG NIR fluorescence intensity was performed using the Quest Research Framework (Quest Medical Imaging, Middenmeer, the Netherlands). The quantification software program generated absolute and normalized time-intensity curves out of all regions of interests (ROIs). The method of normalizing the time-intensity curves was in accordance with a formerly described protocol.²⁰ A total of 10 perfusion parameters were extracted from the time-intensity curves, displayed in Supplemental Figure 1. A postinterventional increase of inflow parameters and decrease of outflow parameters was considered as an improvement of perfusion status.

Angiosome Concept

The ATA from which the dorsalis pedis artery (ADP) originates feeds the complete dorsum of the foot. However, the lateral plantar artery partially feeds the anterior part, the AP the lateral part, and the superficial medial plantar artery the medial part. The ATP supplies the medial ankle and plantar foot consisting out of 2 angiosomes; the lateral and medial plantar angiosome. The lateral plantar artery of the ATP is the most common to feed the hallux. However, the hallux can also be part of the medial plantar angiosome or the dorsalis pedis. The lesser toes are fed by both the lateral plantar angiosome and the dorsalis pedis angiosome with a variability depending on the dominance of the lateral plantar or dorsalis pedis artery. The AP feeds the lateral calcaneal artery which supplies the plantar heel and lateral ankle.^{1,22-24} The selected ROIs were based on the angiosome theory: (1) the hallux angiosome classified as supplied by the ATP, (2) The dorsal angiosome supplied by the ADP (ATA) and (3) the lesser toes defined as the combined angiosome (ADP/ ATP) (Fig. 1).



Supplemental Figure 1. Time-intensity curve with extracted parameters. Abbreviations: a.u., arbitrary units; AUC, Area Under the Curve; Tmax, Time to Max




Figure 1. The classified dorsal angiosomes. Abbreviations: ADP, Arterial Dorsalis Pedis, ATP, Arteria Tibialis Posterior

Statistical Analyses

Preprocedural and postprocedural perfusion parameters were compared in the DR and IR groups for all patients and in the critical limb threatening ischemia (CLTI) subgroup using the Wilcoxon signed-rank test. Percentual changes of the perfusion parameters preinterventionally and postinterventionally comparing the DR and IR groups for all patients and in the CLTI subgroup was performed using the Mann-Whitney U test. A subanalysis based on either endovascular or open surgical revascularization was performed comparing the percentual change of the perfusion parameters after DR and IR using the Mann-Whitney U test. Statistical analysis was performed using IBM SPSS Statistics 25 (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY, USA: IBM Corp.).

3

Results

Patient Characteristics

Preinterventional and postinterventional ICG NIR fluorescence imaging was successfully performed in 52 patients (54 limbs), displayed in Table I. Fontaine stage 2a was classified in 6 limbs (11.1%). Fontaine stage 2b was classified in 26 limbs (48.2%) and CLTI considered as Fontaine stage 3 and 4 was classified in 22 limbs (40.7%). Endovascular revascularization was performed in 26 limbs (48.2%) and surgical intervention in 24 limbs (44.4%). A hybrid intervention was performed in 4 limbs (7.4%). Revascularization on femoral and/or popliteal level was performed in 40 limbs (74.1%) and 12 limbs were treated on aortoiliac level (22.2%). Two limbs were treated on a crural level (3.7%). A total of 157 angiosomes were identified, of which 121 were directly and 36 were indirectly revascularized. Collateral grading was performed in the IR group consisting of 24 limbs (36 angiosomes). Consensus in collateral grading was achieved in all cases. Significant collaterals were scored in 19 limbs, insignificant collaterals in 3 limbs and unclassifiable collaterals in 2 limbs due to insufficient contrast on preoperative CT angiography imaging and/or missing lower limb angiographic images.

Angiosome Characteristics

Characteristics per angiosome are displayed in Table II. Fontaine stage 2b was classified in the limbs of 60 DR angiosomes (49.6%) and 18 IR angiosomes (50.0%) and CLTI was classified in 44 DR and 17 IR angiosomes (36.4% and 47.2%, respectively). Sixty-two DR (51.2%) and 14 IR angiosomes (38.9%) were endovascularly treated. Surgical intervention was performed in 49 DR and 20 IR angiosomes (40.5%, and 55.5%, respectively). Revascularization in the femoral/popliteal area was performed in 84 DR (69.4%) and 32 IR angiosomes (88.9%).

Table I. Patient characteristics *

	Total (n=52, limbs=54)
Age (years)	70.2 ± 7.5
Female – n(%)	24 (46.2)
Body mass index	25.8 ± 4.8
Diabetes – n(%)	15 (28.8)
Active smoking – n(%)	11 (21.2)
Smoking history – n(%)	48 (92.3)
Renal insufficiency – n(%)	1 (1.9)
Hypertension – n(%)	35 (67.3)
Baseline TP (mmHg)	89.6 ± 42.4 ^a
Baseline ABI	0.7 ± 0.3 ^b
Fontaine stage – n(%)	
2a	6 (11.1)
2b	26 (48.2)
3	13 (24.0)
4	9 (16.7)
Type of revascularization – n(%)	
Endovascular	26 (48.2)
Surgical	24 (44.4)
Hybrid	4 (7.4)
Level of revascularization – n(%)	
Aortoiliac	12 (22.2)
Femoral/popliteal	40 (74.1)
Crural	2 (3.7)
Angiosomes – n(%)	
Total	157
DR	121 (77.1)
IR	36 (22.9)

* Plus-minus values are means ± SD

^a Obtained in n=37 limbs

^b Obtained in n=41 limbs

Abbreviations: TP, toe pressure, ABI, ankle-brachial index, DR, direct revascularization, IR, indirect revascularization

Table II. Angiosome characteristics

	DR (n=121)	IR (n=36)
Fontaine stage – n(%)		
2a	17 (14.0)	1 (2.8)
2b	60 (49.6)	18 (50.0)
3	30 (24.8)	8 (22.2)
4	14 (11.6)	9 (25.0)
Type of revascularization –n(%)		
Endovascular	62 (51.2)	14 (38.9)
Surgical	49 (40.5)	20 (55.5)
Hybrid	10 (8.3)	2 (5.6)
Level of revascularization – n(%)		
Aortoiliac	33 (27.3)	3 (8.3)
Femoral/popliteal	84 (69.4)	32 (88.9)
Crural	4 (3.3)	1 (2.8)

Abbreviations: DR, direct revascularization, IR, indirect revascularization

ICG NIR Fluorescence Results

Preprocedural and postprocedural ICG NIR fluorescence parameter values and percentual improvement for the DR and IR groups within all patients are depicted in Table III and within the CLTI subgroup in Table IV. Preinterventional and postinterventional normalized time-intensity curves for the DR and IR groups are displayed in Figure 2. A schematic overview of an endovascular treated patient along with corresponding normalized time-intensity curves is presented in Figure 3. All postprocedural perfusion parameter values changed significantly compared to the preinterventional values for both the DR (P-values <0.001) and IR groups (P-values between <0.001 and 0.007). A median percentual improvement was observed for all 10 perfusion parameters in both the DR and IR groups. When comparing the percentual improvement of perfusion parameters between the DR and IR groups, no significant differences were shown for all perfusion parameters (P-values between 0.253 and 0.881). In the CLTI subgroup analysis, the perfusion parameters changed significantly after both DR and IR (P-values between <0.001 and 0.003, 0.004 and 0.007, respectively). No significant difference was shown when comparing the percentual improvement of the perfusion parameters between the DR and IR groups (P-values between 0.134 and 0.359). Results of the subgroup analysis comparing the percentual improvement of perfusion parameters between the DR and IR groups within the endovascular and surgical procedure subgroups are displayed in Supplementary Table I. In both subgroups, all perfusion parameters displayed positive median percentual improvements and insignificant differences were seen when comparing the percentual improvement between the DR and IR groups (open: P-values between 0.053 and 0.899, endo: P-values between 0.290 and 0.883).

Table III. Pre- and postinterventional values and postinterventional percentual improvement of ICG NIR fluorescence parameters

<i>Parameters</i>	Direct Revascularization			Indirect Revascularization			Percentual improvement		
	Pre	Post	<i>p</i>	Pre	Post	<i>p</i>	DR	IR	<i>p</i>
	<i>Median (quartiles)</i>	<i>Median (quartiles)</i>		<i>Median (quartiles)</i>	<i>Median (quartiles)</i>		<i>Median percentage (quartiles)</i>	<i>Median percentage (quartiles)</i>	
Imax	33.3 (24.9/ 42.9)	46.0 (31.7/ 66.1)	<0.001	39.2 (23.7/ 47.9)	53.8 (41.5/ 68.7)	<0.001	31.3 (0.6/ 83.5)	36.4 (17.3/ 89.7)	0.401
Ingress rate	0.3 (0.2/ 0.7)	1.1 (0.3/ 2.4)	<0.001	0.3 (0.1/ 0.8)	1.2 (0.7/ 2.4)	<0.001	169.8 (10.0/ 580.4)	181.9 (76.9/ 733.1)	0.279
Absolute slope	1.0 (0.5/ 1.9)	2.6 (1.0/ 5.1)	<0.001	1.2 (0.4/ 2.3)	3.2 (1.6/ 4.6)	<0.001	103.7 (4.6/ 349.0)	141.0 (50.9/ 622.5)	0.298
Normalized slope	2.9 (2.1/ 4.4)	5.5 (3.1/ 8.4)	<0.001	3.2 (1.7/ 4.8)	6.1 (3.9/ 8.4)	<0.001	49.7 (4.9/ 159.8)	75.8 (27.1/ 220.3)	0.263
Tmax	105.9 (58.0/ 150.1)	37.7 (21.8/ 105.1)	<0.001	93.5 (47.9/ 174.4)	40.6 (24.7/ 68.2)	<0.001	41.6 (3.4/ 72.7)	57.0 (19.4/ 76.3)	0.253
AUC egress 60	96.3 (93.7/ 97.5)	91.8 (83.0/ 96.5)	<0.001	95.8 (89.9/ 97.3)	92.2 (81.8/ 96.0)	0.007	2.7 (-0.1/ 11.2)	3.8 (0.2/ 11.6)	0.881
AUC egress 120	91.9 (86.3/ 94.3)	83.6 (74.4/ 92.8)	<0.001	91.0 (84.2/ 94.8)	84.7 (72.3/ 90.0)	0.002	5.8 (0.0/ 16.6)	7.2 (0.8/ 16.1)	0.783
AUC egress 180	86.9 (79.3/ 90.1)	75.9 (64.4/ 87.4)	<0.001	85.0 (77.3/ 90.9)	77.9 (65.1/ 84.4)	0.001	7.9 (0.1/ 20.3)	10.3 (0.6/ 19.9)	0.764
AUC egress 240	81.2 (72.7/ 85.7)	69.9 (58.8/ 82.5)	<0.001	80.4 (70.8/ 87.0)	70.6 (59.4/ 79.6)	0.001	8.6 (0.0/ 23.7)	13.5 (-0.7/ 22.6)	0.767
AUC egress 300	75.7 (67.4/ 81.5)	60.5 (50.1/ 75.0)	<0.001	76.2 (64.8/ 83.2)	64.6 (54.7/ 74.1)	0.001	9.1 (0.0/ 25.0)	15.6 (-2.2/ 25.2)	0.698

Abbreviations: Imax, maximum intensity, Tmax, time to max, AUC, area under the curve, DR, direct revascularization, IR, indirect revascularization

Table IV. Pre- and postinterventional values and postinterventional percentual improvement of ICG NIR fluorescence parameters in CLTI patients

<i>Parameters</i>	Direct Revascularization			Indirect Revascularization			Percentual improvement		
	Pre	Post	<i>p</i>	Pre	Post	<i>p</i>	DR	IR	<i>p</i>
	<i>Median (quartiles)</i>	<i>Median (quartiles)</i>		<i>Median (quartiles)</i>	<i>Median (quartiles)</i>		<i>Median percentage (quartiles)</i>	<i>Median percentage (quartiles)</i>	
Imax	33.1 (25.3/ 43.0)	49.4 (32.7/ 60.7)	<0.001	40.2 (23.5/ 48.2)	55.2 (41.2/ 73.1)	0.006	22.5 (-2.5/ 91.6)	36.7 (12.7/ 138.9)	0.311
Ingress rate	0.3 (0.2/ 0.9)	1.3 (0.5/ 3.1)	<0.001	0.6 (0.1/ 1.1)	2.0 (0.8/ 4.2)	0.007	170.1 (-12.3/ 721.2)	267.2 (2.6/ 1912.6)	0.204
Absolute slope	1.1 (0.7/ 2.3)	3.5 (1.5/ 5.8)	<0.001	1.7 (0.4/ 2.6)	4.1 (2.9/ 6.9)	0.005	125.1 (-4.3/ 396.8)	142.9 (40.3/ 877.8)	0.240
Normalized slope	3.3 (2.3/ 5.6)	7.1 (5.0/ 10.4)	<0.001	4.1 (1.7/ 5.6)	7.3 (6.1/ 9.2)	0.004	82.5 (-4.6/ 161.5)	79.4 (14.0/ 353.9)	0.359
Tmax	100.1 (39.9/ 128.7)	35.5 (17.4/ 76.5)	0.003	71.2 (47.1/ 165.3)	26.1 (16.7/ 52.2)	0.004	52.5 (-3.7/ 72.4)	71.9 (4.7/ 88.2)	0.179
AUC egress 60	95.9 (91.7/ 97.8)	90.3 (79.6/ 95.4)	<0.001	96.6 (92.3/ 97.3)	82.2 (76.7/ 94.3)	0.006	4.2 (-0.4/ 12.5)	11.2 (1.9/ 17.3)	0.157
AUC egress 120	91.4 (84.2/ 94.1)	81.7 (68.2/ 91.0)	<0.001	91.0 (84.8/ 94.9)	73.5 (69.3/ 86.5)	0.006	8.3 (-0.3/ 18.9)	14.7 (4.5/ 25.4)	0.157
AUC egress 180	85.5 (77.8/ 89.7)	71.3 (58.5/ 80.0)	<0.001	85.9 (77.6/ 91.5)	67.5 (62.3/ 78.6)	0.004	9.7 (0.0/ 23.7)	17.6 (4.5/ 29.2)	0.135
AUC egress 240	79.9 (71.3/ 85.1)	64.3 (51.0/ 74.8)	<0.001	80.7 (71.3/ 87.9)	62.0 (56.3/ 72.3)	0.006	10.9 (0.0/ 27.0)	19.2 (5.0/ 32.3)	0.152
AUC egress 300	74.1 (62.5/ 80.4)	55.9 (40.1/ 64.8)	<0.001	76.0 (65.6/ 84.6)	56.5 (51.1/ 66.8)	0.006	11.6 (0.0/ 27.9)	21.4 (6.6/ 34.0)	0.134

Abbreviations: Imax, maximum intensity, Tmax, time to max. AUC, area under the curve, DR, direct revascularization, IR, indirect revascularization, CLTI, critical limb threatening ischemia




Figure 2. Pre- and postinterventional normalized time-intensity curves for the direct and indirect revascularized angiosome groups displayed as mean with standard deviations.




Figure 3. Overview of an endovascular revascularized AFS patient with an untreated ATP occlusion resulting in two direct and one indirect treated angiosome. Corresponding pre- and postinterventional normalized time-intensity curves are displayed. Abbreviations: AFS, Arteria Femoralis Superficialis, ATP, Arteria Tibialis Posterior, DR, Direct Revascularization, IR, Indirect Revascularization

Supplementary table I. Percentual improvement ICG NIR fluorescence parameters for open and endovascular procedures

<i>Parameters</i>	Percentual improvement surgical			Percentual improvement endovascular		
	DR	IR		DR	IR	
	<i>Median percentage (quartiles)</i>	<i>Median percentage (quartiles)</i>	<i>p</i>	<i>Median percentage (quartiles)</i>	<i>Median percentage (quartiles)</i>	<i>p</i>
Imax	45.2 (10.3/ 113.6)	44.2 (3.7/ 134.9)	0.899	14.5 (-2.1/ 78.0)	29.2 (20.6/ 49.5)	0.290
Ingress rate	213.3 (20.8/ 661.1)	534.5 (114.9/ 1048.4)	0.152	104.7 (0.0/ 518.3)	83.5 (30.4/ 219.7)	0.883
Absolute slope	142.2 (10.6/ 475.7)	250.2 (94.5/ 860.6)	0.274	74.4 (3.9/ 307.1)	90.8 (15.0/ 145.5)	0.830
Normalized slope	51.4 (0.5/ 226.9)	157.0 (27.8/ 313.3)	0.104	49.5 (6.4/ 124.4)	42.2 (-12.5/ 81.0)	0.414
Tmax	51.6 (11.7/ 72.8)	66.5 (45.5/ 84.9)	0.053	36.2 (0.0/ 71.4)	25.2 (-8.8/ 64.7)	0.380
AUC egress 60	5.1 (0.5/ 14.4)	6.4 (1.0/ 14.5)	0.782	1.9 (-0.8/ 9.0)	1.6 (-6.0/ 11.3)	0.639
AUC egress 120	9.6 (2.2/ 19.4)	12.0 (3.8/ 20.2)	0.640	2.4 (-1.5/ 12.6)	2.7 (-7.9/ 14.7)	0.574
AUC egress 180	14.0 (3.3/ 24.0)	14.7 (6.3/ 23.7)	0.603	3.3 (-1.6/ 14.8)	3.0 (-8.9/ 16.2)	0.503
AUC egress 240	14.6 (3.9/ 26.8)	15.9 (7.5/ 26.0)	0.545	4.3 (-1.5/ 16.3)	1.9 (-8.1/ 16.8)	0.406
AUC egress 300	17.3 (4.2/ 28.1)	18.7 (9.7/ 28.0)	0.510	1.5 (-1.8/ 17.4)	1.1 (-10.6/ 17.0)	0.376

Abbreviations: Imax, maximum intensity, Tmax, time to max, AUC, area under the curve, DR, direct revascularization, IR, indirect revascularization



Discussion

This study shows that perfusion measured with ICG NIR fluorescence significantly increased for both the DR and the IR groups after surgical and endovascular revascularization. Moreover, no difference in the degree of perfusion improvement was found between the DR and IR groups. The same results were seen when solely evaluating patients with CLTI. The significant increase in perfusion after revascularization measured by NIR fluorescence for both DR and IR is in line with the study of Rother et al., which evaluated perfusion changes in bypass surgery patients.⁸ Notwithstanding, others have argued that the angiosome concept might be of greater importance in endovascular revascularizations as during bypass surgery the most optimal outflow artery is selected.^{2,4} Yet the subgroup analysis in this study displays no significant difference comparing DR and IR in both the surgical and endovascular revascularization subgroups. Several studies investigated the clinical outcome in patients undergoing IR, showing similar clinical results compared to patients undergoing DR in case of present collaterals.²⁵⁻²⁷ The high percentage of patients treated with IR and significant collaterals in this study (86.4%), possibly reflecting to the population with LEAD, could therefore explain the significant improvement of perfusion parameters in the IR group. Comparison of perfusion changes using ICG NIR fluorescence in IR patients with and without significant collaterals would be highly relevant; however, it was not feasible in this study due to the limited sample size. Furthermore, the method for grading of collateralization is very heterogeneous in literature. The single-grade scoring system of collaterals in this study was based on 2 articles, which assessed the length, size, and number of collaterals.^{28,29} The use of a validated and unambiguous collateral scoring system is preferable for a more accurate future evaluation of this matter. The presence of choke vessels, as demonstrated by Taylor and Palmer in 1987, should also be taken into account.¹ Choke vessels connect adjacent arterial territories and therefore contribute to the variability in angiosomes borders.

In addition, the anatomical variance and pre- domination of different arteries within patients make it even more complicated to agree on 1 angiosome map.^{22,24,30} These major interpatient differences combined complicate the uniform and precise evaluation of the angiosome concept's value in revascularization decision-making. The results of this study are additionally limited by the sole evaluation of dorsal angiosomes and the small amount of crural revascularizations. Another mentionable topic of consideration is the effect of diabetes on the applicability of the angiosome concept. This patient subgroup is known for prolonged wound healing and inferior collateralization compared to nondiabetics, resulting in a potentially increased clinical benefit from DR.^{6,31,32} ICG NIR fluorescence is a useful tool to evaluate microvascular perfusion status. As microvascular dysfunction usually occurs in diabetic patients, fluorescence imaging is specifically applicable in evaluation of perfusion for this group.³³⁻³⁵

This study population was too small for adequate comparison of the perfusion difference between diabetics and nondiabetics. Several reviews have given an overview of the literature on the usefulness of the angiosome concept in procedural decision-making.³⁻⁷ Although DR seems to be of added value in certain clinical situations and patient groups, the included studies were heterogeneous and the quality of evidence was very low, resulting in the recommendation to further investigate the concept's applicability. Perfusion assessment using ICG NIR fluorescence has already proven its potential as quality control method following revascularization.^{15,17-20} Given this potential and the usability for microvascular perfusion assessment, the evaluation of the angiosome concept using ICG NIR fluorescence seems to be of added value as demonstrated in this study. Patients with claudication and CLTI were included in this study, as the main outcome was to compare perfusion changes within DR and IR. However, the angiosome concept is particularly applicable in CLTI patients with wounds as a revascularization could be targeted toward the angiosome in which a wound is located.

Hence, gaining in-line flow to the foot in patients with wounds is reasonably advocated in revascularization guidelines.³⁶ However, a subgroup analysis in patients with CLTI was performed in this study, which also demonstrated a significant improvement of perfusion in indirectly revascularized angiosomes. Future research is recommended to further evaluate the applicability of the angiosome concept in CLTI patients with the use of ICG NIR fluorescence. We advise that the previously discussed patient-specific factors and other relevant clinical parameters which influence the clinical outcome should be taken into consideration.

Conclusion

This study proves that both DR and IR lead to an improvement of microvascular perfusion measured with ICG NIR fluorescence imaging. This suggests that interventional strategies should not only focus on creating in-line flow to the supplying angiosome. One can argue that the angiosome concept is not applicable in patients with LEAD.

References

1. Taylor GI, Palmer JH. The vascular territories (angiosomes) of the body: experimental study and clinical applications. *British Journal of Plastic Surgery*. 1987 Mar;40(2):113–41.
2. Špillerová K, Settembre N, Biancari F, Albäck A, Venermo M. Angiosome Targeted PTA is More Important in Endovascular Revascularisation than in Surgical Revascularisation: Analysis of 545 Patients with Ischaemic Tissue Lesions. *European Journal of Vascular and Endovascular Surgery*. 2017 Apr;53(4):567–75.
3. Bosanquet DC, Glasbey JCD, Williams IM, Twine CP. Systematic Review and Meta-analysis of Direct Versus Indirect Angiosomal Revascularisation of Infrapopliteal Arteries. *European Journal of Vascular and Endovascular Surgery*. 2014 Jul;48(1):88–97.
4. Jongsma H, Bekken JA, Akkersdijk GP, Hoeks SE, Verhagen HJ, Fiiole B. Angiosome-directed revascularization in patients with critical limb ischemia. *Journal of Vascular Surgery*. 2017 Apr;65(4):1208-1219.e1.
5. Sumpio BE, Forsythe RO, Ziegler KR, Van Baal JG, Lepantalo MJA, Hinchliffe RJ. Clinical implications of the angiosome model in peripheral vascular disease. *Journal of Vascular Surgery*. 2013 Sep;58(3):814–26.
6. Huang TY, Huang TS, Wang YC, Huang PF, Yu HC, Yeh CH. Direct Revascularization With the Angiosome Concept for Lower Limb Ischemia: A Systematic Review and Meta-Analysis. *Medicine*. 2015 Aug;94(34):e1427.
7. Biancari F, Juvonen T. Angiosome-targeted Lower Limb Revascularization for Ischemic Foot Wounds: Systematic Review and Meta-analysis. *European Journal of Vascular and Endovascular Surgery*. 2014 May;47(5):517–22.
8. Rother U, Lang W, Horch RE, Ludolph I, Meyer A, Gefeller O, et al. Pilot Assessment of the Angiosome Concept by Intra-operative Fluorescence Angiography After Tibial Bypass Surgery. *European Journal of Vascular and Endovascular Surgery*. 2018 Feb;55(2):215–21.
9. Špillerová K, Biancari F, Leppäniemi A, Albäck A, Söderström M, Venermo M. Differential Impact of Bypass Surgery and Angioplasty on Angiosome-Targeted Infrapopliteal Revascularization. *European Journal of Vascular and Endovascular Surgery*. 2015 Apr;49(4):412–9.
10. Rother U, Kapust J, Lang W, Horch RE, Gefeller O, Meyer A. The Angiosome Concept Evaluated on the Basis of Microperfusion in Critical Limb Ischemia Patients—an Oxygen to See Guided Study. *Microcirculation*. 2015 Nov;22(8):737–43.
11. Alvelo JL, Papademetris X, Mena-Hurtado C, Jeon S, Sumpio BE, Sinusas AJ, et al. Radiotracer Imaging Allows for Noninvasive Detection and Quantification of Abnormalities in Angiosome Foot Perfusion in Diabetic Patients With Critical Limb Ischemia and Nonhealing Wounds. *Circ Cardiovascular Imaging*. 2018 May;11(5):e006932.
12. Van Den Hoven P, Ooms S, Van Manen L, Van Der Bogt KEA, Van Schaik J, Hamming JF, et al. A systematic review of the use of near-infrared fluorescence imaging in patients with peripheral artery disease. *Journal of Vascular Surgery*. 2019 Jul;70(1):286-297.e1.
13. Schaafsma BE, Mieog JSD, Hutteman M, Van Der Vorst JR, Kuppen PJK, Löwik CWGM, et al. The clinical use of indocyanine green as a near-infrared fluorescent contrast agent for image-guided oncologic surgery. *Journal of Surgical Oncology*. 2011 Sep;104(3):323–32.
14. Muntean MV, Muntean V, Ardelean F, Georgescu A. Dynamic perfusion assessment during perforator flap surgery: An up-to-date. *Medicine and Pharmacy Reports*. 2015 Jul 22;88(3):293–7.
15. Colvard B, Itoga NK, Hitchner E, Sun Q, Long B, Lee G, et al. SPY technology as an adjunctive measure for lower extremity perfusion. *Journal of Vascular Surgery*. 2016 Jul;64(1):195–201.
16. Munabi NCO, Olorunnipa OB, Goltsman D, Rohde CH, Ascherman JA. The ability of intra-operative perfusion mapping with laser-assisted indocyanine green angiography to predict mastectomy flap necrosis in breast reconstruction: A prospective trial. *Journal of Plastic, Reconstructive & Aesthetic Surgery*. 2014 Apr;67(4):449–55.
17. Settembre N, Kauhanen P, Albäck A, Špillerová K, Venermo M. Quality Control of the Foot Revascularization Using Indocyanine Green Fluorescence Imaging. *World j surg*. 2017 Jul;41(7):1919–26.
18. Braun JD, Trinidad-Hernandez M, Perry D, Armstrong DG, Mills JL. Early quantitative evaluation of indocyanine green angiography in patients with critical limb ischemia. *Journal of Vascular Surgery*. 2013 May;57(5):1213–8.

19. Igari K, Kudo T, Toyofuku T, Jibiki M, Inoue Y, Kawano T. Quantitative Evaluation of the Outcomes of Revascularization Procedures for Peripheral Arterial Disease Using Indocyanine Green Angiography. *European Journal of Vascular and Endovascular Surgery*. 2013 Oct;46(4):460–5.
20. Van Den Hoven P, S Weller F, Van De Bent M, Goncalves LN, Ruig M, D Van Den Berg S, et al. Near-infrared fluorescence imaging with indocyanine green for quantification of changes in tissue perfusion following revascularization. *Vascular*. 2022 Oct;30(5):867–73.
21. Van Den Hoven P, Tange F, Van Der Valk J, Nerup N, Putter H, Van Rijswijk C, et al. Normalization of Time-Intensity Curves for Quantification of Foot Perfusion Using Near-Infrared Fluorescence Imaging With Indocyanine Green. *J Endovasc Ther*. 2023 Jun;30(3):364–71.
22. Attinger C, Cooper P, Blume P, Bulan E. The safest surgical incisions and amputations applying the angiosome principles and using the Doppler to assess the arterial-arterial connections of the foot and ankle. *Foot and Ankle Clinics*. 2001 Dec;6(4):745–99.
23. Taylor GI, Minabe T. The Angiosomes of the Mammals and Other Vertebrates: Plastic and Reconstructive Surgery. 1992 Feb;89(2):181–215.
24. Clemens MW, Attinger CE. Angiosomes and Wound Care in the Diabetic Foot. *Foot and Ankle Clinics*. 2010 Sep;15(3):439–64.
25. Zheng XT, Zeng RC, Huang JY, Pan LM, Su X, Wu ZH, et al. The Use of the Angiosome Concept for Treating Infrapopliteal Critical Limb Ischemia through Interventional Therapy and Determining the Clinical Significance of Collateral Vessels. *Annals of Vascular Surgery*. 2016 Apr;32:41–9.
26. Acín F, Varela C, López De Maturana I, De Haro J, Bleda S, Rodríguez-Padilla J. Results of Infrapopliteal Endovascular Procedures Performed in Diabetic Patients with Critical Limb Ischemia and Tissue Loss from the Perspective of an Angiosome-Oriented Revascularization Strategy. *International Journal of Vascular Medicine*. 2014;2014:1–13.
27. Varela C, Acín F, De Haro J, Bleda S, Esparza L, March JR. The Role of Foot Collateral Vessels on Ulcer Healing and Limb Salvage After Successful Endovascular and Surgical Distal Procedures According to an Angiosome Model. *Vasc Endovascular Surg*. 2010 Nov;44(8):654–60.
28. Baumgartner I, Thoeny HC, Kummer O, Roefke C, Skjelsvik C, Boesch C, et al. Leg Ischemia: Assessment with MR Angiography and Spectroscopy. *Radiology*. 2005 Mar;234(3):833–41.
29. Keeling AN, Carroll TJ, McDermott MM, Liu K, Liao Y, Farrelly CT, et al. Clinical correlates of size and number of collateral vessels in peripheral artery disease. *Vasc Med*. 2012 Aug;17(4):223–30.
30. Špillarová K, Sörderström M, Albäck A, Venermo M. The Feasibility of Angiosome-Targeted Endovascular Treatment in Patients with Critical Limb Ischemia and Foot Ulcer. *Annals of Vascular Surgery*. 2016 Jan;30:270–6.
31. Prompers L, Schaper N, Apelqvist J, Edmonds M, Jude E, Mauricio D, et al. Prediction of outcome in individuals with diabetic foot ulcers: focus on the differences between individuals with and without peripheral arterial disease. The EURODIALE Study. *Diabetologia*. 2008 May;51(5):747–55.
32. Schaper NC, Nabuurs-Franssen MH, Huijberts MSP. Peripheral vascular disease and Type 2 diabetes mellitus. *Diabetes Metab Res Rev*. 2000 Sep;16(S1):S11–5.
33. Jörneskog G. Why Critical Limb Ischemia Criteria are Not Applicable to Diabetic Foot and What the Consequences are. *Scand J Surg*. 2012 Jun;101(2):114–8.
34. Apelqvist JAP, Lepäntalo MJA. The ulcerated leg: when to revascularize. *Diabetes Metabolism Res*. 2012 Feb;28(S1):30–5.
35. Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FGR. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *Journal of Vascular Surgery*. 2007 Jan;45(1):S5–67.
36. Conte MS, Bradbury AW, Kolh P, White JV, Dick F, Fitridge R, et al. Global Vascular Guidelines on the Management of Chronic Limb-Threatening Ischemia. *Eur J Vasc Endovasc Surg*. 2019 Jul;58(1S):S1-S109.e33.