



Universiteit
Leiden
The Netherlands

Cell walls of filamentous fungi: challenges and opportunities for biotechnology

Ost, K.J.; Student, M.; Cord-Landwehr, S.; Moerschbacher, B.M.; Ram, A.F.J.; Dirks-Hofmeister, M.E.

Citation

Ost, K. J., Student, M., Cord-Landwehr, S., Moerschbacher, B. M., Ram, A. F. J., & Dirks-Hofmeister, M. E. (2025). Cell walls of filamentous fungi: challenges and opportunities for biotechnology. *Applied Microbiology And Biotechnology*, 109(1).
doi:10.1007/s00253-025-13512-3

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/4309800>

Note: To cite this publication please use the final published version (if applicable).



Cell walls of filamentous fungi – challenges and opportunities for biotechnology

Katharina J. Ost^{1,2} · Mounashree Student² · Stefan Cord-Landwehr² · Bruno M. Moerschbacher² · Arthur F. J. Ram³ · Mareike E. Dirks-Hofmeister¹

Received: 28 February 2025 / Revised: 30 April 2025 / Accepted: 5 May 2025 / Published online: 24 May 2025
© The Author(s) 2025

Abstract

The cell wall of filamentous fungi is essential for growth and development, both of which are crucial for fermentations that play a vital role in the bioeconomy. It typically has an inner rigid core composed of chitin and beta-1,3-/beta-1,6-glucans and a rather gel-like outer layer containing other polysaccharides and glycoproteins varying between and within species. Only a fraction of filamentous fungal species is used for the biotechnological production of enzymes, organic acids, and bioactive compounds such as antibiotics in large amounts on a yearly basis by precision fermentation. Most of these products are secreted into the production medium and must therefore pass through fungal cell walls at high transfer rates. Thus, cell wall mutants have gained interest for industrial enzyme production, although the causal relationship between cell walls and productivity requires further elucidation. Additionally, the extraction of valuable biopolymers like chitin and chitosan from spent fungal biomass, which is predominantly composed of cell walls, represents an underexplored opportunity for circular bioeconomy. Questions persist regarding the effective extraction of these biopolymers from the cell wall and their repurposing in valorization processes. This review aims to address these issues and promote further research on understanding the cell walls in filamentous fungi to optimize their biotechnological use.

Key points

- *The highly complex cell walls of filamentous fungi are important for biotechnology.*
- *Cell wall mutants show promising potential to improve industrial enzyme secretion.*
- *Recent studies revealed enhanced avenues for chitin/chitosan from fungal biomass.*

Keywords Fungal cell walls · Biotechnology · Filamentous fungi · Industrial enzymes · Chitin and chitosan

Introduction

Many bacteria and yeasts can be easily grown in fermenters to produce commercially interesting products, and large-scale fermentation volumes are used to profit

from economies of scale (Demain and Adrio 2008; De Brabander et al. 2023). But sometimes, the metabolism, genetic repertoire, or posttranslational modification potential of prokaryotes and unicellular eukaryotes such as yeasts are not complex enough for target compound production. Therefore, filamentous fungi are used to produce various commodities and specialty products (Nielsen et al. 1995; Meyer 2008; Meyer et al. 2020). Moreover, filamentous fungi outcompete bacteria and yeasts in terms of efficiency of protein secretion, reaching levels of secreted proteins of up to 100 g/L, with the target protein secreted and present in the spent medium (Ward 2012; Cairns et al. 2018). The main filamentous fungal organisms used for biotechnological production are *Aspergillus niger*, *Aspergillus oryzae*, *Trichoderma reesei*, *Penicillium chrysogenum*, *Rhizopus* sp., and *Rhizomucor miehei* (Table 1). Every year, millions of tons of fungal cells are produced

✉ Mareike E. Dirks-Hofmeister
m.dirks-hofmeister@hs-osnabrueck.de

¹ Laboratory for Food Biotechnology, Faculty of Agricultural Sciences and Landscape Architecture, Osnabrück University of Applied Sciences, Oldenburger Landstraße 62, 49090 Osnabrück, Germany

² Institute for Biology and Biotechnology of Plants, University of Münster, Schlossplatz 8, 48143 Münster, Germany

³ Fungal Genetics and Biotechnology, Institute of Biology Leiden, Leiden University, Sylviusweg 72, 2333 BE Leiden, The Netherlands

in commercial biotechnological processes via (precision) fermentation (Allied Market Research 2024; Valuates Reports 2024). Valuable chemicals, food enzymes, antibiotics, and organic acids are produced and secreted, thus requiring passage through the fungal cell wall. In attempts for strain improvement the cell wall and cell wall-related genes increasingly appear as targets. At the same time, cell walls occur as an important by-product in high amounts after fermentation as part of the separated fungal biomass (Nikkilä et al. 2020), and cell wall components are considered as potentially interesting chemical building blocks. Thus, in the interest towards a circular bioeconomy, these biotechnological processes give the chance of recovering and valorizing cell wall components of high purity and reproducibility (Meyer et al. 2016, 2020; Grimm and Wösten 2018; Cairns et al. 2018, 2021b; Wösten 2019).

To provide a tangible impression of the annual quantities of fermentation volumes and fungal cell wall biomass from processes using filamentous fungi from different areas of biotechnology, we accumulated data to come to an approximate estimate as given in Table 1. Considering only fermentation products with the highest estimated yearly production, such as citric acid (Kristiansen et al. 1998; Max et al. 2010; Vandenberghe et al. 2017; Książek 2024), alpha-amylase (Reddy et al. 2003; de Souza and de Oliveira e Magalhães 2010; Balakrishnan et al. 2021; Allied Market Research 2024), glucoamylase (Norouzian et al. 2006; Mendonça et al. 2023) and cellulase enzymes (Ward 2012; Paloheimo et al. 2016; Valuates Reports 2024), penicillin G (Thykaer and Nielsen 2003; Demain 2006; Wenzel and Müller 2013), chymosin, and microbial rennets (Guerrand 2018; Sutay Kocabaş et al. 2022), the yearly fermentation volumes of filamentous fungi can be estimated to exceed 67 million cubic meters. The resulting dry biomass can be assumed to be 4–14% of the fermentation volume, depending on the strain and process (Koutinas et al. 2003; Xie et al. 2020; Bodie et al. 2021), and can then be estimated to sum up to 3–9 million tons of dry weight per year. To date, these side streams are usually either wasted or used in agriculture as fertilizers or feed (Das et al. 2002; Grimm and Wösten 2018; Nikkilä et al. 2020; Meyer et al. 2020; Cairns et al. 2021b; Crosino et al. 2021).

From a biological perspective, hyphal mycelia produced by filamentous fungi are not ideally suited for submerged fermentation. Filamentous hyphae are sensitive to shear stress caused by stirring (Amanullah et al. 2002) and can affect free nutrient diffusion (Olsvik and Kristiansen 1994; El Enshasy 2022). Furthermore, filamentous fungi growing in submerged cultures tend to develop different morphological phenotypes, ranging from dispersed growth to compact pellets, which significantly challenges industrial performance and handling (Grimm et al. 2005; Veiter et al.

2018; El Enshasy 2022). Filamentous fungi are evolutionarily optimized to grow on solid media, developing hyphal tip growth propelled by high turgor pressure (Papagianni 2004; Lew 2019). The cytosol of filamentous fungi tends to be concentrated at the hyphal tips, whereas distant parts are typically empty or filled with vacuoles. The entire mycelium is surrounded by complex cell walls, which are material- and energy-intensive to build. In biotechnological applications, these cell walls influence product secretion and ultimately result in unused and costly biomass waste.

The objective of this review is to discuss the importance of fungal cell walls, particularly those of industrially relevant filamentous species, in biotechnology. This aspect with elaboration on cell walls in the context of industrial utilization of fungi, puts significant novelty in our attempt compared to previous review articles. We (i) briefly describe the current understanding of fungal cell wall composition, architecture, complexity, and flexibility, focusing on biotechnologically relevant filamentous fungal species from different taxonomic groups, excluding yeasts. We then (ii) summarize the fungal cell wall biosynthesis and assembly, remodeling, and integrity pathways, as well as up-to-date data on secretory pathways in fungi. Third, we (iii) focus on the assumably high impact of the cell wall on protein adsorption, morphology, high-secretion stress, and over-all enzyme productivity. Finally, we (iv) will discuss existing and future opportunities and challenges in utilizing fungal biomass, in particular the cell walls, most prominently as an alternative source of the valuable biopolymers chitin and chitosan.

The fungal cell wall

Bacterial cell walls are rather rigid structures that essentially consist of a single, covalently linked peptidoglycan super-molecule enclosing the entire prokaryotic cell (Garde et al. 2021). In contrast, fungal and plant cell walls are more complex and flexible, where different types of polysaccharide polymers are connected by few covalent cross-links and many weak interactions, such as hydrogen bonds. In most plant cell walls, long cellulose fibrils are crosslinked by xyloglucans, and this alkali-insoluble fibrillar network is embedded in an alkali-soluble gel-like matrix composed of hydrated pectins (Cosgrove 2024).

A similar fiber-in-a-matrix architecture has also been suggested for fungal cell walls, as an alkaline insoluble core structure remains after extraction of an alkaline-soluble matrix (Wessels and Sietsma 1981). There is also a strong similarity in the physical structure of the insoluble fibers observed under an electron microscope, as well as in the chemical structure of chitin and cellulose molecules and crystals (Kostag and El Seoud 2021; da Rosa et al.

Table 1 Estimated yearly production and fermentation volumes of major industrial products derived from filamentous fungi fermentation. The source of the underlying data is explained in the footnotes below the last table row

Fungal species as production organism	Product	Annual production (commercial product)	Share of fungal species from the whole production	Assumed content of the ingredient in the products	Annual production of the ingredient by the fungus	Assumed yields of ingredients in industrial fermentations	Annual fermentation volumes
		[× 1000 t/a]			[× 1000 t/a]	[t/m ³]	[× 1000 m ³ /a]
<i>Ascomycetes</i>	<i>Aspergillus</i> sp.	Citric acid	3,300 ^a	80% ^b	100%*	2,640	0.11 ^c
		alpha-Amylases	25,500 ^d	10% ^e	10% ^e	255	0.03 ^f
		Glucoamylase	8,500 ^g	85% ^h	10% ^e	765	0.03 ^f
		Cellulases	549 ⁱ	20% ^e	10% ^e	11	0.03 ^f
		Chymosin	n.r	50% ^j	n.r	n.r	n.r
	<i>Penicillium</i> sp.	Penicillin G	60 ^l	95% ^e	100% ^e	57	0.07 ^m
<i>Mucoromycetes</i>	<i>Trichoderma</i> sp.	Cellulases	549 ⁱ	70% ^e	10% ^e	38	0.10 ⁿ
	<i>Rhizopus</i> sp.	Glucoamylase	8,500	15% ^h	10% ^e	85	0.03 ^f
	<i>Rhizomucor</i> sp.	Microbial rennet	n.r	99% ^k	n.r	n.r	n.r
							66,686

n.r. not relevant to our calculations

*Estimated values

^aKsiążek (2024)

^bVandenbergh et al. (2017)

^cKristiansen et al. (1998) stated that citric acid constitutes up to 15% of the total production, whereas Max et al. (2010) postulated final yields of 110–140 g/L of citric acid; to keep it conservatively, we calculated with 11 tons per m³

^dThe proportion of microbially produced alpha-amylases is estimated to be 25–30% of the industrial enzyme market (Reddy et al. 2003; de Souza and de Oliveira e Magalhães 2010; Balakrishnan et al. 2021). Based on the growing market (Allied Market Research 2024), we assume 30% for our calculations

^eMost industrial enzyme products are standardized to a minimal enzyme activity but mostly state a minimal protein content of 10%

^fCairns et al. (2018) stated that industrial *Aspergillus* strains produce 30 g/L of extracellular enzyme; thus, we used a quotient of 0.03 to calculate the fermentation volume from the annual production

^gThe number was concluded from the Astute Analytica (2024) market research report on “Enzyme Market,” which stated that in 2023, the world total enzyme production by microorganisms was 85 million tons. Mendonça et al. (2023) estimated the share of glucoamylase in the industrial enzyme market as 10%

^hNorouzian et al. (2006) mentioned *Aspergillus* and *Rhizopus* to be the only industrially relevant fungal glucoamylase producers, so we assumed a ratio of 85:15 for these two species

ⁱThe forecast was taken from the Global Cellulase Market Research Report from Valuates Reports (2024)

^jSutay Kocabaş et al. (2022) stated that fermentation-produced chymosin is either produced by *Aspergillus* (Chr. Hansen) or *Kluyveromyces* (DSM)

^kGuerrand (2018) postulated that Chr. Hansen holds 35% of the market share for dairy products and DSM 25–30% including enzymes (\$250 million per year) and 60% of the market of dairy enzymes to be milk clotting enzymes (= \$150 million per year). Additionally, the authors estimated the cost of rennet enzymes to be approximately \$0.05 per liter. From these values, we concluded that the total fermentation volume for rennets was 3,000 million liters, or 3 million m³ per year

^lThykaer and Nielsen (2003) stated that the annual production of penicillin exceeds 60,000 tons

^mDemain (2006) stated commercial *Penicillium* strains to produce 70 g/L penicillin

ⁿWard (2012) reported that *Trichoderma reesei* secretes 100 g/L of cellulase

2024). However, recent developments in analytical techniques, such as enzymatic-mass spectrometric fingerprinting (EMS-FP) and, above all, solid-state nuclear magnetic resonance (ssNMR), which allow investigation of the native

cell wall in situ without prior extraction of individual cell wall components, and the increasing possibility of generating genetically engineered mutants, have provided evidence that tends to challenge this assumption of a simple




Fig. 1 Schematic representation of fungal cell wall structures and biosynthetic proteins. UDP: uridine diphosphate; Glc: glucose; Gal: galactose; GlcNAc: *N*-acetylglucosamine; GalNAc: *N*-acetylgalactosamine; ER: endoplasmic reticulum; RlmA: MADS-type transcrip-

tion factor; MpkA: mitogen-activated protein kinase A; GPI: glycosylphosphatidylinositol (created using BioRender.com and modified from Yoshimi et al. 2016; Kang et al. 2018; Speth et al. 2019; Fontaine and Latgé 2020; Gow and Lenardon 2023)

fiber-in-matrix-architecture (Gow et al. 2017; Chakraborty et al. 2021; Fernando et al. 2023; Sun 2024). However, detailed analyses have only been performed on a few filamentous fungi, including the ascomycetes *Aspergillus fumigatus* and some other Aspergilli, the basidiomycetes *Schizophyllum commune* and *Cryptococcus neoformans*, and a few chitosan-containing mucoromycetes, such as *Rhizopus delemar* (Kang et al. 2018; Ehren et al. 2020; Safeer et al. 2023; Cheng et al. 2024). These results show that fungal cell walls exhibit tremendous variability both between species and within a given species under different environmental conditions or at different developmental stages. Clearly, any generalizations based on the too few examples available today are necessarily preliminary and need to be taken with great caution.

Figure 1 and the remaining of this section provide our view on the current understanding of fungal cell wall composition, architecture, synthesis, and assembly. As mentioned above, we are fully aware that this working model of the fungal cell wall is a snapshot of this work in progress.

Restrictions and improvement of cell wall analytics

As mentioned before, conventional cell wall analysis of fungi and other organisms historically relied on chemical or enzymatic hydrolysis of the polymers and analysis of the degradation products, often preceded by sequential extraction of different polymers (Heux et al. 2000; Wu et al. 2005; Cai et al. 2006; Krake et al. 2024). These techniques have

yielded an inventory of cell wall polymers with known monomer compositions and their glycosidic linkages, but have struggled to resolve their dynamic multilayered architecture (Latgé 2007; Blatzer et al. 2020). Thus, the interactions of polymers to form the complex architecture of the cell wall remain largely speculative. Furthermore, these analyses are prone to errors due to material loss during harsh chemical treatment and/or modifications introduced during the process.

Recent advances in high-resolution techniques, such as ssNMR, have more accurately mapped biomolecular interactions in the cell walls of *A. fumigatus*, *S. commune*, *Rhizopus*, and *Mucor* species, revealing significant deviations from earlier cell wall models (Kang et al. 2018; Ehren et al. 2020; Chakraborty et al. 2021; Ghassemi et al. 2022; Safeer et al. 2023; Lamon et al. 2023; Cheng et al. 2024). However, also reports on polysaccharide and protein composition, as well as the crosslinking of cell wall polysaccharides, differ significantly in the literature, and it remains to be seen whether these differences are the result of the method of analysis, different growth conditions, or proof of the immense flexibility of fungal cell walls.

EMS-FP has helped fine-tune cell wall analyses, for example, providing initial insights into the distribution of acetylated and deacetylated monomeric units within chitin (a polymer of *N*-acetylglucosamine) and chitosan (partially deacetylated chitin) chains (Urs et al. 2023). Solid-state NMR on intact cell walls is well suited for revealing polymer configurations and polymer–polymer interactions, and this has resulted in the realization of additional complexities, such as the existence of up to a dozen subtypes of chitin crystal structures (Cheng et al. 2024). Clearly, advanced cytochemical probes such as antibodies, lectins, or biotechnologically produced affinity proteins, as well as advanced microscopic techniques such as solid-state dynamic nuclear polarization (DNP-) NMR, which has a sensitivity 100–1000 times higher compared to conventional ssNMR, mass spectrometry imaging (MSI), or super-resolution and live-cell imaging techniques such as direct stochastic optical reconstruction microscopy (dSTORM), structured illumination microscopy (SIM), or confocal laser scanning microscopy (CLSM), will contribute to further refining the spatial resolution of fungal cell wall analysis in the near future (Lin et al. 2016; Islam et al. 2017; Götz et al. 2020).

Cell wall components, biosynthesis, and dynamic remodeling

Typically (Fig. 1), fungal cell walls are believed to have two layers: an evolutionary apparently rather conserved rigid skeletal layer of chitin, beta-1,3-glucans, beta-1,6-glucans, and beta-1,3-beta-1,6-glucans and a more variable, partially gel-like and sometimes hydrophobic outer layer varying

between and within species that may include alpha-/beta-glucans and other homo- and heteropolysaccharides as well as (phosphor)glycoproteins, hydrophobins, or melanin, etc. (Latgé 2007, 2010; de Groot et al. 2009; Beauvais et al. 2014; Beauvais and Latgé 2015; Erwig and Gow 2016; Gow et al. 2017; Yoshimi et al. 2017; Ehren et al. 2020; Safeer et al. 2023). This outer layer is probably responsible for mediating interactions with the environment, including a stealth mechanism of pathogenic fungi to cloak the conserved rigid core, thus preventing its recognition by the immune systems of their hosts (Gow et al. 2017; Garcia-Rubio et al. 2020; Student et al. 2024). Studies on the cell wall structure, biosynthesis, and remodeling mechanisms are important, given that cell walls have long been recognized as an ideal target for antifungal therapies, as fungal cell wall polymers are unique and do not occur in humans or, with few exceptions, in plants.

Fungal cell wall polymers are synthesized and assembled through tightly regulated enzymatic pathways in the cytoplasm or at the plasma membrane at the hyphal apex where they integrate into the growing wall (Yoshimi et al. 2016; Kang et al. 2018; Garcia-Rubio et al. 2020). Covalent crosslinks between polymers must be formed extracellularly, possibly by GPI-anchored enzymes on the outer surface of the plasma membrane (Hopke et al. 2018). While some polymers, such as chitin, alpha-1,3-glucan, and beta-1,3-glucans are known to be synthesized by transmembrane glycosyl transferases and directly transported into the extracellular space (Brauer et al. 2023), other cell wall polymers are thought to be synthesized in vesicles and secreted into the cell wall by exocytosis (Henry et al. 2016), but this is more an assumption than supported by experimental evidence. Studies in *Ustilago maydis* and *Zymoseptoria tritici* have shown that chitin synthase and beta-glucan synthase co-localize in the same vesicle which are transported along microtubules to the growing hyphal tip (Schuster et al. 2016, 2020). These studies highlight that the microtubule-dependent co-delivery of cell wall enzymes is a conserved mechanism in both basidiomycetes and ascomycetes, contributing to the efficient synthesis of the fungal cell wall.

Chitin is located in the inner layer and enhances the structural stability and resistance to enzymatic degradation (Bowman and Free 2006; Free 2013; Brown et al. 2019; Garcia-Rubio et al. 2020). Various chitin synthases (Chs) produce chitin polymers with different lengths at specific sites (Mellado et al. 1996; Brauer et al. 2023). Cell wall beta-glucan forms a crucial network for cell shape and mechanical strength and is produced by plasma membrane-bound beta-glucan synthases (Fks) (Beauvais et al. 2001; Ruiz-Herrera and Ortiz-Castellanos 2019; Wagener et al. 2020; Hu et al. 2023). Some fungi also harbor *ags* genes encoding alpha-glucan synthases (Ags), which contribute to both the mobile outer cell wall structure and the inner

structural stability (Yoshimi et al. 2017; Kang et al. 2018; Blatzer et al. 2020; Ehren et al. 2020; Safeer et al. 2023). Additionally, glycoproteins covalently bound to beta-glucans are vital for the cell wall structure and function, serving as adhesins or receptors for environmental and cellular interactions (Vogt et al. 2010; De Groot et al. 2013; Gow et al. 2017; Garcia-Rubio et al. 2020). Other complex polysaccharides with different structures represent cell wall mannans, such as glucomannans, galactomannans (GM), and galactoxylomannan (GXM), and galactosamins and galactosaminogalactans (GAG). These complex polysaccharides were found particularly in opportunistic fungal pathogens from *Aspergillus*, *Candida*, or *Cryptococcus* (Gow et al. 2017), and heterogalactans such as mannogalactans in the cell wall of *S. commune* (Dalonso et al. 2021). Genes involved in the synthesis of most of these polymers have been identified (Wang et al. 2013; Henry et al. 2016, 2019; Onoue et al. 2018; Speth et al. 2019; Hira et al. 2020; Fontaine and Latgé 2020; Kadooka et al. 2022). Cell wall (gluco)mannoproteins can cover the cell wall surface and are shown to be covalently bound to the cell wall core (Klis et al. 2007; Latgé 2007; Tefsen et al. 2012; Henry et al. 2019). In addition to (gluco)mannoproteins, also GAG, produced by the membrane-located glycosyl transferase Gbt3, which uses products from the cytosolic glucose-4-epimerase Uge3, is a recently discovered exopolysaccharide in the outer layer of the *A. fumigatus* cell wall that plays a key role in fungal interactions with the environment and the host immune system by forming a biofilm (Gravelat et al. 2013; Lee et al. 2014; Speth et al. 2019; Le Mauff 2020; Lamon et al. 2023; Le Mauff and Sheppard 2023). And these are just a few examples; it is not meant as a comprehensive overview of all the many complex polysaccharides reported in fungal cell walls.

The ability of fungi to thrive in various environments depends on the integrity and flexibility of their cell walls. To ensure structural stability while facilitating growth and adaptation, fungi employ sophisticated regulatory mechanisms, including the so-called cell wall integrity (CWI) pathway and the mechanisms and enzymes involved in cell wall remodeling (Beauvais et al. 2014; Beauvais and Latgé 2015). The CWI signaling pathways are vital for sensing and responding to cell wall stress. Mechano-sensor proteins, such as wall stress component-type sensors (WscA, WscB), detect changes in the cell wall and initiate signaling cascades to regulate the expression of genes related to cell wall biosynthesis (Rodicio et al. 2008; Yoshimi et al. 2022). These sensors are linked to intracellular signaling pathways, such as the mitogen-activated protein kinase (MAPK) cascade, including MapkA pathways, which are vital for stress signal transduction (Levin 2005, 2011; Perez and Cansado 2010; Dichtl et al. 2012, 2016; Park et al. 2016). Cell wall-modifying enzymes, including

glucanases, chitinases, transglycosylases, mannosidases, peroxidases, deacetylases, and synthases (for chitin, glucans, or other cell wall components), play critical roles in the degradation and remodeling of fungal cell walls and are activated through transcriptional regulators such as RlmA (Damveld et al. 2005; Rocha et al. 2016).

The CWI and cell wall remodeling pathways work together to regulate cell wall biosynthesis, ensuring proper construction, maintenance, and adaptation of the fungal cell wall in response to environmental and physiological changes.

Mechanism of protein secretion and transfer across the fungal cell wall

The ability of filamentous fungi to secrete large amounts of enzymes into the environment is the key reason for their successful application in industrial biotechnology. Their superior secretion performance originates from their lifestyle, with secreted enzymes and organic acids enabling nutrient digestion and absorption from their surroundings (Nevalainen and Peterson 2014; McCotter et al. 2016). Protein secretion in filamentous fungi – as in all organisms – is a complex and highly regulated process involving protein biosynthesis, post-translational modifications, and transport to the extracellular environment (reviewed by Sakekar et al. 2021 and Jadhav et al. 2024). Moreover, the filamentous growth of fungi adds extra complexity because of the relatively long distances for secretion vesicles to be transported to the growing tip of the hyphal cell (Takeshita 2016; Cairns et al. 2021b).

The protein secretion process in filamentous fungi follows the conventional secretion pathway using the ER-Golgi route: extracellular proteins are synthesized at ribosomes, translocated into the ER lumen via an N-terminal signal sequence, folded, and modified post-translationally within the ER (Boppidi et al. 2018; Wang et al. 2020; Jadhav et al. 2024). Folded proteins are packaged into vesicles and sent to the Golgi apparatus for further processing, including sorting (Nevalainen and Peterson 2014; Jadhav et al. 2024). Vesicles with mature secretory proteins or cell wall-remodeling enzymes travel from the Golgi along microtubules and actin filaments to the Spitzenkörper, releasing their contents into the extracellular space upon fusion with the plasma membrane. The classical secretion efficiency is usually much higher for native fungal proteins than for heterologous proteins, often by a factor of 10–1000 (Boppidi et al. 2018; Sakekar et al. 2021). *A. niger* can generate more than 25–30 g/L of homologous glucoamylase in fermentation media (Cairns et al. 2018), while *T. reesei* can secrete up to 100 g/L of its cellulase (Ward 2012), demonstrating their exceptional productivity. More examples of highly valued heterologous protein production in fungal hosts have been reviewed by

Punt et al. (2002), Wei et al. (2021), and Liu et al. (2023). For this reason, filamentous fungi have emerged as versatile platforms for industrial protein production. High levels of protein secretion have been linked to increased biomass formation and altered growth patterns (Hayakawa et al. 2011; Read 2011; Benoit et al. 2015; Fiedler et al. 2018b; Cairns et al. 2019a, 2021a).

Alternative unconventional secretion routes exist next to the conventional secretion pathway, providing a potential alternative for protein export. These unconventional protein secretion (UPS) routes bypass ER quality control and Golgi modifications (Stock et al. 2012). In filamentous fungi, an example of a UPS protein is the chitinase Cst1 in *U. maydis*, which follows a lock-type, unconventional, and signal peptide-independent secretion during cytokinesis (Stock et al. 2012; Reindl et al. 2019; Aschenbroich et al. 2019; Sakekar et al. 2021). Cst1 is crucial for cell wall biosynthesis and regulation, as it hydrolyzes chitin into smaller oligosaccharides for recycling or incorporation into the cell wall matrix (Langner and Göhre 2016). The use of unconventional secretion to co-export heterologous proteins into the culture supernatant is considered an attractive alternative for overcoming the limitations of conventional secretion pathways (Hayakawa et al. 2011; Read 2011; Terfrüchte et al. 2017; Kwon et al. 2017; Cairns et al. 2021a; Liu et al. 2023; Jadhav et al. 2024). In *A. niger*, the aspartic protease PepN has also been suggested to be secreted by an unconventional secretion pathway (Burggraaf et al. 2016). However, the molecular mechanisms underlying the UPS pathway in filamentous fungi are not yet fully understood.

After passing across the cell membrane through any secretory pathway, the secreted fungal proteins still need to pass through the cell wall. Studies on isolated fungal cell walls showed a permeability of 5 to 270 kDa (Trevithick and Metzenberg 1966; De Nobel and Barnett 1991; Jacobson and Ikeda 2005) depending on the species, its degree of melanization, and the fungal stress levels. It has been shown that fungi also have a specific extracellular vesicle transport for macromolecules, including proteins, via the cell wall, which is highly similar to mammalian exosomes and bacterial vesicles (Casadevall et al. 2009; Oliveira et al. 2010; Rodrigues et al. 2015). Although the exact mechanism by which vesicles pass through the cell wall remains unclear, researchers have proposed various potential and partly controversial explanations, such as the existence of pores or channels (De Nobel and Barnett 1991; Casadevall et al. 2009; Rodrigues et al. 2015). Chevalier et al. (2023) describe *A. nidulans* cell walls as dynamic and stable, balanced between secretion and expansion. They showed that localized remodeling occurs at the hyphal tip through secretion of enzymes that assemble and soften the cell wall (Chavelier et al. 2023). This remodeling is spatially confined and regulated via mechanical feedback, where

wall deformation promotes vesicle recruitment, enabling hyphal expansion at the apex. The authors reported that the cell wall at the hyphal apex is thinner and softer than the lateral wall, suggesting that this apical region forms a mechanical barrier through which vesicles can pass. Local softening and decreased thickness correlate with secretory vesicle accumulation, likely facilitating vesicle docking and protein interactions with the cell wall matrix. Further, de Paula et al. (2019) reported that extracellular vesicles transport cellulases in the industrially relevant enzyme producer *Trichoderma reesei*, which shows that vesicle transport via the cell wall is not limited to a communication system for pathogenic species, such as *Cryptococcus* sp., *A. fumigatus*, and *Candida albicans*, but could be a common system in fungi. The extent to which unconventional protein secretion and vesicle transport via the cell wall play a crucial role in high-level industrial protein secretion is not yet evident in the literature; however, it is worth further investigation.

How fungal cell wall structure affects industrial enzyme production

Most industrial hyperproduction strains have historically originated from classical strain engineering without any targeted genetically engineered modifications (Nevalainen and Peterson 2014). For example, high levels of cellulase expression have been generated by random mutagenesis and selection, resulting in *T. reesei* strain Rut-C30 being the parental strain of industrially used *Trichoderma* cellulase producers (Eveleigh and Montenecourt 1979). The industrial *Aspergilli* separate genetically in acid and enzyme producing strain lineages (Schäfer et al. 2020), both developed by separate classical strain development programs.

Improvements such as high-copy insertions, elimination of unwanted side-activities, or deletions of proteases are introduced by genetic engineering (Punt et al. 2008; Heimele 2015), but the underlying chassis strains normally are older “working horses” of the relevant companies like Novonesis (formally Novozymes and Chr. Hansen), DSM, AB Enzymes, and others. Unfortunately, though understandably, publications from industry on these strains are rare, and studies on laboratory fungal strains are not easily transferable to industrial strains, particularly with regard to improvements in protein production. Furthermore, any effect on the cell wall composition or structure by high secretion or vice versa has until now not been a focus of research, and conclusions about its role are difficult to find in the literature.

In this section, we aim to highlight what has been investigated and published regarding the influence of cell wall mutants on enzyme production in industrially relevant filamentous fungi.

Protein transfer through the fungal cell wall is influenced by biochemical interactions

Frequently, enzymes destined for secretion in their filamentous fungal production hosts appear to get stuck in the cell wall and are not released into the media as desired for proper downstream processing (Sato et al. 2011; Zhang et al. 2017a, b). This is more common for heterologous than for homologous proteins. As mentioned earlier, the permeability of filamentous fungal cell walls and the size of the secreted protein have a significant effect on whether it is released into the medium or retained in the periplasmic space (De Nobel and Barnett 1991). The cell wall can be considered a molecular sieve in which the hydrodynamic radius of a secreted molecule, rather than the molecular mass, is important. Hence, glycosylation, as a size increaser, influences transfer via the cell wall, but it also changes the polarity of a protein, and this can also affect the efficiency of proteins secretion into the medium (De Nobel and Barnett 1991). Such absorption effects of proteins and enzymes to the fungal cell wall have been reported for homologous alpha-amylase, heterologous bovine serum albumin, and hen egg lysozyme in *Aspergillus oryzae* (Sato et al. 2011). Recently, it was shown that a catalase produced in an industrially relevant *A. niger* strain was associated with the mycelium, whereas the production of glucoamylase consistently resulted in its secretion into the culture medium (Ram, unpublished data). Moreover, we recently found that two low-molecular-weight (only 20 kDa) luciferases, to be used as reporter genes for secretion, could not be found in the supernatant of *A. niger* cultivation. They remained associated with the biomass in their active, luminescence-producing forms (Ost and Dirks-Hofmeister, unpublished data). Cell walls thus can better be understood as gel matrices in which the density and degree of crosslinking but also the biochemical composition in terms of polarity and charge (e.g., content of chitosan or mannosylphosphates) define the porosity and permeability of secreted compounds (De Nobel et al. 1989, 1990b, 1990a). Clearly, cell-wall-trapped proteins might stick due to affinity reasons or get stuck because of steric hindrance, both limiting passage through the cell wall.

While it is difficult to change the biochemical characteristics of a target enzyme that is to be produced and secreted in filamentous fungi, a few attempts have shown that protein trapping can be circumvented by engineering cell wall-related genes in the production strain. Zhang et al. (2017a, b) reported that alpha-amylase (TAKA-amylase A, TAA) in *A. oryzae* binds to chitin or chitosan in the cell wall and that alpha-1,3-glucans can inhibit this absorption effect. Absorption – as binding of secreted enzymes to cell wall components – may result in the disappearance of enzyme products, especially during later stages of fermentation, and this is likely due to ionic interactions brought about by changes in

the cell wall composition during the fermentation process (Sato et al. 2011). Early studies have shown that this binding is influenced by pH, chemicals, protease treatment, and other factors and that the binding of proteins can also be mediated by isolated cell walls (Tonomura et al. 1963; Yabuki and Fukui 1970).

Productivity of filamentous fungi is related to macromorphology influenced by cell wall composition

Filamentous fungi exhibit diverse morphological phenotypes in submerged cultures that are particularly important for fermentability (reviewed by Papagianni 2004; Krull et al. 2013; Miyazawa et al. 2020; El Enshasy 2022). Macromorphology is known to be one major factor in the handling of filamentous fungi in industrial settings with large-scale bioreactor volumes but, more importantly, morphology seems to impact productivity (Grimm et al. 2005; Veiter et al. 2018). Fungal industrial enzyme-producing strains preferentially exhibit high-branching and dispersed growth phenotypes that enable lower viscosity, better mass transfer, and increased protein secretion (Wongwicharn et al. 1999; McIntyre et al. 2001; Wucherpfennig et al. 2010, 2011; Quintanilla et al. 2015).

The length of a hyphal element between two branching points is referred to as the hyphal growth unit (HGU) (Caldwell and Trinci 1973). A low HGU is preferred for industrial enzyme producers because the hyphal tip is assumed to be more permeable due to higher plasticity of the cell wall (Krull et al. 2013). Fungi growing in dense pellets show an increased number of inactive cores compared to viable hyphal tips, which limits mass transfer (Driouch et al. 2012). Additionally, Papagianni (2004) reviewed autoradiography studies from the 1970s, which showed progressive thickening of the cell walls located more distant from the apex. However, for some bulk chemicals, such as citric acid production in *A. niger*, pellet or clump morphology is advantageous (Papagianni 2004, 2007) while for others, such as penicillin production in *P. chrysogenum*, morphology does not seem to have a clear effect (Nielsen et al. 1995). Zheng et al. (2022) showed that overexpression of the catalytic subunit of cAMP-dependent protein kinase (*pkaC*) in *A. niger* altered pellet surface morphology and promoted hyphal growth at the surface. This leads to a looser outer layer, which improves the mass transfer of nutrients, oxygen, and secreted products, resulting in a 1.87-fold increase in citric acid production.

These examples, however, refer to small metabolites, in contrast to larger proteins and enzymes to be transferred via the cell wall, which could be the reason why morphology and a potentially corresponding cell wall have a stronger impact. A significant difference in the cell wall composition of a dispersed growing fungal strain in contrast to that of a pelleted

growing strain seems plausible but has not yet been studied systematically. Kissler et al. (1980) reported a change in cell wall composition and morphology for citric acid production: the absence of manganese ions in the medium inhibited glycoprotein turnover and resulted in loss of hyphal polarity, increased branching, and chitin biosynthesis in the cell wall. In contrast, hyperbranching induced by deletion of *racA* (Rho-GTPase) in *A. niger* resulted in a modified number of hyphal tips (Kwon et al. 2011, 2014) and led to a four-fold increase in glucoamylase secretion (Meyer et al. 2011; Fiedler et al. 2018a). However, no analyses of the cell wall composition have been performed in these studies. Cairns et al. (2023) found that smaller pellets in *A. niger* were associated with higher protein titres. The analysis showed that a reduced pellet diameter and increased sensitivity to cell wall stress were positively correlated with increased protein secretion. Additionally, factors such as the radial growth rate and fitness under temperature or cell wall stress were described as key predictors of protein productivity. These findings suggest that manipulating pellet size and cell wall integrity could enhance protein yield in submerged cultures.

Further studies have reported targeted genetic modification of the cell wall to change the fungal macromorphology, thereby improving enzyme secretion. DSM patented an *agsE*-deleted strain of *A. niger* as an enzyme producer (Van Peij et al. 2014), and Novozymes (renamed Novonesis in 2023) followed shortly afterwards, stating that deleting more than one *ags*-gene was more beneficial than deleting *agsE* alone (Udagawa 2016). More recent studies on this particular alpha-glucan synthase have shown that its deletion is accompanied by dispersed growth of *A. niger* mutants in submerged cultures (Lyu et al. 2023; Ost et al. 2025). Furthermore, *agsB* is required for normal growth of *A. nidulans* under liquid culture conditions (Yoshimi et al. 2013) and the deletion of two chitin synthases significantly changed the morphology of *A. niger* (Müller et al. 2003). Thus, it can be concluded that the activity of cell wall-modifying enzymes and the macromorphology of filamentous fungi are causally related.

The question remains, if besides changes in morphology, changes in cell wall composition and/or increased enzyme secretion occur in morphological mutants. Some studies have used chitin synthase genes to alter the mycelial morphology to smaller, denser pellets and increase the product yields of penicillin in *P. chrysogenum* (Liu et al. 2013) or citric acid in *A. niger*, accompanied by a 50% reduction in chitin content in the cell wall (Sun et al. 2018). Previous studies have shown that chitin synthesis plays an important role in determining the hyphal morphology during submerged fermentation (Tsuizaki et al. 2013). Recently, Barthel et al. (2024) performed a systematic functional analysis of the entire predicted *chs* gene family (nine genes) in *A. niger*, which showed that the amounts of cell wall chitin

and glucans, macromorphology, and protein secretion were affected by the deletion of specific *chs*-genes. Although the results showed that the deletion of *chsF* resulted in increased titers of extracellular proteins, the authors discussed that there was no obvious link between higher protein secretion and any specific phenotype that was determined in the study.

Deletion of all known *ags*-genes in *A. niger*, together with the whole *crh*-gene family which is involved in cross-linking chitin and glucans in the inner cell wall layer, also affects *A. niger* morphology in submerged cultures, although no changes in the monosaccharide composition of the cell wall or protein secretion have been observed (Ost et al. 2025). In contrast, Lyu et al. (2023) reported increased cellulase production in an *agsE* deletion strain. Moreover, an approach to increase the production of enzymes (cutinase encoding *cutLI*) in *A. oryzae* succeeded by deleting all three alpha-1,3-glucan synthases simultaneously, which decreased pellet size and was accompanied by increased productivity (Miyazawa et al. 2016). A follow-up study showed that combining these *ags*-mutations with the deletion of galactosaminogalactan synthesis resulted in a fully dispersed phenotype with improved fermentative culture rheology and increased productivity of recombinant cutinase (Miyazawa et al. 2019; Ichikawa et al. 2022).

Other reports have shown that enzyme production increases if the morphology is altered only by chemical means, for example, by the addition of inorganic microparticles (Driouch et al. 2011, 2012). Comparably, high agitation in a bioreactor leads to morphologically compact and strong mycelia and provides the cell wall with greater resistance to hydrolases in *A. niger* (Musílková et al. 1981).

These studies suggest that an altered cell wall composition can lead to changes in morphology. Although many morphological mutants have been described in the literature, only a few have analyzed cell wall or protein secretion. Thus, whether morphological changes always go along with changes in the cell wall composition remains to be investigated.

High-secretion stress challenges cell wall integrity and potentially cell wall composition

As described earlier, enzyme secretion from the ER and Golgi apparatus via vesicle exocytosis is correlated with hyphal growth (Wösten et al. 1991; Kwon et al. 2011, 2014, 2017; Read 2011; Fiedler et al. 2018b, 2018a; Cairns et al. 2019b, 2021b, 2021a; Chevalier et al. 2023; Zuriegat et al. 2024). Jadhav et al. (2024) reviewed recently how high-secreting fungal strains are stressed due to their rapidly growing hyphal tips. This consequently not only feedback-stresses the ER machinery but also demands a constantly high supply of nutrients and oxygen with a suitable mass transfer to the inner cell machinery.

At the same time, endocytosis plays a key role in balancing secretion to sustain hyphal growth and cellular maintenance by retrieving excess membrane and recycling secretory components such as cell wall modifying enzymes (Higuchi et al. 2009; Steinberg et al. 2017; Commer and Shaw 2021; Barata-Antunes et al. 2021). Therefore, under high secretion stress, an increase in endocytosis may limit protein secretion efficiency by internalizing proteins before they reach their extracellular targets (Cairns et al. 2019b). During fermentation, secretion is no longer restricted to hyphal tips, but still requires dynamic rebalancing of exo- and endocytosis (Cairns et al. 2019b; Commer and Shaw 2021). Fungal morphology, such as hyphal branching and septation, must adapt to these changes to achieve higher productivity levels. A key regulator of protein secretion and hyphal growth has been identified in studies on the small GTPase Arf in *A. nidulans* and *A. niger* (Lee and Shaw 2008; Meyer et al. 2011; Fiedler et al. 2018b). ArfA is known to be involved in exocytic and endocytic pathways, and its overexpression leads to an increase in both total protein and glucoamylase secretion (Yahara et al. 2001; Meyer et al. 2011; Suda et al. 2018).

To circumvent ER stress, fungi like *Aspergillus* sp. have been shown to induce the expression of genes encoding ER-resident chaperones such as heat-shock proteins (Hsp 70, Hsp90, and Hsp60), foldases such as binding immunoglobulin protein (BiP), protein disulfide isomerase (PDI), calnexin (Clxa), calreticulin (Crt) or glucose-regulated proteins (Grp78), and numerous other proteins, including some involved in cell wall biosynthesis (Krishnan and Askew 2014; Lamothe et al. 2015; Zhang et al. 2017a, b; Rocha et al. 2020b). These proteins are involved in protein secretion as part of the unfolded protein response (UPR), which is initiated when the burden becomes too high to manage cellular homeostasis (reviewed by Heimel 2015; Cairns et al. 2019b). The secretory pathway appears to undergo almost complete restructuring in industrial strains to enhance the cellular architecture for efficient high-volume protein secretion. The interaction between the UPR and CWI pathways was reviewed by Malavazi et al. (2014) and is well discussed in further literature (Harting and Heimel 2020; Rocha et al. 2020a, 2020b). It has been mentioned that many cell wall components (proteins and most polysaccharides) depend on the ER machinery, such that cell wall damage also leads to ER stress and can activate the UPR response in fungi. The loss of UPR function causes cell wall defects in *S. cerevisiae* (Scrimale et al. 2009), and the same is likely to occur in filamentous fungi (Malavazi et al. 2014). This strong and bidirectional dependency of both pathways, UPR and CWI, clearly points to the cell wall, its integrity and composition being influenced by high-secretion stress. Furthermore, a changed cell wall could be a major prerequisite of fungi to

enable UPR. Thus, studies on how the composition and/or crosslinking of the cell wall differs between UPR-stressed and non-stressed strains would be an interesting and relevant topic for future study.

In summary, providing a high secretion capacity for proteins in filamentous fungi demands a complex interaction from the ER machinery with adapted exo- and endocytosis, the CWI pathway maintaining cell walls, and the UPR ensuring proper folding. Thus, impaired responses may lead to cell clustering and reduced viability, which affect fermentation productivity. Consequently, these data suggest that understanding CWI and UPR coordination is essential for improving fungal protein secretion.

Valorization of fungal biomass from biotechnological fermentation

We have already mentioned the enormous amounts of mycelia that accrue as by-products of fungal fermentations. Cell walls account for 20–40% of the dry weight of fungal mycelia (Lamon et al. 2023). Even if our estimated fungal fermentation volumes (Table 1) are partially derived from data with some uncertainties, our calculations still indicate that the estimated annual production of 67 million cubic meters would result in a conservatively estimated yield of six million tons of dry mycelium or two million tons of cell walls. If an average of 10% of this is chitin, which can be converted into more precious chitosans with a yield of approximately 75%, this would amount to 150,000 tons of chitosan annually. With a price tag of approximately 50 €/kg for reasonable quality chitosan, this would be a staggering 7.5 billion € – for the chitin alone! Most spent fungal biomass is used as low-value animal feed or even dumped onto fields as fertilizer, costing the producers a fee. Clearly, there are better uses for precious fungal biopolymers such as chitin and beta-glucans. Corresponding life cycle assessment studies would be very helpful and recommended as soon as the first solid data on resources and costs are available.

Fungal cell walls as feedstock for the biotechnological production of biomolecules

A straightforward option for valorizing fungal cell walls from biotechnological fermentation wastes is to use them as feedstock for producing other commodities in bacteria or fungi. Yeast biomass, for example, a byproduct of beer brewing with *S. cerevisiae*, can be processed into yeast extracts that serve as food or feed additives and N-sources for microbial fermentation (Reed and Nagodawithana 1990; Jacob et al. 2019; Tao et al. 2023). In 1990, Reed and Nagodawithana reported that yeast cell wall mutants

with altered morphologies were used by yeast extract manufacturers because they are highly sensitive to autolytic enzymes, but are also unstable, slow growing, and lyse even during growth. A comparable approach using filamentous fungal biomass as feedstock for microbiology is conceivable but has not been visibly discussed in the scientific literature. Yet, Abasian et al. (2020) showed that *Mucor rouxii* can grow and produce ethanol when the yeast extract in the medium is replaced by a fungal extract obtained by autolysis.

A relevant difference between yeast and filamentous fungi in this respect is the protein content of their biomass. Yeast extracts contain more than 60% protein, which provides accessible nitrogen for microorganisms (reviewed by Tao et al. 2023). The insoluble yeast cell wall fraction is often commercially sold as a dietary supplement under the term *glycan* (or beta-glucans) (Reed and Nagodawithana 1990). In contrast, filamentous fungal biomass after fermentation consists of a higher cell wall portion and a protein content lower than 20% (Isaza-Pérez et al. 2020), though it is noteworthy that specific filamentous fungi used for the production of single cell proteins (SCPs), also called Mycoproteins, may consist of 30–50% protein (Quorn™ as example with 45–54% protein content) (Ritala et al. 2017; Isaza-Pérez et al. 2020). Apart from its use as a nitrogen source, fungal extracts thus could be of interest as inexpensive C-sources (feedstock) for fermentation. It has already been shown that non-competitive bacterial consortia can be constructed to grow on chitin in pretreated fungal biomass and produce lysine (Vortmann et al. 2021). Depolymerization of glucans would even yield glucose as a more universal carbon source. In this respect beta-glucanase containing enzyme cocktails for yeast treatments like Glucanex®, VinoTaste® Pro, Snailase, Lyticase, Protplast F, and Yatalase are well known in the academic community of fungal genetics and biotechnology, as they are efficient for protoplast formation of filamentous fungi before transformation (De Bekker et al. 2009; Daly et al. 2017). A procedure that might enable the production of “fungal extracts” to be used in a manner similar to yeast extracts in biotech and microbiology.

In summary, we suggest that a process in which precious cell wall components (such as chitin) are first extracted and the remaining components of the fungal biomass are further processed (e.g., as feedstock for microorganisms) would be worth further elaboration. In particular, this would valorize and recycle a side-stream in the sense of a circular bioeconomy. To what extent the use of genetically modified strains as sources of the biomass/cell walls affects the potential utilization will likely not only depend on national regulations, but also on the treatment used (e.g. inactivation procedure, purification steps and/or DNA removal).

Chitin/chitosan and beta-glucans are precious biopolymers in fungal cell walls that have applications, e.g., in medicine, food, feed, and agri- or horticulture

As described above, fungal cell walls contain diverse carbohydrate polymers. The most prominent one is chitin, whereas the most abundant ones are beta-glucans. There are many other homo- and heteropolymers containing, e.g., mannose, galactose, or galactosamine residues as monomeric building blocks, but these appear to be less significant in terms of their relative mass contribution to the cell wall, and their presence or absence strongly depends on fungal species and growth conditions.

Among glucans, which account for ca. 50–60% of cell wall dry weight, beta-1,3-glucans account for 65–90% of the total glucans (Bowman and Free 2006). It seems to be the one structural element that is indispensable for fungal vitality, and this is probably the reason why inhibitors of beta-1,3-glucan synthesis are the most successful anti-fungal drugs targeting fungal cell wall biosynthesis. (Ibe and Munro 2021). At the same time, beta-1,3-glucans are among the most prominent fungal pathogen-associated molecular patterns (PAMPs), which are recognized by cell surface receptors of the immune system of animals and humans (Mata-Martínez et al. 2022) and in the non-self-surveillance system of plants (Brown and Gordon 2001; Fesel and Zuccaro 2016). In animals, humans, and plants, both small linear beta-1,3-glucans (Klarzynski et al. 2000; Palma et al. 2006) and branched beta-1,3-beta-1,6-glucan heptamers (Sharp et al. 1984; Adams et al. 2008) appear to trigger immune responses, although the structural details of the PAMP-active oligomers differ. Clearly, such PAMPs have applications as immunostimulatory compounds, triggering “trained immunity” in animals and humans (Quintin et al. 2012) and defense priming in plants (Flors et al. 2024). In particular, branched beta-1,3-beta-1,6-glucan oligomers may offer promising options to support plant/animal/human vaccination strategies. In fact, “Zymosan,” a crude cell wall preparation from *S. cerevisiae* rich in beta-glucans, can be considered the role model for adjuvants in different types of vaccines (Fitzpatrick and DiCarlo 1964; Nihei et al. 2023; Chin et al. 2025), and similar preparations, such as “Cerevisan,” are available to induce disease resistance in crop plants. However, such preparations contain, among other ingredients, a mixture of beta-glucans of different structures that induce different aspects of the immune system (Jin et al. 2018; Rainer et al. 2024). Structurally, slightly different beta-1,3-beta-1,6-glucans from filamentous fungi have been shown to act as either agonists or antagonists of their receptor, Dectin 1, in mouse and human cells (Smith et al. 2018). Similarly, of the hundreds of beta-1,3-beta-1,6-glucan

oligomers, very few are elicitor-active in plants (Shibuya and Minami 2001; Ménard et al. 2004), and their activity varies dramatically with their molecular fine structure. We believe that this level of structural and functional detail for fungal beta-glucans, which far exceeds that from other sources, such as algae or bacteria, offers promising options for further optimizing the value chain from waste mycelial cell walls to functional biopolymers, as discussed below.

Besides glucans, chitin is also a PAMP that triggers immune reactions in humans, animals, and plants (Kaku et al. 2006; Bueter et al. 2013; Hembach et al. 2020). Structurally, chitin is simpler than glucan because it is a linear polymer of beta-1,4-linked *N*-acetylglucosamine residues, some of which may be deacetylated. Currently, most commercially produced chitin is converted into monomeric glucosamine (as the acetyl group is typically lost during chemical depolymerization) for use as an anti-arthritis food and feed supplement (D'Altilio et al. 2007; Conrozier and Lohse 2022), but some are partially deacetylated in concentrated hot alkali to yield chitosans (Younes and Rinaudo 2015). Chitosans are among the most versatile and promising functional biopolymers, which, depending on purity, can fetch prices of up to thousands of euros per kilogram (European Commission 2019; Huq et al. 2022). Clearly, this is an interesting and arguably the most interesting option for the valorization of mycelia waste from fungal fermentation.

However, can chitosan production from fungal chitin complement or compete with conventional chitosan production from crustacean waste? In terms of scale, we estimated that fungal fermentation waste could yield approximately 150,000 tons of chitosan annually. According to the United Nations Food and Agriculture Organization (FAO 2024), approximately 20 million tons of crustaceans were caught in 2022, which could theoretically yield perhaps 2 million tons of chitosan; however taking into account yields for the extraction and conversion steps (Muñoz et al. 2018), 500,000 tons is a more realistic number. While this is more than the estimated potential of fermentation mycelia, the actual global chitosan production from crustacean waste is probably less than ten percent of this potential, as logistics are difficult (Muñoz et al. 2018; Crognale et al. 2022). In addition, sustainability issues and the ecological footprint of chitosan production from crustacean waste have raised increasing concerns (Sebastian et al. 2020; Santos et al. 2020). This is where the sterile mycelial waste from large-scale fermentation with much less batch-to-batch variability may score points. Fungal chitosan is vegan, as compared to the animal origin of conventional chitosans, adding another plus in its favor for sensitive market sectors, such as cosmetics and food additives.

Another point worth mentioning is that true fungal chitosans – as opposed to chitosans produced by chemical deacetylation of fungal chitin – stand out. Some fungi contain

natural chitosan in their cell walls, most prominently mucromycete fungi, to which the industrially exploited *Rhizomucor* sp. and *Rhizopus* sp. belong (Table 1). These fungi enzymatically produce chitosan from chitin using chitin deacetylases (Jeraj et al. 2006; Gauthier et al. 2008; Lindner et al. 2025). To the best of our knowledge, it would seem that this natural chitosan differs in its acetylation pattern from chemically produced chitosans, which may make it unique in both structure and function (Cord-Landwehr and Moerschbacher, unpublished). In addition, the biotechnological production of natural chitosans may be more sustainable than conventional chemical chitosan production from chitin, but this can only be established through a thorough and comprehensive life cycle assessment (Muñoz et al. 2018).

There are many possible applications for chitosans, as mentioned in a plethora of reviews (Shariatnia 2019; Tang et al. 2023; Riseh et al. 2023, just to cite a few recent ones). Therefore, we will not go into detail here. Briefly, some chitosan-based products rely on the superior material properties of the biopolymers, such as their dye- and protein-binding and heavy metal-complexing abilities exploited for waste and drinking water purification or the finishing of papers and textiles (Pal et al. 2021). Here, purity demands – at least in terms of the absence of other biopolymers – may be less stringent than for products based on the biological functionalities of chitosans, such as their antimicrobial, plant-strengthening, or wound-healing activities (Jones et al. 2020; Eickelpasch et al. 2025). However, the latter products are potentially more attractive because chitosans can fetch better prices in these markets. The main problem with such products is that the bioactivities of chitosans critically depend on their molecular fine structure, most importantly their molecular weight and fraction of acetylation, and, as recently revealed, their acetylation pattern; however, structure–function relationships are still only partially understood. Recent advances in structural analysis and industrial production processes of chitosans have led to structurally and functionally well-characterized second-generation or 2G-chitosans with defined molecular weight and fraction of acetylation (Younes and Rinaudo 2015; Cord-Landwehr et al. 2019; Wattjes et al. 2020; Hellmann et al. 2024). These are currently reviving the development of reliable chitosan-based products, such as biostimulants or biopesticides for agriculture or as wound dressings and drug, gene, or vaccine delivery systems in biomedicine. Current research is aimed at developing biotechnological processes for even better-characterized 3G-chitosans with not only tightly controlled molecular weight and fraction of acetylation, but also with defined, non-random acetylation patterns (Basa et al. 2020; Cord-Landwehr et al. 2020; Wattjes et al. 2020; Sreeku-mar et al. 2022; Singh et al. 2023; Hellmann et al. 2024; Richter et al. 2025). These are expected to offer even wider options for the development of chitosan-based products and

applications. Clearly, with an optimistic outlook on future developments, the structural complexity of chitosans can become an asset rather than a problem.

This is where we see a similarity between chitosans and beta-glucans: Their structural complexity is the basis of their functional diversity. Once we understand better the structure–activity relationships of these precious biopolymers from fungal cell walls, we will be able to fine-tune their biological functionalities, producing the “next-generation” designer chitosans and beta-glucans specifically geared towards different uses and different markets. It is a challenge – but one well worth all efforts.

The complexity of cell walls challenges and limits the valorization of filamentous fungal spent mycelia

Industrial-scale fungal chitin and glucan production is of economic interest because of its advantages such as year-round availability, controlled cultivation, and – at least presumed – reduced allergenic risks compared to crustacean-derived chitin. However, the main drawback of fungal chitosan is that chitin is covalently linked to glucans in the fungal cell wall, and potentially also to other components, such as proteins and melanin. Thus, extraction is difficult and purity is a problem. From scientific as well as patent literature, however, processes that show feasible extraction methods, such as alkali or acid treatment of fungal biomass, have been described (Dhillon et al. 2013; Islam et al. 2023). Patents from Cargill, Kitozyme SA, and others show that some industry sectors already have valorization of biomass from industrial fermented filamentous fungi on the horizon (Versali et al. 2002; Anderson et al. 2006; Banner et al. 2011; van Leeuwen et al. 2016).

While many publications report chitin extraction from fungal biomass (Fontaine et al. 2000; Heux et al. 2000; Wu et al. 2005), only some mention its purity. If at all, many studies have indicated the presence of glucose moieties in chitin or chitosan preparations, sometimes up to 50% of the dry weight. Clearly, this is not pure chitin or chitosan but rather a chitin/chitosan-glucan complex (Nwe and Stevens 2002). Depending on the use case, this does not have to be a problem (e.g., for applications relying on the polycationic nature of chitosan which is not impaired by the presence of glucans), but most of the time it will be (namely whenever the precise molecular characteristics of the chitin/chitosan need to be known to ensure reliable performance). However, fungal chitosans are increasingly appearing on the market, and some are surprisingly pure (Cord-Landwehr and Moerschbacher, unpublished).

Yet, to the best of our knowledge, attempts to increase the extractability of fungal biomass by cell wall mutations have not yet been described for filamentous fungi. It would be obvious to combine the advantages of fermentability, such

as changes in morphology, improved fermentation performance, or reduced viscosity, as described for some cell wall mutants in the previous sections, with likely changes in the extractability of chitin/chitosan or beta-glucans. In addition to higher product yields and easier downstream processing, a more porous or loosely packed mycelial network could enhance the accessibility of chemical or enzymatic treatments used in the extraction of chitin and glucans from the biomass. One would assume that, in particular, modified crosslinks between chitin, glucans, and proteins in the fungal cell wall would likely enhance the efficiency of chitin and glucan extraction. Similarly, mutations that reduce the activity of chitin synthases or modify the acetylation pattern of chitin could also influence its solubility and extraction efficiency.

Overall, the use of cell wall mutants provides a promising strategy for improving the valorization of fungal biomass by making the extraction of key biopolymers more efficient and cost-effective. By integrating this approach into industrial fermentation processes for enzyme, protein, or metabolite production, fungal biomass will no longer be a mere byproduct, but a valuable resource for high-value biopolymer production. This circular bioeconomy model ensures that no fungal waste is discarded, thus reducing disposal costs and contributing to sustainable biomanufacturing in industrial biotechnology.

Conclusion

Our literature review and analysis clearly highlight the relevance of cell walls in the biotechnological applications of filamentous fungi. Increasing knowledge of the composition, biosynthesis, and regulation of cell walls is continuously offering new possibilities to develop the cell wall machinery into a powerful set screw to optimize the biotechnological production of enzymes and metabolites. Cell walls play crucial roles in the transfer of proteins to the supernatant of fermentation media and determine fungal macromorphology, both of which have huge impacts on yield and productivity. A detailed understanding of these processes will allow for the design of targeted cell wall modifications by genetic modification and bioprocess engineering. However, the extent and limits of the influence of cell walls on the biotechnological productivity of filamentous fungal systems are yet to be elucidated.

Similarly, cell wall alterations by process and strain engineering offer huge potential for the improvement of valorization routes for fungal biomass as a side-stream of biotechnological production from precision fermentation. The ultimate challenge will be to combine both targets and improve efficiency in both production and valorization.

Acknowledgements We gratefully acknowledge Ludwig Klermund and Peter J. Punt for helpful discussions and feedback

Author contribution KJO contributed to the conception of the manuscript, performed the literature study, drew Fig. 1, drafted, and wrote the manuscript with a focus on Sections 2 (The fungal cell wall) and 3 (How fungal cell wall structure affects industrial enzyme production); MS and SCL performed the literature study with a focus on Section 4 (Valorization of fungal biomass from biotechnological fermentation); BMM contributed to the conception, wrote the manuscript with a focus on Sections 2 and 4, and critically revised the work; AFJR contributed to the conception of the manuscript and critically revised the work; MEDH contributed to the conception, performed the literature study, and created Table 1, contributed to writing with a focus on Sections 1 (Introduction) and 3, and critically revised the manuscript. All the authors have read and approved the final version of the manuscript.

Funding Open Access funding enabled and organized by Projekt DEAL. This research did not receive any specific grants from funding agencies in the public, commercial, or not-for-profit sectors. KJO's position at the Osnabrück University of Applied Sciences was financed by funds from the Lower Saxony Ministry for Science and Culture (MWK) as part of the "Professorinnen für Niedersachsen" research program granted to MEDH.

Data availability The data supporting the findings of this study are available as cited in references. Unpublished data that are used as examples are available from the corresponding authors upon request.

Declarations

Ethics approval This study adhered to ethical standards and did not involve any human or animal studies conducted by the authors.

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abasian L, Shafei Alavijeh R, Satari B, Karimi K (2020) Sustainable and effective chitosan production by dimorphic fungus *Mucor rouxii* via replacing yeast extract with fungal extract. *Appl Biochem Biotechnol* 191:666–678. <https://doi.org/10.1007/s12010-019-03220-w>
- Adams EL, Rice PJ, Graves B, Ensley HE, Yu H, Brown GD, Gordon S, Monteiro MA, Papp-Szabo E, Lowman DW, Power TD, Wempe MF, Williams DL (2008) Differential high-affinity interaction of Dectin-1 with natural or synthetic glucans is dependent upon primary structure and is influenced by polymer chain length and side-chain branching. *J Pharmacol Exp Ther* 325:115–123. <https://doi.org/10.1124/jpet.107.133124>
- Allied Market Research (2024) Amylase market size, share, forecast report, 2033. <https://www.alliedmarketresearch.com/amilase-market-A15505>. Accessed 28 Jan 2025
- Amanullah A, Christensen LH, Hansen K, Nienow AW, Thomas CR (2002) Dependence of morphology on agitation intensity in fed-batch cultures of *Aspergillus oryzae* and its implications for recombinant protein production. *Biotechnol Bioeng* 77:815–826. <https://doi.org/10.1002/bit.10181>
- Anderson T, Zhou S, Bergerud B, Fosdick L, Han T, Vasina J, Vanderhoff A, Trinkle J, Ray B, Bohlmann J (2006) Glucosamine and *N*-acetylglucosamine compositions and methods of making the same fungal biomass. (Patent No. US 8222232 B2) 31 March 2006. <https://worldwide.espacenet.com/patent/search?q=pn%3DUS8222232B2>
- Aschenbroich J, Hussnaetter KP, Stoffels P, Langner T, Zander S, Sandrock B, Bölker M, Feldbrügge M, Schipper K (2019) The germinal centre kinase Don3 is crucial for unconventional secretion of chitinase Cts1 in *Ustilago maydis*. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* 1867:140154. <https://doi.org/10.1016/j.bbapap.2018.10.007>
- Astute Analytica (2024) Enzymes market - industry dynamics, market size, and opportunity forecast to 2035. <https://www.astuteanalytica.com/industry-report/enzymes-market>. Accessed 30 Jan 2025
- Balakrishnan M, Jeevarathinam G, Kumar SKS, Muniraj I, Uthandi S (2021) Optimization and scale-up of α -amylase production by *Aspergillus oryzae* using solid-state fermentation of edible oil cakes. *BMC Biotechnol* 21:33. <https://doi.org/10.1186/s12896-021-00686-7>
- Banner T, Fosmer A, Jessen H, Marasco E, Rush B, Veldhouse J, de Souza M (2011) Microbial bioprocesses for industrial-scale chemical production. In: *Biocatalysis for green chemistry and chemical process development*. Wiley, pp 429–467. <https://doi.org/10.1002/9781118028308.ch16>
- Barata-Antunes C, Alves R, Talaia G, Casal M, Gerós H, Mans R, Paiva S (2021) Endocytosis of nutrient transporters in fungi: the ART of connecting signaling and trafficking. *Comput Struct Biotechnol J* 19:1713–1737
- Barthel L, Cairns T, Duda S, Müller H, Dobbert B, Jung S, Briesen H, Meyer V (2024) Breaking down barriers: comprehensive functional analysis of the *Aspergillus niger* chitin synthase repertoire. *Fungal Biol Biotechnol* 11:3. <https://doi.org/10.1186/s40694-024-00172-7>
- Basa S, Nampally M, Honorato T, Das SN, Podile AR, El Gueddari NE, Moerschbacher BM (2020) The pattern of acetylation defines the priming activity of chitosan tetramers. *J Am Chem Soc* 142:1975–1986. <https://doi.org/10.1021/jacs.9b11466>
- Beauvais A, Bruneau JM, Mol PC, Buitrago MJ, Legrand R, Latgé J-P (2001) Glucan synthase complex of *Aspergillus fumigatus*. *J Bacteriol* 183:2273–2279. <https://doi.org/10.1128/JB.183.7.2273-2279.2001>
- Beauvais A, Fontaine T, Aïmanianda V, Latgé J-P (2014) *Aspergillus* cell wall and biofilm. *Mycopathologia* 178:371–377. <https://doi.org/10.1007/s11046-014-9766-0>
- Beauvais A, Latgé J-P (2015) *Aspergillus* biofilm in vitro and in vivo. *Microbiol Spectr* 3. <https://doi.org/10.1128/microbiolspec.mb-0017-2015>
- Benoit I, van den Esker MH, Patyshakuliyeva A, Mattern DJ, Blei F, Zhou M, Dijksterhuis J, Brakhage AA, Kuipers OP, de Vries RP, Kovács ÁT (2015) *Bacillus subtilis* attachment to *Aspergillus niger* hyphae results in mutually altered metabolism. *Environ Microbiol* 17:2099–2113. <https://doi.org/10.1111/1462-2920.12564>

- Blatzer M, Beauvais A, Henrissat B, Latgé J-P (2020) Revisiting old questions and new approaches to investigate the fungal cell wall construction. In: Latgé J-P (ed) The fungal cell wall. Springer. https://doi.org/10.1007/82_2020_209
- Bodie E, Virag A, Pratt RJ, Leiva N, Ward M, Dodge T (2021) Reduced viscosity mutants of *Trichoderma reesei* with improved industrial fermentation characteristics. J Ind Microbiol Biotechnol 48. <https://doi.org/10.1093/jimb/kuab014>
- Boppidi KR, Ribeiro LFC, Iambamrungs S, Nelson SM, Wang Y, Momany M, Richardson EA, Lincoln S, Srivastava R, Harris SD, Marten MR (2018) Altered secretion patterns and cell wall organization caused by loss of PodB function in the filamentous fungus *Aspergillus nidulans*. Sci Rep 8:11433. <https://doi.org/10.1038/s41598-018-29615-z>
- Bowman SM, Free SJ (2006) The structure and synthesis of the fungal cell wall. BioEssays 28:799–808
- Brauer VS, Pessoni AM, Freitas MS, Cavalcanti-Neto MP, Ries LNA, Almeida F (2023) Chitin biosynthesis in *Aspergillus* species. Journal of Fungi 9:89. <https://doi.org/10.3390/jof9010089>
- Brown GD, Gordon S (2001) A new receptor for β -glucans. Nature 413:36–37. <https://doi.org/10.1038/35092620>
- Brown HE, Esher SK, Alspaugh JA (2019) Chitin: a “hidden figure” in the fungal cell wall. In: Latgé J-P (ed) The fungal cell wall - an armour and weapon for human fungal pathogens. Springer, pp 83–111
- Bueter CL, Specht CA, Levitz SM (2013) Innate sensing of chitin and chitosan. PLoS Pathog 9:e1003080. <https://doi.org/10.1371/journal.ppat.1003080>
- Burggraaf A-M, Punt PJ, Ram AFJ (2016) The unconventional secretion of PepN is independent of a functional autophagy machinery in the filamentous fungus *Aspergillus niger*. FEMS Microbiol Lett 363:fnw152. <https://doi.org/10.1093/femsle/fnw152>
- Cai J, Yang J, Du Y, Fan L, Qiu Y, Li J, Kennedy JF (2006) Enzymatic preparation of chitosan from the waste *Aspergillus niger* mycelium of citric acid production plant. Carbohydr Polym 64:151–157. <https://doi.org/10.1016/j.carbpol.2005.11.004>
- Cairns TC, Nai C, Meyer V (2018) How a fungus shapes biotechnology: 100 years of *Aspergillus niger* research. Fungal Biol Biotechnol 5:13. <https://doi.org/10.1186/s40694-018-0054-5>
- Cairns TC, Feurstein C, Zheng X, Zhang LH, Zheng P, Sun J, Meyer V (2019) Functional exploration of co-expression networks identifies a nexus for modulating protein and citric acid titres in *Aspergillus niger* submerged culture. Fungal Biol Biotechnol 6:18. <https://doi.org/10.1186/s40694-019-0081-x>
- Cairns TC, Zheng X, Zheng P, Sun J, Meyer V (2019) Moulding the mould: understanding and reprogramming filamentous fungal growth and morphogenesis for next generation cell factories. Biotechnol Biofuels 12:77. <https://doi.org/10.1186/s13068-019-1400-4>
- Cairns TC, Barthel L, Meyer V (2021) Something old, something new: challenges and developments in *Aspergillus niger* biotechnology. Essays Biochem 65:213–224. <https://doi.org/10.1042/EBC20200139>
- Cairns TC, de Kanter T, Zheng XZ, Zheng P, Sun J, Meyer V (2023) Regression modelling of conditional morphogene expression links and quantifies the impact of growth rate, fitness and macromorphology with protein secretion in *Aspergillus niger*. Biotechnol Biofuels Bioprod 16:95. <https://doi.org/10.1186/s13068-023-02345-9>
- Cairns TC, Zheng X, Zheng P, Sun J, Meyer V (2021b) Turning inside out: filamentous fungal secretion and its applications in biotechnology, agriculture, and the clinic. Journal of Fungi 7. <https://doi.org/10.3390/jof7070535>
- Caldwell IY, Trinci APJ (1973) The growth unit of the mould *Geotrichum candidum*. Arch Mikrobiol 88:1–10. <https://doi.org/10.1007/BF00408836>
- Casadevall A, Nosanchuk JD, Williamson P, Rodrigues ML (2009) Vesicular transport across the fungal cell wall. Trends Microbiol 17:158–162. <https://doi.org/10.1016/j.tim.2008.12.005>
- Chakraborty A, Fernando LD, Fang W, Dickwella Widanage MC, Wei P, Jin C, Fontaine T, Latgé J-P, Wang T (2021) A molecular vision of fungal cell wall organization by functional genomics and solid-state NMR. Nat Commun 12. <https://doi.org/10.1038/s41467-021-26749-z>
- Cheng Q, Dickwella Widanage MC, Yarava JR, Ankur A, Latgé J-P, Wang P, Wang T (2024) Molecular architecture of chitin and chitosan-dominated cell walls in zygomycetous fungal pathogens by solid-state NMR. Nat Commun 15:8295. <https://doi.org/10.1038/s41467-024-52759-8>
- Chevalier L, Pinar M, Le Borgne R, Durieu C, Peñalva MA, Boudaoud A, Minc N (2023) Cell wall dynamics stabilize tip growth in a filamentous fungus. PLoS Biol 21:e3001981. <https://doi.org/10.1371/journal.pbio.3001981>
- Chin C, Lin Y, Cheng P, Lee P, Chiang B (2025) Effective mucosal adjuvantation of the intranasal enterovirus A71 vaccine with zymosan. Immunology 174:349–362. <https://doi.org/10.1111/imm.13895>
- Commer B, Shaw BD (2021) Current views on endocytosis in filamentous fungi. Mycology 12:1–9. <https://doi.org/10.1080/21501203.2020.1741471>
- Conrozier T, Lohse T (2022) Glucosamine as a treatment for osteoarthritis: what if it's true? Front Pharmacol 13. <https://doi.org/10.3389/fphar.2022.820971>
- Cord-Landwehr S, Richter C, Wattjes J, Sreekumar S, Singh R, Basa S, El Gueddari NE, Moerschbacher BM (2020) Patterns matter part 2: chitosan oligomers with defined patterns of acetylation. React Funct Polym 151:104577. <https://doi.org/10.1016/j.reactfunctpolym.2020.104577>
- Cord-Landwehr S, Niehues A, Wattjes J, Moerschbacher BM (2019) New developments in the analysis of partially acetylated chitosan polymers and oligomers. In: Chitin and Chitosan. Wiley, pp 81–95. <https://doi.org/10.1002/9781119450467.ch4>
- Cosgrove DJ (2024) Structure and growth of plant cell walls. Nat Rev Mol Cell Biol 25:340–358. <https://doi.org/10.1038/s41580-023-00691-y>
- Crognale S, Russo C, Petruccioli M, D'Annibale A (2022) Chitosan production by fungi: current state of knowledge, future opportunities and constraints. Fermentation 8:76. <https://doi.org/10.3390/fermentation8020076>
- Crosino A, Moscato E, Blangetti M, Carotenuto G, Spina F, Bordignon S, Puech-Pagès V, Anfossi L, Volpe V, Prandi C, Gobetto R, Varese GC, Genre A (2021) Extraction of short chain chitooligosaccharides from fungal biomass and their use as promoters of arbuscular mycorrhizal symbiosis. Sci Rep 11. <https://doi.org/10.1038/s41598-021-83299-6>
- D'Altilio M, Peal A, Alvey M, Simms C, Curtsinger A, Gupta RC, Canerdy TD, Goad JT, Bagchi M, Bagchi D (2007) Therapeutic efficacy and safety of undenatured type II collagen singly or in combination with glucosamine and chondroitin in arthritic dogs. Toxicol Mech Methods 17:189–196. <https://doi.org/10.1080/15376510600910469>
- Dalonso N, Petkowicz C, Lugones L, Silveira M, Gern R (2021) Comparison of cell wall polysaccharides in *Schizophyllum commune* after changing phenotype by mutation. An Acad Bras Cienc 93. <https://doi.org/10.1590/0001-3765202120210047>
- Daly P, Slaghek GG, Casado López S, Wiebenga A, Hilden KS, de Vries RP, Mäkelä MR (2017) Genetic transformation of the white-rot fungus *Dichomitus squalens* using a new commercial protoplasting cocktail. J Microbiol Methods 143:38–43. <https://doi.org/10.1016/j.mimet.2017.10.001>
- Damveld RA, Arentshorst M, Franken A, VanKuyk PA, Klis FM, Van Den Hondel CAMJJ, Ram AFJ (2005) The *Aspergillus niger*

- MADS-box transcription factor RlmA is required for cell wall reinforcement in response to cell wall stress. *Mol Microbiol* 58:305–319. <https://doi.org/10.1111/j.1365-2958.2005.04827.x>
- Das KC, Minkara MY, Melear ND (2002) Composting process for the stabilization of an industrial fermentation biomass byproduct. *Transact ASAE* 45:1703–1708. <https://doi.org/10.13031/2013.11033>
- de Bekker C, Wiebenga A, Aguilar G, Wösten HAB (2009) An enzyme cocktail for efficient protoplast formation in *Aspergillus niger*. *J Microbiol Methods* 76:305–306. <https://doi.org/10.1016/j.mimet.2008.11.001>
- de Groot PWJ, Brandt BW, Horiuchi H, Ram AFJ, de Koster CG, Klis FM (2009) Comprehensive genomic analysis of cell wall genes in *Aspergillus nidulans*. *Fungal Genet Biol* 46:S72–S81. <https://doi.org/10.1016/j.fgb.2008.07.022>
- de Groot PWJ, Bader O, de Boer AD, Weig M, Chauhan N (2013) Adhesins in human fungal pathogens: glue with plenty of stick. *Eukaryot Cell* 12:470–481. <https://doi.org/10.1128/EC.00364-12>
- de Paula RG, Antoniêto ACC, Nogueira KMV, Ribeiro LFC, Rocha MC, Malavazi I, Almeida F, Silva RN (2019) Extracellular vesicles carry cellulases in the industrial fungus *Trichoderma reesei*. *Biotechnol Biofuels* 12:146. <https://doi.org/10.1186/s13068-019-1487-7>
- De Nobel JG, Barnett JA (1991) Passage of molecules through yeast cell walls: a brief essay-review. *Yeast* 7:313–323. <https://doi.org/10.1002/yea.320070402>
- De Souza PM, De Oliveira e Magalhães P (2010) Application of microbial α -amylase in industry - a review. *Braz J Microbiol* 41:850–861. <https://doi.org/10.1590/S1517-83822010000400004>
- De Brabander P, Uitterhaegen E, Delmulle T, De Winter K, Soetaert W (2023) Challenges and progress towards industrial recombinant protein production in yeasts: a review. *Biotechnol Adv* 64:108121. <https://doi.org/10.1016/j.biotechadv.2023.108121>
- De Nobel JG, Dijkers C, Hooijberg E, Klis FM (1989) Increased cell wall porosity in *Saccharomyces cerevisiae* after treatment with dithiothreitol or EDTA. *Microbiology (n y)* 135:2077–2084. <https://doi.org/10.1099/00221287-135-7-2077>
- De Nobel JG, Klis FM, Munnik T, Priem J, Van Den Ende H (1990) An assay of relative cell wall porosity in *Saccharomyces cerevisiae*, *Kluyveromyces lactis* and *Schizosaccharomyces pombe*. *Yeast* 6:483–490. <https://doi.org/10.1002/yea.320060605>
- De Nobel JG, Klis FM, Priem J, Munnik T, Van Den Ende H (1990) The glucanase-soluble mannoproteins limit cell wall porosity in *Saccharomyces cerevisiae*. *Yeast* 6:491–499. <https://doi.org/10.1002/yea.320060606>
- Demain AL, Adrio JL (2008) Contributions of microorganisms to industrial biology. *Mol Biotechnol* 38:41–55. <https://doi.org/10.1007/s12033-007-0035-z>
- Demain AL (2006) From natural products discovery to commercialization: a success story. In: *J Ind Microbiol Biotechnol* 486–495. <https://doi.org/10.1007/s12095-005-0076-x>
- Dhillon GS, Kaur S, Brar SK, Verma M (2013) Green synthesis approach: extraction of chitosan from fungus mycelia. *Crit Rev Biotechnol* 33:379–403. <https://doi.org/10.3109/07388551.2012.717217>
- Dichtl K, Helmschrott C, Dirr F, Wagener J (2012) Deciphering cell wall integrity signalling in *Aspergillus fumigatus*: identification and functional characterization of cell wall stress sensors and relevant Rho GTPases. *Mol Microbiol* 83:506–519. <https://doi.org/10.1111/j.1365-2958.2011.07946.x>
- Dichtl K, Samantaray S, Wagener J (2016) Cell wall integrity signalling in human pathogenic fungi. *Cell Microbiol* 18:1228–1238. <https://doi.org/10.1111/cmi.12612>
- Driouch H, Roth A, Dersch P, Wittmann C (2011) Filamentous fungi in good shape: microparticles for tailor-made fungal morphology and enhanced enzyme production. *Bioeng Bugs* 2:100–104. <https://doi.org/10.4161/bbug.2.2.13757>
- Driouch H, Hänsch R, Wucherpfennig T, Krull R, Wittmann C (2012) Improved enzyme production by bio-pellets of *Aspergillus niger*: targeted morphology engineering using titanate microparticles. *Biotechnol Bioeng* 109:462–471. <https://doi.org/10.1002/bit.23313>
- Ehren HL, Appels FVW, Houben K, Renault MAM, Wösten HAB, Baldus M (2020) Characterization of the cell wall of a mushroom forming fungus at atomic resolution using solid-state NMR spectroscopy. *The Cell Surface* 6. <https://doi.org/10.1016/j.tcsu.2020.100046>
- Eickelpasch K, Lemke P, Sreekumar S, Chilukoti N, Moerschbacher BM, Richter C (2025) A bioactivity matrix for antimicrobial activities of chitosans: a review. *Int J Biol Macromol* 299:140740. <https://doi.org/10.1016/j.ijbiomac.2025.140740>
- El Enshasy HA (2022) Fungal morphology: a challenge in bioprocess engineering industries for product development. *Curr Opin Chem Eng* 35:100729. <https://doi.org/10.1016/j.coche.2021.100729>
- Erwig LP, Gow NAR (2016) Interactions of fungal pathogens with phagocytes. *Nat Rev Microbiol* 14:163–176
- European Commission: Directorate-general for research and innovation, Top 20 innovative bio-based products – Task 3 of “study on support to R&I policy in the area of bio-based products and services”, publications office, 2019. <https://doi.org/10.2777/85805>
- Eveleigh DE, Montencourt BS (1979) Increasing yields of extracellular enzymes. *Adv Appl Microbiol* 25:57–74. [https://doi.org/10.1016/S0065-2164\(08\)70146-1](https://doi.org/10.1016/S0065-2164(08)70146-1)
- FAO (2024) The state of world fisheries and aquaculture 2024. In *The state of world fisheries and aquaculture 2024*. FAO. <https://doi.org/10.4060/cd0683en>
- Fernando LD, Zhao W, Gautam I, Ankur A, Wang T (2023) Polysaccharide assemblies in fungal and plant cell walls explored by solid-state NMR. *Structure* 31:1375–1385. <https://doi.org/10.1016/j.str.2023.07.012>
- Fesel PH, Zuccaro A (2016) β -glucan: crucial component of the fungal cell wall and elusive MAMP in plants. *Fungal Genet Biol* 90:53–60. <https://doi.org/10.1016/j.fgb.2015.12.004>
- Fiedler MRM, Barthel L, Kubisch C, Nai C, Meyer V (2018) Construction of an improved *Aspergillus niger* platform for enhanced glucoamylase secretion. *Microb Cell Fact* 17:95. <https://doi.org/10.1186/s12934-018-0941-8>
- Fiedler MRM, Cairns TC, Koch O, Kubisch C, Meyer V (2018b) Conditional expression of the small GTPase ArfA impacts secretion, morphology, growth, and actin ring position in *Aspergillus niger*. *Front Microbiol* 9. <https://doi.org/10.3389/fmicb.2018.00878>
- Fizpatrick FW, DiCarlo FJ (1964) ZYMOSAN. *Ann N Y Acad Sci* 118:235–261. <https://doi.org/10.1111/j.1749-6632.1964.tb33982.x>
- Flors V, Kyndt T, Mauch-Mani B, Pozo MJ, Ryu C-M, Ton J (2024) Enabling sustainable crop protection with induced resistance in plants. *Frontiers in Science* 2. <https://doi.org/10.3389/fsci.2024.1407410>
- Fontaine T, Latgé J-P (2020) Galactomannan produced by *Aspergillus fumigatus*: an update on the structure, biosynthesis and biological functions of an emblematic fungal biomarker. *Journal of Fungi* 6:283. <https://doi.org/10.3390/jof6040283>
- Fontaine T, Simenel C, Dubreucq G, Adam O, Delepierre M, Lemoine J, Vorgias CE, Diaquin M, Latgé J-P (2000) Molecular organization of the alkali-insoluble fraction of *Aspergillus fumigatus* cell wall. *J Biol Chem* 275:27594–27607. <https://doi.org/10.1074/jbc.M909975199>
- Free SJ (2013) Fungal cell wall organization and biosynthesis. In: Friedmann T, Dunlap J, Goodwin S (eds) *Advances in genetics*. Academic Press, pp 33–82

- Garcia-Rubio R, de Oliveira HC, Rivera J, Trevijano-Contador N (2020) The fungal cell wall: *Candida*, *Cryptococcus*, and *Aspergillus* species. *Front Microbiol* 10. <https://doi.org/10.3389/fmicb.2019.02993>
- Garde S, Chodiseti PK, Reddy M (2021) Peptidoglycan: structure, synthesis, and regulation. *EcoSal Plus* 9. <https://doi.org/10.1128/ecosalplus.ESP-0010-2020>
- Gauthier C, Clerisse F, Dommes J, Jaspars-Versali M (2008) Characterization and cloning of chitin deacetylases from *Rhizopus circinans*. *Protein Expr Purif* 59:127–137. <https://doi.org/10.1016/j.pep.2008.01.013>
- Ghassemi N, Poulhazan A, Deligey F, Mentink-Vigier F, Marcotte I, Wang T (2022) Solid-state NMR investigations of extracellular matrices and cell walls of algae, bacteria, fungi, and plants. *Chem Rev* 122:10036–10086
- Götz R, Panzer S, Trinks N, Eilts J, Wagener J, Turrà D, Di Pietro A, Sauer M, Terpitz U (2020) Expansion microscopy for cell biology analysis in fungi. *Front Microbiol* 11. <https://doi.org/10.3389/fmicb.2020.00574>
- Gow NAR, Latgé J-P, Munro CA (2017) The fungal cell wall: structure, biosynthesis, and function. *Microbiol Spectr* 5. <https://doi.org/10.1128/microbiolspec.funk-0035-2016>
- Gow NAR, Lenardon MD (2023) Architecture of the dynamic fungal cell wall. *Nat Rev Microbiol* 21:248–259. <https://doi.org/10.1038/s41579-022-00796-9>
- Gravelat FN, Beauvais A, Liu H, Lee MJ, Snarr BD, Chen D, Xu W, Kravtsov I, Hoareau CMQ, Vanier G, Urb M, Campoli P, Al Abdallah Q, Lehoux M, Chabot JC, Ouimet MC, Baptista SD, Fritz JH, Nierman WC, Latgé J-P, Mitchell AP, Filler SG, Fontaine T, Sheppard DC (2013) *Aspergillus* galactosaminogalactan mediates adherence to host constituents and conceals hyphal β -glucan from the immune system. *PLoS Pathog* 9. <https://doi.org/10.1371/journal.ppat.1003575>
- Grimm D, Wösten HAB (2018) Mushroom cultivation in the circular economy. *Appl Microbiol Biotechnol* 102:7795–7803
- Grimm LH, Kelly S, Krull R, Hempel DC (2005) Morphology and productivity of filamentous fungi. *Appl Microbiol Biotechnol* 69:375–384
- Guerrand D (2018) Economics of food and feed enzymes: status and perspectives. In: *Enzymes in human and animal nutrition: Principles and perspectives*. Elsevier Inc, pp 487–514. <https://doi.org/10.1016/B978-0-12-805419-2.00026-5>
- Harting R, Heimel K (2020) Genetics of the unfolded protein response in fungi. In: Benz JP, Schipper K (eds) *Genetics and Biotechnology. The Mycota*, vol 2. Springer, Cham., 49–76. https://doi.org/10.1007/978-3-030-49924-2_3
- Hayakawa Y, Ishikawa E, Shoji Ya J, Nakano H, Kitamoto K (2011) Septum-directed secretion in the filamentous fungus *Aspergillus oryzae*. *Mol Microbiol* 81:40–55. <https://doi.org/10.1111/j.1365-2958.2011.07700.x>
- Heimel K (2015) Unfolded protein response in filamentous fungi – implications in biotechnology. *Appl Microbiol Biotechnol* 99:121–132
- Hellmann MJ, Gillet D, Trombotto S, Raetz S, Moerschbacher BM, Cord-Landwehr S (2024) Heterogeneously deacetylated chitosans possess an unexpected regular pattern favoring acetylation at every third position. *Nat Commun* 15:6695. <https://doi.org/10.1038/s41467-024-50857-1>
- Hembach L, Bonin M, Gorzelanny C, Moerschbacher BM (2020) Unique subsite specificity and potential natural function of a chitosan deacetylase from the human pathogen *Cryptococcus neoformans*. *Proc Natl Acad Sci* 117:3551–3559. <https://doi.org/10.1073/pnas.1915798117>
- Henry C, Fontaine T, Heddergott C, Robinet P, Aïmanianda V, Beau R, Beauvais A, Mouyna I, Prevost M-C, Fekkar A, Zhao Y, Perlín D, Latgé J-P (2016) Biosynthesis of cell wall mannan in the conidium and the mycelium of *Aspergillus fumigatus*. *Cell Microbiol* 18:1881–1891. <https://doi.org/10.1111/cmi.12665>
- Henry C, Li J, Danion F, Alcazar-Fuoli L, Mellado E, Beau R, Jou-vion G, Latgé J-P, Fontaine T (2019) Two KTR mannosyltransferases are responsible for the biosynthesis of cell wall mannans and control polarized growth in *Aspergillus fumigatus*. *mBio* 10. <https://doi.org/10.1128/mBio.02647-18>
- Heux L, Brugnerotto J, Desbrières J, Versali M-F, Rinaudo M (2000) Solid state NMR for determination of degree of acetylation of chitin and chitosan. *Biomacromol* 1:746–751. <https://doi.org/10.1021/bm000070y>
- Higuchi Y, Shoji J, Arioka M, Kitamoto K (2009) Endocytosis is crucial for cell polarity and apical membrane recycling in the filamentous fungus *Aspergillus oryzae*. *Eukaryot Cell* 8:37–46. <https://doi.org/10.1128/EC.00207-08>
- Hira D, Onoue T, Oka T (2020) Structural basis for the core-mannan biosynthesis of cell wall fungal-type galactomannan in *Aspergillus fumigatus*. *J Biol Chem* 295:15407–15417. <https://doi.org/10.1074/jbc.RA120.013742>
- Hopke A, Brown AJP, Hall RA, Wheeler RT (2018) Dynamic fungal cell wall architecture in stress adaptation and immune evasion. *Trends Microbiol* 26:284–295. <https://doi.org/10.1016/j.tim.2018.01.007>
- Hu X, Yang P, Chai C, Liu J, Sun H, Wu Y, Zhang M, Zhang M, Liu X, Yu H (2023) Structural and mechanistic insights into fungal β -1,3-glucan synthase FKS1. *Nature* 616:190–198. <https://doi.org/10.1038/s41586-023-05856-5>
- Huq T, Khan A, Brown D, Dhayagude N, He Z, Ni Y (2022) Sources, production and commercial applications of fungal chitosan: a review. *J Bioresour Bioprod* 7:85–98. <https://doi.org/10.1016/j.jobab.2022.01.002>
- Ibe C, Munro CA (2021) Fungal cell wall: an underexploited target for antifungal therapies. *PLoS Pathog* 17:e1009470. <https://doi.org/10.1371/journal.ppat.1009470>
- Ichikawa H, Miyazawa K, Komeiji K, Susukida S, Zhang S, Muto K, Orita R, Takeuchi A, Kamachi Y, Hitosugi M, Yoshimi A, Shintani T, Kato Y, Abe K (2022) Improved recombinant protein production in *Aspergillus oryzae* lacking both α -1,3-glucan and galactosaminogalactan in batch culture with a lab-scale bioreactor. *J Biosci Bioeng* 133:39–45. <https://doi.org/10.1016/j.jbiosc.2021.09.010>
- Isaza-Pérez F, Ramírez-Carmona M, Rendón-Castrillón L, Ocampo-López C (2020) Potential of residual fungal biomass: a review. *Environ Sci Pollut Res* 27:13019–13031. <https://doi.org/10.1007/s11356-020-08193-6>
- Islam N, Hoque M, Taharat SF (2023) Recent advances in extraction of chitin and chitosan. *World J Microbiol Biotechnol* 39:28. <https://doi.org/10.1007/s11274-022-03468-1>
- Islam KT, Shah DM, El-Mounadi K (2017) Live-cell imaging of fungal cells to investigate modes of entry and subcellular localization of antifungal plant defensins. *Journal of Visualized Experiments* 2017. <https://doi.org/10.3791/55995>
- Jacob FF, Striegel L, Rychlik M, Hutzler M, Methner F-J (2019) Spent yeast from brewing processes: a biodiverse starting material for yeast extract production. *Fermentation* 5:51. <https://doi.org/10.3390/fermentation5020051>
- Jacobson ES, Ikeda R (2005) Effect of melanization upon porosity of the cryptococcal cell wall. *Med Mycol* 43:327–333. <https://doi.org/10.1080/13693780412331271081>
- Jadhav R, Mach RL, Mach-Aigner AR (2024) Protein secretion and associated stress in industrially employed filamentous fungi. *Appl Microbiol Biotechnol* 108:1–14
- Jeraj N, Kunič B, Lenasi H, Breskvar K (2006) Purification and molecular characterization of chitin deacetylase from *Rhizopus nigricans*. *Enzyme Microb Technol* 39:1294–1299. <https://doi.org/10.1016/j.enzmtec.2006.03.017>

- Jin Y, Li P, Wang F (2018) β -glucans as potential immunoadjuvants: a review on the adjuvanticity, structure-activity relationship and receptor recognition properties. *Vaccine* 36:5235–5244. <https://doi.org/10.1016/j.vaccine.2018.07.038>
- Jones M, Kujundzic M, John S, Bismarck A (2020) Crab vs. mushroom: a review of crustacean and fungal chitin in wound treatment. *Mar Drugs* 18:64. <https://doi.org/10.3390/md18010064>
- Kadooka C, Hira D, Tanaka Y, Chihara Y, Goto M, Oka T (2022) Mnt1, an α -(1 $\rightarrow$ 2)-mannosyltransferase responsible for the elongation of *N*-glycans and *O*-glycans in *Aspergillus fumigatus*. *Glycobiology* 32:1137–1152. <https://doi.org/10.1093/glycob/cwac049>
- Kaku H, Nishizawa Y, Ishii-Minami N, Akimoto-Tomiyama C, Dohmae N, Takio K, Minami E, Shibuya N (2006) Plant cells recognize chitin fragments for defense signaling through a plasma membrane receptor. *Proc Natl Acad Sci* 103:11086–11091. <https://doi.org/10.1073/pnas.0508882103>
- Kang X, Kirui A, Muszyński A, Widanage MCD, Chen A, Azadi P, Wang P, Mentink-Vigier F, Wang T (2018) Molecular architecture of fungal cell walls revealed by solid-state NMR. *Nat Commun* 9. <https://doi.org/10.1038/s41467-018-05199-0>
- Kisser M, Kubicek CP, Röhr M (1980) Influence of manganese on morphology and cell wall composition of *Aspergillus niger* during citric acid fermentation. *Arch Microbiol* 128:26–33. <https://doi.org/10.1007/BF00422301>
- Klarzynski O, Plesse B, Joubert J-M, Yvin J-C, Kopp M, Kloareg B, Fritig B (2000) Linear β -1,3 glucans are elicitors of defense responses in tobacco. *Plant Physiol* 124:1027–1038. <https://doi.org/10.1104/pp.124.3.1027>
- Klis FM, de Jong M, Brul S, de Groot PWJ (2007) Extraction of cell surface-associated proteins from living yeast cells. *Yeast* 24:253–258. <https://doi.org/10.1002/yea.1476>
- Kostag M, El Seoud OA (2021) Sustainable biomaterials based on cellulose, chitin and chitosan composites - a review. *Carbohydrate Polymer Technologies and Applications* 2:100079. <https://doi.org/10.1016/j.carpta.2021.100079>
- Koutinas AA, Wang R, Webb C (2003) Estimation of fungal growth in complex, heterogeneous culture. *Biochem Eng J* 14:93–100. [https://doi.org/10.1016/S1369-703X\(02\)00154-7](https://doi.org/10.1016/S1369-703X(02)00154-7)
- Krake S, Conzelmann C, Heuer S, Dyballa M, Zibek S, Hahn T (2024) Production of chitosan from *Aspergillus niger* and quantitative evaluation of the process using adapted analytical tools. *Biotechnol Bioprocess Eng* 29:942–954. <https://doi.org/10.1007/s12257-024-00124-3>
- Krishnan K, Askew DS (2014) Endoplasmic reticulum stress and fungal pathogenesis. *Fungal Biol Rev* 28:29–35
- Kristiansen B, Linden J, Matthey M (1998) Citric acid biotechnology. In: Kristiansen B, Linden J, Matthey M (eds). CRC Press
- Krull R, Wucherpfennig T, Esfandabadi ME, Walisko R, Melzer G, Hempel DC, Kampen I, Kwade A, Wittmann C (2013) Characterization and control of fungal morphology for improved production performance in biotechnology. *J Biotechnol* 163:112–123. <https://doi.org/10.1016/j.jbiotec.2012.06.024>
- Książek E (2023) Citric acid: Properties, microbial production, and applications in industries. *Molecules* 29:22. <https://doi.org/10.3390/molecules29010022>
- Kwon MJ, Arentshorst M, Roos ED, van den Hondel CAMJJ, Meyer V, Ram AFJ (2011) Functional characterization of Rho GTPases in *Aspergillus niger* uncovers conserved and diverged roles of Rho proteins within filamentous fungi. *Mol Microbiol* 79:1151–1167. <https://doi.org/10.1111/j.1365-2958.2010.07524.x>
- Kwon MJ, Arentshorst M, Fiedler M, de Groen FLM, Punt PJ, Meyer V, Ram AFJ (2014) Molecular genetic analysis of vesicular transport in *Aspergillus niger* reveals partial conservation of the molecular mechanism of exocytosis in fungi. *Microbiology* (n y) 160:316–329. <https://doi.org/10.1099/mic.0.074252-0>
- Kwon HS, Kawaguchi K, Kikuma T, Takegawa K, Kitamoto K, Higuchi Y (2017) Analysis of an acyl-CoA binding protein in *Aspergillus oryzae* that undergoes unconventional secretion. *Biochem Biophys Res Commun* 493:481–486. <https://doi.org/10.1016/j.bbrc.2017.08.166>
- Lamon G, Lends A, Valsecchi I, Wong SSW, Duprès V, Lafont F, Tolchard J, Schmitt C, Mallet A, Grélard A, Morvan E, Dufourc EJ, Habenstein B, Guijarro JI, Aimaniananda V, Loquet A (2023) Solid-state NMR molecular snapshots of *Aspergillus fumigatus* cell wall architecture during a conidial morphotype transition. *Proceedings of the National Academy of Sciences* 120. <https://doi.org/10.1073/pnas.2212003120>
- Lamoth F, Juvvadi PR, Soderblom EJ, Moseley MA, Steinbach WJ (2015) Hsp70 and the cochaperone StiA (Hop) orchestrate Hsp90-mediated caspofungin tolerance in *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 59(8):4727–4733. <https://doi.org/10.1128/aac.00946-15>
- Langner T, Göhre V (2016) Fungal chitinases: function, regulation, and potential roles in plant/pathogen interactions. *Curr Genet* 62:243–254
- Latgé J-P (2007) The cell wall: a carbohydrate armour for the fungal cell. *Mol Microbiol* 66:279–290
- Latgé J-P (2010) Tasting the fungal cell wall. *Cell Microbiol* 12:863–872
- Le Mauff F (2020) Exopolysaccharides and biofilms. In: Latgé J-P (ed) The fungal cell wall - an armour and a weapon for human fungal pathogens. Springer, pp 225–254
- Le Mauff F, Sheppard DC (2023) Understanding *Aspergillus fumigatus* galactosaminogalactan biosynthesis: a few questions remain. *The Cell Surface* 9:100095. <https://doi.org/10.1016/j.tcs.2023.100095>
- Lee SC, Shaw BD (2008) Localization and function of ADP ribosylation factor A in *Aspergillus nidulans*. *FEMS Microbiol Lett* 283:216–222. <https://doi.org/10.1111/j.1574-6968.2008.01174.x>
- Lee MJ, Gravelat FN, Cerone RP, Baptista SD, Campoli PV, Choe S-I, Kravtsov I, Vinogradov E, Creuzenet C, Liu H, Berghuis AM, Latgé J-P, Filler SG, Fontaine T, Sheppard DC (2014) Overlapping and distinct roles of *Aspergillus fumigatus* UDP-glucose 4-epimerases in galactose metabolism and the synthesis of galactose-containing cell wall polysaccharides. *J Biol Chem* 289:1243–1256. <https://doi.org/10.1074/jbc.M113.522516>
- van Leeuwen J, Norton G, Ndlela S, Rudnick D (2016) Processes for isolating chitin and chitosan from fungal biomass. (Patent No. US 9,249,235 B2) 14 March 2014. <https://worldwide.espacenet.com/patent/search?q=pn%3DUS9249235B2>
- Levin DE (2005) Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* 69:262–291. <https://doi.org/10.1128/MMBR.69.2.262-291.2005>
- Levin DE (2011) Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the cell wall integrity signaling pathway. *Genetics* 189:1145–1175. <https://doi.org/10.1534/genetics.111.128264>
- Lew RR (2019) Biomechanics of hyphal growth. *Biology of the Fungal Cell*. Springer International Publishing, Cham, pp 83–94
- Lin J, Wester MJ, Graus MS, Lidke KA, Neumann AK (2016) Nanoscopic cell-wall architecture of an immunogenic ligand in *Candida albicans* during antifungal drug treatment. *Mol Biol Cell* 27:1002–1014. <https://doi.org/10.1091/mbc.E15-06-0355>
- Lindner S, Bonin M, Hellmann MJ, Moerschbacher BM (2025) Three intertwining effects guide the mode of action of chitin deacetylase de- and *N*-acetylation reactions. *Carbohydr Polym* 347:122725. <https://doi.org/10.1016/j.carbpol.2024.122725>
- Liu H, Zheng Z, Wang P, Gong G, Wang L, Zhao G (2013) Morphological changes induced by class III chitin synthase gene silencing could enhance penicillin production of *Penicillium chrysogenum*. *Appl Microbiol Biotechnol* 97:3363–3372. <https://doi.org/10.1007/s00253-012-4581-3>

- Liu D, Garrigues S, de Vries RP (2023) Heterologous protein production in filamentous fungi. *Appl Microbiol Biotechnol* 107:5019–5033
- Liu J-J, Hou Y-K, Wang X, Zhou X-T, Yin J-Y, Nie S-P (2024) Recent advances in the biosynthesis of fungal glucan structural diversity. *Carbohydr Polym* 329:121782. <https://doi.org/10.1016/j.carbpol.2024.121782>
- Lyu J, Torchia C, Post H, Moran Torres JP, Altelaar AFM, de Cock H, Wösten HAB (2023) The α -(1,3)-glucan synthase gene *agsE* impacts the secretome of *Aspergillus niger*. *Antonie Van Leeuwenhoek* 116:867–882. <https://doi.org/10.1007/s10482-023-01853-w>
- Malavazi I, Goldmann GH, Brown NA (2014) The importance of connections between the cell wall integrity pathway and the unfolded protein response in filamentous fungi. *Brief Funct Genomics* 13:456–470. <https://doi.org/10.1093/bfpg/elu027>
- Mata-Martínez P, Bergón-Gutiérrez M, del Fresno C (2022) Dectin-1 signaling update: new perspectives for trained immunity. *Front Immunol* 13. <https://doi.org/10.3389/fimmu.2022.812148>
- Max B, Salgado JM, Rodríguez N, Cortés S, Converti A, Manuel Domínguez J (2010) Biotechnological production of citric acid. *Braz J Microbiol* 41:862–875
- McCotter SW, Horianopoulos LC, Kronstad JW (2016) Regulation of the fungal secretome. *Curr Genet* 62:533–545
- McIntyre M, Müller C, Dynesen J, Nielsen J (2001) Metabolic engineering of the morphology of *Aspergillus*. In: *Advances in biochemical engineering/ biotechnology*, pp 103–128. https://doi.org/10.1007/3-540-45300-8_6
- Mellado E, Aufaivre-Brown A, Gow NAR, Holden DW (1996) The *Aspergillus fumigatus* *chsC* and *chsG* genes encode Class III chitin synthases with different functions. *Mol Microbiol* 20:667–679. <https://doi.org/10.1046/j.1365-2958.1996.55710.84.x>
- Ménard R, Alban S, de Ruffray P, Jamois F, Franz G, Fritig B, Yvin J-C, Kauffmann S (2004) β -1,3 Glucan sulfate, but not β -1,3 glucan, induces the salicylic acid signaling pathway in *Tobacco* and *Arabidopsis*. *Plant Cell* 16:3020–3032. <https://doi.org/10.1105/tpc.104.024968>
- Mendonça APS, dos Reis KL, Barbosa-Tessmann IP (2023) *Aspergillus clavatus* UEM 04: an efficient producer of glucoamylase and α -amylase able to hydrolyze gelatinized and raw starch. *Int J Biol Macromol* 249:125890. <https://doi.org/10.1016/j.ijbiomac.2023.125890>
- Meyer V (2008) Genetic engineering of filamentous fungi - Progress, obstacles and future trends. *Biotechnol Adv* 26:177–185
- Meyer V, Wanka F, van Gent J, Arentshorst M, van den Hondel CAMJJ, Ram AFJ (2011) Fungal gene expression on demand: an inducible, tunable, and metabolism-independent expression system for *Aspergillus niger*. *Appl Environ Microbiol* 77:2975–2983. <https://doi.org/10.1128/AEM.02740-10>
- Meyer V, Andersen MR, Brakhage AA, Braus GH, Caddick MX, Cairns TC, de Vries RP, Haarmann T, Hansen K, Hertz-fowler C, Krappmann S, Mortensen UH, Peñalva MA, Ram AFJ, Head RM (2016) Current challenges of research on filamentous fungi in relation to human welfare and a sustainable bio-economy: a white paper. *Fungal Biol Biotechnol* 3:1–17. <https://doi.org/10.1186/s40694-016-0024-8>
- Meyer V, Basenko EY, Benz JP, Braus GH, Caddick MX, Csukai M, De Vries RP, Endy D, Frisvad JC, Gunde-Cimerman N, Haarmann T, Hadar Y, Hansen K, Johnson RI, Keller NP, Kraševac N, Mortensen UH, Perez R, Ram AFJ, Record E, Ross P, Shapaval V, Steiniger C, Van Den Brink H, Van Munster J, Yarden O, Wösten HAB (2020) Growing a circular economy with fungal biotechnology: a white paper. *Fungal Biol Biotechnol* 7
- Miyazawa K, Yoshimi A, Zhang S, Sano M, Nakayama M, Gomi K, Abe K (2016) Increased enzyme production under liquid culture conditions in the industrial fungus *Aspergillus oryzae* by disruption of the genes encoding cell wall α -1,3-glucan synthase. *Biosci Biotechnol Biochem* 80:1853–1863. <https://doi.org/10.1080/09168451.2016.1209968>
- Miyazawa K, Yoshimi A, Abe K (2020) The mechanisms of hyphal pellet formation mediated by polysaccharides, α -1,3-glucan and galactosaminogalactan, in *Aspergillus* species. *Fungal Biol Biotechnol* 7:10. <https://doi.org/10.1186/s40694-020-00101-4>
- Miyazawa K, Yoshimi A, Sano M, Tabata F, Sugahara A, Kasahara S, Koizumi A, Yano S, Nakajima T, Abe K (2019) Both galactosaminogalactan and α -1,3-Glucan contribute to aggregation of *Aspergillus oryzae* hyphae in liquid culture. *Front Microbiol* 10. <https://doi.org/10.3389/fmicb.2019.02090>
- Müller C, Hansen K, Szabo P, Nielsen J (2003) Effect of deletion of chitin synthase genes on mycelial morphology and culture viscosity in *Aspergillus oryzae*. *Biotechnol Bioeng* 81:525–534. <https://doi.org/10.1002/bit.10491>
- Muñoz I, Rodríguez C, Gillet DM, Moerschbacher B (2018) Life cycle assessment of chitosan production in India and Europe. *Int J Life Cycle Assess* 23:1151–1160. <https://doi.org/10.1007/s11367-017-1290-2>
- Musílková M, Ujcová E, Pláček J, Fencel Z, Seichert L (1981) Release of protoplasts from a fungus as a criterion of mechanical interactions in the fermentor. *Biotechnol Bioeng* 23:441–446. <https://doi.org/10.1002/bit.260230219>
- Nevalainen H, Peterson R (2014) Making recombinant proteins in filamentous fungi – are we expecting too much? *Front Microbiol* 5:75
- Nielsen J, Johansen CL, Jacobsen M, Krabben P, Villadsen J (1995) Pellet formation and fragmentation in submerged cultures of *Penicillium chrysogenum* and its relation to Penicillin production. *Biotechnol Prog* 11:93–98. <https://doi.org/10.1021/bp00031a013>
- Nihei Y, Higashiyama M, Miyauchi K, Haniuda K, Suzuki Y, Kubo M, Kitamura D (2023) Subcutaneous immunisation with zymosan generates mucosal IgA-eliciting memory and protects mice from heterologous influenza virus infection. *Int Immunol* 35:377–386. <https://doi.org/10.1093/intimm/dxad013>
- Nikkilä I, Waldén M, Maina NH, Tenkanen M, Mikkonen KS (2020) Fungal cell biomass from enzyme industry as a sustainable source of hydrocolloids. *Frontiers in Chemical Engineering* 2. <https://doi.org/10.3389/fceng.2020.574072>
- Norouzzian D, Akbarzadeh A, Scharer JM, Moo Young M (2006) Fungal glucoamylases. *Biotechnol Adv* 24:80–85
- Nwe N, Stevens WF (2002) Production of fungal chitosan by solid substrate fermentation followed by enzymatic extraction. *Biotechnol Lett* 24:131–134. <https://doi.org/10.1023/A:1013850621734>
- Oliveira DL, Nakayasu ES, Joffe LS, Guimarães AJ, Sobreira TJP, Nosanchuk JD, Cordero RJB, Frases S, Casadevall A, Almeida IC, Nimrichter L, Rodrigues ML (2010) Characterization of yeast extracellular vesicles: evidence for the participation of different pathways of cellular traffic in vesicle biogenesis. *PLoS ONE* 5:1–13. <https://doi.org/10.1371/journal.pone.0011113>
- Olsvik E, Kristiansen B (1994) Rheology of filamentous fermentations. *Biotechnol Adv* 12:1–39. [https://doi.org/10.1016/0734-9750\(94\)90288-7](https://doi.org/10.1016/0734-9750(94)90288-7)
- Onoue T, Tanaka Y, Hagiwara D, Ekino K, Watanabe EA, Ohta K, Kamei K, Shibata N, Goto M, Oka T (2018) Identification of two mannosyltransferases contributing to biosynthesis of the fungal-type galactomannan α -core-mannan structure in *Aspergillus fumigatus*. *Sci Rep* 8:16918. <https://doi.org/10.1038/s41598-018-35059-2>
- Ost KJ, Arentshorst M, Moerschbacher BM, Dirks-Hofmeister ME, Ram AFJ (2025) Comprehensive phenotypic analysis of multiple gene deletions of α -glucan synthase and Crh-transglycosylase gene families in *Aspergillus niger* highlighting the versatility of

- the fungal cell wall. *The Cell Surface* 13:100141. <https://doi.org/10.1016/j.tcs.2025.100141>
- Pal P, Pal A, Nakashima K, Yadav BK (2021) Applications of chitosan in environmental remediation: a review. *Chemosphere* 266:128934. <https://doi.org/10.1016/j.chemosphere.2020.128934>
- Palma AS, Feizi T, Zhang Y, Stoll MS, Lawson AM, Díaz-Rodríguez E, Campanero-Rhodes MA, Costa J, Gordon S, Brown GD, Chai W (2006) Ligands for the β -glucan receptor, Dectin-1, assigned using “designer” microarrays of oligosaccharide probes (neoglycolipids) generated from glucan polysaccharides. *J Biol Chem* 281:5771–5779. <https://doi.org/10.1074/jbc.M511461200>
- Paloheimo M, Haarmann T, Mäkinen S, Vehmaanperä J (2016) Production of industrial enzymes in *Trichoderma reesei*. In: Schmoll M, Dattenböck C (eds) *Gene expression systems in fungi: advancements and applications*. Springer International Publishing, Cham, pp 23–57
- Papagianni M (2004) Fungal morphology and metabolite production in submerged mycelial processes. *Biotechnol Adv* 22:189–259
- Papagianni M (2007) Advances in citric acid fermentation by *Aspergillus niger*: biochemical aspects, membrane transport and modeling. *Biotechnol Adv* 25:244–263. <https://doi.org/10.1016/j.biotechadv.2007.01.002>
- Park J, Hulsman M, Arentshorst M, Breeman M, Alazi E, Lagendijk EL, Rocha MC, Malavazi I, Nitsche BM, Hondel CAMJJ, Meyer V, Ram AFJ (2016) Transcriptomic and molecular genetic analysis of the cell wall salvage response of *Aspergillus niger* to the absence of galactofuranose synthesis. *Cell Microbiol* 18:1268–1284. <https://doi.org/10.1111/cmi.12624>
- van Peij NNME, Beishuizen M, van de Vondervoort PJI (2014) AgsE-deficient strain (Patent No. WO 2014/013074 A1) 23 January 2014. <https://worldwide.espacenet.com/patent/search?q=pn%3DWO2014013074A1>
- Perez P, Cansado J (2010) Cell integrity signaling and response to stress in fission yeast. *Curr Protein Pept Sci* 11:680–692. <https://doi.org/10.2174/138920310794557718>
- Punt PJ, van Biezen N, Conesa A, Albers A, Mangnus J, van den Hondel C (2002) Filamentous fungi as cell factories for heterologous protein production. *Trends Biotechnol* 20:200–206. [https://doi.org/10.1016/S0167-7799\(02\)01933-9](https://doi.org/10.1016/S0167-7799(02)01933-9)
- Punt PJ, Schuren FHJ, Lehmbeck J, Christensen T, Hjort C, van den Hondel CAMJJ (2008) Characterization of the *Aspergillus niger* *prrT*, a unique regulator of extracellular protease encoding genes. *Fungal Genet Biol* 45:1591–1599. <https://doi.org/10.1016/j.fgb.2008.09.007>
- Quintanilla D, Hagemann T, Hansen K, Gernaey KV (2015) Fungal morphology in industrial enzyme production – modelling and monitoring. *Adv Biochem Eng Biotechnol* 149:29–54. https://doi.org/10.1007/10_2015_309
- Quintin J, Saeed S, Martens JHA, Giamarellos-Bourboulis EJ, Ifrim DC, Logie C, Jacobs L, Jansen T, Kullberg B-J, Wijmenga C, Joosten LAB, Xavier RJ, van der Meer JWM, Stunnenberg HG, Netea MG (2012) *Candida albicans* infection affords protection against reinfection via functional reprogramming of monocytes. *Cell Host Microbe* 12:223–232. <https://doi.org/10.1016/j.chom.2012.06.006>
- Rainer H, Goretzki A, Lin Y-J, Schiller HR, Krause M, Döring S, Strecker D, Junker A-C, Wolfheimer S, Toda M, Scheurer S, Schülke S (2024) Characterization of the immune-modulating properties of different β -glucans on myeloid dendritic cells. *Int J Mol Sci* 25:9914. <https://doi.org/10.3390/ijms25189914>
- Read ND (2011) Exocytosis and growth do not occur only at hyphal tips. *Mol Microbiol* 81:4–7
- Reddy NS, Nimmagadda A, Rao KRSS (2003) An overview of the microbial α -amylase family. *Afr J Biotechnol* 2:645–648
- Reed G, Nagodawithana TW (1990) *Yeast technology*. Springer, Netherlands, Dordrecht
- Reindl M, Hänsch S, Weidtkamp-Peters S, Schipper K (2019) A potential lock-type mechanism for unconventional secretion in fungi. *Int J Mol Sci* 20:460. <https://doi.org/10.3390/ijms20030460>
- Richter C, Cord-Landwehr S, Singh R, Ryll J, Moerschbacher BM (2025) Dissecting and optimizing bioactivities of chitosans by enzymatic modification. *Carbohydr Polym* 349:122958. <https://doi.org/10.1016/j.carbpol.2024.122958>
- Riseh RS, Vazvani MG, Kennedy JF (2023) The application of chitosan as a carrier for fertilizer: a review. *Int J Biol Macromol* 252:126483. <https://doi.org/10.1016/j.ijbiomac.2023.126483>
- Ritala A, Häkkinen ST, Toivari M, Wiebe MG (2017) Single cell protein-state-of-the-art, industrial landscape and patents 2001–2016. *Front Microbiol* 8:2009. <https://doi.org/10.3389/fmicb.2017.02009>
- Rocha MC, Fabri JHTM, de Godoy KF, de Castro PA, Hori JI, da Cunha AF, Arentshorst M, Ram AFJ, van den Hondel CAMJJ, Goldman GH, Malavazi I (2016) *Aspergillus fumigatus* MADS-Box transcription factor *rlmA* is required for regulation of the cell wall integrity and virulence. *G3: Genes, Genomes, Genetics* 6:2983–3002. <https://doi.org/10.1534/g3.116.031112>
- Rocha MC, Fabri JHTM, Simões IT, Silva-Rocha R, Hagiwara D, da Cunha AF, Goldman GH, Cánovas D, Malavazi I (2020) The cell wall integrity pathway contributes to the early stages of *Aspergillus fumigatus* asexual development. *Appl Environ Microbiol* 86:e02347–e2419. <https://doi.org/10.1128/AEM.02347-19>
- Rocha MC, Minari K, Fabri JHTM, Kerkaert JD, Gava LM, da Cunha AF, Cramer RA, Borges JC, Malavazi I (2020) *Aspergillus fumigatus* Hsp90 interacts with the main components of the cell wall integrity pathway and cooperates in heat shock and cell wall stress adaptation. *Cell Microbiol* 23(2):e13273. <https://doi.org/10.1111/cmi.13273>
- Rodicio R, Buchwald U, Schmitz H-P, Heinisch JJ (2008) Dissecting sensor functions in cell wall integrity signaling in *Kluyveromyces lactis*. *Fungal Genet Biol* 45:422–435. <https://doi.org/10.1016/j.fgb.2007.07.009>
- Rodrigues ML, Godinho RMC, Zamith-Miranda D, Nimrichter L (2015) Traveling into outer space: Unanswered questions about fungal extracellular vesicles. *PLoS Pathog* 11:e1005240. <https://doi.org/10.1371/journal.ppat.1005240>
- da Rosa RR, Fernandes SN, Mitov M, Godinho MH (2024) Cellulose and chitin twisted structures: from nature to applications. *Adv Funct Mater* 34. <https://doi.org/10.1002/adfm.202304286>
- Ruiz-Herrera J, Ortiz-Castellanos L (2019) Cell wall glucans of fungi. *A Review the Cell Surface* 5:100022. <https://doi.org/10.1016/j.tcs.2019.100022>
- Safeer A, Kleijburg F, Bahri S, Beriashvili D, Veldhuizen EJA, van Neer J, Tegelaar M, de Cock H, Wösten HAB, Baldus M (2023) Probing cell-surface interactions in fungal cell walls by high-resolution 1H-detected solid-state NMR spectroscopy. *Chem A Eur J* 29. <https://doi.org/10.1002/chem.202202616>
- Sakekar AA, Gaikwad SR, Puneekar NS (2021) Protein expression and secretion by filamentous fungi. *J Biosci* 46:5. <https://doi.org/10.1007/s12038-020-00120-8>
- Santos VP, Marques NSS, Maia PCSV, Lima MAB, Franco LO, Campos-Takaki GM (2020) Seafood waste as attractive source of chitin and chitosan production and their applications. *Int J Mol Sci* 21:4290. <https://doi.org/10.3390/ijms21124290>
- Sato H, Toyoshima Y, Shintani T, Gomi K (2011) Identification of potential cell wall component that allows Taka-amylase A adsorption in submerged cultures of *Aspergillus oryzae*. *Appl Microbiol Biotechnol* 92:961–969. <https://doi.org/10.1007/s00253-011-3422-0>
- Schäfer D, Schmitz K, Weuster-Botz D, Benz JP (2020) Comparative evaluation of *Aspergillus niger* strains for endogenous pectin-depolymerization capacity and suitability for d-galacturonic acid

- production. *Bioprocess Biosyst Eng* 43:1549–1560. <https://doi.org/10.1007/s00449-020-02347-z>
- Schuster M, Martin-Urdiroz M, Higuchi Y, Hacker C, Kilaru S, Gurr SJ, Steinberg G (2016) Co-delivery of cell-wall-forming enzymes in the same vesicle for coordinated fungal cell wall formation. *Nat Microbiol* 1(11):1–11. <https://doi.org/10.1038/nmicrobiol.2016.149>
- Schuster M, Guiu-Aragones C, Steinberg G (2020) Class V chitin synthase and β -(1,3)-glucan synthase co-travel in the same vesicle in *Zyloseptoria tritici*. *Fungal Genet Biol* 135:103286. <https://doi.org/10.1016/j.fgb.2019.103286>
- Scrimale T, Didone L, de Mesy Bentley KL, Krysan DJ (2009) The unfolded protein response is induced by the cell wall integrity mitogen-activated protein kinase signaling cascade and is required for cell wall integrity in *Saccharomyces cerevisiae*. *Mol Biol Cell* 20:164–175. <https://doi.org/10.1091/mbc.e08-08-0809>
- Sebastian J, Rouissi T, Brar SK (2020) Fungal chitosan: prospects and challenges. In: *Handbook of Chitin and Chitosan*. Elsevier, pp 419–452. <https://doi.org/10.1016/B978-0-12-817970-3.00014-6>
- Shariatnia Z (2019) Pharmaceutical applications of chitosan. *Adv Colloid Interface Sci* 263:131–194. <https://doi.org/10.1016/j.cis.2018.11.008>
- Sharp JK, McNeil M, Albersheim P (1984) The primary structures of one elicitor-active and seven elicitor-inactive hexa(beta-D-glucopyranosyl)-D-glucitols isolated from the mycelial walls of *Phytophthora megasperma* f. sp. *glycinea*. *J Biol Chem* 259:11321–11336. [https://doi.org/10.1016/S0021-9258\(18\)90865-3](https://doi.org/10.1016/S0021-9258(18)90865-3)
- Shibuya N, Minami E (2001) Oligosaccharide signalling for defence responses in plant. *Physiol Mol Plant Pathol* 59:223–233. <https://doi.org/10.1006/pmpp.2001.0364>
- Singh R, Smiatek J, Moerschbacher BM (2023) Unraveling the impact of acetylation patterns in chitosan oligomers on Cu²⁺ ion binding: Insights from DFT calculations. *Int J Mol Sci* 24:13792. <https://doi.org/10.3390/ijms241813792>
- Smith AJ, Graves B, Child R, Rice PJ, Ma Z, Lowman DW, Ensley HE, Ryter KT, Evans JT, Williams DL (2018) Immunoregulatory activity of the natural product laminarin varies widely as a result of its physical properties. *J Immunol* 200:788–799. <https://doi.org/10.4049/jimmunol.1701258>
- Speth C, Rambach G, Lass-Flörl C, Howell PL, Sheppard DC (2019) Galactosaminogalactan (GAG) and its multiple roles in *Aspergillus* pathogenesis. *Virulence* 10:976–983. <https://doi.org/10.1080/21505594.2019.1568174>
- Sreekumar S, Wattjes J, Niehues A, Mengoni T, Mendes AC, Morris ER, Goycoolea FM, Moerschbacher BM (2022) Biotechnologically produced chitosans with nonrandom acetylation patterns differ from conventional chitosans in properties and activities. *Nat Commun* 13:7125. <https://doi.org/10.1038/s41467-022-34483-3>
- Steinberg G, Peñalva MA, Riquelme M, Wösten HA, Harris SD (2017) Cell biology of hyphal growth. *Microbiol Spectr* 5. <https://doi.org/10.1128/microbiolspec.FUNK-0034-2016>
- Stock J, Sarkari P, Kreibich S, Brefort T, Feldbrügge M, Schipper K (2012) Applying unconventional secretion of the endochitinase Cts1 to export heterologous proteins in *Ustilago maydis*. *J Biotechnol* 161:80–91. <https://doi.org/10.1016/j.jbiotec.2012.03.004>
- Student M, Hellmann MJ, Cord-Landwehr S, Moerschbacher BM (2024) Chitins and chitosans – a tale of discovery and disguise, of attachment and attainment. *Curr Opin Plant Biol* 82:102661. <https://doi.org/10.1016/j.pbi.2024.102661>
- Suda Y, Kurokawa K, Nakano A (2018) Regulation of ER-Golgi transport dynamics by GTPases in budding yeast. *Front Cell Dev Biol* 5. <https://doi.org/10.3389/fcell.2017.00122>
- Sun X, Wu H, Zhao G, Li Z, Wu X, Liu H, Zheng Z (2018) Morphological regulation of *Aspergillus niger* to improve citric acid production by *chsC* gene silencing. *Bioprocess Biosyst Eng* 41:1029–1038. <https://doi.org/10.1007/s00449-018-1932-1>
- Sun W (2024) Fungal mycelia: from innovative materials to promising products: Insights and challenges. *Biointerphases* 19. <https://doi.org/10.1116/6.0003441>
- Sutay Kocabaş D, Lyne J, Ustunol Z (2022) Hydrolytic enzymes in the dairy industry: applications, market and future perspectives. *Trends Food Sci Technol* 119:467–475
- Takeshita N (2016) Coordinated process of polarized growth in filamentous fungi. *Biosci Biotechnol Biochem* 80:1693–1699. <https://doi.org/10.1080/09168451.2016.1179092>
- Tang W, Wang J, Hou H, Li Y, Wang J, Fu J, Lu L, Gao D, Liu Z, Zhao F, Gao X, Ling P, Wang F, Sun F, Tan H (2023) Review: application of chitosan and its derivatives in medical materials. *Int J Biol Macromol* 240:124398. <https://doi.org/10.1016/j.ijbio.2023.124398>
- Tao Z, Yuan H, Liu M, Liu Q, Zhang S, Liu H, Jiang Y, Huang D, Wang T (2023) Yeast extract: characteristics, production, applications and future perspectives. *J Microbiol Biotechnol* 33:151–166
- Tefsen B, Ram AFJ, Van Die I, Routier FH (2012) Galactofuranose in eukaryotes: aspects of biosynthesis and functional impact. *Glycobiology* 22:456–469
- Terfrüchte M, Reindl M, Jankowski S, Sarkari P, Feldbrügge M, Schipper K (2017) Applying unconventional secretion in *Ustilago maydis* for the export of functional nanobodies. *Int J Mol Sci* 18:937. <https://doi.org/10.3390/ijms18050937>
- Thykaer J, Nielsen J (2003) Metabolic engineering of β -lactam production. *Metab Eng* 5:56–69. [https://doi.org/10.1016/S1096-7176\(03\)00003-X](https://doi.org/10.1016/S1096-7176(03)00003-X)
- Tonomura K, Futai F, Tanabe O (1963) Binding of α -amylase to the cell wall of *Aspergillus oryzae*. *Biochim Biophys Acta* 78:802–805. [https://doi.org/10.1016/0006-3002\(63\)91068-0](https://doi.org/10.1016/0006-3002(63)91068-0)
- Trevithick JR, Metzner RL (1966) Molecular sieving by *Neurospora* cell walls during secretion of invertase isozymes. *J Bacteriol* 92:1010–1015. <https://doi.org/10.1128/jb.92.4.1010-1015.1966>
- Tsuizaki M, Ohta A, Horiuchi H (2013) Myosin motor-like domain of class VI chitin synthase CsmB of *Aspergillus nidulans* is not functionally equivalent to that of class V chitin synthase CsmA. *Biosci Biotechnol Biochem* 77:369–374. <https://doi.org/10.1271/bbb.120822>
- Udagawa H (2016) Filamentous fungal double-mutant host cells (Patent No. WO 2016/066690 A1) 6 May 2016. <https://worldwide.espacenet.com/patent/search?q=pn%3DWO2016066690A1>
- Urs MJ, Moerschbacher BM, Cord-Landwehr S (2023) Quantitative enzymatic-mass spectrometric analysis of the chitinous polymers in fungal cell walls. *Carbohydr Polym* 301:120304. <https://doi.org/10.1016/j.carbpol.2022.120304>
- Valuates Reports (2024) Global cellulase market research report 2024 <https://reports.valuates.com/market-reports/QYRE-Auto-1P12307/global-cellulase>. Accessed 5 Jan 2025
- Vandenberghe LPS, Rodrigues C, de Carvalho JC, Medeiros ABP, Soccol CR (2017) Production and application of citric acid. In: Pandey A, Negi S, Soccol CR (eds) *Current Developments in Biotechnology and Bioengineering*. Elsevier, pp 557–575
- Veiter L, Rajamanickam V, Herwig C (2018) The filamentous fungal pellet—relationship between morphology and productivity. *Appl Microbiol Biotechnol* 102:2997–3006. <https://doi.org/10.1007/s00253-018-8818-7>
- Versali M, Clerisse F, Bruyere J, Gautier S (2002) Cell wall derivatives form biomass and preparation thereof (CA2475258C) 12 February 2003. <https://worldwide.espacenet.com/patent/search?q=pn%3DCA2475258C>

- Vogt S, Varón Silva D, Mösch H-U, Essen L-O (2010) Structural base for the transfer of GPI-anchored glycoproteins into fungal cell walls. *PNAS* 117:22061–22067. <https://doi.org/10.1073/pnas.2010661117/-DCSupplemental>
- Vortmann M, Stumpf AK, Sgobba E, Dirks-Hofmeister ME, Krehenbrink M, Wendisch VF, Philipp B, Moerschbacher BM (2021) A bottom-up approach towards a bacterial consortium for the biotechnological conversion of chitin to l-lysine. *Appl Microbiol Biotechnol* 105:1547–1561. <https://doi.org/10.1007/s00253-021-11112-5>
- Wagener J, Striegler K, Wagener N (2020) α - and β -1,3-glucan synthesis and remodeling. In: Latgé J-P (ed) *The fungal cell wall - an armour and a weapon for human fungal pathogens*. Springer, pp 53–82
- Wang Y, Mortimer JC, Davis J, Dupree P, Keegstra K (2013) Identification of an additional protein involved in mannan biosynthesis. *Plant J* 73:105–117. <https://doi.org/10.1111/tpj.12019>
- Wang Q, Zhong C, Xiao H (2020) Genetic engineering of filamentous fungi for efficient protein expression and secretion. *Front Bioeng Biotechnol* 8:293. <https://doi.org/10.3389/fbioe.2020.00293>
- Ward OP (2012) Production of recombinant proteins by filamentous fungi. *Biotechnol Adv* 30:1119–1139. <https://doi.org/10.1016/j.biotechadv.2011.09.012>
- Wattjes J, Sreekumar S, Richter C, Cord-Landwehr S, Singh R, El Gueddari NE, Moerschbacher BM (2020) Patterns matter part 1: Chitosan polymers with non-random patterns of acetylation. *React Funct Polym* 151:104583. <https://doi.org/10.1016/j.reactfunctpolym.2020.104583>
- Wei H, Wu M, Fan A, Su H (2021) Recombinant protein production in the filamentous fungus *Trichoderma*. *Chin J Chem Eng* 30:74–81. <https://doi.org/10.1016/j.cjche.2020.11.006>
- Wenzel SC, Müller R (2013) Antibiotika. In: Sahm H, Antranikian G, Stahmann KP, Takors R (eds) *Industrielle Mikrobiologie*. Springer Spektrum, Berlin, Heidelberg. https://doi.org/10.1007/978-3-8274-3040-3_8
- Wessels JGH, Sietsma JH (1981) Fungal cell walls: A survey. *Plant carbohydrates II*. Springer, Berlin Heidelberg, Berlin, Heidelberg, pp 352–394
- Wongwicharn A, McNeil B, Harvey LM (1999) Effect of oxygen enrichment on morphology, growth, and heterologous protein production in chemostat cultures of *Aspergillus niger* B1-D. *Biotechnol Bioeng* 65:416–424. [https://doi.org/10.1002/\(SICI\)1097-0290\(19991120\)65:4%3C416::AID-BIT6%3E3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1097-0290(19991120)65:4%3C416::AID-BIT6%3E3.0.CO;2-Z)
- Wösten HAB (2019) Filamentous fungi for the production of enzymes, chemicals and materials. *Curr Opin Biotechnol* 59:65–70
- Wösten HAB, Moukha SM, Sietsma JH, Wessels JGH (1991) Localization of growth and secretion of proteins in *Aspergillus niger*. *J Gen Microbiol* 137:2017–2023. <https://doi.org/10.1099/00221287-137-8-2017>
- Wu T, Zivanovic S, Draughon FA, Conway WS, Sams CE (2005) Physicochemical properties and bioactivity of fungal chitin and chitosan. *J Agric Food Chem* 53:3888–3894. <https://doi.org/10.1021/jf048202s>
- Wucherpennig T, Hestler T, Krull R (2011) Morphology engineering - osmolality and its effect on *Aspergillus niger* morphology and productivity. *Microb Cell Fact* 10:58. <https://doi.org/10.1186/1475-2859-10-58>
- Wucherpennig T, Kiep KA, Driouch H, Wittmann C, Krull R (2010) Morphology and rheology in filamentous cultivations. In: *Advances in applied microbiology*. Academic Press Inc, pp 89–136. [https://doi.org/10.1016/S0065-2164\(10\)72004-9](https://doi.org/10.1016/S0065-2164(10)72004-9)
- Xie H, Ma Q, Wei D, Wang F (2020) Metabolic engineering of an industrial *Aspergillus niger* strain for itaconic acid production. *3 Biotech* 10:113. <https://doi.org/10.1007/s13205-020-2080-2>
- Yabuki M, Fukui S (1970) Presence of binding site for α -amylase and of masking protein for this site on mycelial cell wall of *Aspergillus oryzae*. *J Bacteriol* 104:138–144. <https://doi.org/10.1128/jb.104.1.138-144.1970>
- Yahara N, Ueda T, Sato K, Nakano A (2001) Multiple roles of Arf1 GTPase in the yeast exocytic and endocytic pathways. *Mol Biol Cell* 12:221–238. <https://doi.org/10.1091/mbc.12.1.221>
- Yoshimi A, Sano M, Inaba A, Kokubun Y, Fujioka T, Mizutani O, Hagiwara D, Fujikawa T, Nishimura M, Yano S, Kasahara S, Shimizu K, Yamaguchi M, Kawakami K, Abe K (2013) Functional analysis of the α -1,3-glucan synthase genes *agsA* and *agsB* in *Aspergillus nidulans*: AgsB is the major α -1,3-glucan synthase in this fungus. *PLoS ONE* 8:e54893. <https://doi.org/10.1371/journal.pone.0054893>
- Yoshimi A, Miyazawa K, Abe K (2016) Cell wall structure and biogenesis in *Aspergillus* species. *Biosci Biotechnol Biochem* 80:1700–1711. <https://doi.org/10.1080/09168451.2016.1177446>
- Yoshimi A, Miyazawa K, Abe K (2017) Function and biosynthesis of cell wall α -1,3-glucan in fungi. *Journal of Fungi* 3:63. <https://doi.org/10.3390/jof3040063>
- Yoshimi A, Miyazawa K, Kawauchi M, Abe K (2022) Cell wall integrity and its industrial applications in filamentous fungi. *Journal of Fungi* 8:435. <https://doi.org/10.3390/jof8050435>
- Younes I, Rinaudo M (2015) Chitin and chitosan preparation from marine sources. Structure, properties and applications. *Mar Drugs* 13:1133–1174. <https://doi.org/10.3390/md13031133>
- Zhang S, Sato H, Ichinose S, Tanaka M, Miyazawa K, Yoshimi A, Abe K, Shintani T, Gomi K (2017) Cell wall α -1,3-glucan prevents α -amylase adsorption onto fungal cell in submerged culture of *Aspergillus oryzae*. *J Biosci Bioeng* 124:47–53. <https://doi.org/10.1016/j.jbiosc.2017.02.013>
- Zhang S, Zheng H, Chen Q, Chen Y, Wang S, Lu L, Zhang S (2017) The lectin chaperone calnexin is involved in the endoplasmic reticulum stress response by regulating Ca^{2+} homeostasis in *Aspergillus nidulans*. *Appl Environ Microbiol* 83:e00673–e717. <https://doi.org/10.1128/AEM.00673-17>
- Zheng X, Cairns TC, Ni X, Zhang L, Zhai H, Meyer V, Zheng P, Sun J (2022) Comprehensively dissecting the hub regulation of PkaC on high-productivity and pellet macromorphology in citric acid producing *Aspergillus niger*. *Microb Biotechnol* 15(6):1867–1882. <https://doi.org/10.1111/1751-7915.14020>
- Zuriegat Q, Abubakar YS, Wang Z, Chen M, Zhang J (2024) Emerging roles of exocyst complex in fungi: a review. *Journal of Fungi* 10:614. <https://doi.org/10.3390/jof10090614>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.